

Corrigendum

Micronutrient-fortified infant cereal improves Hb status and reduces iron-deficiency anaemia in Indian infants: an effectiveness study – Corrigendum

Shally Awasthi, Narayan U Reddy, Monjori Mitra, Shweta Singh, Sanjeev Ganguly, Ivana Jankovic, Dominik Grathwohl, Colin I. Cercamondi and Apurba Ghosh

DOI: [10.1017/S0007114519003386](https://doi.org/10.1017/S0007114519003386) (First published online 3 January 2020)

The authors apologise for an error in Table 3 The geometric mean for Serum ferritin (ng/ml) was incorrectly given as 40.0 when the correct value is 36.0. All other values remain the same. The corrected Table is given below.

Correct Table 3. Iron status indices and C-reactive protein concentration of infants in the intervention group at 6 and 12 months of age and in the static cross-sectional control group at 12 months of age.

(Mean values and standard deviations; geometric mean and -1 standard deviation and +1 standard deviation)

Parameter	IC (n 64)				CG (n 80)		P*
	6 months		12 months		12 months		
	Geometric Mean	-1 SD, +1 SD	Geometric Mean	-1 SD, +1 SD	Geometric Mean	-1 SD, +1 SD	
Hemoglobin (g/l)							
Mean	113.4		118.1		109.5		<0.001
SD	8.7		10.2		16.4		
Serum ferritin (ng/ml)	36.0	16.4, 78.7	27.0	13.4, 54.4	20.3	7.5, 55.0	0.085
Serum soluble transferrin receptor (mg/l)	1.68	1.22, 2.31	1.70	1.19, 2.43	2.07	1.29, 3.33	0.014
Serum C-reactive protein (mg/l)	0.46	0.16, 1.34	0.84	0.20, 3.60	0.77	0.17, 3.39	0.741

CG, static cross-sectional control group; IC, intervention group; SD, standard deviation.

* P-value for between group comparison (ANCOVA model correcting for sex, exclusively breastfeeding until 6 months of age (Y/N), Kuppaswamy socioeconomic status⁽¹⁾, birth weight, gestational age, and study site).

