

## ON REDUCIBILITY OF PROVABILITY IN THE PRIMITIVE LOGIC [LO]

JIRO ITO

*Dedicated to Professor Katuzi Ono on his 60th birthday*

### Introduction.

In the present paper, we would like to show a theorem concerning with *reducibility of provability* in the primitive logic. This theorem seems to suggest a procedure to find the proof-note of a given proposition which is provable in the primitive logic.

The formulations of the primitive logic **LO** and [LO] have been introduced in Ono [1], [2], [3], and [5]. The primitive logic is the logic having only two logical constants IMPLICATION  $\rightarrow$  and UNIVERSAL QUANTIFICATION ( $\forall$ ), and has an interesting property that any logic belonging to intuitionistic series or to classical series can be faithfully interpreted in it (Ono [3], [4]).

In the proof-note of the logic [LO], there are some propositions enclosed in pairs of brackets. Any proposition in proof-notes of [LO] is said to be CLAD or BARE according as it is enclosed or is not enclosed in a pair of brackets. The followings are the inference rules of the logic [LO]:

[F]: *The step  $\mathfrak{A}$  can be deduced from the step  $[\mathfrak{A}]$ .*

[I]: *The step  $[\mathfrak{B}]$  can be deduced from the steps  $\mathfrak{A}$  and  $[\mathfrak{A} \rightarrow \mathfrak{B}]$ .*

[I\*]: *The step  $\mathfrak{A} \rightarrow \mathfrak{B}$  can be deduced from the fact that  $\mathfrak{B}$  is deducible from  $[\mathfrak{A}]$ .*

[U]: *The step  $[\mathfrak{A}(t)]$  can be deduced from the step  $[(x)\mathfrak{A}(x)]$  as far as  $\mathfrak{A}(u)$  contains no free variable  $x$  at all.*

[U\*]: *The step  $(x)\mathfrak{A}(x)$  can be deduced from the fact that  $\mathfrak{A}(t)$  is deducible for any variable  $t$  whatever, i.e. from the fact that the step  $\mathfrak{A}(t)$  is deducible from the step  $\forall t : .$*

---

Received March 31, 1969

Any proof-note is understood as a practical discription of steps of reasonings. Every step is introduced and denoted by its introductory index-word. Any index-word is a sequence of letters, such as  $\mathbf{A}, \mathbf{b}, \mathbf{c}, \dots$  assuming the usual alphabetical order between them, including the null sequence. We denote a sequence of letters by an underlined single letter such as  $\underline{\mathbf{p}}$  or  $\underline{\mathbf{t}}$ , and especially null sequence by  $\emptyset$ .

Here, we must further refer to [5] on some technical terms such as ORDER (Natural, Fundamental and Basis) of steps, REFERENCE STEP and ASSUMPTION STEP. In [5], K. Ono showed a characteristic feature of the logic [LO] by proving the theorem:

*Any proposition occurring in a wasteless proof-note of a proposition  $\mathfrak{P}$  is a subformula of the proposition  $\mathfrak{P}$ .*

In this paper, we shall show that this theorem may be described more precisely from a view-point on reducibility of provability.

In section 1, we shall introduce some expressions on *LO-formulas* such as REGULAR PARSING-FORM,  $\tau$ -KERNEL and  $\sigma$ -CONSTRUCTION with their PARSING-FACTORS and KERNEL.

In section 2, we shall prove the main theorem on reducibility of provability such as the theorem:

*The provability of a given proposition  $\mathfrak{A}$  is reducible to the fact that there are finite sequences of factors  $\{\mu\}$  and  $\{\nu\}$  which satisfy the following conditions; 1) the  $\tau$ -kernel of  $\mathfrak{A}$  by factors  $\{\mu\}$  is deducible from the assumptions  $\{\mu\}$ , 2)  $\{\mu\}$  contains a  $\sigma$ -construction by factors  $\{\nu\}$  with that  $\tau$ -kernel of  $\mathfrak{A}$  by factors  $\{\mu\}$  as its kernel, and 3) any formula of  $\{\nu\}$  is deducible from the assumptions  $\{\mu\}$ .*

### 1. Parsing expression.

Any formula which has no logical constants other than *implication* and *universal quantification* is called an *LO-formula*.

Firstly, we would like to introduce the parsing expressions.

#### (1. 1) Regular parsing-form

We introduce a PARSING-FORM with its FACTOR and KERNEL as follows;

An expression  $\langle \lambda \rangle \mathfrak{A}$  which is called a *parsing-form* by a  $\pi$ -factor  $\lambda$  with kernel  $\mathfrak{A}$ , is defined by

- 1) if  $\lambda$  is a formula  $\mathfrak{C}$ ,  $\langle \mathfrak{C} \rangle \mathfrak{A} \equiv^1 \mathfrak{C} \rightarrow \mathfrak{A}$ :

---

<sup>1)</sup>  $P \equiv Q$  means “ $P$  is defined by  $Q$ ”.

- 2) if  $\lambda$  is a variable  $x$ ,  $\langle x \rangle \mathfrak{A} \equiv (x)\mathfrak{A}(x)$ ;
- 3) otherwise,  $\langle \lambda \rangle \mathfrak{A}$  is undefined.

Especially, if  $\lambda$  is empty or denoted by  $\lambda_0$ ,  $\langle \ \ \rangle \mathfrak{A}$  or  $\langle \lambda_0 \rangle \mathfrak{A} \equiv \mathfrak{A}$  is called a parsing-form by the *empty-factor* or  $\lambda_0$  with kernel  $\mathfrak{A}$ .

For any  $i (i \geq 1)$ , a parsing-form with kernel  $\mathfrak{A}$  by  $\pi$ -factors  $\{\lambda_0, \dots, \lambda_i\}$ ,  $\langle \lambda_0, \lambda_1, \dots, \lambda_i \rangle \mathfrak{A} \equiv \langle \lambda_i \rangle (\langle \lambda_0, \dots, \lambda_{i-1} \rangle \mathfrak{A})$  recursively, where  $\lambda_0$  is empty and each of  $\lambda_i (i \geq 1)$  is a formula or a variable, respectively. In the case that  $\lambda_i$  is a formula each of  $\lambda_i$  is called a  $\pi$ -*formula*, otherwise a  $\pi$ -*variable*.

From the definition of parsing-forms we have the following lemmas.

LEMMA 1. *Any parsing-form which has an LO-formula as its kernel is also an LO-formula.*

LEMMA 2. *Any LO-formula is expressible in a parsing-form by  $\pi$ -factors.*

Accordingly, corresponding to any LO-formula  $\mathfrak{B}$ , there is one and only one parsing-form which has an *elementary* formula as its kernel. This expression is called a *regular parsing-form* of  $\mathfrak{B}$ .

(1. 2)  $\tau$ -Kernel.

Let  $\mathfrak{A}$  be a formula whose regular parsing-form is  $\langle \lambda_0, \dots, \lambda_n \rangle E$ . The expression  $\tau \langle \mu \rangle \mathfrak{A}$  which is called a  $\tau$ -kernel of  $\mathfrak{A}$  by a  $\tau$ -factor  $\mu$ , is defined as follows:

- 1) If  $\mu$  is a formula  $\mathfrak{C}$ , and  $\lambda_n$  is  $\mathfrak{C}$ ,  $\tau \langle \mathfrak{C} \rangle (\langle \mathfrak{C} \rangle \mathfrak{B}) \equiv \mathfrak{B}$ ,
- 2) If  $\mu$  is a free variable  $t$  and  $\lambda_n$  is a variable  $x$ ,  $\tau \langle t \rangle (\langle x \rangle \mathfrak{B}(x)) \equiv \mathfrak{B}(t)$ ,
- 3) otherwise,  $\tau \langle \mu \rangle \mathfrak{A}$  is undefined,

where  $\mathfrak{B}$  or  $\mathfrak{B}(x)$  denotes  $\langle \lambda_0, \dots, \lambda_{n-1} \rangle E$ . The  $\tau$ -factor  $\mu$  is called a  $\tau$ -*formula* or a  $\tau$ -*variable* according as  $\mu$  is a formula or a variable, respectively.

Especially, if  $\mu$  is empty or denoted by  $\mu_0$ ,  $\tau \langle \ \ \rangle \mathfrak{A}$  or  $\tau \langle \mu_0 \rangle \mathfrak{A} \equiv \mathfrak{A}$ , is called a  $\tau$ -kernel of  $\mathfrak{A}$  by the *empty-factor* or  $\mu_0$ .

For any  $i (i \geq 1)$ , a  $\tau$ -kernel of  $\mathfrak{A}$  by  $\tau$ -factors  $\{\mu_0, \dots, \mu_i\}$

$$\tau \langle \mu_0, \mu_1, \dots, \mu_i \rangle \mathfrak{A} \equiv \tau \langle \mu_i \rangle (\tau \langle \mu_0, \dots, \mu_{i-1} \rangle \mathfrak{A})$$

recursively, where  $\mu_0$  is empty and each of  $\mu_j (i \geq j \geq 1)$  is a  $\tau$ -*formula* or a  $\tau$ -*variable* according as  $\lambda_{n-j-1}$  is a formula or a variable, respectively.

(1. 3) *σ-construction.*

An expression  $\sigma\langle\nu\rangle\mathfrak{A}$  which is called a *σ-construction* with *kernel*  $\mathfrak{A}$  by a *σ-factor*  $\nu$ , is defined as follows;

- 1) If  $\nu$  is a formula  $\mathfrak{C}$ ,  $\sigma\langle\mathfrak{C}\rangle\mathfrak{A} \equiv \langle\mathfrak{C}\rangle\mathfrak{A}$ .
- 2) If  $\nu$  is a variable  $t$ ,  $\sigma\langle t\rangle\mathfrak{A}(t) \equiv \langle x\rangle\mathfrak{A}(x)$

where  $\mathfrak{A}(x)$  denotes the derived formula from  $\mathfrak{A}(t)$  replacing  $t$  in its position by  $x$  which has no occurrence in  $\mathfrak{A}(t)$ . The *σ-factor*  $\nu$  is called *σ-formula* or *σ-variable* according as  $\nu$  is a formula or a variable, respectively.

Especially, if  $\nu$  is empty or denoted by  $\nu_0$ ,  $\sigma\langle \rangle\mathfrak{A}$  or  $\sigma\langle\nu_0\rangle\mathfrak{A} \equiv \mathfrak{A}$  is called a *σ-construction* with kernel  $\mathfrak{A}$  by the *empty-factor* or  $\nu_0$ . For any  $i$  ( $i \geq 1$ ), a *σ-construction* with kernel  $\mathfrak{A}$  by *σ-factors*  $\{\nu_0, \dots, \nu_i\}$ ,

$$\sigma\langle\nu_0, \nu_1, \dots, \nu_i\rangle\mathfrak{A} \equiv \sigma\langle\nu_i\rangle(\sigma\langle\nu_0, \dots, \nu_{i-1}\rangle\mathfrak{A})$$

recursively, where  $\nu_0$  is empty and each of  $\nu_j$  ( $j \geq 1$ ) is a *σ-formula* or a *σ-variable*.

(1. 4) *Modulation of inference rules.*

Using our expressions the inference rules of the logic [LO] are modulated as follows;

- [F]: The step  $\mathfrak{A}$  can be deduced from the step  $[\mathfrak{A}]$ .
- [I]: The step  $\langle\mathfrak{B}\rangle$  can be deduced from the steps  $\mathfrak{A}$  and  $[\sigma\langle\mathfrak{A}\rangle\mathfrak{B}]$ .
- [I\*]: The step  $\langle\mathfrak{A}\rangle\mathfrak{B}$  can be deduced from the fact that  $\tau\langle\mathfrak{A}\rangle(\langle\mathfrak{A}\rangle\mathfrak{B})$  is deducible from  $[\mathfrak{A}]$ .
- [U]: The step  $\mathfrak{A}(t)$  can be deduced from the step  $[\sigma\langle t\rangle\mathfrak{A}(t)]$ .
- [U\*]: The step  $\langle x\rangle\mathfrak{A}(x)$  can be deduced from the fact that  $\tau\langle t\rangle(\langle x\rangle\mathfrak{A}(x))$  is deducible for any variable  $t$  whatever, i.e., from the fact that the step  $\tau\langle t\rangle(\langle x\rangle\mathfrak{A}(x))$  is deducible from the step  $[t]$ .<sup>2)</sup>

Let  $\Gamma$  be an ordered set of clad formulas or variables which belong to assumption steps of a step  $\mathfrak{s}$ , arranged its index-word in the fundamental order. Then, the step  $\mathfrak{s}: \mathfrak{s}$   $\mathfrak{A}$  or  $[\mathfrak{A}]$  is said to be *deducible* from the *assumption*  $\Gamma$ . We would like to denote this by  $\Gamma \vdash \mathfrak{A}$  or  $\Gamma \vdash [\mathfrak{A}]$ , respectively. Especially, if  $\mathfrak{s}$  is  $\emptyset$ , the formula of  $\mathfrak{s}$  is bare and  $\Gamma$  is empty. We denote this by  $\vdash \mathfrak{A}$ .

<sup>2)</sup> We will use clad variable  $[t]$  instead of  $\forall t$ ; because in a proof-note, denominating quantifier  $\forall t$ : as well as clad formula  $[\mathfrak{A}]$  has an assumptional character.

Any proposition  $\mathfrak{B}$  is said to be *provable* in  $[LO]$  if and only if the step  $\mathfrak{B}$  is deducible in  $[LO]$ .

**2. Reducibility of provability.**

(2. 1) *Lemmas.*

Now, we are going to prove the following lemmas preparatory to the main theorem.

LEMMA 3.  $\vdash \langle \lambda \rangle \mathfrak{A}$  if and only if  $[\mu] \vdash \tau \langle \mu \rangle \langle \lambda \rangle \mathfrak{A}$ , where  $\mu$  is a  $\tau$ -variable which has no occurrence in  $\langle \lambda \rangle \mathfrak{A}$  or a  $\tau$ -formula according as  $\lambda$  is a variable or a formula, respectively.

*Proof.* Firstly we assume  $\vdash \langle \lambda \rangle \mathfrak{A}$ . In this proof-note, the assumption step of the step  $\emptyset$  is empty, therefore the inference rule for the step  $\emptyset$  is  $[U^*]$  or  $[I^*]$ . Accordingly, there are steps  $A) [\mu]$  and  $\epsilon) \tau \langle \mu \rangle \langle \lambda \rangle \mathfrak{A}$  in this proof-note, where  $\mu$  is a  $\tau$ -variable which has no occurrence in  $\langle \lambda \rangle \mathfrak{A}$  or a  $\tau$ -formula according as  $\lambda$  is a variable or a formula. Thus we have  $[\mu] \vdash \tau \langle \mu \rangle \langle \lambda \rangle \mathfrak{A}$ . Conversely, we assume  $[\mu] \vdash \tau \langle \mu \rangle \langle \lambda \rangle \mathfrak{A}$  with the  $\mu$ -condition. Adding to this proof-note the step  $\emptyset) \langle \lambda \rangle \mathfrak{A}$ , we have  $\vdash \langle \lambda \rangle \mathfrak{A}$  by  $[U^*]$  or  $[I^*]$  according as  $\lambda$  is a variable or a formula, respectively.

In the following lemmas, let  $\langle \lambda_0, \dots, \lambda_n \rangle E$  be the regular parsing-form of  $\mathfrak{A}$ .

LEMMA 4. For any  $k$  ( $n \geq k \geq 1$ ), if  $[\mu_1], \dots, [\mu_k] \vdash \tau \langle \mu_0, \dots, \mu_k \rangle \mathfrak{A}$ , then  $\vdash \mathfrak{A}$ , where the  $\mu$ -condition:  $\mu_i$  ( $k \geq i \geq 1$ ) is a  $\tau$ -variable which has no occurrence in any  $\tau \langle \mu_0, \dots, \mu_{j-1} \rangle \mathfrak{A}$  ( $i \geq j \geq 1$ ) or a  $\tau$ -formula according as  $\lambda_{n-i+1}$  is a variable or a formula, respectively, holds.

*Proof.* For  $k = 1$ , the case is the last part of lemma 3.

Now, we assume this lemma for any number less than  $k$ , and prove it for  $k$  by induction. In this proof-note, there are the step  $\mathfrak{s}) \tau \langle \mu_0, \dots, \mu_k \rangle \mathfrak{A}$  and its assumption step  $\mathfrak{uA}) [\mu_k]$  with the  $\mu$ -condition. Therefore, adding the step  $\mathfrak{u}) \tau \langle \mu_0, \dots, \mu_{k-1} \rangle \mathfrak{A}$  to this proof-note, we have  $[\mu_1], \dots, [\mu_{k-1}] \vdash \tau \langle \mu_0, \dots, \mu_{k-1} \rangle \mathfrak{A}$  which leads up to  $\vdash \mathfrak{A}$  by assumption of induction.

LEMMA 5.  $\vdash \mathfrak{A}$  if and only if there is a sequence of factors  $\{\mu_0, \dots, \mu_k\}$  ( $n \geq k \geq 1$ ) such as

$$[\tau \langle \mu_0, \dots, \mu_k \rangle \mathfrak{A}] \in \{[\mu_1], \dots, [\mu_k]\}$$

$$\text{or } [\mu_1], \dots, [\mu_k] \vdash [\tau \langle \mu_0, \dots, \mu_k \rangle \mathfrak{A}]$$

where the  $\mu$ -condition holds.

*Proof.* Firstly by assuming  $\vdash \mathfrak{A}$ , we have  $[\mu_1] \vdash \tau\langle \mu_0, \mu_1 \rangle \mathfrak{A}$  by lemma 3. If the inference for the step  $\mathfrak{s}$ )  $\tau\langle \mu_0, \dots, \mu_{m-1} \rangle \mathfrak{A}$  is not  $[F]$ , it must be  $[I^*]$  or  $[U^*]$  and then there are steps  $\mathfrak{s}A$ )  $[\mu_m]$  and  $\mathfrak{s}t$ )  $\tau\langle \mu_0, \dots, \mu_m \rangle \mathfrak{A}$  in this proof-note, where the  $\mu$ -condition holds. However,  $m$  can not exceed  $n$  so that there is  $k$  ( $k \leq n$ ) such as the inference for the step  $\mathfrak{p}$ )  $\tau\langle \mu_0, \dots, \mu_k \rangle \mathfrak{A}$  is  $[F]$ . And therefore,  $[\tau\langle \mu_0, \dots, \mu_k \rangle \mathfrak{A}]$  belongs to the set of assumption steps of the step  $\mathfrak{p}$  or is deducible from the assumption steps of the step  $\mathfrak{p}$ .

Conversely, by assuming the condition, from the step  $\mathfrak{s}$ )  $[\tau\langle \mu_0, \dots, \mu_k \rangle \mathfrak{A}]$  we have the step  $\mathfrak{p}$ )  $\tau\langle \mu_0, \dots, \mu_k \rangle \mathfrak{A}$  by  $[F]$ , and the assumption steps of the step  $\mathfrak{s}$  is also those of the step  $\mathfrak{p}$  so that we have  $\vdash \mathfrak{A}$  by lemma 4.

LEMMA 6.  $\Gamma \vdash [\mathfrak{B}]$  if and only if there is a sequence of factor  $\{\nu_0, \dots, \nu_m\}$  such as

$$1) [\sigma\langle \nu_0, \dots, \nu_m \rangle \mathfrak{B}] \in \Gamma$$

and 2) for any  $\sigma$ -formula  $\nu_j$  ( $1 \leq j \leq m$ )  $\Gamma \vdash \nu_j$ .

*Proof.* Firstly we assume  $\Gamma \vdash [\mathfrak{B}]$ , and let the final step of this proof-note which is arranged in the fundamental order of its index-words be  $\mathfrak{s}$  (i.e.,  $\mathfrak{s}$ ) $[\mathfrak{B}]$ ). We may prove the conclusion by induction referring to the step  $\mathfrak{s}$ .

If the step  $\mathfrak{s}$  is the first step, the step  $\mathfrak{s}$  is the assumption step of itself and the  $\sigma$ -construction with kernel  $\mathfrak{B}$  by the empty factor i.e.,  $\sigma\langle \ \rangle \mathfrak{B}$ . In this case,  $m = 0$  and the condition 2) is omitted.

Otherwise, we assume the assertion for any step  $\mathfrak{r}$  which takes precedence of  $\mathfrak{s}$  in the fundamental order.

The formula of the step  $\mathfrak{s}$  is clad so that the step  $\mathfrak{s}$  is either an assumption step of itself (case 1), or a step deduced from a step  $\mathfrak{u}$  by  $[U]$  (case 2), or from steps  $\mathfrak{u}$  and  $\mathfrak{v}$  by  $[I]$  (case 3).

In the case 1, the assertion holds evidently.

In the case 2, the step  $\mathfrak{u}$  is a clad formula and is a  $\sigma$ -construction with kernel  $\mathfrak{B}$  by a  $\sigma$ -variable i.e.,  $\mathfrak{u}$ )  $[\sigma\langle \nu \rangle \mathfrak{B}]$ . And the step  $\mathfrak{u}$  takes precedence of  $\mathfrak{s}$  in the fundamental order so that there is an assumption step  $\mathfrak{p}$  of  $\mathfrak{u}$  which is a  $\sigma$ -construction with kernel  $\sigma\langle \nu \rangle \mathfrak{B}$ , by our assumption of induction. The step  $\mathfrak{p}$  is an assumption step of  $\mathfrak{s}$ , and a  $\sigma$ -construction

with kernel  $\sigma\langle\nu\rangle\mathfrak{B}$  is also a  $\sigma$ -construction with kernel  $\mathfrak{B}$ . Thus the assertion holds in the step  $\mathfrak{s}$ .

In the case 3, one of the steps  $\mathfrak{u}$  and  $\mathfrak{v}$  is a clad formula and the other is a bare formula by [I], and both of them take precedences of  $\mathfrak{s}$  in the fundamental order.

We would like to suppose the step  $\mathfrak{u}$  is clad, then there is an assumption step of  $\mathfrak{s}$  whose formula is  $\sigma$ -construction with kernel  $\mathfrak{B}$  as we had it in the case 1, where  $\nu$  is a  $\sigma$ -formula in this case. And then, the step  $\mathfrak{v}$  is the same bare formula as  $\nu$  and is deduced from its assumption steps. The assumption steps of  $\mathfrak{v}$  are the assumption steps of  $\mathfrak{s}$  too. Thus the assertion also holds in this case.

Conversely, we have  $\Gamma \vdash [\mathfrak{B}]$ , by assuming the fact that there is a sequence of factor  $\{\nu_0, \dots, \nu_m\}$  such as the conditions 1) and 2) hold for any formula  $\mathfrak{B}$ , by induction as follows;

If  $m = 0$ , the assertion is trivial.

We assume the assertion holds for any  $i$  ( $i < m$ ).

If  $\nu_m$  is a free variable, we can deduce the step  $\mathfrak{s}[\sigma\langle\nu_0, \dots, \nu_{m-1}\rangle\mathfrak{B}]$  from the step  $\mathfrak{u}[\sigma\langle\nu_0, \dots, \nu_{m-1}, \nu_m\rangle\mathfrak{B}]$  of  $\Gamma$  by [U].

If  $\nu_m$  is a formula, according to the condition 2), we have a step  $\mathfrak{v}$   $\nu_m$  which is deduced from its assumption steps  $\Gamma$ , and the step  $\mathfrak{u}[\sigma\langle\nu_0, \dots, \nu_{m-1}, \nu_m\rangle\mathfrak{B}]$  of  $\Gamma$  by the condition 1). Then we can deduce the step  $\mathfrak{s}[\sigma\langle\nu_0, \dots, \nu_{m-1}\rangle\mathfrak{B}]$  from the steps  $\mathfrak{u}$  and  $\mathfrak{v}$  by [I]. In the both cases of  $\nu_m$ , the steps of  $\Gamma$  are also the assumption steps of  $\mathfrak{s}$ .

Thus we have the conclusion.

(2. 2) *Theorem.*

From the preceding Lemmas 5 and 6 we have the main theorem and its corollaries.

**THEOREM.** *Let  $\mathfrak{A}$  be any LO-formula which has its regular parsing form by  $\pi$ -factors  $\{\lambda_0, \dots, \lambda_n\}$ . In the primitive logic [LO], the provability of a given proposition  $\mathfrak{A}$  is reducible to the fact that there are  $\tau$ -factors  $\{\mu_0, \dots, \mu_k\}$  ( $k \leq n$ ) and  $\sigma$ -factors  $\{\nu_0, \dots, \nu_m\}$  which satisfy the following conditions;*

- 1) *The  $\tau$ -kernel of  $\mathfrak{A}$  by  $\tau$ -factors  $\{\mu_0, \dots, \mu_k\}$  which satisfies the  $\mu$ -condition of lemma 4, is deducible from the assumptions  $\{[\mu_1], \dots, [\mu_k]\}$ ,*
- 2)  *$\{[\mu_1], \dots, [\mu_k]\}$  contains a  $\sigma$ -construction by  $\sigma$ -factors  $\{\nu_0, \dots, \nu_m\}$  with that  $\tau$ -kernel of  $\mathfrak{A}$  by  $\tau$ -factors  $\{\mu_0, \dots, \mu_k\}$  as its kernel,*

3) Any  $\sigma$ -formula  $\nu_j$  such as  $1 \leq j \leq m$ , is deducible from the assumptions  $\{[\mu_1], \dots, [\mu_k]\}$ .

COROLLARY 1. *If there is no  $\sigma$ -construction of  $\tau$ -kernel of  $\mathfrak{A}$  in the set of  $\tau$ -factors of  $\mathfrak{A}$ ,  $\mathfrak{A}$  is unprovable.*

COROLLARY 2. *Without assumption, any elementary proposition is unprovable.*

This theorem seems to suggest a procedure to find a proof-note of a given proposition which is provable in the primitive logic [LO]. But we would like to mention about it in our later paper ([1]-part 3).

Finally, I would like to thank Professor Katuzi Ono for his guidance and his helpful suggestions.

#### REFERENCES

- K. Ono, [ 1 ] On a practical way of describing formal deductions, *Nagoya Math. J.*, **21** (1962), 115–121.  
 [ 2 ] A certain kind of formal theories, *Nagoya Math. J.*, **25** (1965), 59–86.  
 [ 3 ] On universal character of the primitive logic, *Nagoya Math. J.*, **27** (1966), 331–353.  
 [ 4 ] Reduction of logics to the primitive logic, *J. Math. Soc. Jap.*, Vol. **19**, No. 3 (1967), 384–398.  
 [ 5 ] A study on formal deductions in the primitive logic, *Nagoya Math. J.*, **31** (1968), 1–14.  
 J. Ito, [ 1 ] Studies on mechanical deductions part 1, *Jour. Toyota Technical College*, Vol. **1** (1968), 41–49.

*Toyota Technical College*