

Merging of the superbanana plateau and $\sqrt{\nu}$ transport regimes in nearly quasisymmetric stellarators

Peter J. Catto ^{1,†}, Elizabeth A. Tolman ^{2,3} and Felix I. Parra ⁴

¹Plasma Science and Fusion Center, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

²Institute for Advanced Study, Princeton, NJ 08540, USA

³Center for Computational Astrophysics, Flatiron Institute, 162 Fifth Avenue, New York, NY 10010, USA

⁴Princeton Plasma Physics Laboratory, Princeton University, Princeton, NJ 08540, USA

(Received 18 August 2022; revised 20 December 2022; accepted 21 December 2022)

Alpha particle confinement is one of the most demanding issues for stellarators. It now seems clear that it is possible to design optimized stellarators that confine the background plasma at near tokamak radial transport levels. Moreover, adequate collisionless alpha particle confinement is possible in the core of a highly optimized stellarator. Here, the collisional confinement of barely trapped alphas in an optimized stellarator is considered by accounting for the resonance due to the reversal in direction of the drift within a flux surface and investigating the sensitive role of magnetic shear in keeping this resonance close to the passing boundary in some nearly quasisymmetric stellarator configurations. The treatment relies on a narrow collisional boundary layer formulation that combines the responses of both these resonant pitch angle alphas and the remaining barely trapped alphas. A novel merged regime treatment leads to explicit expressions for the energy diffusivity for both superbanana plateau (or resonant plateau) and $\sqrt{\nu}$ transport in the large aspect ratio limit for a slowing down tail alpha distribution function, where ν is the effective pitch angle scattering collision frequency of the trapped alphas off the background ions. Depending on the details of the optimization scheme and the sign of the magnetic shear, modest magnetic shear can be used to reduce superbanana (or resonant) plateau transport to below the $\sqrt{\nu}$ transport level. In addition, a quasilinear equation retaining spatial diffusion is derived for a general alpha distribution function that allows the radial alpha transport to modify the distribution so it is no longer isotropic in velocity space.

Key words: fusion plasma, plasma confinement

1. Introduction

Recent stellarator optimization efforts (Landreman & Sengupta 2018, 2019; Landreman 2019; Landreman, Sengupta & Plunk 2019; Plunk *et al.* 2019) extending the earlier near magnetic axis formalism of Garren & Boozer (1991*a,b*) demonstrate that many optimized nearly quasisymmetric and/or omnigenous (Boozer 1983; Nührenberg & Zille 1988; Cary

† Email address for correspondence: catto@psfc.mit.edu

& Shasharina 1997a,b; Parra *et al.* 2015) configurations are possible and can be efficiently constructed. For an exactly quasisymmetric (QS) magnetic flux surface in a stellarator a drift kinetic canonical angular momentum (Boozer 1983; Landreman & Catto 2011) exists that is a constant of the motion as in a tokamak. Optimized nearly QS stellarators provide good collisional confinement for the background ions and electrons (Beidler *et al.* 2011; Landreman & Sengupta 2019). However, the alpha particles produced by fusion must also be well confined so that most of their energy heats the background plasma before they are lost by collisional radial heat transport. As this is a more difficult task, collisional radial transport at low collisionality in nearly optimized QS stellarators remains a focus of magnetic fusion efforts (Gates *et al.* 2017; Henneberg *et al.* 2019), with a proxy sometimes used in place of full collisional treatments (Bader *et al.* 2019; Velasco *et al.* 2021) to reduce computational costs. Importantly, recent work by Landreman & Paul (2022) demonstrates that quasiaxisymmetric (QAS) and quasihelically symmetric (QHS) configurations with excellent collisionless alpha confinement are within reach.

Here, earlier work on weakly collisional superbanana (or resonant) plateau and $\sqrt{\nu}$ alpha particle transport (Catto 2019b) is extended to retain magnetic shear, where ν is the pitch angle scattering collision frequency for the trapped alphas. Superbanana plateau transport occurs when the tangential drift vanishes so the alphas encounter a drift resonance. The definition of tangential drift is such that there is no parallel streaming along the tangential drift direction within a flux surface. The resonance enhances the transport caused by any small radial drift departure from quasisymmetry. Because the trapped alpha response must vanish at the passing velocity space boundary, $\sqrt{\nu}$ transport also arises as a result of the same symmetry breaking radial drift. Even though shear is often weak to modest in a stellarator, it is shown to enter the results in a sensitive way. Indeed, shear can result in a merging of the superbanana plateau (Galeev *et al.* 1969; Shaing 2015) and $\sqrt{\nu}$ (Galeev *et al.* 1969; Ho & Kulsrud 1987) transport regimes. The sensitivity arises because shear alters the location in velocity space of the sign change in the tangential drift frequency. The reversal of the tangential drift is responsible for the resonant alphas associated with the superbanana or resonant plateau transport. Depending on the details of the QS stellarator configuration, shear can move the resonance closer to the trapped–passing boundary, where the vanishing of the passing response provides the boundary condition for $\sqrt{\nu}$ transport. It is this feature that causes the merging of the two regimes and allows the two separate regimes to be treated as a single merged regime close to the trapped–passing boundary. The key new aspect of the work presented here is a method for treating the merging of these two regimes in a unified manner. The narrow collisional boundary layer treatment demonstrates that superbanana plateau transport need not dominate over $\sqrt{\nu}$ regime transport (Beidler *et al.* 2011; Calvo *et al.* 2017). Depending on the details of the QS configuration, the shear is typically restricted in sign to justify the narrow boundary layer assumption employed here. The sensitivity to shear arises when it causes the resonant alphas to be so close to the passing boundary that the pitch angle variation of the drift becomes strong enough to reduce the number of resonant alphas.

The narrow boundary layer treatment also allows the formulation of a quasilinear (QL) treatment whereby the radial transport of the alphas alters the background alpha distribution function. This QL equation indicates that a substantial departure from QS is needed for radial transport to modify the unperturbed alpha distribution function from its usual isotropic form.

The details of the treatments in the sections that follow are somewhat involved kinetic theory calculations with the added complication of stellarator geometry. Consequently, it is useful to keep in mind that the departure from QS only matters in the radial drift. Unperturbed QS alpha trajectories are used everywhere else. In addition, the full

alpha collision operator with electron and ion drag and pitch angle scattering by the background ions matters for determining the unperturbed distribution function. However, radial transport introduces velocity space anisotropy, thereby requiring the retention of pitch angle scattering off the background ions to resolve narrow boundary layers associated with the perturbed alpha distribution function.

The superbanana or resonant plateau regime is particularly interesting since it occurs because the tangential drift vanishes, thereby allowing a drift resonance to occur. The occurrence of this resonant plateau behaviour is similar to the effect of the small parallel velocity particles that are responsible for the usual plateau regime of neoclassical transport in a tokamak. Indeed, the superbanana plateau radial diffusivity is independent of collision frequency even though the process responsible for the transport is collisional. To understand how this happens its diffusivity can be estimated by considering the tangential magnetic drift frequency ω_α and the radial drift speed V_r . In a large aspect ratio torus the QS tangential magnetic drift is of order $\omega_\alpha \sim qv_0^2/\Omega_0 Rr$, with v_0 the alpha birth speed, Ω_0 the on axis alpha gyrofrequency, R and r the major and minor radii and q the safety factor. The radial drift due to a departure from QS is of order $V_r \sim \delta qv_0^2/\Omega_0 r$, where $\delta \ll \varepsilon = r/R$ is the error or QS breaking magnetic field amplitude normalized by the on axis magnetic field (Calvo *et al.* 2013, 2014a,b). Diffusive collisions result in a narrow boundary layer of dimensionless width w_{sbp} in pitch angle such that the effective collision frequency in the superbanana plateau regime is $\nu_{sbp} \sim \nu/w_{sbp}^2$, with ν the effective collision frequency for pitch angle scattering of the trapped alphas by the background ions. A resonance occurs because the tangential drift vanishes at some pitch angle with collisions resulting in a spread out drift resonance with an effective width given by $\omega_\alpha w_{sbp}$. Balancing the effective drift and collisions ($\omega_\alpha w_{sbp} \sim \nu_{sbp}$) gives the dimensionless width (Catto & Tolman 2021) to be $w_{sbp} \sim (\nu/\omega_\alpha)^{1/3} \ll 1$, and thereby an enhanced collision frequency of $\nu_{sbp} \sim \nu/w_{sbp}^2 \gg \nu \sim \nu_{\alpha i}/\varepsilon$, where $\nu_{\alpha i}$ is the pitch angle scattering frequency of the alphas by the background ions. The resulting radial step size is then $\Delta = V_r/\nu_{sbp}$. Only the trapped particles in the narrow collisional boundary layer contribute to transport. Defining the inverse aspect ratio as $\varepsilon = r/R$, then this fraction is $w_{sbp}\varepsilon^{1/2}$. Consequently, the diffusivity is seen to be independent of the collision frequency and given by

$$D_{sbp} \sim (w_{sbp}\varepsilon^{1/2})V_r^2/\nu_{sbp} \sim \varepsilon^{1/2}V_r^2/\omega_\alpha \sim \delta^2qv_0^2/\varepsilon^{1/2}\Omega_0, \tag{1.1}$$

due to a cancelation between the boundary layer trapped fraction and the effective collision frequency. In some QAS and QHS configurations it will be shown that magnetic shear can reduce w_{sbp} by a factor that increases exponentially with shear. These configurations are the ones of interest here and for them the superbanana plateau diffusivity is reduced to

$$D_{sbp} \sim \varepsilon^{1/2}V_r^2/\gamma\omega_\alpha \sim \delta^2qv_0^2/\gamma\varepsilon^{1/2}\Omega_0, \tag{1.2}$$

as the shear reduction factor γ is large. The reduction occurs because very few barely trapped alphas are able to resonate with the drift once the drift varies strongly with pitch angle very near the trapped–passing boundary.

The estimate for the $\sqrt{\nu}$ regime is a bit more straightforward to obtain since the tangential drift ω_α does not vanish so the effective collision frequency is $\nu_{\sqrt{\nu}} \sim \nu/w_{\sqrt{\nu}}^2$. The dimensionless width $w_{\sqrt{\nu}}$ is found from $\omega_\alpha \sim \nu/w_{\sqrt{\nu}}^2$ to be $w_{\sqrt{\nu}} \sim (\nu/\omega_\alpha)^{1/2} \ll 1$. In this case, the trapped fraction in the narrow boundary layer is $w_{\sqrt{\nu}}\varepsilon^{1/2}$, and results in the smaller diffusivity of

$$D_{\sqrt{\nu}} \sim (w_{\sqrt{\nu}}\varepsilon^{1/2})V_r^2/\nu_{\sqrt{\nu}} \sim w_{\sqrt{\nu}}\varepsilon^{1/2}V_r^2/\omega_\alpha \sim \gamma w_{\sqrt{\nu}}D_{sbp}. \tag{1.3}$$

Magnetic shear will be shown to have little effect on this estimate, but as the shear reduction factor is large $D_{sbp} \gg D_{\sqrt{\nu}}$ may no longer hold.

Results very similar to the preceding will be recovered in the merged regime treatment by assuming that radial transport does not modify the unperturbed distribution function significantly.

The sections that follow begin by introducing notation and briefly presenting some background for general stellarators in § 2 and QS stellarators in § 3. In § 4 an average over the tangential angle variable α is introduced and used to obtain the energy and particle balance equations with radial transport retained. Section 5 uses the α average to derive the QL equation for the unperturbed distribution function and the linearized equation for the perturbed distribution function. Simplification of the linearized kinetic equation for a narrow boundary layer is performed in § 6 and, except for the retention of finite drift effects to retain magnetic shear, is similar to the procedure used by Catto (2019b). A unified solution of the merged superbanana plateau and $\sqrt{\nu}$ regime is presented in § 7. Section 8 derives the QL velocity space dependent radial diffusivity needed to complete the derivation of the full QL equation for the unperturbed distribution function. In § 9 the alpha particle energy diffusivity is evaluated for the merged superbanana plateau and $\sqrt{\nu}$ regime by using the usual slowing down tail alpha distribution function. The results indicate that magnetic shear in some QS configurations can be used to reduce superbanana plateau transport below the $\sqrt{\nu}$ transport level. The last section summarizes and discusses the results.

2. General stellarator properties

A general stellarator magnetic field depends on the poloidal flux function ψ_p , and the poloidal and toroidal angles ϑ and ζ , respectively. Introducing the angle variable

$$\alpha = \zeta - q\vartheta = \zeta - \ell^{-1}\vartheta, \quad (2.1)$$

the Boozer (1981) and Clebsch representations for a general stellarator magnetic field are

$$\mathbf{B} = B\mathbf{b} = \nabla\alpha \times \nabla\psi_p = K_p\nabla\psi_p + G\nabla\vartheta + I\nabla\zeta, \quad (2.2)$$

with $\mathbf{B} \cdot \nabla\psi_p = 0 = \mathbf{B} \cdot \nabla\alpha$, q the safety factor and ℓ the rotational transform flux functions, and

$$\mathbf{B} \cdot \nabla\zeta = q\nabla\psi_p \times \nabla\vartheta \cdot \nabla\zeta = q\mathbf{B} \cdot \nabla\vartheta = qB^2/(qI + G). \quad (2.3)$$

The coefficients I and G are flux functions and related to the toroidal current enclosed by a magnetic field line or flux surface

$$\int_{tor} d^2\mathbf{r} \cdot \mathbf{J} = \int_0^{\psi_p} d\psi_p \int_0^{2\pi} d\vartheta \frac{\nabla\zeta \cdot \mathbf{J}}{\mathbf{B} \cdot \nabla\vartheta} = \frac{cG}{2}, \quad (2.4)$$

and the poloidal current outside the flux surface

$$\int_{pol} d^2\mathbf{r} \cdot \mathbf{J} = \int_0^{\psi_p} d\psi_p \int_0^{2\pi} d\zeta \frac{\nabla\vartheta \cdot \mathbf{J}}{\mathbf{B} \cdot \nabla\vartheta} = \frac{cI}{2}. \quad (2.5)$$

In the preceding, unlike in usual stellarator notation, I is used with $\nabla\zeta$ to more closely conform to tokamak notation.

3. Quasisymmetric stellarator properties

The magnetic field $B = |\mathbf{B}|$ of a QS stellarator depends on the flux and is periodic in a single angle variable

$$\eta = M\vartheta - N\zeta, \tag{3.1}$$

where $M \geq 0$ and $N \geq 0$ with $N=0$ and $M=1$ for a tokamak. On a QS flux surface a point on a constant B contour closes on itself after ϑ increases by $2\pi N$ and ζ increases by $2\pi M$. Defining the helical magnetic flux function (Boozer 1983; Landreman & Catto 2011) as

$$\psi_h = M\psi_p - N\psi_t, \tag{3.2}$$

with the poloidal flux function ψ_p and toroidal flux function ψ_t related by

$$\partial\psi_t/\partial\psi_p = q = \varphi^{-1}, \tag{3.3}$$

then

$$\nabla\psi_h = (M - qN)\nabla\psi_p. \tag{3.4}$$

The special case of $M=0$ is often referred to as quasi-poloidal symmetry (QPS), the $N=0$ case is referred to as quasi-axisymmetry (QAS includes, but is not limited to, strict axisymmetry), and the general $N \neq 0 \neq M$ case is referred to as quasi-helical symmetry (QHS). The preceding imply that the Boozer (1981) representations for a QS stellarator magnetic field may also be written as

$$\mathbf{B} = B\mathbf{b} = (M - qN)^{-1}\nabla\alpha \times \nabla\psi_h = K_h(\psi_h, \vartheta, \zeta)\nabla\psi_h + G\nabla\vartheta + I\nabla\zeta, \tag{3.5}$$

with $K_p = (M - qN)K_h$

$$\mathbf{B} \cdot \nabla\eta = (M - qN)\mathbf{B} \cdot \nabla\vartheta = (M - qN)B^2/(qI + G), \tag{3.6}$$

and

$$\mathbf{B} \cdot \nabla\zeta = q(M - qN)^{-1}\nabla\psi_h \times \nabla\vartheta \cdot \nabla\zeta = q\mathbf{B} \cdot \nabla\vartheta = qB^2/(qI + G). \tag{3.7}$$

In terms of α and η

$$\vartheta = (N\alpha + \eta)/(M - qN), \tag{3.8}$$

and

$$\zeta = (M\alpha + q\eta)/(M - qN). \tag{3.9}$$

For fixed α , η changes by 2π when $\vartheta \rightarrow \vartheta + 2\pi/(M - qN)$ and $\zeta \rightarrow \zeta + 2\pi q/(M - qN)$.

In a QS stellarator the drift kinetic canonical angular momentum constant of the motion (Boozer 1983; see Appendix A of Landreman & Catto 2011) is

$$\psi_* = \psi_h - I_h v_{||}/\Omega, \tag{3.10}$$

where $\Omega = ZeB/M_\alpha c$ and

$$I_h = MI + NG, \tag{3.11}$$

with Z the alpha charge number, M_α the alpha mass, e the charge on a proton and c the speed of light. As in a tokamak, the drift kinetic angular momentum in ψ_h, ϑ, ζ variables

satisfies

$$(v_{\parallel} \mathbf{b} + \mathbf{v}_d) \cdot \nabla \psi_* = 0, \tag{3.12}$$

where the combined magnetic and electric drifts used are

$$\mathbf{v}_d = \Omega^{-1} \mathbf{b} \times (\mu \nabla B + v_{\parallel}^2 \mathbf{b} \cdot \nabla \mathbf{b}) + cB^{-1} \mathbf{b} \times \nabla \Phi \simeq \Omega^{-1} v_{\parallel} \nabla \times (v_{\parallel} \mathbf{b}), \tag{3.13}$$

in magnetic moment, $\mu = v_{\perp}^2/2B$, and total energy variables

$$E = v^2/2 + Ze\Phi/M_{\alpha}, \tag{3.14}$$

for which $v_{\parallel}^2 = 2(E - Ze\Phi/M_{\alpha} - \mu B)$, with Φ the electrostatic potential. The parallel component of the last form of \mathbf{v}_d is negligible compared with parallel streaming so $\mathbf{v}_d \simeq \Omega^{-1} v_{\parallel} \nabla_{\perp} \times (v_{\parallel} \mathbf{b})$ may be employed.

4. Alpha transport for nearly QS stellarators

In E , μ and φ (= gyrophase) velocity space variables the drift kinetic equation for alpha particles is simply

$$(v_{\parallel} \mathbf{b} + \mathbf{v}_d) \cdot \nabla f = C\{f\} + \frac{S(\psi_p)\delta(v - v_0)}{4\pi v^2}, \tag{4.1}$$

where the flux function $S = n_T n_D \langle \sigma v \rangle_{DT}$ is the birth rate of the alphas born isotropically in velocity space at the birth speed v_0 , and

$$C\{f\} = \frac{1}{\tau_s} \nabla_v \cdot \left[\left(\frac{v^3 + v_c^3}{v^3} \right) \mathbf{v} f + \frac{v_{\lambda}^3}{2v^3} (v^2 \vec{I} - \mathbf{v} \mathbf{v}) \cdot \nabla_v f \right], \tag{4.2}$$

is the collision operator retaining electron and ion drag and pitch angle scattering by ions, with

$$\tau_s = \frac{3M_i T_e^{3/2}}{4(2\pi m_e)^{1/2} Z_i^2 e^4 n_e \ell n \Lambda}, \tag{4.3}$$

the slowing down time, v_c the critical speed defined by

$$v_c^3 = \frac{3\pi^{1/2} T_e^{3/2}}{(2m_e)^{1/2} n_e} \sum_i \frac{Z_i^2 n_i}{M_i}, \tag{4.4}$$

and $\tau_s v^3 v_{\lambda}^{-3}$ the pitch angle scattering time, where

$$v_{\lambda}^3 = \frac{3\pi^{1/2} T_e^{3/2}}{(2m_e)^{1/2} M_{\alpha} n_e} \sum_i Z_i^2 n_i. \tag{4.5}$$

The electron temperature T_e and the electron and ion densities n_e and n_i are taken to be flux functions and quasineutrality ($n_e \simeq \sum_i Z_i n_i$) is employed. The masses of the background ions and electrons are M_i and m_e , respectively, and Z_i is the charge number of the ions with \sum_i denoting a sum over all background ion species. The alpha particle collision operator is obtained by expanding unlike collision operators for $v_i \ll v_{\lambda} \sim v_c \ll v_0 \ll v_e$, making energy scattering of alphas by ions and electrons small because $v^2 \gg v_i^2 = 2T_i/M_i$ and $v^2 \ll v_e^2 = 2T_e/m_e$, respectively.

In an imperfect stellarator B , and therefore v_{\parallel} , depends on α as well as η and the flux. Using α and η from (3.8) and (3.9) in the Boozer form of (3.5) along with (3.4) and (3.6) gives

$$\begin{aligned} \mathbf{v}_d \cdot \nabla \psi_h &= \frac{v_{\parallel}}{B} \left[\nabla \alpha \times \nabla \psi_p \cdot \nabla \left(\frac{I_h v_{\parallel}}{\Omega} \right) + (qI + G) \nabla \eta \times \nabla \psi_p \cdot \nabla \left(\frac{v_{\parallel}}{\Omega} \right) \right] \\ &= v_{\parallel} \mathbf{b} \cdot \nabla \left(\frac{I_h v_{\parallel}}{\Omega} \right) - (M - qN) v_{\parallel} B \frac{\partial}{\partial \alpha} \left(\frac{v_{\parallel}}{\Omega} \right). \end{aligned} \tag{4.6}$$

In the final form of (4.6), the last term proportional $\partial/\partial\alpha$ is the drive term that arises from the lack of QS and the first term proportional to $\mathbf{b} \cdot \nabla$ is the neoclassical drive term. As the departure from QS is assumed small, the QS trajectories may be used to evaluate all other terms in the drift kinetic equation. Consequently, departures from QS must be retained in the radial drift, while unperturbed QS trajectories are employed elsewhere.

As the departure from QS is assumed small, the lowest order alpha distribution function f_0 cannot depend on α . Consequently,

$$f = f(\psi_p, \eta, \alpha, E, \mu, \sigma) = f_0(\psi_p, \eta, E, \mu, \sigma) + f_1 + \dots, \tag{4.7}$$

with $f_0 \gg f_1$. Then, compared with parallel streaming, the radial and tangential drifts are small and the source term and collisions are weak, giving to lowest order

$$v_{\parallel} \mathbf{b} \cdot \nabla f_0 = v_{\parallel} \mathbf{b} \cdot \nabla \eta \partial f_0 / \partial \eta = 0. \tag{4.8}$$

Therefore, for small departures from QS, $\partial f_0 / \partial \eta = 0$ as well as $\partial f_0 / \partial \alpha = 0$, leading to

$$f_0 = f_0(\psi_p, E, \mu), \tag{4.9}$$

with no dependence on σ as the alphas are born isotropically.

Going to next order by retaining $\partial f_1 / \partial \psi_p$ to allow radial transport to modify f_0 , the kinetic equation becomes

$$\begin{aligned} (v_{\parallel} \mathbf{b} + \mathbf{v}_d) \cdot \nabla f_1 + \mathbf{v}_d \cdot \nabla \psi_h \frac{\partial f_0}{\partial \psi_h} &= C\{f_0 + f_1\} + \frac{S\delta(v - v_0)}{4\pi v^2} \\ &= \frac{v_{\parallel}}{B} \nabla \cdot \left[\frac{B}{v_{\parallel}} (v_{\parallel} \mathbf{b} + \mathbf{v}_d) (f_0 + f_1) \right], \end{aligned} \tag{4.10}$$

where the alternate form on the far right follows by using

$$B v_{\parallel}^{-1} \mathbf{v}_d \cdot \nabla f_1 = \nabla \cdot (B v_{\parallel}^{-1} \mathbf{v}_d f_1), \tag{4.11}$$

and $C\{f_0 + f_1\} = C\{f_0\} + C\{f_1\}$. The usual slowing down distribution function

$$f_0 = \frac{S\tau_s H(v_0 - v)}{4\pi(v^3 + v_c^3)}, \tag{4.12}$$

with H the Heaviside step function, satisfies $4\pi v^2 C\{f_0\} + S\delta(v - v_0) = 0$. However, strong radial losses in (4.10) cause a departure from the isotropic behaviour of (4.12).

The right most form of the drift kinetic operator is convenient for forming flux surface averaged moments. To do so, use $d^3v \rightarrow 2\pi \sum_{\sigma} B dE d\mu / |v_{\parallel}| = 2\pi \sum_{\sigma} B v^2 dv d\lambda / B_0 |\xi|$,

with \sum_σ a reminder to sum over both signs of $\sigma = v_{||}/|v_{||}|$, $\xi = v_{||}/v$, $\lambda = 2\mu B_0/v^2$, and the flux surface average of an arbitrary function A as defined by

$$\langle A \rangle = \oint \frac{d\vartheta d\zeta A}{\mathbf{B} \cdot \nabla \vartheta} / \oint \frac{d\vartheta d\zeta}{\mathbf{B} \cdot \nabla \vartheta} = \oint \frac{d\eta d\alpha A}{\mathbf{B} \cdot \nabla \eta} / \oint \frac{d\eta d\alpha}{\mathbf{B} \cdot \nabla \eta}, \tag{4.13}$$

where $d\eta d\alpha = (M - qN) d\vartheta d\zeta$. Then allowing for a collisional loss sink L at small speeds where the alphas become ash due to ion drag, the continuity equation leads to the expected result that radial particle loss reduces the amount of ash

$$\frac{1}{V'} \frac{\partial}{\partial \psi_p} \left[V' \left\langle \int d^3 v f_1 \mathbf{v}_d \cdot \nabla \psi_p \right\rangle \right] = S - L, \tag{4.14}$$

where $V' = \oint d\vartheta d\zeta / \mathbf{B} \cdot \nabla \vartheta$, (3.13), $\langle \int d^3 v f_0 \mathbf{v}_d \cdot \nabla \psi_p \rangle = \langle \Omega^{-1} \int d^3 v v_{||} \nabla \cdot (f_0 v_{||} \mathbf{b} \times \nabla \psi_p) \rangle = 0$ and

$$\langle \nabla \cdot \mathbf{A} \rangle = \frac{1}{V'} \frac{\partial}{\partial \psi_p} (V' \langle \mathbf{A} \cdot \nabla \psi_p \rangle), \tag{4.15}$$

are employed. The loss into a small velocity space sphere about the origin is

$$L \equiv - \left\langle \int d^3 v C \{f_0 + f_1\} \right\rangle = -\tau_s^{-1} \oint d^3 v \nabla_v \cdot \left[\mathbf{v} \left(\frac{v^3 + v_c^3}{v^3} \right) f \right] \xrightarrow[\text{xpert}]{no} S, \tag{4.16}$$

where $\nabla_v \cdot (\mathbf{v} v^{-3} f) \xrightarrow[v \rightarrow 0]{} -4\pi \delta(\mathbf{v}) f(\psi, v = 0)$. In the absence of alpha transport loss $L = S$ as all the alphas become ash.

A related, but more important, result holds for energy conservation

$$\frac{1}{V'} \frac{\partial}{\partial \psi_p} \left[V' \left\langle \frac{M_\alpha}{2} \int d^3 v f_1 v^2 \mathbf{v}_d \cdot \nabla \psi_p \right\rangle \right] = \frac{1}{2} M_\alpha v_0^2 S - E, \tag{4.17}$$

with E the alpha energy loss to the background ions and electrons

$$E \equiv -\frac{M_\alpha}{2} \left\langle \int d^3 v v^2 C \{f\} \right\rangle = \frac{M_\alpha}{\tau_s} \int d^3 v v^{-1} (v^3 + v_c^3) f \xrightarrow[\text{xpert}]{no} \frac{1}{2} M_\alpha v_0^2 S. \tag{4.18}$$

To avoid severe depletion, radial alpha energy losses must be kept small compared with $M_\alpha v_0^2 S/2$.

5. Quasilinear theory for nearly QS stellarators

Quasilinear theory allows the unperturbed distribution to be altered if the radial transport is large. To obtain this QL feature the form of the drift kinetic equation must be simplified further. To do so it is necessary to introduce the α average of an arbitrary function A as

$$\begin{aligned} \langle A \rangle_\alpha &= \oint d\alpha A / \oint d\alpha \equiv \int_\alpha^{\alpha+2\pi(M-qN)} d\alpha' A / \int_\alpha^{\alpha+2\pi(M-qN)} d\alpha' \\ &= \int_\alpha^{\alpha+2\pi(M-qN)} d\alpha' A / 2\pi(M - qN), \end{aligned} \tag{5.1}$$

where the integration on a fixed flux surface is over a closed loop on a constant B contour that closes back on itself when $\vartheta \rightarrow \vartheta + 2\pi N$ and $\zeta \rightarrow \zeta + 2\pi M$ (causing $\alpha \rightarrow \alpha + 2\pi(M - qN)$). Such B contours exist by definition for a QS magnetic field.

The drive term (4.6) in the kinetic equation is the sum of two contributions

$$\mathbf{v}_d \cdot \nabla \psi_h = \mathbf{v}_d \cdot \nabla \psi_h|_{nc} + \mathbf{v}_d \cdot \nabla \psi_h|_{im}, \tag{5.2}$$

with

$$\mathbf{v}_d \cdot \nabla \psi_h|_{nc} = v_{||} \mathbf{b} \cdot \nabla (I_h v_{||} / \Omega), \tag{5.3}$$

the usual QS neoclassical drive term, and

$$\mathbf{v}_d \cdot \nabla \psi_h|_{im} = -(M - qN) v_{||} B \frac{\partial}{\partial \alpha} \left(\frac{v_{||}}{\Omega} \right), \tag{5.4}$$

the drive term due to the departure from QS. Therefore, f_1 is written as the sum of the neoclassical (nc) and non-QS (im) contributions

$$f_1 = f_1^{nc} + f_1^{im}, \tag{5.5}$$

with $f_1^{nc} \equiv \langle f_1 \rangle_\alpha$ and $\langle f_1^{im} \rangle_\alpha = 0$. Then the neoclassical portion satisfies

$$v_{||} \mathbf{b} \cdot \nabla f_1^{nc} + \mathbf{v}_d \cdot \nabla \psi_h|_{nc} \frac{\partial f_0}{\partial \psi_h} = C\{f_1^{nc}\} = v_{||} \mathbf{b} \cdot \nabla \left(f_1^{nc} + \frac{I_h v_{||}}{\Omega} \frac{\partial f_0}{\partial \psi_h} \right), \tag{5.6}$$

where $\partial f_1^{nc} / \partial \alpha = 0$ and the $\mathbf{v}_d \cdot \nabla \eta \partial f_1^{nc} / \partial \eta$ term is ignored as negligible. The solution to the neoclassical equation for alphas in a QS stellarator does not involve any resonant particles and can be obtained from Hsu *et al.* (1990) and Catto (2018) using the isomorphism of stellarators with tokamaks (Boozer 1983; Landreman & Catto 2011). It is of no concern here.

Subtracting the neoclassical equation from the full equation and continuing to keep $\partial f_1 / \partial \psi_h$, leaves

$$\frac{v_{||}}{B} \nabla \cdot \left[\frac{B}{v_{||}} (v_{||} \mathbf{b} + \mathbf{v}_d) f_1^{im} \right] + \mathbf{v}_d \cdot \nabla \psi_h|_{im} \frac{\partial (f_0 + f_1^{nc})}{\partial \psi_h} = C\{f_0 + f_1^{im}\} + \frac{S\delta(v - v_0)}{4\pi v^2}, \tag{5.7}$$

where the ash sink is implicit as $\nabla_v \cdot (\mathbf{v} v^{-3} f_0) \rightarrow -4\pi \delta(\mathbf{v}) f_0(\psi, v = 0)$ and $\int d^3 v \delta(\mathbf{v}) = 1$. Departures from a QS magnetic field are assumed to satisfy stellarator symmetry. Upon Fourier decomposition they may be written as proportional to $\cos(m\vartheta - n\zeta)$ with $m \neq M$ and $n \neq N$. As a result, the non-QS drive terms are proportional to $\sin(m\vartheta - n\zeta)$, where

$$\chi \equiv m\vartheta - n\zeta = [(mN - nM)\alpha + (m - qn)\eta] / (M - qN). \tag{5.8}$$

Then f_1^{im} is the portion of f_1 periodic in α such that $\langle f_1^{im} \rangle_\alpha = 0$, $\langle \partial f_1^{im} / \partial \alpha \rangle_\alpha = 0$ and $\langle \mathbf{v}_d \cdot \nabla \psi_h|_{im} / v_{||} \mathbf{b} \cdot \nabla \eta \rangle_\alpha = 0$ (recall $\mathbf{b} \cdot \nabla \eta \propto B$). Consequently, writing out the divergence on the left of (5.7) in ψ_h, η, α variables, dividing by $v_{||} \mathbf{b} \cdot \nabla \eta$, and averaging over α as in (5.1), leaves

$$\frac{1}{\oint d\alpha} \frac{\partial}{\partial \psi_h} \oint d\alpha \frac{f_1^{im} \mathbf{v}_d \cdot \nabla \psi_h|_{im}}{v_{||} \mathbf{b} \cdot \nabla \eta} + \frac{\partial}{\partial \eta} \left\langle \frac{f_1^{im} (v_{||} \mathbf{b} + \mathbf{v}_d) \cdot \nabla \eta}{v_{||} \mathbf{b} \cdot \nabla \eta} \right\rangle_\alpha = \left\langle \frac{C\{f_0\}}{v_{||} \mathbf{b} \cdot \nabla \eta} \right\rangle_\alpha + \left\langle \frac{S\delta(v - v_0)}{4\pi v^2 v_{||} \mathbf{b} \cdot \nabla \eta} \right\rangle_\alpha, \tag{5.9}$$

where for narrow collisional boundary layers $\langle B^{-1} v_{||}^{-1} C\{f_1^{im}\} \rangle_\alpha \simeq B^{-1} v_{||}^{-1} C\{f_1^{im}\} = 0$ is assumed.

A final simplification follows by integrating η over a full bounce for the trapped and over a fixed α path for which η changes by 2π for the passing. Periodicity in η is then used to obtain the QL equation to evaluate f_0 once the solution f_1^{im} is found, namely,

$$\frac{1}{\oint d\alpha} \frac{\partial}{\partial \psi_p} \oint d\alpha \oint_{\alpha} \frac{d\eta f_1^{im} \mathbf{v}_d \cdot \nabla \psi_p|_{im}}{v_{||} \mathbf{b} \cdot \nabla \eta} = \left\langle \oint_{\alpha} \frac{d\eta C\{f_0\}}{v_{||} \mathbf{b} \cdot \nabla \eta} \right\rangle_{\alpha} + \frac{S\delta(v - v_0)}{4\pi v^2} \left\langle \oint_{\alpha} \frac{d\eta}{v_{||} \mathbf{b} \cdot \nabla \eta} \right\rangle_{\alpha}. \tag{5.10}$$

The subscript on the integral is a reminder that it is performed at fixed α . This QL equation for f_0 implies that depletion due to strong ψ_p variation of $f_1^{im} \mathbf{v}_d \cdot \nabla \psi_p|_{im}$ caused by α variation can alter the alpha distribution function for large enough departures from QS. The radial transport term on the left must remain unimportant to control alpha particle losses in a stellarator.

Subtracting the α average of (5.7) from the full equation yields the f_1^{im} equation

$$\begin{aligned} (v_{||} \mathbf{b} + \mathbf{v}_d) \cdot \nabla \eta \frac{\partial f_1^{im}}{\partial \eta} + \mathbf{v}_d \cdot \nabla \alpha \frac{\partial f_1^{im}}{\partial \alpha} + \mathbf{v}_d \cdot \nabla \psi_{h|nc} \frac{\partial f_1^{im}}{\partial \psi_h} - C\{f_1^{im}\} + \mathbf{v}_d \cdot \nabla \psi_{h|im} \frac{\partial f_0}{\partial \psi_h} \\ \simeq v_{||} \mathbf{b} \cdot \nabla \eta \left\langle \frac{\mathbf{v}_d \cdot \nabla \psi_{h|im}}{v_{||} \mathbf{b} \cdot \nabla \eta} \frac{\partial f_1^{im}}{\partial \psi_h} \right\rangle_{\alpha} - \mathbf{v}_d \cdot \nabla \psi_{h|im} \frac{\partial f_1^{im}}{\partial \psi_h}, \end{aligned} \tag{5.11}$$

where $\partial f_1^{nc} / \partial \psi_h \ll \partial f_0 / \partial \psi_h$ is assumed. In a standard QL treatment the terms on the right side of this equation are ignored. The neglect corresponds to an assumption that the departure from QS is small. Therefore, only the remaining linear equation,

$$(v_{||} \mathbf{b} + \mathbf{v}_d) \cdot \nabla f_1^{im} - C\{f_1^{im}\} = -\mathbf{v}_d \cdot \nabla \psi_{h|im} \frac{\partial f_0}{\partial \psi_h} = v_{||} B \frac{\partial}{\partial \alpha} \left(\frac{v_{||}}{\Omega} \right) \frac{\partial f_0}{\partial \psi_p}, \tag{5.12}$$

is to be solved. The passing alphas trace out flux surfaces and, as $v_{||} \mathbf{b} \cdot \nabla f_1^{im} \gg \mathbf{v}_d \cdot \nabla f_1^{im} \gg C\{f_1^{im}\}$ for all passing particles, little to no transport occurs. The trapped alphas are unable to trace out flux surfaces. Moreover, the tangential drift in a flux surface is known to change sign at some pitch angle. Consequently, it is these trapped alphas that give the dominant contribution to the radial transport. Therefore, the next task is to further simplify this kinetic equation for the trapped alphas by using the unperturbed QS trajectories on the left side based on the approximations already outlined.

6. Formulation of the linearized drift kinetic equation for nearly QS stellarators

As the unperturbed QS trajectories are adequate on the left side, $(v_{||} \mathbf{b} + \mathbf{v}_d) \cdot \nabla \psi_* = 0$ is used there. Then, to retain radial drift departures from QS it is convenient to use the variables ψ_*, η and

$$\alpha_* \equiv \zeta - q_* \vartheta = \zeta - \ell_*^{-1} \vartheta, \tag{6.1}$$

along with E and μ , with $q_* = q_*(\psi_*)$, $q = q(\psi_h)$ and $d\psi_h = (M - qN) d\psi_p$. Using (3.12)

$$\begin{aligned} \omega_{\alpha} \equiv (v_{||} \mathbf{b} + \mathbf{v}_d) \cdot \nabla \alpha_* = \mathbf{v}_d \cdot \nabla \zeta - q_* \mathbf{v}_d \cdot \nabla \vartheta - (q_* - q) v_{||} \mathbf{b} \cdot \nabla \vartheta \\ \simeq \frac{B v_{||}}{(qI + G)} \left[\frac{\partial}{\partial \psi_p} \left(\frac{G v_{||}}{\Omega} \right) + q \frac{\partial}{\partial \psi_p} \left(\frac{I v_{||}}{\Omega} \right) + \frac{I_h v_{||}}{(M - qN) \Omega} \frac{\partial q}{\partial \psi_p} - (M - qN) \frac{\partial}{\partial \eta} \left(\frac{K_h v_{||}}{\Omega} \right) \right], \end{aligned} \tag{6.2}$$

where magnetic shear effects are retained (unlike Catto 2019b), and $q_* \mathbf{v}_d \cdot \nabla \vartheta \simeq q \mathbf{v}_d \cdot \nabla \vartheta$ and

$$q_* \simeq q(\psi_h) + (\psi_* - \psi_h) \partial q / \partial \psi_h = q(\psi_h) - I_h \Omega^{-1} v_{\parallel} \partial q / \partial \psi_h, \tag{6.3}$$

are employed. Using α_* instead of α removes the awkward secular term $\vartheta \mathbf{v}_d \cdot \nabla q \neq 0$ as $\vartheta (v_{\parallel} \mathbf{b} + \mathbf{v}_d) \cdot \nabla q_* = 0$ in ω_α . Consequently, the drift kinetic equation for the trapped alphas with finite orbit effects retained becomes

$$v_{\parallel} \mathbf{b} \cdot \nabla \eta \frac{\partial f_1^{im}}{\partial \eta} + \omega_\alpha \frac{\partial f_1^{im}}{\partial \alpha_*} - C\{f_1^{im}\} \simeq v_{\parallel} B \frac{\partial}{\partial \alpha} \left(\frac{v_{\parallel}}{\Omega} \right) \frac{\partial f_0}{\partial \psi_p} = -\frac{v^2}{\Omega} \left(1 - \frac{\lambda B}{2B_0} \right) \frac{\partial B}{\partial \alpha} \frac{\partial f_0}{\partial \psi_p}, \tag{6.4}$$

where the drift correction to the streaming term is neglected as small and now

$$f_1^{im} = f_1^{im}(\psi_*, \eta, \alpha_*, v, \lambda). \tag{6.5}$$

For alphas the electrostatic potential is unimportant and it is convenient to introduce the pitch angle variable

$$\lambda = 2\mu B_0 / v^2, \tag{6.6}$$

with

$$B_0^2 = \langle B^2 \rangle, \tag{6.7}$$

so that

$$v_{\parallel}^2 = v^2 (1 - \lambda B / B_0). \tag{6.8}$$

As the drive term due to the departure from QS is small and parallel streaming dominates the left side of the equation, to lowest order $\partial f_1^{im} / \partial \eta = 0$. Letting $f_1^{im} = \bar{f}_1^{im} + \tilde{f}_1^{im}$ with $\partial \tilde{f}_1^{im} / \partial \eta = 0$ and $f_1^{im}(\psi_p, \alpha, v, \lambda) \gg \tilde{f}_1^{im}$, the next order equation is

$$v_{\parallel} \mathbf{b} \cdot \nabla \eta \frac{\partial \tilde{f}_1^{im}}{\partial \eta} + \omega_\alpha \frac{\partial \tilde{f}_1^{im}}{\partial \alpha_*} - C\{\tilde{f}_1^{im}\} \simeq v_{\parallel} B \frac{\partial}{\partial \alpha} \left(\frac{v_{\parallel}}{\Omega} \right) \frac{\partial f_0}{\partial \psi_p}, \tag{6.9}$$

where the distinction between α_* and α no longer matters to the requisite order, collisions are weak and $v_{\parallel} \mathbf{b} \cdot \nabla \eta \gg \omega_\alpha$ is assumed.

To annihilate the streaming term integration along the characteristics associated with the alpha trajectories at fixed α is employed by defining the total trajectory derivatives using

$$d\eta(\tau) / d\tau = v_{\parallel} \mathbf{b} \cdot \nabla \eta \simeq v_{\parallel} (M - qN) B / qI, \tag{6.10}$$

with $\eta(\tau = 0) = \eta$ and $\eta(\tau_0) = 0$ at the closed minimum B contour of the lowest order QS magnetic field (recall that unperturbed QS trajectories are being used on the left side of the kinetic equation). Transit or bounce averaging over a full bounce for the trapped alphas leads to the reduced kinetic equation

$$\left(\oint_{\alpha} d\tau \omega_{\alpha} \right) \frac{\partial \bar{f}_1^{im}}{\partial \alpha_*} - \oint_{\alpha} d\tau C\{\bar{f}_1^{im}\} \simeq \frac{qI}{M - qN} \frac{\partial}{\partial \alpha} \left(\oint_{\alpha} \frac{d\eta v_{\parallel}}{\Omega} \right) \frac{\partial f_0}{\partial \psi_p}, \tag{6.11}$$

where $qI \gg G$ is employed and the transit average subscript is a reminder to hold α fixed. The usual superbanana plateau resonance and \sqrt{v} behaviour occurs for the trapped when $\oint_{\alpha} d\tau \omega_{\alpha}$ changes sign and near the passing boundary, respectively. Consequently, trapped particle contributions to transport will always occur.

From here on the focus is on evaluating the superbanana plateau and \sqrt{v} transport for the trapped alphas. To proceed analytically it is convenient to assume a large aspect ratio stellarator by using the approximations $B_t \simeq B_0$ and $I \simeq RB_0$, where B_0 is a constant equal to the lowest order toroidal magnetic field while the constant R is the nominal major radius. Then, to lowest order, $\psi_t \simeq B_0 a(r)$, where $a = a(r)$ is the area enclosed by the flux surface labelled by r , and $\partial/\partial\psi_p = q\partial/\partial\psi_t \simeq (q/B_0 a')\partial/\partial r$ with $a' = da/dr \sim r$ and $a^2 \sim r^2$. Using $G/qI \ll 1$, $NG \ll MI$ and $\Omega_0 \equiv ZeB_0/M_\alpha c$,

$$\begin{aligned} \omega_\alpha + v_{||} \mathbf{b} \cdot \nabla_\eta \frac{\partial}{\partial \eta} \left(\frac{K_h v_{||}}{\Omega} \right) \\ \simeq \frac{B_0}{2\Omega_0} \left[\frac{\partial v_{||}^2}{\partial \psi_p} + \frac{2I_h v_{||}^2}{(M - qN)Iq} \frac{\partial q}{\partial \psi_p} \right] \simeq \frac{q}{2\Omega_0 a'} \left[\frac{\partial v_{||}^2}{\partial r} + \frac{2M v_{||}^2}{(M - qN)q} \frac{\partial q}{\partial r} \right]. \end{aligned} \tag{6.12}$$

Continuing to assume large aspect ratio by taking

$$B = B_0 \left[1 - \varepsilon \cos(M\vartheta - N\zeta) + \sum_{m,n} \delta_m^n \cos(m\vartheta - n\zeta) \right], \tag{6.13}$$

with $1 \gg \varepsilon \simeq r/R \gg \delta_m^n$, and \sum denoting the sums over $n \neq N \geq 0$ and $m \neq M \geq 0$ ($n = N$ and $m \neq M$ or $n \neq N$ and $m = M$ represent field departures from QS). Then

$$\frac{\partial B}{\partial \alpha} = -B_0 \sum_{m,n} \frac{mN - nM}{M - qN} \delta_m^n \sin \chi. \tag{6.14}$$

For the unperturbed QS trajectories using $B \simeq B_0(1 - \varepsilon \cos \eta)$ gives

$$\omega_\alpha \simeq \frac{qv^2}{2\Omega_0 R a'} \left[\lambda \cos \eta + \frac{2sM}{(M - qN)\varepsilon} (1 - \lambda + \lambda \varepsilon \cos \eta) \right] - v_{||} \mathbf{b} \cdot \nabla_\eta \frac{\partial}{\partial \eta} \left(\frac{K_h v_{||}}{\Omega} \right), \tag{6.15}$$

with the magnetic shear defined as

$$s \equiv \frac{a' \partial q}{q \partial r}. \tag{6.16}$$

To evaluate the transit averages it is convenient to define

$$\xi = v_{||}/v = \sqrt{1 - \lambda B/B_0} = \sqrt{1 - \lambda(1 - \varepsilon \cos \eta)}, \tag{6.17}$$

and use the QS trapped results by letting $\sin(\eta/2) = \kappa \sin x$ to find

$$\begin{aligned} \oint_\alpha d\eta/\xi &= 8(2\varepsilon\lambda)^{-1/2} \int_0^{\pi/2} dx/\sqrt{1 - \kappa^2 \sin^2 x} \\ &= 8(2\varepsilon)^{-1/2} \sqrt{1 - \varepsilon + 2\varepsilon\kappa^2} K(\kappa) \simeq 8(2\varepsilon)^{-1/2} K(\kappa), \end{aligned} \tag{6.18}$$

$$\oint_\alpha d\eta\xi = \frac{8\sqrt{2\varepsilon\kappa^2}}{\sqrt{1 - \varepsilon + 2\varepsilon\kappa^2}} \int_0^{\pi/2} \frac{dx \cos^2 x}{\sqrt{1 - \kappa^2 \cos^2 x}} \simeq 8\sqrt{2\varepsilon} [E(\kappa) - (1 - \kappa^2)K(\kappa)], \tag{6.19}$$

and

$$\oint_{\alpha} d\eta \cos \eta / \xi = 8(2\varepsilon\lambda)^{-1/2} \int_0^{\pi/2} dx \frac{(1 - 2\kappa^2 \sin^2 x)}{(1 - \kappa^2 \sin^2 x)^{1/2}}$$

$$= 8(2\varepsilon)^{-1/2} \sqrt{1 - \varepsilon + 2\varepsilon\kappa^2} [2E(\kappa) - K(\kappa)], \tag{6.20}$$

where $\kappa^2 = [1 - (1 - \varepsilon)\lambda] / 2\varepsilon\lambda$. Then $\mathbf{b} \cdot \nabla \eta \simeq (M - qN) / qR$ leads to $\oint_t d\tau \xi \simeq 2\pi qR / |M - qN|v$,

$$\tau_t = \oint_{\alpha} d\tau = \oint_{\alpha} \frac{d\eta}{v_{||} |\mathbf{b} \cdot \nabla \eta|} \simeq \frac{qR}{|M - qN|v} \oint_{\alpha} \frac{d\eta}{\xi} \simeq \frac{8qRK(\kappa)}{|M - qN|v\sqrt{2\varepsilon}}, \tag{6.21}$$

and

$$\oint_{\alpha} d\tau \omega_{\alpha} = \oint_{\alpha} \frac{d\eta \omega_{\alpha}(\eta)}{v_{||} |\mathbf{b} \cdot \nabla \eta|}$$

$$\simeq \frac{4q^2 v}{|M - qN| \Omega_0 a' \sqrt{2\varepsilon}} \left\{ 2E(\kappa) - K(\kappa) + \frac{4sM}{M - qN} [E(\kappa) - (1 - \kappa^2)K(\kappa)] \right\}. \tag{6.22}$$

In the absence of shear and a radial electric field a superbanana plateau resonance ($\oint_{\alpha} d\tau \omega_{\alpha} = 0$) occurs at $2E = K$ (Shaing, Sabbagh & Chu 2009; Catto 2019b), corresponding to $\kappa^2 \simeq 0.83$. In a tokamak, $s > 0$ moves the resonance closer to the trapped-passing boundary. Stellarators tend to have weaker shear but can have $s > 0$ or $s < 0$. Shear is unimportant in QPS stellarators ($M = 0$), while for QAS and QHS stellarators the limiting forms of the trapped tangential drift follow from

$$2E(\kappa) - K(\kappa) + \frac{4sM}{M - qN} [E(\kappa) - (1 - \kappa^2)K(\kappa)]$$

$$\rightarrow \begin{cases} \pi \left[1 - \left(1 + \frac{4sM}{qN - M} \right) \kappa^2 / 2 \right] / 2 & \kappa^2 \ll 1 \\ -\ell n(4 / \sqrt{1 - \kappa^2}) + 2 \left(\frac{2sM}{M - qN} + 1 \right) & \kappa^2 \rightarrow 1 \end{cases}. \tag{6.23}$$

Magnetic shear tends to be weak or modest in a stellarator, keeping q between any lower rational surfaces ($M \sim 1 \sim N$). A deeply trapped $\kappa^2 \ll 1$ resonance is less likely as it requires $4sM / (qN - M) \gg 1$ and often $M = 1$ (Landreman & Paul 2022). More importantly, the resonance as $\kappa^2 \rightarrow 1$ is exponentially sensitive to even modest shear and leads to shear reduction factor γ to be introduced in the next section that also appears in (1.2). It is these barely trapped alphas that give a superbanana plateau resonance when

$$1 - \kappa_{res}^2 = 16 e^{-4 - 8sM / (M - qN)} = 16 / \gamma \ll 1, \tag{6.24}$$

although (6.24) is less accurate for $s = 0$. To keep $1 - \kappa_{res}^2 \ll 1$ requires either $s > 0$ for $M > qN$, or $s < 0$ for $M < qN$. In addition, as $sM / (M - qN) > 0$ increases the resonance moves even closer to the passing boundary, merging the superbanana plateau and \sqrt{v} transport regimes. The behaviour when the two regions merge is the focus of much of the remaining sections, with only QAS (with $N = 0$ and $s > 0$) and QHS stellarators (with

$sM/(M - qN) > 0$) of interest as the resonance for QPS stellarators is insensitive to shear. These are two QS configurations recently shown by Landreman & Paul (2022) to offer the possibility of good alpha confinement.

The form of the kinetic equation for the trapped alphas suggests a solution of the form

$$f_1^{im} = \text{Im} \sum_{m,n} h_m^n(\psi_p, v, \lambda) e^{i(mN-nM)/(M-qN)\alpha}, \tag{6.25}$$

where $\partial h_m^n / \partial \eta = 0$, which leads to

$$ih_m^n \oint_{\alpha} d\tau \omega_{\alpha} = \frac{M - qN}{mN - nM} \oint_{\alpha} d\tau C\{h_m^n\} + \frac{M_{\alpha} c v^2}{Ze} \left(1 - \frac{\lambda}{2}\right) \frac{\partial f_0}{\partial \psi_h} \delta_m^n \oint_{\alpha} d\tau e^{i((m-qn)/(M-qN))\eta(\tau)}. \tag{6.26}$$

Next, the solution of (6.26) is given when the superbanana plateau and \sqrt{v} regimes merge.

7. Merged solution for the trapped alphas in nearly QS stellarators

In the presence of shear, superbanana plateau and \sqrt{v} transport collisions only matter in narrow layers near the trapped-passing boundary. Consequently, in (6.26)

$$\bar{\omega}_{\alpha} \equiv \frac{\oint_{\alpha} d\tau \omega_{\alpha}}{\oint_{\alpha} d\tau} \xrightarrow{\kappa \rightarrow 1} \frac{-qv^2}{2\Omega_0 R a'} \left\{ 1 - \frac{2[1 + 2sM/(M - qN)]}{\ell n(4/\sqrt{1 - \kappa^2})} \right\}, \tag{7.1}$$

where $\partial \bar{\omega}_{\alpha} / \partial \lambda = (\partial \kappa^2 / \partial \lambda) \partial \bar{\omega}_{\alpha} / \partial \kappa^2 \xrightarrow{\kappa \rightarrow 1} -qv^2[1 + 2sM/(M - qN)]/4\Omega_0 a' r(1 - \kappa^2)\ell n^2(4/\sqrt{1 - \kappa^2})$ is responsible for the sensitivity to magnetic shear that enters via the shear reduction factor

$$\gamma \equiv e^{4+8sM/(M-qN)} \gg 1. \tag{7.2}$$

As only pitch angle scattering need be retained in the collision operator in (6.26),

$$C\{h_m^n\} \rightarrow \frac{2v_{\lambda}^3 B_0}{\tau_s v^5 B} v_{\parallel} \frac{\partial}{\partial \lambda} \left(\lambda v_{\parallel} \frac{\partial h_m^n}{\partial \lambda} \right), \tag{7.3}$$

gives

$$\begin{aligned} \frac{\oint_{\alpha} d\tau C\{h_m^n\}}{\oint_{\alpha} d\tau} &\rightarrow \frac{2v_{\lambda}^3 B_0}{\tau_s v^3 \oint_{\alpha} d\tau} \frac{\partial}{\partial \lambda} \left[\lambda \left(\oint_{\alpha} d\tau \frac{\xi^2}{B} \right) \frac{\partial h_m^n}{\partial \lambda} \right] \\ &\simeq \frac{4\varepsilon v_{\lambda}^3}{\tau_s v^3 K(\kappa)} \frac{\partial}{\partial \lambda} \left\{ [E(\kappa) - (1 - \kappa^2)K(\kappa)] \frac{\partial h_m^n}{\partial \lambda} \right\} \xrightarrow{\kappa \rightarrow 1} \frac{v_{\lambda}^3}{4\varepsilon \tau_s v^3 \ell n(4/\sqrt{1 - \kappa^2})} \frac{\partial^2 h_m^n}{\partial \kappa^2}. \end{aligned} \tag{7.4}$$

In addition, the phase factor in the drive term of (6.26), which enters as

$$\Theta \equiv \oint_{\alpha} d\tau e^{i((m-qn)/(M-qN))\eta(\tau)} / \oint_{\alpha} d\tau \simeq \oint_{\alpha} d\eta \xi^{-1} e^{i((m-qn)/(M-qN))\eta} / \oint_{\alpha} d\eta \xi^{-1}, \tag{7.5}$$

can be taken to be approximately equal to one as long as $2\pi(m - qn)/(M - qN) < 1$. Keeping Θ allows the recovery of ripple modifications in a tokamak ($M = 1, N = 0, n \gg 1 \sim m$).

Using the preceding simplifies (6.26) to

$$\frac{i(nM - mN)qv^2}{2(M - qN)\Omega_0 Ra'} \left[1 - \frac{\ell n(\sqrt{\gamma})}{\ell n(4/\sqrt{1 - \kappa^2})} \right] h_m^n - \frac{v_\lambda^3}{4\varepsilon\tau_s v^3 \ell n(4/\sqrt{1 - \kappa^2})} \frac{\partial^2 h_m^n}{\partial \kappa^2} \tag{7.6}$$

$$\simeq \frac{(mN - nM)}{(M - qN)} \frac{B_0 \delta_m^n v^2}{2\Omega_0} \frac{\partial f_0}{\partial \psi_p} \Theta.$$

To consider the merged trapped regime it is convenient to first rewrite the kinetic equation by introducing the new variable

$$\zeta = (1 - \kappa^2)/16 \simeq (1 - \kappa)/8, \tag{7.7}$$

to obtain

$$\frac{1}{\ell n(\zeta)} \frac{\partial^2 h_m^n}{\partial \zeta^2} + i2W \left[1 + \frac{\ell n(\gamma)}{\ell n(\zeta)} \right] h_m^n = 2V, \tag{7.8}$$

where

$$W \equiv \frac{32q(nM - mN)\varepsilon\tau_s v^5}{(M - qN)v_\lambda^3 \Omega_0 Ra'}, \tag{7.9}$$

and

$$V \equiv -\frac{32(nM - mN)B_0 \varepsilon \delta_m^n \tau_s v^5}{(M - qN)\Omega_0 v_\lambda^3} \frac{\partial f_0}{\partial \psi_p} \Theta. \tag{7.10}$$

As $\zeta \rightarrow 0$, $h_m^n \rightarrow -iV/W$, provided $\ell n(1/\zeta) \gg \ell n(\gamma)$. However, a solution is required that vanishes at the trapped–passing boundary. An approximate matched asymptotic solution in \sqrt{v} regime is given by (Catto 2019a)

$$h_m^n \Big|_{\sqrt{v}} \simeq \frac{iV}{W} [e^{-(1\pm i)\zeta\sqrt{|W|\ell n(1/\zeta)}} - 1] \simeq \frac{iV}{W} [e^{-(1\pm i)\zeta\sqrt{|W|\ell n(|W|^{1/2})}} - 1], \tag{7.11}$$

with the upper (lower) sign for $W > 0$ ($W < 0$) and where the \sqrt{v} boundary layer width is

$$w_{\sqrt{v}} \sim |W|^{-1/2} \ll 1. \tag{7.12}$$

To carefully capture the preceding features in a merged superbanana plateau and \sqrt{v} evaluation and demonstrate that the two contributions are additive, an expansion in the vicinity of $\gamma\zeta = 1$ is employed by writing $\ell n(\gamma\zeta) \simeq (\gamma\zeta - 1) + \dots$ to be able to consider $\kappa^2 \simeq 1 - 16/\gamma$ close to one when $\gamma \gg 1$. Then letting $x = \gamma\zeta$ the kinetic equation can be approximated by

$$\frac{\partial^2 h_m^n}{\partial x^2} + i\frac{2W}{\gamma^2}(x - 1)h_m^n = -\frac{2V}{\gamma^2}\ell n\gamma. \tag{7.13}$$

This merged limit is the situation of interest and $x = \gamma\zeta \sim 1$ implies that $\zeta = (1 - \kappa^2)/16 \sim 1/\gamma$, giving $\partial\bar{\omega}_\alpha/\partial\lambda \xrightarrow{\kappa \rightarrow 1} -qv^2\gamma/64\Omega_0 v^3 r \ell n(\gamma) \sim \bar{\omega}_\alpha\gamma/\varepsilon \ell n(\gamma) \sim \bar{\omega}_\alpha\gamma/\varepsilon$, as used to obtain estimate (1.2). Using this estimate for the merged case for which the superbanana plateau boundary layer extends to the trapped–passing boundary gives $C\{h_m^n\} \sim v h_m^n/w_{sbp}^2 \sim \bar{\omega}_\alpha\gamma\varepsilon^{-1}w_{sbp}h_m^n$, and a width $w_{sbp} \sim (v/\gamma\omega_\alpha)^{1/3} \ll 1$, where $v \sim v_\lambda^3/\varepsilon v_0^3 \tau_s \sim v_{\alpha i}/\varepsilon$ is the effective pitch angle scattering frequency of the trapped alphas

with the background ions. This width becomes narrower than the $\sqrt{\nu}$ boundary layer width $w_{\sqrt{\nu}} \sim (\nu/\omega_\alpha)^{1/2} \ll 1$ when shear becomes strong, that is when $\gamma > (\omega_\alpha/\nu)^{1/2}$.

Letting

$$z = \pm \frac{(2|W|)^{1/3}}{\gamma^{2/3}}(x - 1) = \pm \frac{x - 1}{\gamma w_{sbp}}, \tag{7.14}$$

with the upper (lower) sign for $W > 0$ ($W < 0$) and w_{sbp} the superbanana plateau boundary layer width, more precisely defined as

$$w_{sbp} = 1/|2W\gamma|^{1/3} \ll 1, \tag{7.15}$$

leads to the Su & Oberman (1968) form

$$\frac{\partial^2 h_m^n}{\partial z^2} + izh_m^n \simeq -\frac{2V\ell n(\gamma)}{(2|W|\gamma)^{2/3}} = -2Vw_{sbp}^2 \ell n(\gamma). \tag{7.16}$$

As $W \sim \omega_\alpha/\nu$, this more precise definition of w_{sbp} is consistent with the rougher estimates. The superbanana plateau solution is

$$h_m^n|_{sbp} = \frac{2V\ell n(\gamma)}{(2|W|\gamma)^{2/3}} \int_0^\infty d\tau e^{iz\tau - \tau^3/3} \xrightarrow{|z| \gg 1} \frac{i2V\ell n(\gamma)}{(2|W|\gamma)^{2/3}z}. \tag{7.17}$$

The resonance $\gamma\zeta = x = 1$ remains near the trapped–passing boundary as long as $16/\gamma \ll 1$, a condition that is marginally satisfied even for $s=0$. Using (7.12) and (7.15) gives

$$w_{sbp}/w_{\sqrt{\nu}} \sim |W|^{1/6}/\gamma^{1/3}, \tag{7.18}$$

indicating w_{sbp} will become smaller than $w_{\sqrt{\nu}}$ as the shear becomes stronger.

To estimate the magnetic shear level at which the narrow boundary layer treatment fails consider a D-T plasma with $v_0 \simeq 1.3 \times 10^9$ cm sec⁻¹, $R = 10$ m, $B = 10$ T, $T = 10$ keV and $n_e = 10^{14}$ cm⁻³, to find $\tau_s \simeq 0.6$ sec, $\rho_0 \simeq 1$ cm, $v_0\tau_s/R \sim 10^6$, $R/\rho_0 \sim 10^3$, $R/a = 10$ and $v_0/v_\lambda \sim 3$. These parameters give $|W| \sim 10^6$ if $|(nM - mN)/(M - qN)| \sim 1$, indicating that, for $\gamma \sim 10^3$, the boundary layers are of comparable width. For the modest shear ($0 < sM/(M - qN) < 1$) case of interest here, $w_{\sqrt{\nu}} < w_{sbp} \ll 1$. Once $\gamma \gg |W|^{1/2}$, the superbanana plateau regime gives negligible transport.

The merged regime solution about to be presented is valid when $1 \ll \gamma \ll (\omega_\alpha/\nu)^{1/2}$. It is obtained by adding the appropriate homogeneous solution to the Su & Oberman (1968) form to make the total solution vanish at the trapped–passing boundary. To do so it is convenient to consider

$$\partial^2 \Upsilon / \partial z^2 + iz\Upsilon = -1, \tag{7.19}$$

then the merged solution is given by

$$h_m^n = \frac{2V\ell n(\gamma)}{(2|W|\gamma)^{2/3}} \Upsilon = -q^{-1} \delta_m^n \ell n(\gamma) B_0 R a' \frac{\partial f_0}{\partial \psi_p} \frac{\Theta W}{u|W|} \Upsilon, \tag{7.20}$$

with

$$u = \gamma / (2|W|\gamma)^{1/3} = \gamma w_{sbp}. \tag{7.21}$$

A particular solution to this inhomogeneous Airy equation is the Su & Oberman form,

$$\Upsilon_{SO} = \int_0^\infty d\tau e^{iz\tau - \tau^3/3} \xrightarrow{|z| \gg 1} i/z. \tag{7.22}$$




FIGURE 1. The solution for Υ given by (7.28) is plotted for various values of the shear reduction factor γ . The $\Upsilon_{\sqrt{v}}$ contribution from the truncated tail is small and nearer $\kappa = 1$, while the larger Υ_{SO} feature moves toward the trapped–passing boundary and narrows as γ increases.

For the homogenous solutions it is convenient to use Airy functions by letting $y = iz$ to obtain

$$\partial^2 \Upsilon / \partial y^2 - y \Upsilon = 1. \tag{7.23}$$

Then the convenient homogenous solutions are $Ai(y)$ and $Ai(ye^{-i2\pi/3})$ for large $z > 0$ (upper sign) and $Ai(y)$ and $Ai(ye^{i2\pi/3})$ for $z < 0$ with $-z$ large (lower sign) (Abramowitz & Stegun 1964). Asymptotically these give

$$Ai(y) = Ai(\pm z e^{\pm i\pi/2}) \propto e^{-2} e^{\pm i3\pi/4} (\pm z)^{3/2/3} \rightarrow \infty, \tag{7.24}$$

while for $z > 0$ with $|z| \gg 1$

$$Ai(ye^{-i2\pi/3}) = Ai(ze^{-i\pi/6}) \propto e^{-2} e^{-i\pi/4} z^{3/2/3} \rightarrow 0, \tag{7.25}$$

and for $z < 0$ with $|z| \gg 1$

$$Ai(ye^{i2\pi/3}) = Ai(-ze^{i\pi/6}) \propto e^{-2} e^{-i\pi/4} (-z)^{3/2/3} \rightarrow 0. \tag{7.26}$$

Consequently, the solution vanishing for the barely trapped alphas is

$$\Upsilon = \Upsilon_{SO}(z) + aAi(\pm z e^{\mp i\pi/6}), \tag{7.27}$$

where the constant a is determined by the need to satisfy $\Upsilon(z = \mp 1/u) = 0$ at the trapped–passing boundary. Hence, once the two regimes merge

$$\Upsilon = \Upsilon_{SO}(z) - \Upsilon_{SO}(z = \mp 1/u) Ai(\pm z e^{\mp i\pi/6}) / Ai(-e^{\mp i\pi/6} / u) \equiv \Upsilon_{SO}(z) + \Upsilon_{\sqrt{v}}(z), \tag{7.28}$$

is the solution from which the QL diffusivity and alpha heat flux can be evaluated, with $\Upsilon_{\sqrt{v}}(z)$ the homogenous contribution needed to satisfy the trapped–passing boundary condition. The solution Υ in figure 1 shows the $\Upsilon_{\sqrt{v}}$ contribution from the truncated tail nearer $\kappa = 1$ is small and the larger Υ_{SO} feature moves toward the trapped–passing boundary and narrows as γ increases.

8. The QL alpha diffusivity in nearly QS stellarators

At large aspect ratio the QL equation (5.10) for f_0 becomes

$$\begin{aligned} \frac{S\delta(v-v_0)}{4\pi v^2} + \frac{1}{\tau_s v^2} \frac{\partial}{\partial v} [(v^3 + v_c^3)f_0] + \frac{2v_\lambda^3}{\tau_s v^3 \oint_\alpha d\eta \xi^{-1}} \frac{\partial}{\partial \lambda} \left[\lambda \left(\oint_\alpha d\eta \xi \right) \frac{\partial f_0}{\partial \lambda} \right] \\ \simeq \frac{\frac{\partial}{\partial \psi_p} [qR \langle \oint_\alpha d\eta \xi^{-1} f_1^{im} \mathbf{v}_d \cdot \nabla \psi_p |_{im} \rangle_\alpha]}{qR \oint_\alpha d\eta \xi^{-1}}, \end{aligned} \tag{8.1}$$

where the sink term is contained in $v_c^3 \int d^3 v v^{-2} \partial f_0 / \partial v \xrightarrow{v \rightarrow 0} -4\pi v_c^3 f_0(\psi, v = 0)$.

To evaluate the diffusivity in the QL equation for f_0 and the alpha energy flux, the remaining angle integral is performed by first using $\lambda \simeq 1$ and

$$2 \langle e^{i(mN-nM)\alpha/(M-qN)} \sin \chi \rangle_\alpha = i e^{-i(m-qn)\eta/(M-qN)}, \tag{8.2}$$

to find

$$\begin{aligned} \left\langle \oint_\alpha d\eta \xi^{-1} f_1^{im} \mathbf{v}_d \cdot \nabla \psi_p |_{im} \right\rangle_\alpha &\simeq \frac{v^2}{2\Omega_0} \text{Im} \sum_{m,n} h_m^n \oint_\alpha d\eta \xi^{-1} \left\langle e^{i(mN-nM)\alpha/(M-qN)} \frac{\partial B}{\partial \alpha} \right\rangle_\alpha \\ &\simeq -\frac{B_0 v^2}{2\Omega_0} \text{Im} \sum_{m,n} h_m^n \sum_{m',n'} \frac{m'N - n'M}{M - qN} \delta_{m'}^{n'} \oint_\alpha d\eta \xi^{-1} \langle e^{i(mN-nM)\alpha/(M-qN)} \sin \chi' \rangle_\alpha \\ &\simeq -\frac{B_0 v^2}{4\Omega_0} \text{Im} \sum_{m,n} i h_m^n \frac{mN - nM}{M - qN} \delta_m^n \Theta^* \oint_\alpha d\eta \xi^{-1}, \end{aligned} \tag{8.3}$$

where only $m\vartheta - n\zeta$ in $\chi' = m'\vartheta - n'\zeta$ contributes.

Inserting h_m^n with $u = \gamma w_{sbp}$, (8.3) becomes

$$\left\langle \oint_\alpha d\eta \xi^{-1} f_1^{im} \mathbf{v}_d \cdot \nabla \psi_p |_{im} \right\rangle_\alpha = \frac{\ell n(\gamma) a' v^2}{4q\Omega_0 |M - qN|} \left[\sum_{m,n} |mN - nM| (\delta_m^n)^2 |\Theta|^2 \frac{Re\Upsilon}{u} \right] B_0^2 R \frac{\partial f_0}{\partial \psi_p} \oint_\alpha \frac{d\eta}{\xi}. \tag{8.4}$$

The delta function behaviour of Υ_{SO} occurs upon integration over pitch angle λ (Catto & Tolman 2021). Using

$$\frac{\gamma Re\Upsilon_{SO}}{32\pi u \varepsilon} = \frac{\gamma}{32\pi u \varepsilon} \int_0^\infty d\tau e^{-\tau^3/3} \cos(\tau\tau), \tag{8.5}$$

with $d\lambda \simeq -2\varepsilon d\kappa^2$ and $\gamma d\kappa^2 \simeq -16u dz$ gives

$$\frac{\gamma}{32\pi u \varepsilon} \int_{1/(1+\varepsilon)}^{1/(1-\varepsilon)} d\lambda Re\Upsilon_{SO} \simeq \frac{1}{\pi} \int_0^\infty \frac{d\tau}{\tau} e^{-\tau^3/3} \int_{-1/\gamma w_{sbp}}^{(\gamma-16)/16\gamma w_{sbp}} dz \frac{d}{dz} \sin(\tau\tau) \simeq 1, \tag{8.6}$$

due to the superbanana or resonant plateau contribution.

Using $\partial/\partial\psi_p \simeq (q/B_0a')\partial/\partial r$ and $\varepsilon \simeq r/R$, and defining the spatial diffusivity

$$D \equiv \frac{q\ell n(\gamma)v^2}{4\varepsilon\Omega_0|M - qN|} \left[\sum_{m,n} |nM - mN|(\delta_m^n)^2 |\Theta|^2 \frac{Re\Upsilon}{\gamma^w w_{sbp}} \right], \tag{8.7}$$

the QL equation (5.10) becomes

$$\begin{aligned} & \frac{S\delta(v - v_0)}{4\pi v^2} + \frac{1}{\tau_s v^2} \frac{\partial}{\partial v} [(v^3 + v_c^3)f_0] + \frac{2v_\lambda^3}{\tau_s v^3 \oint_\alpha d\eta\xi^{-1}} \frac{\partial}{\partial \lambda} \left[\lambda \left(\oint_\alpha d\eta\xi \right) \frac{\partial f_0}{\partial \lambda} \right] \\ & \simeq - \frac{1}{|\oint_\alpha d\eta\xi^{-1}|a'} \frac{\partial}{\partial r} \left(D \left| \oint_\alpha d\eta\xi^{-1} \right| r \frac{\partial f_0}{\partial r} \right), \end{aligned} \tag{8.8}$$

where $\oint_\alpha d\eta\xi/\oint_\alpha d\eta\xi^{-1} = 2\varepsilon[E(\kappa) - (1 - \kappa^2)K(\kappa)]/K(\kappa)$ and $\oint_\alpha d\eta\xi^{-1} \propto \varepsilon^{-1/2}$.

In addition to retaining alpha birth and slowing down (including the ash sink) on the left, the preceding QL equation retains the radial loss of alphas on the right. The strong dependence of the spatial diffusivity on pitch angle allows pitch angle scattering to enter on the left side of (8.8). Notice that $D \propto v^2$ implies that radial loss becomes negligible at low speed where f_0 becomes isotropic.

Negative definite entropy production is associated with both the radial transport and pitch angle scattering terms as multiplying by $\ell n f_0 |\oint_\alpha d\eta\xi^{-1}|$ and integrating over velocity space and cross section $da \simeq dr a'$ out to some reference flux surface label r_0 leads to the expression

$$\begin{aligned} & \int d^3v \int_0^{r_0} dr \ell n f_0 \left\{ \frac{\partial}{\partial r} \left[D \left| \oint_\alpha d\eta\xi^{-1} \right| r \frac{\partial f_0}{\partial r} \right] + \frac{2a'v_\lambda^3}{\tau_s v^3 \oint_\alpha d\eta\xi^{-1}} \frac{\partial}{\partial \lambda} \left[\lambda \left(\oint_\alpha d\eta\xi \right) \frac{\partial f_0}{\partial \lambda} \right] \right\} \\ & = - \int d^3v \int_0^{r_0} \frac{dr}{f_0} \left[r D \left| \oint_\alpha d\eta\xi^{-1} \right| \left(\frac{\partial f_0}{\partial r} \right)^2 + \frac{2a'v_\lambda^3 \lambda}{\tau_s v^3} \left| \oint_\alpha d\eta\xi \right| \left(\frac{\partial f_0}{\partial \lambda} \right)^2 \right]. \end{aligned} \tag{8.9}$$

The entropy production is associated with the alphas attempting to remove radial and pitch angle gradients to become the slowing down distribution function of (4.12) as implied by the remaining source and drag terms in (8.8).

9. The QL alpha heat flux in nearly QS stellarators

The heat flux evaluation involves performing velocity space integrals in addition to the angle integral evaluated in the last section. Rewriting

$$\begin{aligned} & \left\langle \int d^3v j_1^{im} v^2 \mathbf{v}_d \cdot \nabla \psi_p |_{im} \right\rangle \simeq \int_0^{v_0} dv v^4 \int_{1/(1+\varepsilon)}^{1/(1-\varepsilon)} d\lambda \left\langle \oint_{\xi} \frac{d\eta}{\xi} f_1^{im} \mathbf{v}_d \cdot \nabla \psi_p |_{im} \right\rangle_\alpha \\ & = \frac{B_0 R \ell n(\gamma)}{4\Omega_0(M - qN)} \sum_{m,n} \left[(mN - nM)(\delta_m^n)^2 \int_0^{v_0} dv v^6 \frac{\partial f_0}{\partial r} \int_{1/(1+\varepsilon)}^{1/(1-\varepsilon)} d\lambda \left(\frac{Re\Upsilon}{w} \right) \left(\oint_{\xi} d\eta\xi^{-1} \right) |\Theta|^2 \right], \end{aligned} \tag{9.1}$$

where $\oint d\eta = 2\pi$. To simplify further recall that the superbanana plateau resonance is at $\gamma\zeta = 1$, and the \sqrt{v} boundary layer width estimate from (7.11) is $\zeta = (1 - \kappa^2)/16 \sim W^{-1/2}$, while the superbanana plateau boundary layer width estimate is $\zeta =$

$(1 - \kappa^2)/16 \sim \gamma^{-1}$. As these small ζ are very close to the trapped–passing boundary they imply

$$\oint_t d\eta \xi^{-1} \simeq 8(2\varepsilon)^{1/2} \ell n \left(\frac{4}{\sqrt{1 - \kappa^2}} \right) \simeq 8(2\varepsilon)^{1/2} \begin{cases} \ell n(\sqrt{\gamma}) & sbp \\ \ell n(W_0^{1/4}) & \sqrt{v} \end{cases} \simeq \begin{cases} \oint_t d\eta \xi^{-1}|_{sbp} \\ \oint_t d\eta \xi^{-1}|_{\sqrt{v}} \end{cases}, \tag{9.2}$$

where

$$W_0 \equiv \frac{32q(nM - mN)\tau_s v_0^5}{(M - qN)v_\lambda^3 \Omega_0 R^2} > 0. \tag{9.3}$$

Moreover, the barely trapped ($\kappa \rightarrow 1$) evaluation in Appendix A of Catto (2019b) suggests

$$|\Theta|^2 \simeq \cos^2 \left[\left(\frac{qn - m}{M - qN} \right) \pi \right] \frac{\ell n^2 \left[4 / \left(1 + \left| \frac{qn - m}{M - qN} \right| \right) \sqrt{1 - \kappa^2} \right]}{\ell n^2(4/\sqrt{1 - \kappa^2})} \\ \simeq \cos^2 \left[\left(\frac{qn - m}{M - qN} \right) \pi \right] \begin{cases} \frac{\ell n^2 \left[\sqrt{\gamma} / \left(1 + \left| \frac{qn - m}{M - qN} \right| \right) \right]}{\ell n^2(\sqrt{\gamma})} sbp \\ \frac{\ell n^2 \left[W_0^{1/4} / \left(1 + \left| \frac{qn - m}{M - qN} \right| \right) \right]}{\ell n^2(W_0^{1/4})} \sqrt{v} \end{cases} \equiv \begin{cases} |\Theta|_{sbp}^2 \\ |\Theta|_{\sqrt{v}}^2 \end{cases}, \tag{9.4}$$

then $|\Theta|^2 \rightarrow 1$ as required for $|(qn - m)/(M - qN)| \ll 1$, but $|\Theta|^2 < 1$ otherwise. When $|(qn - m)/(M - qN)| \gg 1$ the rapid oscillation of the Θ phase factor in (7.5) cannot be ignored as it acts to reduce the effective step size of a trapped alpha (for a tokamak $N = 0$, $M = 1$ and $n \gg m \sim 1$). Using the preceding approximations indicates the pitch angle dependence of $|\Theta|^2$ and $\oint_t d\eta \xi^{-1}$ only enters logarithmically and suggests writing

$$\left\langle \int d^3 v f_1^{im} v^2 \mathbf{v}_d \cdot \nabla \psi_p |_{im} \right\rangle \simeq \frac{B_0 R \ell n(\gamma)}{4\Omega_0(M - qN)} \sum_{m,n} \{(mN - nM)(\delta_m^n)^2 \\ \left[|\Theta|_{sbp}^2 \oint_t d\eta \xi^{-1}|_{sbp} \int_0^{v_0} dv v^6 \frac{\partial f_0}{\partial r} \int_{1/(1+\varepsilon)}^{1/(1-\varepsilon)} d\lambda \left(\frac{\text{Re} \Upsilon_{SO}}{u} \right) \right. \\ \left. + |\Theta|_{\sqrt{v}}^2 \oint_t d\eta \xi^{-1}|_{\sqrt{v}} \int_0^{v_0} dv v^6 \frac{\partial f_0}{\partial r} \int_{1/(1+\varepsilon)}^{1/(1-\varepsilon)} d\lambda \left(\frac{\text{Re} \Upsilon_{\sqrt{v}}}{u} \right) \right] \Bigg\}, \tag{9.5}$$

where defining $w_{\sqrt{v}} = 1/(2|W|)^{1/2}$ leads to $u = \gamma w_{sbp} = \gamma^{2/3} w_{\sqrt{v}}^{2/3}$. The Υ_{SO} term has already been evaluated in (8.6). For the $\Upsilon_{\sqrt{v}}$ term only the large z limit is required giving

$$\frac{\gamma}{32\pi u \varepsilon} \int_{1/(1+\varepsilon)}^{1/(1-\varepsilon)} d\lambda \text{Re} \Upsilon_{\sqrt{v}} \simeq -\frac{1}{\pi} \text{Re} \left[\frac{\Upsilon_{SO}(z = \mp 1/u)}{Ai(-e^{\mp i\pi/6}/u)} \int_{-1/u}^{\gamma^{1/6} u} dz Ai(\pm z e^{\mp i\pi/6}) \right]. \tag{9.6}$$

Using the $u \ll 1$ asymptotic forms (Abramowitz & Stegun 1964)

$$Ai(-e^{\mp i\pi/6}/u) \simeq \frac{e^{\pm i\pi/24} u^{1/4} \sin[2^{1/2}(1 - i)/3u^{3/2} + \pi/4]}{\pi^{1/2}} + \dots, \tag{9.7}$$

and

$$\begin{aligned} \int_{-1/u}^{\gamma/16u} dz Ai(\pm z e^{\mp i\pi/6}) &= \int_0^{\gamma/16u} dz Ai(\pm z e^{\mp i\pi/6}) + \int_0^{1/u} dz Ai(\mp z e^{\mp i\pi/6}) \\ &= e^{\pm i\pi/6} \left[\int_0^{\gamma e^{\mp i\pi/6}/16u} dt Ai(t) \pm \int_0^{e^{\mp i\pi/6}/u} dt Ai(-t) \right] \\ &\simeq e^{\pm i\pi/6} \left[1 \mp \frac{e^{\pm i\pi/8} u^{3/4} \cos[2^{1/2}(1-i)/3u^{3/2} + \pi/4]}{\pi^{1/2}} + \dots \right], \end{aligned} \tag{9.8}$$

leads to

$$\frac{\int_{-1/u}^{\gamma/16u} dz Ai(\pm z e^{\mp i\pi/6})}{Ai(-e^{\mp i\pi/6}/u)} \simeq -e^{\pm i\pi/4} u^{1/2} \cot[2^{1/2}(1-i)/3u^{3/2} + \pi/4] \simeq -ie^{\pm i\pi/4} u^{1/2}. \tag{9.9}$$

Therefore, using $\Upsilon_{SO}(\mp 1/u) \simeq \mp iu$,

$$\frac{\gamma}{32\pi u \varepsilon} \int_{1/(1+\varepsilon)}^{1/(1-\varepsilon)} d\lambda \text{Re} \Upsilon_{\sqrt{v}} \simeq \frac{u^{3/2}}{2^{1/2}\pi}, \tag{9.10}$$

and

$$\frac{\gamma}{32\pi u \varepsilon} \int_{1/(1+\varepsilon)}^{1/(1-\varepsilon)} d\lambda \text{Re} \Upsilon \simeq 1 + \frac{u^{3/2}}{2^{1/2}\pi}. \tag{9.11}$$

Amazingly, $u^{3/2} = \gamma/\sqrt{2|W|} \propto \tau_s^{-1/2}$, and is just the \sqrt{v} regime contribution. Consequently, the merged procedure formulated here has the virtue of recovering both the \sqrt{v} and superbanana plateau regimes in a unified and additive manner when $u \ll 1$. Only the $u \ll 1$ limit is considered here. Once $\gamma \gg \sqrt{2|W|}$ is satisfied, presumably \sqrt{v} transport will dominate since so few trapped alphas experience the tangential drift resonance.

Inserting the preceding results into the heat flux leads to

$$\begin{aligned} \left\langle \int d^3 v f_1^{im} v^2 \mathbf{v}_d \cdot \nabla \psi_p |_{im} \right\rangle &\simeq \frac{8\pi B_0 \ell n(\gamma) r}{\Omega_0 |M - qN| \gamma} \sum_{m,n} \{|mN - nM| (\delta_m^n)^2 \\ &\left[|\Theta|_{sbp}^2 \oint_t d\eta \xi^{-1} |_{sbp} \int_0^{v_0} dv v^6 \frac{\partial f_0}{\partial r} + |\Theta|_{\sqrt{v}}^2 \oint_t d\eta \xi^{-1} |_{\sqrt{v}} \int_0^{v_0} dv v^6 \frac{\partial f_0}{\partial r} \frac{u^{3/2}}{2^{1/2}\pi} \right] \}. \end{aligned} \tag{9.12}$$

Then using the approximation that near the birth speed $v^3 \partial f_0 / \partial r \simeq (4\pi)^{-1} \partial(S\tau_s) / \partial r$,

$$\begin{aligned} \left\langle \frac{M_\alpha q}{2B_0 r} \int d^3 v f_1^{im} v^2 \mathbf{v}_d \cdot \nabla \psi_p |_{im} \right\rangle &\simeq \frac{qv_0^2 \ell n(\gamma)}{2\Omega_0 |M - qN| \gamma} \left(\frac{M_\alpha v_0^2}{2} \right) \sum_{m,n} \{|mN - nM| (\delta_m^n)^2 \\ &\left[|\Theta|_{sbp}^2 \oint_t d\eta \xi^{-1} |_{sbp} + \frac{4\gamma}{3\pi W_0^{1/2}} |\Theta|_{\sqrt{v}}^2 \oint_t d\eta \xi^{-1} |_{\sqrt{v}} \right] \} \frac{\partial(S\tau_s)}{\partial r}, \end{aligned} \tag{9.13}$$

where for the slowing down distribution the alpha density is $n_\alpha \simeq S\tau_s \ell n(v_0/v_c)$. Importantly, in the presence of magnetic shear with $sM/(qN - M) > 0$, superbanana

plateau transport no longer dominates over $\sqrt{\nu}$ regime transport when $\gamma \rightarrow W_0^{1/2} \sim (\omega_\alpha/\nu)^{1/2}$ as suggested by the estimate of (7.18). The merged solution found here fails in this limit, but by then superbanana plateau transport is expected to be as small or smaller than $\sqrt{\nu}$ transport.

The preceding leads to the heat transport as being the sum of the superbanana plateau and $\sqrt{\nu}$ heat diffusivities (D_{sbp} and $D_{\sqrt{\nu}}$, respectively)

$$D_{sbp} = \frac{32[1 + 2sM/(M - qN)]^2 qv_0^2}{\sqrt{2\varepsilon}\Omega_0|M - qN|} e^{-4[1+2sM/(M-qN)]} \sum_{m,n} |mN - nM|(\delta_m^n)^2 |\Theta|_{sbp}^2, \tag{9.14}$$

and

$$D_{\sqrt{\nu}} = \frac{16[1 + 2sM/(M - qN)]qv_0^2}{3\pi\sqrt{2\varepsilon}\Omega_0|M - qN|} \ell n(W_0) \sum_{m,n} |mN - nM|(\delta_m^n)^2 \frac{|\Theta|_{\sqrt{\nu}}^2}{W_0^{1/2}}, \tag{9.15}$$

where in some stellarator cases $|\Theta|_{sbp}^2 \simeq 1$ and $|\Theta|_{\sqrt{\nu}}^2 \simeq 1$ may be adequate approximations in

$$\begin{aligned} \frac{1}{V'} \frac{\partial}{\partial \psi_p} \left[V' \left\langle \frac{M_\alpha}{2} \int d^3 v f_1 v^2 \mathbf{v}_d \cdot \nabla \psi_p \right\rangle \right] &\simeq \frac{1}{a'} \frac{\partial}{\partial r} \left[\left\langle \frac{M_\alpha q}{2B_0} \int d^3 v f_1 v^2 \mathbf{v}_d \cdot \nabla \psi_p \right\rangle \right] \\ &= \frac{1}{a'} \frac{\partial}{\partial r} \left[r \frac{M_\alpha v_0^2}{2} (D_{sbp} + D_{\sqrt{\nu}}) \frac{\partial (S\tau_s)}{\partial r} \right]. \end{aligned} \tag{9.16}$$

The preceding results retain magnetic shear for $sM/(qN - M) > 0$ with $\gamma \equiv e^{4+8sM/(M-qN)} \gg 1$ and confirm the estimates in the Introduction of

$$D_{sbp} \sim (\delta_m^n)^2 qv_0^2 / \gamma \sqrt{\varepsilon} \Omega_0, \tag{9.17}$$

and

$$D_{\sqrt{\nu}}/D_{sbp} \sim \gamma/W_0^{1/2} \sim (\Omega_0 R/v_0)^{1/2} (v_\lambda^3 R/v_0^4 \tau_s)^{1/2} e^{4+8sM/(M-qN)}. \tag{9.18}$$

The general expressions (9.14) and (9.15) indicate that shear can significantly reduce superbanana plateau transport but has little effect on $\sqrt{\nu}$ regime transport. As $\gamma \equiv e^{4+8sM/(M-qN)}$ increases the superbanana plateau boundary layer narrows and the diffusivity is strongly reduced because of the rapid variation the transit average tangential drift as seen in (7.1) and noted there. As a result, fewer trapped alphas are able participate in the drift resonance, thereby reducing the transport. The shear at the edge of a tokamak means that superbanana transport is unlikely to ever be of concern, and the small ripple makes $\sqrt{\nu}$ transport weak as well (Catto 2019a). The $\sqrt{\varepsilon}$ and q factors found here are consistent with the shearless limit (Catto 2019b). However, the result here is larger, presumably because the barely trapped asymptotic limits of the elliptic integrals are used here to treat finite shear. This last approximation works well for finite shear, but is less accurate in the absence of shear.

The superbanana plateau diffusivity can be rewritten in an alternate approximate form. Using the resonance condition $\kappa_{res}^2 = 1 - 16/\gamma$ in $\kappa^2 = [1 - (1 - \varepsilon)\lambda]/2\varepsilon\lambda$ gives the

resonant pitch angle to be

$$\lambda_{res} = 1 - (1 - 32\gamma^{-1})\varepsilon. \quad (9.19)$$

Then taking advantage of the implicit delta function in $Re\mathcal{Y}$ by letting

$$Re\mathcal{Y} \rightarrow 32\pi u\varepsilon\gamma^{-1}\delta(\lambda - \lambda_{res}), \quad (9.20)$$

this result is used to obtain an approximate superbanana plateau D for the QL operator (8.8) of the form

$$D = \frac{8\pi q\ell n(\gamma)v^2}{\Omega|M - qN|\gamma} \delta(\lambda - \lambda_{res}) \sum_{m,n} |nM = mN|(\delta_m^n)^2, \quad (9.21)$$

where $|\Theta|^2 \simeq 1$ is also assumed. In this form collisions no longer appear even though the derivation is based on a collisional boundary layer treatment. The diffusivity can also be rewritten using $\delta(\lambda - \lambda_{res}) = \delta(\kappa - \kappa_{res})/4\varepsilon$ since $d\lambda \simeq -4\varepsilon d\kappa$. The absence of an explicit collisional dependence is characteristic of resonant plateau regime behaviour to lowest order.

The strong pitch angle dependence of the diffusivity D_{sbp} means that ignoring the λ dependence of f_0 when evaluating the evaluation of the radial heat transport is not justified for large departures from QS.

10. Discussion

Superbanana plateau and \sqrt{v} transport of alpha particles in a nearly QS, large aspect ratio stellarator, having a sheared magnetic field with $sM/(M - qN) > 0$ is evaluated by solving a kinetic equation retaining both processes in the same narrow collisional boundary layer. The radial superbanana plateau alpha energy flux is reduced by magnetic shear s satisfying $sM/(M - qN) > 0$, with M and N the integers associated with the toroidal and poloidal variation of the QS field, respectively, for the QS angle variable $\eta = M\vartheta - N\zeta$ (for which a B contour closes on itself after a $2\pi N$ change in ϑ and a $2\pi M$ change in ζ). As $M \geq 0$ and $N \geq 0$ the shear in a QHS stellarator needs to be positive for $M > qN$ and negative for $M < qN$. In a QAS stellarator ($N = 0$) the shear must be positive. In a QPS stellarator ($M = 0$) shear has no significant effect. The corresponding superbanana plateau and \sqrt{v} energy diffusivities are given by (9.14) and (9.15), respectively, for a slowing down tail alpha distribution function, with the alpha energy balance equation given by (9.16). The key new result is the significant reduction of superbanana plateau transport due to magnetic shear for the cases just enumerated. Shear acts to move the tangential drift resonance of the alphas very close to the trapped-passing boundary where the strong pitch angle variation of the drift reduces the number of alphas that are able to resonate, thereby reducing the transport. Even at modest shear levels it is possible to substantially reduce superbanana plateau transport. Although the solution presented here fails in this limit, it suggests that superbanana plateau transport will become as small or smaller than \sqrt{v} transport at rather modest shear. Similar behaviour is expected for the background ions when the superbanana plateau and \sqrt{v} regimes merge.

In addition, a QL description of radial alpha particle transport is derived to find (8.8) with the QL diffusivity given by (8.7). The QL formulation demonstrates how transport acts to deplete the resonant pitch angle and thereby introduce a pitch angle modification to the usual isotropic slowing down tail alpha distribution.

As the departure from QS becomes smaller and smaller, the perturbed radial drift term in the kinetic equation (5.12) can no longer be ignored. Once it matters collisional

detrapping and retrapping in the wells due to imperfect QS is expected to lead to transport becoming linear in the collision frequency due to collisional changes in the second adiabatic invariant. In this superbanana regime the collision operator acts to place a constraint on how the constants of the drift motion and the first and second adiabatic invariant dependences are allowed to change the non-Maxwellian features (Hazeltine & Catto 1981; Shaing 2015; d’Herbemont *et al.* 2022). The stellarator case is discussed in detail in this last reference for background ions.

The alpha particle case differs from the background ion case because the magnetic tangential drift matters rather than the tangential $\mathbf{E} \times \mathbf{B}$ drift. Accounting for this difference allows an estimate of when a superbanana plateau regime treatment is expected to remain a valid description. As the trapped alphas drift radially at speed $V_r \sim \delta\omega_\alpha R$ into a different magnetic field the pitch angle of the turning point of the trapped, $\lambda = B_0/B$, does not change, but the bounce points change by $\Delta r/R \sim \Delta B/B_0$ thereby shifting the trapped–passing boundary $\lambda = 1/(1 + \varepsilon)$ by $\Delta\lambda \sim \Delta r/R$ relative to the drift reversal layer location. Consequently, the nonlinear or finite orbit radial drift term neglected in (5.12) enters once this boundary change becomes comparable to the boundary layer width, $\Delta\lambda \sim w_{sbp} \sim \Delta r/R$. Therefore, the radial and tangential drift terms will compete once

$$\frac{V_r \partial f_1^{im} / \partial r}{\omega_\alpha \partial f_1^{im} / \partial \alpha} \sim \frac{\delta \omega_\alpha R / w_{sbp} R}{\gamma \omega_\alpha w_{sbp}} \sim \frac{\delta}{\gamma w_{sbp}^2} \sim 1, \tag{10.1}$$

where $\partial f_1^{im} / \partial r \sim f_1^{im} / w_{sbp} R$ and recall that expanding about drift reversal layer near the trapped–passing boundary gives the estimate $\omega_\alpha \rightarrow \gamma \omega_\alpha w_{sbp}$. As a result, a superbanana plateau treatment for the alphas using unperturbed trajectories assumes

$$\left(\frac{v}{\omega_\alpha}\right)^{2/3} \sim \left(\frac{vR}{v_0}\right)^{2/3} \left(\frac{a\Omega_0}{qv_0}\right)^{2/3} > \frac{\delta}{\gamma^{1/3}}, \tag{10.2}$$

where shear enters through γ . For the numbers given below (7.18) this requires highly optimized fields with $\delta < 10^{-3}$ for $s = 0$ (for which $\gamma^{1/3} \sim 4$). When $sM/(M - qN) > 0$ this requirement is relaxed, while for $sM/(M - qN) < 0$ it is more stringent (and the description here begins to fail as $\gamma \rightarrow 1$). In addition, to the preceding, $\partial f_1^{im} / \partial r < \partial f_0 / \partial r$, requires $f_1^{im} / f_0 \sim w_{sbp} / \varepsilon \ll 1$. Combining this with (10.2) leads to the restriction

$$(\delta/\varepsilon)^{1/2} \ll (\varepsilon\gamma)^{1/2}, \tag{10.3}$$

further confirming that $\delta/\varepsilon \ll 1$ is required.

Based on the QL (8.8), $\tau_s D_{sbp} / a^2$ must be small to avoid superbanana plateau transport from becoming a concern. For $|\Theta|_{sbp}^2 |mN - nM| / |M - qN| \sim 1$, (9.14) gives the estimate of (1.2), namely $D_{sbp} \sim \delta^2 q v_0^2 / \gamma \varepsilon^{1/2} \Omega_o \sim \delta^2 \omega_\alpha R a / \gamma \varepsilon^{1/2}$. Using it with the preceding estimate $\delta < \gamma w_{sbp}^2$ and $r \sim a$, leads to an estimate of the condition for small alpha loss while the transport still remains in the superbanana plateau regime

$$\left(\frac{\delta}{\varepsilon}\right)^{1/2} \frac{v \tau_s}{\varepsilon \gamma^{1/2}} < \frac{\tau_s D_{sbp}}{a^2} \ll 1. \tag{10.4}$$

As $v \tau_s / \gamma^{1/2} \ll 1$, as long as $(\delta/\varepsilon)^{1/2} \ll 1$, then for any reasonable aspect ratio it seems likely that superbanana plateau alpha transport will remain small. In particular, for $\delta \sim 10^{-3}$ and $\varepsilon \sim 0.1$, large alpha losses are unlikely to be an issue as long as shear is favourable or unimportant. Indeed, as superbanana plateau transport is always expected to be larger than superbanana transport, well optimized stellarators are expected to provide adequate alpha confinement as long as magnetic shear is weak or favourable.

Acknowledgements

The United States Government retains a non-exclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.

Editor Per Helander thanks the referees for their advice in evaluating this article.

Declaration of interests

The authors report no conflict of interest.

Funding

This work was supported by the U.S. Department of Energy under contract numbers DE-FG02-91ER-54109 and DE-AC02-09CH11466, research at the Institute for Advanced Study was supported by the W. M. Keck Foundation Fund, and research at the Flatiron Institute is supported by the Simons Foundation.

REFERENCES

- ABRAMOWITZ, M. & STEGUN, I.A. 1964 *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, 10th printing, pp. 446–449. National Bureau of Standards.
- BADER, A., DREVLAK, M., ANDERSON, D.T., FABER, B.J., HEGNA, C.C., LIKIN, K.M., SCHMITT, J.C. & TALMADGE, J.N. 2019 Stellarator equilibria with reactor relevant energetic particle losses. *J. Plasma Phys.* **85**, 905850508; 18pp.
- BEIDLER, C.D., ALLMAIER, K., ISAEV, M.Y., KASLOV, S.V., KERNBICHLER, W., LEITOLD, G.O., MAAßBERG, H., MIKKELSEN, D.R., MURAKAMI, S., SCHMIDT, M., SPONG, D.A., TRIBALDOS, V. & WAKASA, A. 2011 Benchmarking of the mono-energetic transport coefficients - results from the International Collaboration on Neoclassical Transport in Stellarators (ICNTS). *Nucl. Fusion* **51**, 076001; 28pp.
- BEIDLER, C.D. & D'HAESELEER, W.D. 1995 A general solution of the ripple-averaged kinetic equation (GSRAGE). *Plasma Phys. Control. Fusion* **37**, 463–490.
- BOOZER, A.H. 1981 Plasma equilibrium with rational magnetic surfaces. *Phys. Fluids* **24**, 1999–2003.
- BOOZER, A.H. 1983 Transport and isomorphic equilibria. *Phys. Fluids* **26**, 496–499.
- BOOZER, A.H. 1995 Quasi-helical symmetry in stellarators. *Plasma Phys. Control. Fusion* **37**, A103–A117.
- CALVO, I., PARRA, F.I., ALONSO, J.A. & VELASCO, J.L. 2014a Optimizing stellarators for large flows. *Plasma Phys. Control. Fusion* **56**, 094003; 11pp.
- CALVO, I., PARRA, F.I., VELASCO, J.L. & ALONSO, J.A. 2013 Stellarators close to quasisymmetry. *Plasma Phys. Control. Fusion* **55**, 125014; 28pp.
- CALVO, I., PARRA, F.I., VELASCO, J.L. & ALONSO, J.A. 2014b Flow damping in stellarators close to quasisymmetry. *Plasma Phys. Control. Fusion* **57**, 014014; 7pp.
- CALVO, I., PARRA, F.I., VELASCO, J.L. & ALONSO, J.A. 2017 The effect of tangential drifts on neoclassical transport in stellarators close to omnigenity. *Plasma Phys. Control. Fusion* **59**, 055014; 19pp.
- CARY, J.R. & SHASHARINA, S.G. 1997a Helical plasma confinement devices with good confinement properties. *Phys. Rev. Lett.* **78**, 674–677.
- CARY, J.R. & SHASHARINA, S.G. 1997b Omnigenity and quasihelicity in helical plasma confinement systems. *Phys. Plasmas* **4**, 3323–3333.
- CATTO, P.J. 2018 Ripple modifications to alpha transport in tokamaks. *J. Plasma Phys.* **84**, 905840508; 39pp.
- CATTO, P.J. 2019a Collisional alpha transport in a weakly rippled magnetic field. *J. Plasma Phys.* **85**, 905850203; 16pp.
- CATTO, P.J. 2019b Collisional alpha transport in a weakly non-quasisymmetric stellarator magnetic field. *J. Plasma Phys.* **85**, 905850213; 20pp & 945850501; 1p.

- CATTO, P.J. & TOLMAN, E.A. 2021 Collisional broadening of nonlinear resonant wave-particle interactions. *J. Plasma Phys.* **87**, 905870606; 15pp.
- D'HERBEMONT, V., PARRA, F.I., CALVO, I. & VELASCO, J.L. 2022 Finite orbit width effects in large aspect ratio stellarators. *J. Plasma Phys.* **88**, 905880507; 80pp.
- GALEEV, A.A., SAGDEEV, R.Z., FURTH, H.P. & ROSENBLUTH, M.N. 1969 Plasma diffusion in a toroidal stellarator. *Phys. Rev. Lett.* **22**, 511–514.
- GARREN, D.A. & BOOZER, A.H. 1991a Magnetic field strength of toroidal plasma equilibria. *Phys. Fluids B* **3**, 2805–2821.
- GARREN, D.A. & BOOZER, A.H. 1991b Existence of quasihelically symmetric stellarators. *Phys. Fluids B* **3**, 2822–2834.
- GATES, D.A., BOOZER, A.H., BROWN, T., BRESLAU, J., CURRELI, D., LANDREMAN, M., LAZERSON, S.A., LORE, J., MYNICK, H., NEILSON, G.H., POMPHREY, N., XANTHOPOULOS, P. & ZOLFAGHARI, A. 2017 Recent advances in stellarator optimization. *Nucl. Fusion* **57**, 126064; 9pp.
- HAZELTINE, R.D. & CATTO, P.J. 1981 Bumpy torus transport in the low collision frequency limit. *Phys. Fluids* **24**, 290–306.
- HENNEBERG, S.A., DREVLAK, M., NÜHRENBURG, C., BEIDLER, C.D., TURKIN, Y., LOIZU, J. & HELANDER, P. 2019 Properties of a new quasi-axisymmetric configuration. *Nucl. Fusion* **59**, 026014; 11pp.
- HO, D.D.-M. & KULSRUD, R.M. 1987 Neoclassical transport in stellarators. *Phys. Fluids* **30**, 442–461.
- HSU, C.T., CATTO, P.J. & SIGMAR, D.J. 1990 Neoclassical transport of isotropic fast ions. *Phys. Fluids B* **2**, 280; 11pp.
- LANDREMAN, M. 2019 Optimized quasi-symmetric stellarators are consistent with the Garren-Boozer construction. *Plasma Phys. Control. Fusion* **61**, 075001; 8pp.
- LANDREMAN, M. & CATTO, P.J. 2011 Effects of the radial electric field in a quasisymmetric stellarator. *Plasma Phys. Control. Fusion* **53**, 015004; 28pp.
- LANDREMAN, M. & PAUL, E. 2022 Magnetic fields with precise quasisymmetry for plasma confinement. *Phys. Rev. Lett.* **128**, 035001-6.
- LANDREMAN, M. & SENGUPTA, W. 2018 Direct construction of optimized stellarator shapes. Part 1. Theory in cylindrical coordinates. *J. Plasma Phys.* **84**, 905840616; 22pp.
- LANDREMAN, M. & SENGUPTA, W. 2019 Constructing stellarators with quasisymmetry to high order. *J. Plasma Phys.* **85**, 815850601; 43pp.
- LANDREMAN, M., SENGUPTA, W. & PLUNK, G. 2019 Direct construction of optimized stellarator shapes. Part 2. Numerical quasisymmetric solutions. *J. Plasma Phys.* **85**, 905850103; 22pp.
- NÜHRENBURG, J. & ZILLE, R. 1988 Quasi-helically symmetric toroidal stellarators. *Phys. Lett. A* **129**, 113–117.
- PARRA, F.I., CALVO, I., HELANDER, P. & LANDREMAN, M. 2015 Less constrained omnigenous stellarators. *Nucl. Fusion* **55**, 033005; 5pp.
- PARRA, F.I. & CATTO, P.J. 2008 Limitations of gyrokinetics on transport time scales. *Plasma Phys. Control. Fusion* **50**, 065014; 23pp.
- PLUNK, G., LANDREMAN, M., SENGUPTA, W. & HELANDER, P. 2019 Direct construction of optimized stellarator shapes. Part 3. Omnigenity near the magnetic axis. *J. Plasma Phys.* **85**, 905850602; 24pp.
- PYTTE, A. & BOOZER, A.H. 1981 Neoclassical transport in helically symmetric plasmas. *Phys. Fluids* **24**, 88–92.
- SHAING, K.C. 2015 Superbanana and superbanana plateau transport in finite aspect ratio tokamaks with broken symmetry. *J. Plasma Phys.* **81**, 905810203; 12pp.
- SHAING, K.C., SABBAGH, S.A. & CHU, M.S. 2009 Neoclassical toroidal plasma viscosity in the superbanana plateau for tokamaks. *Plasma Phys. Control. Fusion* **51**, 035009; 9pp.
- SU, C.H. & OBERMAN, C. 1968 Collisional damping of a plasma echo. *Phys. Rev. Lett.* **20**, 427–429.
- TOLMAN, E.A. & CATTO, P.J. 2021 Drift kinetic theory of alpha transport by tokamak perturbations. *J. Plasma Phys.* **87**, 855870201; 44pp.
- VELASCO, J.L., CALVO, I., MULAS, S., SÁNCHEZ, E., PARRA, F.I. & CAPPÀ, Á. AND THE W7-X TEAM 2021 A model for the fast evaluation of prompt losses of energetic ions in stellarators. *Nucl. Fusion* **61**, 116059; 26pp.
- WHITE, R.B. 2001 *The Theory of Toroidally Confined Plasmas*, 2nd edn, pp. 298–302. Imperial College Press.