

Iron deficiency is uncommon among lactating women in urban Nepal, despite a high risk of inadequate dietary iron intake

Sigrun Henjum^{1*}, Mari Manger², Eli Skeie², Manjeswori Ulak³, Andrew L. Thorne-Lyman^{4,5}, Ram Chandyo^{2,3}, Prakash S. Shrestha³, Lindsey Locks⁴, Rune J. Ulvik^{6,7}, Wafaie W. Fawzi^{4,8,9} and Tor A. Strand^{2,10}

¹Oslo and Akershus University College of Applied Sciences, PO Box 4, St Olavs Plass, N-0130 Oslo, Norway

²Centre for International Health, University of Bergen, PO Box 7800, 5020 Bergen, Norway

³Department of Child Health, Institute of Medicine, Tribhuvan University, Kathmandu, Nepal

⁴Department of Nutrition, Harvard School of Public Health, 665 Huntington Avenue, Boston, MA 02115, USA

⁵Center on Globalization and Sustainable Development, the Earth Institute, Columbia University, New York, NY, USA

⁶Department of Clinical Science, University of Bergen, 5020 Bergen, Norway

⁷Laboratory of Clinical Biochemistry, Haukeland University Hospital, 5021 Bergen, Norway

⁸Department of Epidemiology, Harvard School of Public Health, 665 Huntington Avenue, Boston, MA 02115, USA

⁹Department of Global Health and Population, Harvard School of Public Health, 665 Huntington Avenue, Boston, MA 02115, USA

¹⁰Division of Medical Services, Innlandet Hospital Trust, 2629 Lillehammer, Norway

(Submitted 3 July 2013 – Final revision received 5 February 2014 – Accepted 21 February 2014 – First published online 8 April 2014)

Abstract

The main objective of the present study was to examine the association between dietary Fe intake and dietary predictors of Fe status and Hb concentration among lactating women in Bhaktapur, Nepal. We included 500 randomly selected lactating women in a cross-sectional survey. Dietary information was obtained through three interactive 24 h recall interviews including personal recipes. Concentrations of Hb and plasma ferritin and soluble transferrin receptors were measured. The daily median Fe intake from food was 17.5 mg, and 70% of the women were found to be at the risk of inadequate dietary Fe intake. Approximately 90% of the women had taken Fe supplements in pregnancy. The prevalence of anaemia was 20% (Hb levels < 123 g/l) and that of Fe deficiency was 5% (plasma ferritin levels < 15 µg/l). In multiple regression analyses, there was a weak positive association between dietary Fe intake and body Fe (β 0.03, 95% CI 0.014, 0.045). Among the women with children aged < 6 months, but not those with older infants, intake of Fe supplements in pregnancy for at least 6 months was positively associated with body Fe (P for interaction < 0.01). Due to a relatively high dietary intake of non-haem Fe combined with low bioavailability, a high proportion of the women in the present study were at the risk of inadequate intake of Fe. The low prevalence of anaemia and Fe deficiency may be explained by the majority of the women consuming Fe supplements in pregnancy.

Key words: Iron deficiency: Lactating women: Iron intakes: Plasma ferritin: Soluble transferrin receptors

Fe deficiency is the most prevalent micronutrient deficiency globally. It is an important cause of anaemia⁽¹⁾ that affects 42% of pregnant women and nearly one-third of all non-pregnant women of reproductive age⁽²⁾. Low dietary Fe content, often combined with low bioavailability from plant-based diets, are important causes of Fe deficiency, especially in pregnant women with increased Fe requirements⁽³⁾.

The degree of Fe absorption depends on the form of Fe consumed and by the presence of dietary enhancers or inhibitors of its absorption. Highly bioavailable haem Fe is likely to

be consumed infrequently and in small amounts by people in resource-poor settings. The majority of dietary Fe consumed in such countries is in the form of non-haem Fe, which is often poorly absorbed due to the presence of dietary inhibitors such as phytate and tannins.

The present study was carried out in Bhaktapur municipality in semi-urban communities in the Kathmandu valley. A previous survey carried out in this population has reported an anaemia prevalence of 12% and that 54% consumed inadequate amounts of Fe⁽⁴⁾. The 2011 Demographic Health

Abbreviations: EAR, estimated average requirement; FCT, food composition table; R:F, soluble serum transferrin receptor:plasma ferritin ratio; RNI, recommended nutrient intake; TfR, transferrin receptor.

* **Corresponding author:** S. Henjum, email sigrun.henjum@hioa.no

Survey showed that 39% of lactating women in Nepal (15–49 years old) were anaemic⁽⁵⁾. Another study conducted among pregnant women in the south-eastern plains of Nepal reported that dietary intake of haem Fe was significantly associated with a lower risk for Fe deficiency without anaemia⁽⁶⁾. However, few studies carried out in developing-country settings have related the measures of dietary intake to the biochemical measures of Fe status⁽⁷⁾. Furthermore, a systematic review on dietary micronutrient intakes of women in resource-poor settings has concluded that there is a need for more documentation of the risk of inadequate micronutrient intakes among women living in low-income settings⁽⁸⁾.

Therefore, the main objective of the present study was to examine the association between dietary Fe intake and dietary predictors of Fe status and Hb concentration in a representative sample among lactating women in Bhaktapur, Nepal.

Methods

Study area and population

A cross-sectional survey was carried out among 500 randomly selected healthy lactating women (17–44 years old) from Bhaktapur municipality, Nepal. Bhaktapur is an urban area located 15 km east of the capital Kathmandu and was chosen because of the socio-economic diversity of this population, which gave us a unique opportunity to explore dietary variation. It has a total population of approximately 75 000, predominantly of the Newari ethnic group, and mostly farmers, semi-skilled or unskilled labourers and daily wage earners.

Study design and participants

From a public health perspective, we wanted to detect deficiencies of micronutrients such as Zn, Fe and vitamin B₁₂ with a prevalence of >25%. It was calculated that 450 mothers would be adequate to detect this prevalence with an absolute precision of 4%, i.e. with a 95% CI ranging from 21 to 29%. Assuming incomplete sampling from approximately 10% of these women, we calculated a final desired sample size of 500 women. In the first stage of sampling, we used a population-proportional-to-size method to select sixty-six of 160 geographical areas ('toles'). In the second stage, we obtained the census lists of all women living in the sixty-six toles and selected the subjects randomly from these lists. We approached 582 women in order to enrol 500 women in the study (Fig. 1). A total of 500 lactating (encompassing both exclusive and partial) women were enrolled in the study and completed the first 24 h dietary recall. Due to the dropouts, the sample sizes for the second and third 24 h recalls were 487 and 477, respectively. A total of eleven women were excluded due to errors in the interactive 24 h recalls, and thus the final sample size consisted of 466 lactating women who had completed three 24 h recalls.

Women came to the hospital to receive physical examinations, dietary interviews and blood draws. The first woman was enrolled in January 2008 and the last in February 2009. The inclusion criteria were that they were lactating, had no



Fig. 1. Flow chart of the recruitment of the study subjects.

self-reported on-going infections and were able to provide household information. Women with anaemia (Hb levels < 123 g/l) were offered free treatment with Fe supplements according to the national guidelines. All women gave written informed consent before the start of the study, which was approved by the ethical review board of the Institute of Medicine, Tribhuvan University. Women were classified as literate if they could read and write, and illiterate if they could do neither or only one of the two. Schooling was defined as primary school (1st–3rd grade), secondary school (4th–10th grade), school-leaving certificate, intermediate school or Bachelor's degree.

Dietary assessment

Nepali-speaking, trained fieldworkers performed the interactive 24 h recalls, and each woman participated in three interactive 24 h recalls⁽⁹⁾. Every fieldworker in the present survey received training by a dietitian for a period of 2 months. The fieldworkers were trained in interview techniques, how to use the electronic scales, how to estimate the volume of different foods, how to collect recipes, how to use the food codes, how to handle difficult situations and how to calibrate the weights before every interview. They were not only trained as a group, but also practised 24 h recalls on each other and on at least five women from the community before conducting the recalls for the present study. To ensure that the days represented normal intake and to minimise interviewer biases, the recalls were obtained to represent three different weekdays with each recall period separated by 2–11 d, and conducted by three different fieldworkers. Saturdays (weekends) were excluded. The 24 h recalls were

collected during 1 year in order to cover the seasonal variation in food supply across the participants in the study area. The same procedure was used to collect the 24 h recalls for each of the 3 d. First, the participants were asked to name all the food and drinks consumed during the preceding day, including anything consumed outside the home and the time of consumption. Second, they were asked to describe the ingredients and cooking methods for each recipe. Third, the amounts of foods and dishes were estimated using an electronic scale (Philips) with a precision of 1 g and a maximum capacity of 5 kg. The scales were calibrated daily. Cooked rice was used for estimating the volume of rice, vegetable stew ('tarkari') and pickles, and water was used to estimate the volume of lentils ('dal'). Fresh vegetables were used to measure the size and quantity of vegetables used in the recipes. The amounts of meat, fish, bread and fruits were estimated by food models and pictures made exclusively for the present study according to Gibson & Ferguson⁽⁹⁾. Clay models were used for estimating the portions of meat and fish, wooden models were used for bread, whereas pictures were used for estimating the amounts of fruits consumed. Finally, the participants were asked to recall snacks consumed between meals during the last 24 h from a list of snack foods, made especially for the present study. The personal recipes of the participants were collected. Standard recipes were made for tea, spices (masala), lentils, bread, vegetable stew (tarkari) and pickles, and were used when the participants had bought ready-made food or when the food had been eaten at someone else's place. The standard recipes were developed from a collection of recipes in a pilot study, and the average of the ingredients from at least twelve recipes for each dish was calculated. Information on the consumption of fortified foods was not collected. Most of the fortified food available in the study area were designed for infants and preschool children, and thus were not commonly consumed by adults.

Intake of iron supplements in pregnancy

Information on the consumption of Fe supplements in pregnancy was collected through questionnaires. The women were encouraged at the hospital to take Fe supplements during pregnancy, and bought Fe supplements from the hospital or from local drug stores. In the questionnaire, the women were asked for how many months they consumed Fe supplements in pregnancy and in which trimester they started to have the supplements. Information on the frequency of consumption, the dosage or the brand name of Fe supplements was not collected, and thus the intake of supplements was not included in the analysis of dietary Fe intake. The common practice for the administration of Fe supplements was 60 mg of elemental iron sulphate from the second trimester of pregnancy.

Nutrient analysis

Because there is no standard food composition table (FCT) available for Nepal, a FCT was compiled for the present study ('Composite Bhaktapur FCT'). In this FCT, nutrient

values and foods were derived from WorldFood 2⁽¹⁰⁾, the 'Nutritive Value of Indian Foods'⁽¹¹⁾ and, where necessary, from the Thai FCT⁽¹²⁾ or the US FCT⁽¹³⁾. The three 24 h dietary recalls and the 'Composite Bhaktapur FCT' were entered in nutrient analysis software designed exclusively for the present study. The source of phytate data was the Indian FCT⁽¹¹⁾. Intake of haem Fe was calculated based on the assumption that haem Fe makes up 40% of Fe in meat, poultry or fish. The usual intake distributions were calculated by the multiple source method, which is characterised by a two-part shrinkage technique applied to the residuals of two regression models: one for the positive daily intake data and another for the event of consumption^(14,15).

Estimation of adequacy of dietary iron intake

Estimation of dietary Fe availability in this population was made according to the method described by Murphy *et al.*⁽¹⁶⁾, using quantitative data on the intake of haem Fe, non-haem Fe, coffee and tea, as well as on the amount of ascorbic acid and protein from meat, fish and poultry per 4184 kJ (1000 kcal) of energy consumed. The average bioavailability of dietary Fe intake in this population was calculated to be 4.5%. The bioavailability of dietary Fe was also calculated according to the algorithm developed by Bhargava *et al.*⁽¹⁷⁾. This algorithm includes estimation of Fe stores, intake of fish and meat, ascorbic acid, phytate, non-haem Fe and haem Fe, which was developed for Bangladeshi women, a population that might be equalled to that of the present study. This algorithm gave an estimated average bioavailability of 1.8%. The WHO recommends the use of the calculation of 5 or 10% Fe absorption in developing countries depending on the diet⁽¹⁸⁾. We therefore used the 5% bioavailability assumption for the evaluation of Fe intake in the present study.

The 2004 Estimated Average Requirements (EAR) and Recommended Nutrient Intake (RNI) of the FAO/WHO were used to evaluate the nutritional adequacy of the women's diets depending on post-partum. The RNI of Fe for lactating women (0–3 months) and women aged 19–50 years is 30 and 58.8 mg, respectively, when the bioavailability is 5%. The risk of inadequate dietary Fe intake for women who had been lactating for <3 months was classified using the following definitions: very high risk (average dietary Fe intake below the EAR); moderate risk (average dietary Fe intake between the EAR and the RNI); low risk (average dietary Fe intake higher than the RNI)⁽⁸⁾. A full probability approach was used⁽¹⁹⁾ that estimated the risk of inadequate dietary Fe intake as a total product of the probability of inadequacy for a given range of intake multiplied by the percentage of women with intakes in that range.

Anthropometric measurements

Weight was measured using a UNICEF weighing scale (Seca; Salter). Height was measured with a locally made board in the clinic and calibrated weekly. Maternal BMI was calculated as weight/(height)² (kg/m²). BMI < 18.5 kg/m² was



considered as underweight, $18.5 \text{ kg/m}^2 > \text{BMI} < 25 \text{ kg/m}^2$ as normal weight and $\text{BMI} \geq 25 \text{ kg/m}^2$ as overweight⁽²⁰⁾.

Laboratory tests

During the first hospital visit, the first of the three 24 h recalls was performed, and a venous blood sample was collected from a cubital vein into a micronutrient-free, heparinised polypropylene tube (Sarstedt). Hb concentration was measured immediately by HemoCue (Vedbæk)⁽²¹⁾, which was regularly calibrated as recommended by the manufacturer. After centrifuging the sample at 760g for 10 min at room temperature, plasma was transferred to micronutrient-free polypropylene vials (Eppendorf, Hinz, Germany) and stored at -70°C before transport on dry ice to Norway, and then further stored at -80°C until analysis was performed at the Laboratory of Clinical Biochemistry, Haukeland University Hospital, Bergen. The plasma concentration of biochemical components was determined on a Modular Analytics System by Roche Diagnostics (Roche Diagnostics GmbH), with the analytical CV being 5% for each test. Plasma ferritin concentration was analysed by an electrochemiluminescence immunoassay, while soluble transferrin receptor (TfR) concentration was analysed by immunoturbidimetry.

Cut-off limits for the analytical test

After adjusting for the altitude of the study area (1400 m), anaemia in this population was defined as Hb levels $< 123 \text{ g/l}$ ⁽²²⁾. Mild anaemia was defined as Hb levels between 103 and 122 g/l, moderate anaemia as Hb levels between 73 and 102 g/l and severe anaemia as Hb levels $< 73 \text{ g/l}$ ⁽²²⁾. Fe deficiency, expressed as depleted Fe stores, was defined as plasma ferritin levels $< 15 \mu\text{g/l}$ ⁽¹⁾. Increased need of Fe in the erythropoietic bone marrow and peripheral tissues was defined as TfR levels $> 4.4 \text{ mg/l}$ ⁽²³⁾. As increased concentration of C-reactive protein is a sensitive marker of inflammation, women with C-reactive protein levels $> 5 \text{ mg/l}$ were excluded from the analyses in which plasma ferritin was involved.

Calculation of body iron stores

Body Fe assessed as surplus (positive value) or deficit (negative value) of Fe in the tissues was calculated by using the formula described by Cook *et al.*⁽²⁴⁾. The formula was derived from a close linear relationship that was found between the logarithm of the ratio of the concentration of soluble serum TfR and plasma ferritin (R:F ratio, $\mu\text{g}/\mu\text{g}$) and body Fe expressed as mg Fe/kg body weight, corrected for the absorption of dietary Fe:

$$\text{mg Fe/kg} = -(\log(\text{R:F ratio}) - 2.8229)/0.1207.$$

Since analysis of TfR refers to the in-house ELISA developed by Flowers *et al.*⁽²⁵⁾, the results obtained for TfR by the Roche method were converted by using the regression

equation presented by Pfeiffer *et al.*⁽²⁶⁾:

$$\text{TfR-Flowers} = 1.5 \times \text{TfR-Roche} + 0.35.$$

Data processing and statistical analysis

Data were analysed using SPSS version 17 (SPSS, Inc.), STATA version 12 (Stata Corporation) and R version 2.16 (r-project.org). Continuous data that were not normally distributed are presented as medians and 25th (P₂₅) and 75th (P₇₅) percentiles. Two-tailed tests with a significance level of 5% were used for all analyses. We measured the association of relevant independent

Table 1. Demographic and anthropometric characteristics of 500 lactating women in Bhaktapur, Nepal

(Mean values and standard deviations; number of subjects and percentages)

	Mean	SD
Demographic information		
Age (years)	25.8	4.2
Employed*		
n	103	
%	21.0	
Years of schooling†	2.5	1.0
Literate		
n	379	
%	81.5	
BMI (kg/m ²)‡	22.5	3.1
< 18.5 kg/m ²		
n	23	
%	4.6	
18.5–25 kg/m ²		
n	366	
%	78.7	
> 25 kg/m ²		
n	84	
%	17.0	
Parity	1.9	0.9
One		
n	203	
%	41.0	
Two		
n	206	
%	41.0	
Three or more		
n	90	
%	18.0	
Intake of Fe supplements in pregnancy		
n	449	
%	89.8	
Months of Fe supplementation (n 449)	4.8	1.9
Started in the first trimester		
n	144	
%	29.0	
Started in the second trimester		
n	262	
%	52.4	
Started in the third trimester		
n	43	
%	8.6	

* Farmers, carpet factory workers, daily wage earners, self-employed or service workers.

† Primary school (1st–3rd grade); secondary school (4th–10th grade); school-leaving certificate; intermediate school; or Bachelor's degree.

‡ BMI $< 18.5 \text{ kg/m}^2$ defined as underweight; BMI 18.5–25 kg/m² defined as normal weight; BMI $> 25 \text{ kg/m}^2$ defined as overweight.

Table 2. Dietary intakes* of energy, nutrients, foods and constituents influencing iron absorption in 466 lactating women in Bhaktapur, Nepal (Mean values and standard deviations; median values and 25th–75th percentiles)

Variables	Median	25th–75th percentile
Energy intake (kcal)		
Mean	2024	
SD	312	
Energy intake (kJ)		
Mean	8468	
SD	1305	
Dietary Fe (mg)	17.5	13.3–24.7
Haem (mg)	0.3	0.0–0.6
Available Fe (mg)†	0.8	0.6–1.3
Vitamin C (mg)	55.8	42.0–71.7
Ca (mg)	461.0	334.3–632.4
Meat, fish and poultry (g)‡	54.0	37.9–72.5
Protein meat, fish and poultry (g)‡	18.35	7.9–23.7
Dry tea (g)§	2.1	1.5–2.6
Phytate (mg)		
Mean	3304	
SD	776	
Phytate:Fe molar ratio	15.9	10.4–24.3

* Usual intakes based on three 24 h recalls per women.

† 5% bioavailability calculated according to Murphy *et al.*⁽¹⁶⁾.

‡ *n* 327.

§ *n* 425.

|| Gibson & Ferguson⁽⁹⁾.

variables with dependent variables (Hb and body Fe) using multiple linear regression models. The variables that were known to influence Hb concentration and body Fe as well as selected socio-economic variables were included in initial crude models. Candidate variables included dietary Fe intake, vitamin C intake, vitamin A intake (β -carotene), phytate intake, use of Fe supplements in pregnancy (for at least 6 months), mother's age, mother's BMI, mother's literacy, parity and infant's age at the time of the baseline visit. All covariates showing a linear association ($P < 0.10$) in the crude regression models were included in a preliminary multiple regression model. The variables that were still significantly associated in this model ($P < 0.10$) were retained in the final model⁽²⁷⁾. Analysis of the residuals was performed in order to examine the fit of the model. In the final model, the following interactions between the independent variables were assessed and included if the interaction term was significant ($P < 0.10$): (Dietary Fe intake \times vitamin C (dichotomous variable, intake P_{25} (42 mg/d) and P_{75} (72 mg/d)) and time since birth (dichotomous variable, cut-off 6 months) \times prenatal Fe supplement for at least 6 months). We explored and depicted the linearity of the associations between the independent and dependent variables in generalised additive models⁽²⁸⁾. We adjusted for clustering of outcomes due to the sampling design using the SVY group of commands for complex survey data in STATA.

Results

The mean age of the 500 enrolled women was 25.8 years, and the majority were literate, had a healthy BMI and less than three children. The intake of Fe supplements in pregnancy was reported by 90% of the women, the mean duration of

which was 4.8 months, and more than half of them started the supplementation during the second trimester (Table 1).

Intakes of dietary iron and risk of inadequate intakes of iron

The intake of dietary Fe and other dietary factors with the potential to affect Fe absorption are presented in Table 2. The median daily Fe intake from food was 17.5 mg and almost all the Fe consumed was in the form of non-haem Fe (98%). The daily dietary intake of phytate (3304 mg/d) was well above the level known to adversely affect Fe absorption, and the main sources were rice, dal, potato and whole-wheat flour. Enhancers of Fe absorption (meat, fish and poultry, and vitamin C) were consumed in moderate amounts. The principal sources of dietary Fe in this population were mustard leaves, rice flakes ('beaten rice'), turnip leaves, rice and whole-wheat flour (Table 3). Based on the EAR of the WHO/FAO for women consuming a diet with low bioavailability (5%), 72.7% of the women who were ≤ 3 months post-partum (*n* 33) were considered to have a very high risk of inadequate dietary Fe intake. Only 15% of the women who were ≤ 3 months post-partum were at a low risk of inadequate dietary Fe intake (Table 4). Using a full probability approach for women who were > 3 months post-partum (*n* 432), the total prevalence of inadequate intake of dietary Fe was estimated to be 78%.

Iron status and anaemia

After adjusting for the altitude of the study area (1400 m), the prevalence of anaemia (Hb levels < 123 g/l) was 20% and mild anaemia (Hb levels 103–122 g/l) was 17%. Of the women, only 5% (*n* 26) had depleted Fe stores with plasma ferritin levels < 15 μ g/l, while 15% (*n* 73) had TfR levels > 4.4 mg/l, indicating an insufficient supply of Fe to the erythropoietic bone marrow and peripheral tissues. Of the women with plasma ferritin levels < 15 μ g/l, 69% (*n* 18) also had TfR levels > 4.4 mg/l as a sign of empty Fe stores with accompanying Fe-deficient erythropoiesis evidenced by a mean Hb concentration of 112 (SD 1.3) g/l. Thus, the prevalence in the whole study group of true Fe-deficient anaemia was 3.6%. The remaining eight subjects with plasma ferritin levels < 15 μ g/l and no increase in the levels of TfR had a sufficient supply of Fe to the tissues despite depleted Fe stores.

Table 3. Main sources of dietary iron in Bhaktapur, Nepal

Food items	Contribution to Fe intake (%)	Fe content (mg/100 g food)*
Mustard leaves	25.4	16.3
Rice flakes	50.0	20.0†
Turnip leaves	10.2	28.4
Rice	8.1	0.2
Whole-wheat flour	4.9	4.9

* Value obtained from the Indian food composition table⁽¹¹⁾.

† When using a value from Suma *et al.*⁽³³⁾, the contribution to iron intake was 18% and the iron content was 5.2 mg.

Table 4. Proportion of different risk groups of inadequate iron intake for thirty-three lactating women (≤ 3 months post-partum) consuming a diet with 5% iron bioavailability* in Bhaktapur, Nepal (Number of subjects and percentages)

Risk of inadequate dietary Fe intake	Dietary Fe intake (mg)	Proportion	
		n	%
Very high†	≤ 23.4	24	72.7
Moderate‡	23.4–30.0	4	12.1
Low§	≥ 30.0	5	15.2

* Iron bioavailability calculated according to Murphy *et al.*⁽¹⁶⁾ and WHO recommendations⁽¹⁸⁾.

† Average dietary iron intake is less than or equal to the estimated average requirement for lactating women.

‡ Average dietary iron intake is greater than the estimated average requirement and less than or equal to the recommended nutrient intake for lactating women.

§ Average dietary iron intake is greater than the recommended nutrient intake for lactating women.

Of the study participants, 103 had plasma ferritin levels between 15 and 35 $\mu\text{g/l}$, which is compatible with depleted or very small Fe stores in many women⁽²⁹⁾. In this subgroup, twenty-one women had TfR levels $> 4.4 \text{ mg/l}$, indicating restricted synthesis of Hb due to an insufficient supply of Fe to the bone marrow. Since low levels of plasma ferritin may be the evidence of a negative Fe balance, it is conceivable that thirty-nine (53%, with plasma ferritin levels $< 35 \text{ mg/l}$) of the seventy-three women with increased TfR values could be explained by a restricted supply of Fe to the tissues.

Body Fe calculated from the logarithm of the geometric mean of the R:F ratios (95.6) was 7.0 (SD 3.3) mg Fe/kg body weight with a range between -8.9 and 14.5 mg Fe/kg . Of the women, fifteen (3%) had negative values indicating tissue Fe deficiency, with a mean of -2.8 (SD 3.4) mg Fe/kg. The rest of the group ($n 485$, i.e. 97%) had a mean body Fe stores of 7.3 (SD 5.5) mg Fe/kg (Table 5).

Dietary predictors of iron status and anaemia

In the linear regression models (Table 6), intake of Fe supplements in pregnancy predicted Hb concentration. Dietary Fe intake and intake of Fe supplements in pregnancy predicted body Fe. Intake of Fe supplements in pregnancy was associated with a higher Hb concentration of 0.29 (95% CI 0.04, 0.54) g/l ($P=0.03$). Dietary Fe intake and potential enhancers and inhibitors of Fe absorption were not associated with Hb concentration. There was a weak positive association between dietary intake of Fe and body Fe ($\beta 0.03$, 95% CI 0.014, 0.045). In addition, for women with children aged < 6 months, but not for those with older infants (P for interaction < 0.01), intake of Fe supplements in pregnancy was positively associated with body Fe stores. We identified the same predictors in the multiple linear regression models using ferritin and TfR concentrations as the dependent variables. The R^2 values in the models with ferritin and TfR were 0.22 and 0.09, respectively. The association of dietary Fe intake with biochemical markers is also depicted in the graphs obtained from generalised additive models (Fig. 2).

Discussion

In the population of lactating women in Bhaktapur, we found that more than 98% of dietary Fe intake consumed was in the form of non-haem Fe and that the intake of phytate was high. Therefore, $> 70\%$ of these women were estimated to be at the risk of inadequate Fe intake. At the same time, only 5% of the women had plasma ferritin levels indicative of Fe deficiency and 15% of the women had elevated levels of TfR mainly indicative of Fe-deficient erythropoiesis. We demonstrated that dietary Fe intake and intake of Fe supplements in pregnancy predicted body Fe. Furthermore, intake of Fe supplements in pregnancy, but not dietary variables, showed a strong positive association with Hb concentration.

Table 5. Hb, plasma ferritin and transferrin receptor concentrations among 500 lactating women in Bhaktapur, Nepal

(Mean values and standard deviations or ranges; number of subjects and percentages)

	Mean	SD
Hb (g/l)	131	13
Anaemia, $< 120 \text{ g/l}$		
n	61	
%	12	
Anaemia, $< 123 \text{ g/l}^*$		
n	100	
%	20	
Mild anaemia, 103–122 g/l*		
n	87	
%	17.4	
Moderate anaemia, 73–102 g/l*		
n	13	
%	2.6	
Severe anaemia, $< 73 \text{ g/l}^*$		
n	0	
%	0	
Plasma ferritin ($\mu\text{g/l}$)†	68.8	46.2
$< 15 \mu\text{g/l}^\ddagger$		
n	26	
%	5.3	
$< 35 \mu\text{g/l}^\ddagger$		
n	129	
%	25.8	
Plasma transferrin receptor (TfR, mg/l)	3.4	1.5
$> 4.4 \text{ mg/l}^\ddagger$		
n	73	
%	15	
Ferritin $< 15 \mu\text{g/l}$ and TfR $> 4.4 \text{ mg/l}$		
n	18	
%	3.6	
C-reactive protein (CRP) $> 5.0 \text{ mg/l}$		
n	24	
%	5	
Body Fe (mg Fe/kg body weight)§	7.0	3.3
Range	-8.9 to 14.5	
Tissue Fe deficit	-2.8	3.4
n	15	
%	3	
Storage Fe present	7.3	5.5
n	485	
%	97	

* Threshold for defining anaemia is adjusted for altitude⁽²²⁾.

† $n 476$; twenty-four women excluded due to the elevated WHO cut-off levels of CRP for the depletion of iron stores⁽²¹⁾.

‡ Cut-off point for iron-deficient erythropoiesis⁽²³⁾.

§ Negative value indicates a quantitative measure of tissue iron deficit (i.e. lack of stored iron).

Table 6. Multiple linear regression models of the relationship between dietary iron intake, Hb concentration and body iron (n 500) (β Coefficients and 95 % confidence intervals)

	Multiple adjusted* β	95 % CI	P	Standardised β coefficients
Model 1: Hb (R^2 0.02)				
Intake of Fe supplements in pregnancy†‡	0.29	0.04, 0.54	0.03	0.11
Model 2: body Fe (mg Fe/kg body weight), R^2 0.20				
Dietary Fe (mg)	0.029	0.014, 0.045	<0.01	0.16
Time since birth \times Fe in pregnancy§	2.69	1.54, 3.84	<0.01	
Intake of Fe supplements in pregnancy				
Time since birth < 6 months	2.72	1.79, 3.65	<0.01	0.40
Time since birth \geq 6 months	0.03	-0.67, 0.78	0.93	0.00
Time since birth < 6 months				
Intake of Fe supplements in pregnancy	2.24	1.36, 3.13	<0.01	0.33
Non-intake of Fe supplements in pregnancy	-0.45	-0.29, 1.18	0.23	-0.07
Mother's age	0.21	0.14, 0.28	<0.01	0.26
Literacy¶	0.81	0.25, 1.39	<0.01	0.12
Ownership of land¶	0.74	0.19, 1.30	<0.01	0.11

* Both models included mother's age, parity, literacy and child's age.

† Intake of Fe supplements in pregnancy at least 6 months, dichotomous variable (yes/no).

‡ No significant interaction between time since birth (dichotomous variable, cut-off 6 months) and iron supplements in pregnancy for at least 6 months.

§ Interaction between time since birth (dichotomous variable, cut-off 6 months) and iron supplements in pregnancy for at least 6 months.

|| Continuous variable.

¶ Dichotomous variable (yes/no).

Dietary iron intake

The Fe consumed by these Nepali women was almost exclusively in the form of non-haem Fe. It should be noted that no fortified food staples were available to this population at the time of data collection. The main contributors of dietary Fe intake were cooked green leaf relishes, unrefined whole-meal bread 'roti' and rice flakes, all of which contained significant amounts of phytate based on available food composition data^(10,11), and therefore impaired the bioavailability of Fe. In addition, meat that enhances the uptake of non-haem Fe⁽³⁰⁾ was not commonly consumed. In contrast, intakes of vitamin C were about 70 % of the RNI for lactating women and obtained from vegetable relishes commonly eaten with meals, which may have had a positive impact on the bioavailability of Fe. Tea consumption is unlikely to have had any substantial negative impact on the bioavailability of Fe because it is typically not consumed with meals.

The dietary Fe sources were similar to that of a previous study among non-pregnant, non-lactating women in Bhaktapur; however, that study reported lower mean intakes of Fe (8.4 mg/d)⁽⁴⁾. This may in part be due to the different FCT used. The previous study used exclusively data from the WorldFood Dietary Assessment System⁽¹⁰⁾, whereas we used primarily the Indian food composition data⁽¹¹⁾ because Indian foods were more specific to food items consumed in Nepal. The Fe content of mustard leaves, turnip leaves, rice flakes, refined rice and whole-wheat flour was consistently higher in the Indian FCT, compared with similar substitute foods in WorldFood. These values may be higher than the true Fe content of these common Nepali foods. However, a study in the Kathmandu valley found that dietary Fe intakes based on chemical analysis of 24 h diet composites were three times higher than the calculated dietary intakes for the same 24 h period using the US Department of Agriculture

database⁽³¹⁾. Therefore, it is possible that the Fe content of the foods consumed in Bhaktapur is higher than estimated. There is a clear need for a Nepal-specific FCT.

Probability of inadequacy

Based on plasma ferritin and TfR levels, we found the prevalence of Fe deficiency to be 5.3 and 14.6 %, respectively. Yet, we estimated that 70 % of the women were at the risk of inadequate dietary Fe intake. There are several possible reasons for this discrepancy. First, by using a cut-off value of plasma ferritin <15 μ g/l for depleted Fe stores, we have probably underestimated the prevalence of Fe deficiency as signalled by this biomarker (the prevalence of Fe deficiency was 25.8 % by using the cut-off values of plasma ferritin <35 μ g/l). As shown in the study by Hallberg *et al.*⁽²⁹⁾, at this cut-off value, the diagnostic sensitivity and specificity of plasma ferritin are 75 and 98 %, respectively. They have found that the Fe stores could be negligible at plasma ferritin concentrations from 35 μ g/l and below. Second, 90 % of the women had used Fe supplements in pregnancy and more than half of them initiated the supplementation in the second trimester. Thus, temporary Fe supplementation in pregnancy is likely to have had a positive effect on Fe stores during the first few months of lactation, but probably not later⁽⁵⁾ when Fe loss increased due to the return of menstruation and probably because Fe supplementation was stopped. Third, we may have overestimated the probability of inadequate dietary Fe intake because we used the EAR for all non-pregnant, non-lactating women who were >3 months post-partum. Many of these women may still have had lactation amenorrhoea and therefore had lower Fe requirements than what was the basis of the EAR used. However, the risk of inadequate dietary Fe intake for women \leq 3 months post-partum, for whom lactation amenorrhoea was taken into

account, was similar to those who had given birth earlier. Fourth, as discussed above, the food composition data used may not have accurately reflected the true Fe, phytate and vitamin C contents of Nepali foods. It is possible that the Fe content of several foods is higher and/or the phytate content is lower than what was estimated based on the available data.

The calculation of body Fe based on the ratio between the concentrations of TfR and plasma ferritin gives a more precise picture of the tissue Fe content than what is achieved by using the cut-off limits of the biochemical tests. The present results in Fe-replete and Fe-deficient women were somewhat different from what was reported by Cook *et al.*⁽²⁴⁾ in their study of women between 20 and 45 years of age. In women with normal Fe status, they have found mean Fe stores of 5.5 (SD 3.35) *v.* 7.3 (SD 5.5) mg Fe/kg in the present study, and in women with Fe deficiency, they have found a mean deficit in tissue Fe of -3.9 (SD 3.23) *v.* -2.8 (SD 3.4) mg/kg in the present study. As discussed above, the Fe status of women in the present study may have benefited from Fe supplementation and less Fe loss due to amenorrhoea.

Dietary predictors of iron status and anaemia

We found a significant, albeit weak, positive association between body Fe and dietary Fe intake, and also demonstrated that this association was linear. The lack of an association between dietary Fe intake and Hb concentration may in part be due to the fact that only a small proportion of anaemia in this population was Fe-deficiency anaemia; 82 and 63% of the anaemic women did not have low plasma ferritin or elevated TfR levels, respectively. However, the generalised additive model curve revealed a linear association between dietary Fe intake and Hb concentration at lower dietary Fe intakes (Fig. 2). However, this effect is small compared with the strong association between the intake of Fe supplements in pregnancy and Hb concentration in the present study. As discussed earlier, this effect persists in the lactation period. In addition, the 2011 Nepal Demographic and Health Survey showed that 80% of Nepali women took Fe supplementation during last pregnancy and 41% consumed Fe supplements post-partum⁽⁵⁾. Lastly, other micronutrient deficiencies may also have been responsible for lower Hb concentrations observed in the present study, such as folate, vitamin B₁₂ and vitamin A⁽³²⁾, but their relationships to Hb were not studied.

Strengths and weaknesses

The present study had a number of strengths. We had a representative sample of lactating women, with a relatively large sample when compared with that used in most other dietary studies. We had three interactive 24 h recalls with personal recipes adapted to local foods. We trained local staff to perform the recall interviews and the interviews were done throughout the year, covering the seasonal variation in food intake at a group level. Using plasma ferritin and TfR as biomarkers, the near-linear association of their relevant changes with dietary Fe intake indicates strong relative validity of our




Fig. 2. Association between daily iron intake and concentrations of (a) Hb, (b) plasma ferritin and (c) plasma transferrin receptor in lactating women (*n* 466) in Bhaktapur, Nepal. The graph was constructed using the generalised additive models in R. The solid curves depict the estimated dose-response curve; the shaded areas represent the 95% CI. The small vertical lines on the x-axis show the distribution of the observations.

adapted interactive 24 h recall method. Inflammation will obscure the interpretation of plasma ferritin, but not TfR, which can explain the difference in the prevalence of Fe deficiency based on plasma ferritin and TfR. However, chronic and acute illness was an exclusion criterion in the present study and very few (*n* 24 women) had elevated C-reactive protein levels. Therefore, we believe that the difference in prevalence is not due to inflammation. The primary weakness of the present study was the reliance of external FCT, which may not reflect the true nutrient composition of local Nepali foods. Our findings may have also been adversely affected by bias caused by the fact that participating women knew they were coming in for a dietary interview and may have altered their diet. However, the fact that few women had insufficient Fe intake may suggest that potential bias would have been towards the overestimation, rather than underestimation, of Fe intake.

Conclusion

Probably due to a high dietary intake of non-haem Fe combined with low bioavailability, a high proportion of the lactating women in the present study were at the risk of inadequate intake of Fe. The low prevalence of anaemia and Fe deficiency may be explained by the majority of the women consuming Fe supplements in pregnancy.

Acknowledgements

The present study was funded by a grant from the Norwegian Council of Universities' Committee for Development Research, Education (project no. NUFUSM-2007/10177), Research Council of Norway (project no. 172226), and a grant from the GCRieber Funds and Feed the Future Food Security Innovation Lab: Collaborative Research on Nutrition which is funded by the United States Agency for International Development. A. L. T.-L. was supported by the National Institutes of Health training grant T32 #DK 007703.

The contributions of the authors were as follows: T. A. S., R. C., S. H., W. W. F. and P. S. S. designed the research; R. C., M. U., E. S. and S. H. conducted the research; S. H., A. T.-L., R. J. U., L. L. and T.A.S analysed the data; S. H. and M. M. wrote the paper; S. H. and T. A. S. had primary responsibility for the final content. All authors read and approved the final manuscript.

None of the authors has any conflict of interest to declare.

References

- World Health Organization, United Nations Children's Fund, United Nations University (2001) *Iron Deficiency Anaemia Assessment, Prevention, and Control. A Guide For Programme Managers*. Geneva: WHO. http://www.who.int/nutrition/publications/micronutrients/anaemia_iron_deficiency/WHO_NHD_01.3/en/index.html
- McLean E, Cogswell M, Egli I, *et al.* (2009) Worldwide prevalence of anaemia, WHO Vitamin and Mineral Nutrition Information System, 1993–2005. *Public Health Nutr* **12**, 444–454.
- Zimmermann MB & Hurrell RF (2007) Nutritional iron deficiency. *Lancet* **370**, 511–520.
- Chandyo RK, Strand TA, Ulvik RJ, *et al.* (2007) Prevalence of iron deficiency and anemia among healthy women of reproductive age in Bhaktapur, Nepal. *Eur J Clin Nutr* **61**, 262–269.
- NDHS: Ministry of Health and Population (Nepal) (2012) *Nepal Demographic Health Survey – 2011*. Kathmandu, Nepal: New ERA and ICF International.
- Makhoul Z, Taren D, Duncan B, *et al.* (2012) Risk factors associated with anemia, iron deficiency and iron deficiency anemia in rural Nepali pregnant women. *Southeast Asian J Trop Med Public Health* **43**, 735–746.
- Backstrand JR, Allen LH, Black AK, *et al.* (2002) Diet and iron status of nonpregnant women in rural Central Mexico. *Am J Clin Nutr* **76**, 156–164.
- Torheim LE, Ferguson EL, Penrose K, *et al.* (2010) Women in resource-poor settings are at risk of inadequate intakes of multiple micronutrients. *J Nutr* **140**, 2051S–2058S.
- Gibson RS & Ferguson EL (2008) *An Interactive 24-Hour Recall for Assessing the Adequacy of Iron and Zinc Intakes in Developing Countries*. HarvestPlus Technical Monograph 8. Washington, DC: International Food Policy Research Institute: International Food Policy Research Institute (IFPRI) and International Center for Tropical Agriculture (CIAT).
- Food and Agriculture Organization of the United Nations (2010) Food Composition: Overview of the WorldFood Dietary Assessment System. http://typo3.fao.org/fileadmin/templates/food_composition/documents/World_food_dietary_assessment_system/Overview_of_the_WorldFood_Dietary_Assessment_System.pdf
- Gopalan C, Rama Sastri B & Balasubramanian S (1989) *Nutritive Value of Indian Foods*. Hyderabad, India: National Institute of Nutrition.
- Institute of Nutrition Mahidol University (2002) *Food Composition Database for INMUCAL Program*. Thailand: Institute of Nutrition.
- United States Department of Agriculture, Agricultural Research Service (2012) Search the USDA National Nutrient Database for Standard Reference. <http://ndb.nal.usda.gov/>
- Haubrock J, Nothlings U, Volatier JL, *et al.* (2011) Estimating usual food intake distributions by using the multiple source method in the EPIC-Potsdam Calibration Study. *J Nutr* **141**, 914–920.
- Harttig U, Haubrock J, Knuppel S, *et al.* (2011) The MSM program: web-based statistics package for estimating usual dietary intake using the Multiple Source Method. *Eur J Clin Nutr* **65**, Suppl. 1, S87–S91.
- Murphy SP, Beaton GH & Calloway DH (1992) Estimated mineral intakes of toddlers: predicted prevalence of inadequacy in village populations in Egypt, Kenya, and Mexico. *Am J Clin Nutr* **56**, 565–572.
- Bhargava A, Bouis HE & Scrimshaw NS (2001) Dietary intakes and socioeconomic factors are associated with the hemoglobin concentration of Bangladeshi women. *J Nutr* **131**, 758–764.
- World Health Organization, Food and Agriculture Organization of the United Nations (2002) *Human Vitamin and Mineral Requirements*. Rome, Italy: Food and Agriculture Organization of the United Nations (FAO).
- World Health Organization, Food and Agriculture Organization of the United Nations (2006) *Guidelines for Food Fortification with Micronutrients* [LH Allen, B de Benoist, O Dary and R Hurrell, editors]. Geneva: World Health Organization, Food and Agriculture Organization of the United Nations.
- WHO (2006) *Global Database on Body Mass Index: An Interactive Surveillance Tool for Monitoring Nutrition Transition*. Geneva: WHO.
- Hemocue Corporate (2011) Our Products – Hemoglobin Portfolio. Available at <http://www.hemocue.com/index.php?page=51>
- Goodman RA (1989) Current trends: CDC criteria for anemia in children and child bearing-aged women. *MMWR Morb Mortality Wkly Rep* **38**, 400–404.
- Kolbe-Busch S, Lotz J, Hafner G, *et al.* (2002) Multicenter evaluation of a fully mechanized soluble transferrin receptor assay on the Hitachi and Cobas Integra analysers. The determination of reference ranges. *Clin Chem Lab Med* **40**, 529–536.
- Cook JD, Flowers CH & Skikne BS (2003) The quantitative assessment of body iron. *Blood* **101**, 3359–3364.
- Flowers CH, Skikne BS, Covell AM, *et al.* (1989) The clinical measurement of serum transferrin receptor. *J Lab Clin Med* **114**, 368–377.
- Pfeiffer CM, Cook JD, Mei Z, *et al.* (2007) Evaluation of an automated soluble transferrin receptor (sTfR) assay



- on the Roche Hitachi analyzer and its comparison to two ELISA assays. *Clin Chim Acta* **382**, 112–116.
27. Hosmer DW & Lemeshow S (2000) *Applied Logistic Regression*, 2nd ed. New York: John Wiley & Sons.
 28. Wood SN (2000) Modelling and smoothing parameter estimation with multiple quadratic penalties. *J R Statist Soc B* **62**, 413–428.
 29. Hallberg L, Bengtsson C, Lapidus L, *et al.* (1993) Screening for iron deficiency: an analysis based on bone-marrow examinations and serum ferritin determinations in a population sample of women. *Br J Haematol* **85**, 787–798.
 30. Bothwell TH, Charlton RW, Cook JD, *et al.* (1979) *Iron Metabolism in Man*. Oxford, UK: Blackwell Scientific Publications.
 31. Moser PB, Reynolds RD, Acharya S, *et al.* (1988) Copper, iron, zinc, and selenium dietary intake and status of Nepalese lactating women and their breast-fed infants. *Am J Clin Nutr* **47**, 729–734.
 32. WHO (2004) *Assessing the Iron Status of Populations: A Report of a Joint World Health Organization/Centers for Disease Control Technical Consultation on the Assessment of Iron Status at the Population Level*. Geneva: WHO.
 33. Suma RC, Sheetal G, Jyothi LA, *et al.* (2007) Influence of phytin phosphorous and dietary fibre on *in vitro* iron and calcium bioavailability from rice flakes. *Int J Food Sci Nutr* **58**, 637–643.