

Research Article

Cite this article: Robertson HE, Schiffer PH, Telford MJ (2018). The mitochondrial genomes of the mesozoans *Intoshia linei*, *Dicyema* sp. and *Dicyema japonicum*. *Parasitology Open* 4, e16, 1–9. <https://doi.org/10.1017/pao.2018.12>

Received: 9 May 2018

Revised: 15 June 2018

Accepted: 22 June 2018

Key words:

Mesozoa; mitochondrial genomes; mitochondrial mini-circles

Author for correspondence:

Maximilian J. Telford,

E-mail: m.telford@ucl.ac.uk

The mitochondrial genomes of the mesozoans *Intoshia linei*, *Dicyema* sp. and *Dicyema japonicum*

Helen E. Robertson, Philipp H. Schiffer and Maximilian J. Telford

Department of Genetics, Evolution and Environment, Centre for Life's Origins and Evolution, University College London, Darwin Building, Gower Street, London, WC1E 6BT, UK

Abstract

The Dicyemida and Orthonectida are two groups of tiny, simple, vermiform parasites that have historically been united in a group named the Mesozoa. Both Dicyemida and Orthonectida have just two cell layers and appear to lack any defined tissues. They were initially thought to be evolutionary intermediates between protozoans and metazoans but more recent analyses indicate that they are protostomian metazoans that have undergone secondary simplification from a complex ancestor. Here we describe the first almost complete mitochondrial genome sequence from an orthonectid, *Intoshia linei*, and describe nine and eight mitochondrial protein-coding genes from *Dicyema* sp. and *Dicyema japonicum*, respectively. The 14 247 base pair long *I. linei* sequence has typical metazoan gene content, but is exceptionally AT-rich, and has a unique gene order. The data we have analysed from the Dicyemida provide very limited support for the suggestion that dicyemid mitochondrial genes are found on discrete mini-circles, as opposed to the large circular mitochondrial genomes that are typical of the Metazoa. The *cox1* gene from dicyemid species has a series of conserved, in-frame deletions that is unique to this lineage. Using *cox1* genes from across the genus *Dicyema*, we report the first internal phylogeny of this group.

Introduction

The Mesozoa is a historic name given to two different groups of very small, vermiform and morphologically simple parasitic animals: the Dicyemida (=Rhombozoa), whose adults are made up of approximately 40 cells (Furuya and Tsuneki, 2003); and the Orthonectida, which possess just a few hundred cells. As there is uncertainty over whether Dicyemida and Orthonectida form a monophyletic group, the Mesozoa grouping is now generally used informally rather than as a formal taxonomic assignment as a phylum. Both dicyemids and orthonectids are parasites of various marine animals: the dicyemids live in the renal tissue of cephalopods, whilst orthonectids occupy the internal body spaces of a variety of marine invertebrates, including brittle stars, bivalve molluscs, nemerteans and polychaetes.

Adults of both dicyemids and orthonectids have just two cell layers. Although they are multicellular, they lack defined complex tissues and organs and there is no evidence for the presence of true ectoderm or endoderm (Margulis and Chapman, 2009). Adult mesozoans have an external layer of multiciliated cells, which facilitate movement, and at least one reproductive cell. Analysis of the orthonectid *Intoshia linei* found evidence for a simple nervous system, comprising just 10–12 nerve cells, and a simple muscular system, composed of four longitudinal and 9–11 circular muscle cells (Slyusarev and Starunov, 2016).

In the 19th century, their simple body organization led to the idea that members of the Mesozoa – as their name suggests – were an evolutionary intermediate between the protozoans and the metazoans. More recent reassessment of mesozoan species indicates that they are in fact bilaterian metazoans that have undergone extreme simplification from a more complex ancestor (Dodson, 1956). *In situ* hybridizations for 16 diverse genes in different life stages of *Dicyema japonicum* suggested the presence of multiple different cell types, providing further support for the idea of a complex ancestor of Dicyemida, followed by extreme simplification of body organization (Ogino *et al.*, 2011).

Two significant contributions for understanding the evolutionary history of the Mesozoa are the recent publication of the nuclear genome of *Intoshia linei* (Mikhailov *et al.*, 2016), which represents one of approximately 20 species of this genus; and a transcriptome of *Dicyema japonicum* (Lu *et al.*, 2017). The genomic sequence of *I. linei* is 43 Mbp in length and encodes just ~9000 genes, including those essential for the development and activity of muscular and nervous systems. Neither a phylogenomic analysis based on 500 orthologous groups nor an analysis of transcriptomic data from *D. japonicum* and a dataset compiled from 29 taxa and >300 gene orthologues, could confidently place the Orthonectida or Dicyemida in a precise position within the Lophotrochozoa. However, analysis by Lu *et al.* (2017) found strong statistical support for a grouping of the Dicyemida with the Orthonectida in the phylum Mesozoa. A new and taxonomically broader phylogenomic analysis, however, found *Intoshia* to be nested within the annelids (Schiffer *et al.*, 2018). While the

© Cambridge University Press 2018. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

CAMBRIDGE
UNIVERSITY PRESS

position of *Dicyema* within the Lophotrochozoa could not be unambiguously resolved, it seems clear that Orthonectids and Dicyemida/Rhombzoa are not joined in a single taxon. Nevertheless, both of these groups can be regarded as examples of the extreme morphological and genomic simplification often found in parasites.

One resource that has not been extensively investigated to study the evolution and biology of the mesozoans is their mitochondrial genomes. Interestingly, the limited mitochondrial data analysed from Dicyemida to date suggest a highly unusual mitochondrial gene structure. The three mitochondrial genes that have been sequenced from *D. misakiense* (*cox1*, 2 and 3) appear to be on individual mini-circles of DNA, rather than being part of a typical circular mitochondrial genome (Watanabe *et al.*, 1999; Awata *et al.*, 2005; Catalano *et al.*, 2015). Although the vast majority of metazoan mitochondrial genomes are found as a single circular molecule, a few multipartite circular genomes (that is, genomes where mitochondrial genes are found on more than one closed-circle molecule) have been reported across the Bilateria. The majority of these appear to come from parasitic species and multipartite mitochondrial genomes have been described, for example, from several species of lice (Shao *et al.*, 2005; Cameron *et al.*, 2011; Dong *et al.*, 2014) and parasitic nematodes (Hunt *et al.*, 2016; Phillips *et al.*, 2016). The Dicyemida could, therefore, represent another example of a parasitic organism with a multipartite mitochondrial genome.

Cox1 genes have been deposited in GenBank for a number of Dicyemida species, (*Dicyema koinonum*; *Dicyema acuticephalum*; *Dicyema vincentense*; *Dicyema multimegalum*; *Dicyema coffinense*), but these have not been used in a published molecular systematic study.

In order to provide new mitochondrial gene data from these species, we looked for mitochondrial sequences in publicly available short read sequence data from members of the genus *Dicyema* (*D. japonicum* and *Dicyema* sp.) and Orthonectida (*Intoshia linei*). We compared features of the mitochondrial genome of *I. linei* to other metazoan members to shed additional light on its possible rapid and extreme simplification and used mitochondrial protein-coding gene data from *Dicyema* sp. and *D. japonicum* to investigate the internal phylogeny of the dicyemids.

Materials and methods

Genome and transcriptome assemblies

Genomic (*I. linei*: SRR4418796, SR4418797) and transcriptomic (*Dicyema* sp.: SRR827581; *D. japonicum*: DRR057371) data were downloaded from the NCBI short read archive. Adapter sequences and low-quality bases were removed from the sequencing reads using Trimmomatic (Bolger *et al.*, 2014). The *I. linei* genome was re-assembled using the CLC assembly cell (v.5.0 <https://www.qiagenbioinformatics.com/>). The CLC assembly cell (v.5.0) and Trinity pipeline (Haas *et al.*, 2013) (v.2.3.2) were used to assemble the *Dicyema* sp. and *D. japonicum* transcriptomes, with default settings.

Mitochondrial genome fragment identification and annotation

Mitochondrial protein-coding gene sequences from flatworms were used as tblastn queries to search for mitochondrial fragments in the *I. linei* genome assembly and *Dicyema* sp. and *D. japonicum* Trinity RNA-Seq assemblies, using NCBI translation table 5 'invertebrate mitochondrial'. Positively identified sequences were blasted against the NCBI nucleotide database in order to detect possible contaminant sequences from *Dicyema* host species. For each *Dicyema* sp. gene-bearing contig, a number of other contigs were found that had very high sequence similarity

to *Octopus* or other cephalopods, indicating a degree of host species contamination in the RNA-Seq data. These sequences were discarded from the subsequent analysis.

A 14 176 bp mitochondrial contig was identified from initial blast queries against the reassembled *I. linei* genome. An additional mitochondrial contig was found that overlapped with this 14 176 bp contig to extend the mitochondrial sequence of *I. linei* to 14 247 bp long. This final contig was annotated using MITOS (Bernt *et al.*, 2013). The locations of MITOS-predicted protein-coding genes were manually verified by aligning to orthologous protein-coding gene sequences taken from the published mitochondrial genomes of lophotrochozoan taxa. Where possible, the locations of protein-coding genes were inferred to start from the first in-frame start codon (ATN) and the C-terminal of the protein-coding genes inferred to be the first in-frame stop codon (TAA, TAG or TGA). Contigs from *Dicyema* sp. and *D. japonicum* identified as containing mitochondrial protein-coding genes were verified and annotated in the same way.

The secondary structure of tRNAs identified in the *I. linei* mitochondrial genome were inferred using the Mitfi program within MITOS. MITOS was also used to screen the *Dicyema* sp. and *D. japonicum* contigs which had been positively-identified as containing mitochondrial genes using blast. Using MITOS, we identified one reliable tRNA sequence for trnV on the same contig as *cox3* from *Dicyema* sp. The secondary structure of this tRNA was also inferred using Mitfi in MITOS.

Dicyema internal phylogeny

cox1 nucleotide sequences from *Dicyema* sp. and *D. japonicum* were aligned to publicly available *cox1* nucleotide sequences from other *Dicyema* species, and outgroup sequences from diverse lophotrochozoan taxa (see Supplementary Table S2). Sequences were aligned using Muscle v.3.8.31 (Edgar, 2004) visualized in Mesquite v.3.31 (<http://mesquiteproject.org>) and trimmed to remove uninformative residues with trimal v1.4.rev15. Maximum Likelihood inference was carried out for two trimmed alignments, one with all species, one with only the *Dicyema* species, using IQ-Tree (v.1.6.1) (Nguyen *et al.*, 2015), letting the implemented model testing (ModelFinder (Kalyaanamoorthy *et al.*, 2017)) pick the best-fitting phylogenetic model (TVM + I + G4 and TIM + F + I + G4, respectively), and performing 1000 bootstrap replicates (UFBoot (Hoang *et al.*, 2018)). The trees were visualized using Seaview v.4.4.2 (Gouy *et al.*, 2010) and annotated with Inkscape.

Data availability

All mitochondrial genome data presented in this study have been submitted to NCBI GenBank, under accession numbers: *Intoshia linei* mitochondrial genome sequence, MG839537; *Dicyema* sp. genes, MG839520-MG839528; *D. japonicum* genes, MG839529-MG839536.

Results

Intoshia linei mitochondrial genome composition

We assembled an *I. linei* mitochondrial genome sequence of 14 247 base pairs length but were unable to close the circular genome with paired-end reads. This could be attributed to the missing sequence being an AT-rich repetitive region, making it difficult to resolve. Furthermore, the *I. linei* mitochondrial genome we report is very AT-rich (83.40% AT).

Using MITOS and manual verification, we were able to predict the full reading frames of 12 protein-coding genes, 20 tRNAs and the small subunit rRNA (*rrnS*). No sequences resembling *atp8*,

trnQ or *trnR* could be found in the final sequence. It is possible that *rrnL* is found between *trnC* and *trnM* based on a weak prediction using MITOS, but this could not be confirmed by aligning to known *rrnL* sequences. Genes in the *I. linei* mitochondrial genome are found in two blocks of opposing transcriptional polarity: those in the first 3104 base pairs are found on the reverse strand (*trnK-trnV-trnT-trnY-nad4-trnS2-cox1-trnF*); all other genes are found on the forward strand, where the forward strand is defined as that containing a greater portion of the protein-coding sequences (Fig. 1, Table 1). All protein-coding genes have standard initiation codons (ATA × 8; ATG × 1; and ATT × 3). Ten of the protein-coding genes have a standard termination codon (TAA). *nad5* and *nad6* appear to have a truncated stop codon (T- and TA-, respectively) (Table 1).

Protein-coding genes account for 70.9% of the 14 247 base pair long sequence (allowing for overlap between genes); tRNAs 8.13%; rRNAs (including the uncertain prediction for *rrnL*) 13.83%; and non-coding DNA 6.85%. Four regions of non-coding sequence greater than 100 base pairs are found in the *I. linei* genome: a 143 bp-long region between *trnS2* and *cox1*; 230 bp between *trnF* and *nad1* (where the two genes are found on opposite strands); 112 bp between *trnW* and *nad3*; and 112 bp following the *nad5* at the end of the genomic sequence. There is very little overlap between coding sequences: *rrnS* and the best prediction for the *trnM* location overlap by 48 nucleotides, and there are eight incidences of overlap of coding sequences of fewer than 10 nucleotides across the sequence.

In total 20 out of 22 typical tRNAs were identified in the *I. linei* mitochondrial genome (see Supplementary Fig. S3). All predicted tRNAs have an amino-acyl acceptor stem composed of seven or eight base pairs and an anticodon stem composed of four or five base pairs, with the exception of *trnA* and *trnV*, which appear to have truncated acceptor stems. The structure of the DHU arm is, for the most part, consistent with standard tRNA secondary structure: 14 of the predicted tRNAs have a typical four or five base pair DHU stem; four tRNAs have a truncated or modified DHU arm (*trnI*, *trnL1*, *trnL2* and *trnP*); and two tRNAs appear to have lost the DHU arm entirely (*trnS1* and *trnS2*). More unusually, the TC arm in almost all of the predicted tRNAs is either truncated or replaced by a TV loop. The only tRNA found to have a 'typical' TC arm structure is *trnS2*.

Gene order

The protein-coding and rRNA gene order of *I. linei* was compared with a number of other metazoan taxa using CREx (Bernt *et al.*, 2007). Our analysis included representatives from across the Lophotrochozoa, Ecdysozoa and Deuterostomia (see Supplementary Table S1). CREx analysis aims to identify common gene intervals between different mitochondrial genomes, and infer the reversals, transpositions and reverse transpositions required to obtain an observed gene order from the mitochondrial gene orders of other species.

Compared with the other taxa included for analysis, *I. linei* has a highly divergent mitochondrial gene order. The species with the most similar gene order was the carmine spider mite *Tetranychus cinnabarinus* (Arthropoda; Chelicerata) (Fig. 2). However, the similarity in gene order between these genomes is low, with the two sharing just four short gene 'blocks' and with a degree of variation in gene order even within these conserved regions. Of the lophotrochozoan species included for analysis, the species with the highest similarity to *I. linei* was found to be the nemertean *Paranemertes peregrina* (Fig. 2). Both species share the common arrangement of *nad1-nad6-cob*, and the adjacency of *rrnL-rrnS-cox2-atp6* and *nad4-nad5*, but with variation in the order of these two blocks. Overall, conservation of gene order

between *I. linei* and *P. peregrina* is very low given the number of possible shared gene boundaries. It is clear that the gene order of *I. linei* is novel and very divergent compared with other published metazoan mitochondrial genomes.

Dicyema sp. and *D. japonicum* mitochondrial genes

Using BLAST and manual sequence verification of contigs assembled using Trinity (Haas *et al.*, 2013) and the CLC assembly cell, we were able to identify nine reconstructed mitochondrial transcripts containing protein-coding genes from *Dicyema* sp. (*cox1*, 2, 3; *cob*; and *nad1*, 2, 3, 4 and 5). In *D. japonicum* we found contigs for eight mitochondrial transcripts of protein-coding genes (*cox1*, 2, 3; *nad1*, 3, 4 and 5; and *cob*). The identification of *cob* and *nad3*, *nad4* and *nad5* are novel for this taxon. All of the complete Dicyemida protein-coding genes identified have full initiation and termination codons, with the exception of *D. japonicum nad1*, which has a truncated TA stop codon.

All mitochondrial protein-coding genes found for *Dicyema* sp. and *D. japonicum* were located on individual contigs. In no instance were two or more protein-coding genes found on the same reconstructed transcript. However, each reconstructed mitochondrial transcript did contain non-coding sequence in addition to protein-coding gene sequence (Fig. 3). The length and location (5' and/or 3') of non-coding sequence in the reconstructed transcripts was variable between the two species and between genes (Fig. 3). For each protein-coding gene, we compared contigs assembled using both the CLC-assembler and Trinity in order to identify reconstructed mitochondrial transcripts with the longest stretch of protein-coding sequence. Of the 17 dicyemid genes we report, seven were derived from CLC-assembled contigs (*cox1*, 2, 3, *nad3*, 5 from *D. japonicum* and *nad2*, *nad5* from *Dicyema* sp.) and ten from Trinity-assembled contigs (*cob*, *nad1*, 4 from *D. japonicum* and *cox1*, 2, 3, *cob*, *nad1*, 3, 4 from *Dicyema* sp.). The best reconstructed mitochondrial contig we identified for *nad1* (*Dicyema* sp.) was found to contain duplicated stretches of the identical protein-coding gene sequence. We attribute this to an assembly artefact rather than speculating about this providing potential evidence for mitochondrial gene mini-circles. We also identified and corrected a frameshift in the coding sequence for *nad2* from *Dicyema* sp. in the longest reconstructed mitochondrial contig from this species (Fig. 3).

By screening the *Dicyema* sp. gene-containing contigs with MITOS standalone software, we were able to predict one reliable tRNA sequence for *trnV* adjacent to *cox3* (Fig. 3). *Dicyema* sp. *trnV* has an eight base pair acceptor stem; a five base pair anti-codon stem; and a four base pair DHU stem (Supplementary Information S3). As was found in *I. linei*, the TC arm appears to be modified from a standard 'cloverleaf' structure. In no other cases did we find any sequence from more than one gene on a single contig.

Dicyemida internal phylogeny using *cox1*

Cox1 is commonly used as a species 'barcoding' gene and can be used in phylogenetic inference to discriminate between closely related species (Hebert *et al.*, 2003). Given that a number of *cox1* genes have been sequenced from different Dicyemida species, we used publicly available *cox1* sequences along with the two new *cox1* sequences found in this study for phylogenetic inference to determine the relationship of *Dicyema* sp. and *D. japonicum* to other dicyemids. After aligning the *cox1* nucleotide sequences from dicyemids to other metazoans, we found that all Dicyemida *cox1* sequences had several conserved deletions, not present in other metazoan *cox1* sequences included in our alignment (Fig. 4). These comprise in-frame deletions of two, five, four and two amino acids moving from the N-terminus to the C-terminus of the protein. These deletions appear to be unique




Fig. 1. Overview of the mitochondrial sequence resolved for *Intoshia linei*. Genes not drawn to scale. Numbers beneath the sequences show intergenic spaces (positive values) or intergenic overlap (negative values). Protein-coding genes are denoted by three letter abbreviations; ribosomal genes by four-letter abbreviations. tRNAs are shown by single uppercase letters (for recognized codons of L1, L2, S1 and S2 see Table 1). Genes found on the negative (reverse) strand are coloured green; genes found on the positive (forward) strand are coloured blue. The unreliable prediction for *rrnS* is shown in orange.

Table 1. Organization of the *I. linei* mitochondrial genome. Uncertain position of *rrnL* shown in brackets. Recognized codons for tRNAs L1, L2, S1 and S2 indicated in brackets

Feature	Strand	Start	Stop	Length (nucleotides)	Length (AA)	Start codon	Stop codon	Intergenic region
<i>trnK</i>	-	99	157	59				0
<i>trnV</i>	-	158	209	52				2
<i>trnT</i>	-	212	271	60				-2
<i>trnY</i>	-	270	330	61				-3
<i>nad4</i>	-	328	1509	1182	394	ATA	TAA	70
<i>trnS2</i> (TGA)	-	1580	1637	58				143
<i>cox1</i>	-	1781	3175	1395	465	ATA	TAA	1
<i>trnF</i>	-	3177	3237	61				230
<i>nad1</i>	+	3468	4358	891	297	ATG	TAA	60
<i>trnE</i>	+	4419	4470	52				8
<i>trnP</i>	+	4479	4539	61				48
<i>nad6</i>	+	4588	5028	441	147	ATT	TA-	9
<i>trnL2</i> (TAA)	+	5038	5098	61				47
<i>cob</i>	+	5146	6168	1023	341	ATA	TAA	-1
<i>trnW</i>	+	6168	6224	57				112
<i>nad3</i>	+	6337	6687	351	117	ATA	TAA	-1
<i>trnI</i>	+	6687	6749	63				0
<i>trnC</i>	+	6750	6804	55				-1
(<i>rrnL</i>)	+	6804	8031	1228				1
<i>trnM</i>	+	8033	8096	64				-48
<i>rrnS</i>	+	8049	8790	742				-9
<i>trnL1</i> (TAG)	+	8782	8838	57				0
<i>trnD</i>	+	8839	8894	56				68
<i>cox2</i>	+	8963	9580	618	206	ATA	TAA	4
<i>trnN</i>	+	9585	9649	65				1
<i>trnH</i>	+	9651	9705	55				2
<i>atp6</i>	+	9708	10 394	687	229	ATA	TAA	7
<i>nad4l</i>	+	10 402	10 677	276	92	ATT	TAA	-1
<i>trnS1</i> (TCT)	+	10 677	10 728	52				0
<i>trnA</i>	+	10 729	10 781	53				0
<i>trnG</i>	+	10 782	10 837	56				-4
<i>cox3</i>	+	10 834	11 613	780	260	ATA	TAA	24
<i>nad2</i>	+	11 638	12 507	870	290	ATT	TAA	41
<i>nad5</i>	+	12 549	14 135	1587	529	ATA	T--	

to members of the Dicyemida and were present in all Dicyemida species included in the alignment.

Maximum likelihood phylogenetic analysis was carried out using *cox1* sequences from publicly available *Dicyema* species; the

D. japonicum and *Dicyema* sp. *cox1* sequences assembled in this study; and *cox1* sequences from a diverse representation of lophotrochozoans as outgroup taxa (Supplementary Table S2). As anticipated, the dicyemids included in analysis form their own branch on




Fig. 2. Gene order comparison between the mitochondrial genomes of *Intoshia linei*, the chelicerate *Tetranychus cinnabarinus* and the nemertean *Paranemertes peregrina*, as analysed using CREx. Genes on the minus strand are denoted by a minus sign (-X). Conserved blocks of genes between *I. linei* and *T. cinnabarinus* are shown in orange (*nad1*), blue (*nad6*, *cob*), pink (*nad3*, *rrnL*), green (*rrnS*, *cox2*), yellow (*nad4l*, *cox3*) and red (*nad2*, *nad5*). Variation in gene order and translocation between strands is found within these blocks between the two species. Conserved blocks of genes between *I. linei* and *P. peregrina* are shown in grey (*nad1*, *nad6*, *cob*), purple (*rrnL*, *rrnS*, *cox2*, *atp6*) and light blue (*nad4*, *nad5*). As before, variation in gene order and translocation between strands is evident between these common intervals. For both comparisons, solid or dashed-line boxes show larger regions of the genomes that encompass the same genes, but with no conserved gene order.

the tree. The topology of the subtree containing the Dicyemida species and the unrooted tree inferred from Dicyemida sequences alone is identical (Fig. 5). The analyses found a close affinity between *D. japonicum* with *D. misakiense*, with 97% bootstrap support at this node. At the nucleotide level, sequences from *D. japonicum* and *D. misakiense* are ~98% identical. The remaining *Dicyema* species branch into four groups, with the *Dicyema* sp. sequence from our analysis being an outgroup to the other six species. As found for *D. japonicum* and *D. misakiense*, the *cox1* sequences from *D. multimegalum* and *D. coffinense* are ~98% identical at the nucleotide level (Fig. 5).

Discussion

Although the mitochondrial genome we found from *I. linei* is not a closed circular molecule, this represents the first mitochondrial genome from an orthonectid species and includes 12 protein-coding genes, 20 tRNAs and (possibly) both ribosomal RNAs. *Atp8* was not found, but this gene has been lost from the mitochondrial genomes of taxa in many different metazoan lineages (Boore, 1999) and so its absence from our assembly may be real rather than representing missing data. Compared with the drastically reduced *I. linei* nuclear genome, its mitochondrial genome has a gene complement that is fairly standard across the Metazoa (Mikhailov *et al.*, 2016). Genes in the *I. linei* mitochondrial genome are clustered into two blocks of opposite transcriptional polarity. The block comprising *trnK-trnV-trnT-trnY-nad4-trnS2-cox1-trnF* at the 'start' of the sequence is found on the negative strand, whilst all other genes are transcribed from the positive strand, suggesting an inversion event (Fig. 1).

The ~84% A + T content found in the *I. linei* mitochondrial genome is high compared with other invertebrate mitochondrial genomes – for example, the chelicerate *Limulus polyphemus* (A + T content = 67.6%) (Lavrov *et al.*, 2000) and the annelid *Lumbricus terrestris* (A + T content = 61.6%) (Boore and Brown, 1995). The very high A + T content of the mitochondrial

genome is even higher than the high A + T content of the *I. linei* nuclear genome (73%), and provides evidence for the very fast rate of mitochondrial evolution in this species.

The small proportion of non-coding mtDNA we found for *I. linei* is typical of mitochondrial genomes. Although there is very little gene overlap in the sequence found for *I. linei*, the sequence for *trnM* is predicted with significant overlap with *rrnS*. Whilst it is possible that this is a mis-prediction, large gene overlaps have been reported in other mitochondrial genomes and this overlap could result from selection to minimize mitochondrial genome size (Robertson *et al.*, 2017).

Mitochondrial gene order can be a useful tool for inferring phylogenetic relationships. Gene order in the mitochondrial genomes of different lineages are largely stable, with the rearrangement of protein-coding genes occurring relatively infrequently. Where rearrangement events do occur, they are thought to be a result of tandem duplications and multiple random deletions. In this model, a portion of the mitochondrial genome is erroneously duplicated, and the subsequent random loss of one copy of a gene (by deletion or the accumulation of mutations) results in a novel gene order (Boore, 2000). Comparing mitochondrial gene order between different taxa has been informative not only for the study of larger-scale evolutionary lineages (Boore *et al.*, 1998) but also for understanding the phylogeny of, for example, parasitic flatworms (Le *et al.*, 2000; Liu *et al.*, 2013).

Our analysis demonstrated that *I. linei* has a highly divergent mitochondrial gene order in comparison with other published metazoan mitochondrial genomes. Of the species included – chosen as a broad representation of different metazoan lineages (see Supplementary Table S1) – the closest similarity was found to be to the chelicerate *T. cinnabarinus* (Fig. 2). However, of the highest possible conserved gene order score of 204 (that is, when two mitochondrial genomes have identical gene orders for the 12 protein-coding genes included for analysis (*atp8* is not present in *I. linei*) and two rRNAs included in the CREx matrix calculation), similarity of gene order between *I. linei* and *T. cinnabarinus*

Overview of reconstructed mitochondrial contigs




Fig. 3. Overview of the reconstructed mitochondrial transcripts of protein-coding genes found in *Dicyema japonicum* and *Dicyema* sp. Transcripts are not to scale. Regions coloured blue indicate protein-coding sequence; regions coloured white indicate non-coding sequence. Numbers under each transcript correspond to the length of each respective coding or non-coding region, in base pairs. For *cox3* in *Dicyema* sp. the location of the putative *trnV* sequence is shown in green (nucleotides 1130–1200). For *nad2* in *Dicyema* sp., the location of a frameshift at position 463 in the longest coding sequence assembly is denoted by an asterisk (*).




Fig. 4. Conserved deletions in the amino acid sequence of *cox1* taken from publicly available dicyemid sequences (*D. vincentense*, *D. multimegalum*, *D. coffinense*, *D. furuyi*, *D. misakiense*, *D. koinonum*, *D. papuceum*) and the *Dicyema* species presented in this analysis (*D. japonicum* and *Dicyema* sp.), in alignment with other lophotrochozoan *cox1* sequences. The location of the deletions (two amino acids; five amino acids; four amino acids and two amino acids) moving from the N-terminus to the C-terminus of the protein, are shown by red stars. Colours in the alignment correspond to amino acid colours as used by Mesquite v.3.31.




Fig. 5. Phylogenetic analysis of *cox1* genes from dicyemids using Maximum Likelihood (IQ-Tree), including the two *Dicyema* species presented in this analysis (*Dicyema* sp. and *D. japonicum*). *Cox1* sequences from diverse lophotrochozoan taxa included as outgroups (see Supplementary Fig. S4 for a tree based on the *Dicyema* species alone). Bootstrap support is shown at relevant nodes. Phylogenetic inference shows a split of dicyemid species into two groups, one containing *D. japonicum* and *D. misakiense* (green box) and the other containing all other dicyemid species included in analysis (pink box). Within the second group, *Dicyema* sp. is found on a separate branch from the rest of the included dicyemids. Very short branch lengths between both *D. coffinense* and *D. multimegalum* and *D. japonicum* and *D. misakiense*, indicate that they are very closely related or could be identical species (dashed-line boxes).

was still poor, with a score of just 32. In light of the proposed affinity of *I. linei* to the Lophotrochozoa, the highest-scoring similarity to the lophotrochozoan species included in the analysis was with the nemertean *P. peregrina*. Again, these two species share comparatively little gene order conservation: just three common

gene blocks were identified, one of which (*rrnL-rrnS-cox2-atp6*) has further rearrangement therein (Fig. 2). Of the common intervals identified, the block of *nad1-nad6-cob* is conserved between *I. linei* and *P. peregrina* and it is possible that this arrangement is plesiomorphic within the Lophotrochozoa.

Interestingly, previous analysis of five mitochondrial genomes from early branching annelids found that gene order was highly variable between these species (Weigert *et al.*, 2016). Other studies of various lophotrochozoan mitochondrial genomes – including Brachiopoda (*Lingula anatina*) (Luo *et al.*, 2015); various *Schistosoma* species (Webster and Littlewood, 2012); and nemerteans (Podsiadlowski *et al.*, 2009) – also indicate that extensive gene order rearrangements have occurred in different lophotrochozoan lineages. It is possible that the divergent gene order in *I. linei* can be associated with the parasitic lifestyle and rapid rate of evolution seen for this species: studies in a number of other parasitic taxa indicate that an accelerated rate of gene rearrangement in mitochondrial genomes could be associated with this lifestyle. For example, various *Schistosoma* species have mitochondrial genomes with a unique gene order (Le *et al.*, 2000) and this has also been observed in the ectoparasitic louse *Heterodoxus macropus* (Shao *et al.*, 2001), various species of mosquito (Beard *et al.*, 1993) and parasitic hymenopterans (Dowton and Austin, 1999), amongst others.

All of the tRNAs predicted from *I. linei* and the one tRNA found in *Dicyema* sp. have deviations from the ‘standard’ secondary structure of the TC arm (Supplementary Information S3). Although it is unusual amongst typical metazoan mitochondrial genomes to have such consistent modifications to one element of the tRNA cloverleaf structure, a great deal of variation can be found in tRNA structures across the Metazoa. Mitochondrial genomes with almost all tRNAs lacking either the TC arm or DHU arm – termed ‘minimal functional tRNAs’ – have been reported in a number of different lineages. In nematodes, analysis of tRNAs with TV loops in place of a TC arm suggests that an ‘L-shaped’ tRNA, analogous to a typical cloverleaf-structure tRNA – can maintain normal tertiary interactions and remain functional. Furthermore, it is likely that the tRNAs reported in this analysis are functional, as the acceptor stems and anti-codon stems are, for the most part, complete, and it is highly likely that they would have accumulated mutations should they have lost functionality. Instead, the reduction we observe in tRNA secondary structure could be a result of selective pressure to reduce the TC arm and provide another example of minimally functional tRNAs, in addition to those already found across the Metazoa.

Previous analyses had suggested that mitochondrial genes (*cox1*, *cox2* and *cox3*) in dicyemids are found on individual mini-circles in somatic cells, as opposed to being located on a larger circular mitochondrial genome (Watanabe *et al.*, 1999; Catalano *et al.*, 2015). Mitochondrial mini-circles – although rare across the Metazoa – do appear to be most prevalent in parasitic species. A number of studies have reported the presence of mini-circle mitochondrial genomes, fragmented to various degrees, in a number of lice and nematode species (Shao *et al.*, 2009; Cameron *et al.*, 2011; Dong *et al.*, 2014; Hunt *et al.*, 2016; Phillips *et al.*, 2016).

We identified a mitochondrial contig for *nad1* in *Dicyema* sp. that contained repeated *nad1* protein-coding sequence. This could be an assembly artefact resulting from sequencing a circular *nad1* molecule, but the question of the presence of mini-circles remains unresolved in our analysis. Further investigating the validity of mitochondrial mini-circles was outside of the scope of the present study, but future approaches involving long-range polymerase chain reaction or long-read sequencing should be conducted to resolve this question.

All dicyemid *cox1* sequences were found to have a series of four in-frame deletions within a region of the gene that was well-aligned with *cox1* sequences taken from other invertebrate species (Fig. 4). Insertions and deletions (indels) in genes are rare genomic changes that can be used to infer common evolutionary history (Belinky *et al.*, 2010), but this set of conserved deletions are

so far known only from the *Dicyema* genus. No such deletions are found in the *I. linei* *cox1* gene, or in the same protein-coding sequence taken from across the Lophotrochozoa.

The internal phylogeny of dicyemids was inferred using Maximum Likelihood reconstruction based on *cox1* gene sequences. Based on our analysis it is possible that *D. coffinense* and *D. multimegalum* are the same species, with both dicyemids isolated from Australian *Sepia* species (Catalano, 2013) and with a greater than 98% sequence similarity at the nucleotide sequence level. Our analysis also found a very close affinity for *D. japonicum* with *D. misakiense*, with 97% bootstrap support. As found for *D. coffinense* and *D. multimegalum* these *cox1* sequences are >98% identical at the nucleotide level and it is thus likely that they are very closely related or the same species, both found in octopus living in the North West Pacific off the coast of Honshu Island, Japan. The *Dicyema* sp., despite being isolated from *Sepia* living off the coast of Florida, is a close sister to *D. coffinense* and *D. multimegalum* based on the similarities of the *cox 1* assembled in this analysis, hinting at long-range dispersal with the host species. Further investigation into dicyemid members isolated from hosts in other geographical locations could help to inform whether the phylogenetic structure of the parasitic dicyemids is reflective of dispersal of the host octopus (Tobias *et al.*, 2017).

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/pao.2018.12>.

Acknowledgements. We thank members of the Telford and Oliveri labs at UCL for helpful comments on the analyses and manuscript.

Financial support. The research was funded by a European Research Council grant (ERC-2012-AdG 322790) to MJT.

Conflicts of interest. None.

Ethical standards. Not applicable.

References

- Awata H, Noto T and Endoh H (2005) Differentiation of somatic mitochondria and the structural changes in mtDNA during development of the dicyemid *Dicyema japonicum* (Mesozoa). *Molecular Genetics and Genomics* **273**, 441–449.
- Beard CB, Hamm DM and Collins FH (1993) The mitochondrial genome of the mosquito *Anopheles gambiae*: DNA sequence, genome organization, and comparisons with mitochondrial sequences of other insects. *Insect Molecular Biology* **2**, 103–124.
- Belinky F, Cohen O and Huchon D (2010) Large-scale parsimony analysis of metazoan indels in protein-coding genes. *Molecular Biology and Evolution* **27**, 441–451.
- Bernt M, Merkle D, Ramsch K, Fritsch G, Perseke M, Bernhard D, Schlegel M, Stadler PF and Middendorf M (2007) CREx: inferring genomic rearrangements based on common intervals. *Bioinformatics (Oxford, England)* **23**, 2957–2958.
- Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritsch G, Pütz J, Middendorf M and Stadler PF (2013) MITOS: improved de novo metazoan mitochondrial genome annotation. *Molecular Phylogenetics and Evolution* **69**, 313–319.
- Bolger AM, Lohse M and Usadel B (2014) Trimmomatic: a flexible trimmer for illumina sequence data. *Bioinformatics (Oxford, England)* **30**, 2114–2120.
- Boore JL (1999) Animal mitochondrial genomes. *Nucleic Acids Research* **27**, 1767–1780.
- Boore JL (2000) The duplication/random loss model for gene rearrangement exemplified by mitochondrial genomes of deuterostome animals. In Sankoff D and Nadeau JH (eds), *Comparative Genomics: Empirical and Analytical Approaches to Gene Order Dynamics, Map Alignment and the Evolution of Gene Families*, Netherlands, Dordrecht: Springer, pp. 133–147.
- Boore JL and Brown WM (1995) Complete sequence of the mitochondrial DNA of the annelid worm *Lumbricus terrestris*. *Genetic* **141**, 305–319.

- Boore JL, Lavrov DV and Brown WM** (1998) Gene translocation links insects and crustaceans. *Nature* **392**, 667–668.
- Cameron SL, Yoshizawa K, Mizukoshi A, Whiting MF and Johnson KP** (2011) Mitochondrial genome deletions and minicircles are common in lice (Insecta: Phthiraptera). *BMC Genomics* **12**, 394–394.
- Catalano SR** (2013) Five new species of dicyemid mesozoans (Dicyemida: Dicyemidae) from two Australian cuttlefish species, with comments on dicyemid fauna composition. *Systematic Parasitology* **86**, 125–151.
- Catalano SR, Whittington ID, Donnellan SC, Bertozzi T and Gillanders BM** (2015) First comparative insight into the architecture of COI mitochondrial minicircle molecules of dicyemids reveals marked inter-species variation. *Parasitology* **142**, 1066–1079.
- Dodson EO** (1956) A note on the systematic position of the Mesozoa. *Systematic Zoology* **5**, 37–40.
- Dong W-G, Song S, Guo X-G, Jin D-C, Yang Q, Barker SC and Shao R** (2014) Fragmented mitochondrial genomes are present in both major clades of the blood-sucking lice (suborder Anoplura): evidence from two Hoplopleura rodent lice (family Hoplopleuridae). *BMC Genomics* **15**, 751.
- Dowton M and Austin AD** (1999) Evolutionary dynamics of a mitochondrial rearrangement ‘hotspot’ in the Hymenoptera. *Molecular Biology and Evolution* **16**, 298–309.
- Edgar RC** (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* **32**, 1792–1797.
- Furuya H and Tsuneki K** (2003) Biology of dicyemid mesozoans. *Zoological Science* **20**, 519–532.
- Gouy M, Guindon S and Gascuel O** (2010) Seaview version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Molecular Biology and Evolution* **27**, 221–224.
- Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD and Bowden J** (2013) De novo transcript sequence reconstruction from RNA-seq using the trinity platform for reference generation and analysis. *Nature Protocols* **8**, 1494–1512.
- Hebert PDN, Ratnasingham S and deWaard JR** (2003) Barcoding animal life: cytochrome c oxidase subunit I divergences among closely related species. *Proceedings of the Royal Society B: Biological Sciences* **270**, S96–S99.
- Hoang DT, Chernomor O, von Haeseler A, Minh BQ and Vinh LS** (2018) UFBoot2: improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* **35**, 518–522.
- Hunt VL, Tsai IJ, Coghlan A, Reid AJ, Holroyd N, Foth BJ, Tracey A, Cotton JA, Stanley EJ, Beasley H, Bennett HM, Brooks K, Harsha B, Kajitani R, Kulkarni A, Harbecke D, Nagayasu E, Nichol S, Ogura Y, Quail MA, Randle N, Xia D, Brattig NW, Soblik H, Ribeiro DM, Sanchez-Flores A, Hayashi T, Itoh T, Denver DR, Grant W, Stoltzfus JD, Lok JB, Murayama H, Wastling J, Streit A, Kikuchi T, Viney M and Berriman M** (2016) The genomic basis of parasitism in the Strongyloides clade of nematodes. *Nature Genetics* **48**, 299–307.
- Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A and Jermin LS** (2017) ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods* **14**, 587.
- Lavrov DV, Boore JL and Brown WM** (2000) The complete mitochondrial DNA sequence of the horseshoe crab *Limulus polyphemus*. *Molecular Biology and Evolution* **17**, 813–824.
- Le TH, Blair D, Agatsuma T, Humair P-F, Campbell NJH, Iwagami M, Littlewood DTJ, Peacock B, Johnston DA, Bartley J, Rollinson D, Herniou EA, Zarlenga DS and McManus DP** (2000) Phylogenies inferred from mitochondrial gene orders—A cautionary tale from the parasitic flatworms. *Molecular Biology and Evolution* **17**, 1123–1125.
- Liu G-H, Shao R, Li D-H, Li H and Zhu X-Q** (2013) The complete mitochondrial genomes of three parasitic nematodes of birds: a unique gene order and insights into nematode phylogeny. *BMC Genomics* **14**, 414.
- Lu T-M, Kanda M, Satoh N and Furuya H** (2017) The phylogenetic position of dicyemid mesozoans offers insights into spiralian evolution. *Zoological Letters* **3**, 6.
- Luo Y-J, Satoh N and Endo K** (2015) Mitochondrial gene order variation in the brachiopod *Lingula anatina* and its implications for mitochondrial evolution in lophotrochozoans. *Marine Genomics* **24**, Part 1, 31–40.
- Margulis L and Chapman MJ** (2009) Chapter three – Animalia. In *Kingdoms and Domains*. London: Academic Press, pp. 231–377.
- Mikhailov KV, Slyusarev GS, Nikitin MA, Logacheva MD, Penin AA and Aleoshin VV** (2016) The genome of *Intoshia linei* affirms orthonectids as highly simplified spiralian. *Current Biology* **26**, 1768–1774.
- Nguyen L-T, Schmidt HA, von Haeseler A and Minh BQ** (2015) IQ-TREE: a fast and effective stochastic algorithm for estimating Maximum-likelihood phylogenies. *Molecular Biology and Evolution* **32**, 268–274.
- Ogino K, Tsuneki K and Furuya H** (2011) Distinction of cell types in *Dicyema japonicum* (phylum Dicyemida) by expression patterns of 16 genes. *Journal of Parasitology* **97**, 596–601.
- Phillips WS, Brown AMV, Howe DK, Peetz AB, Blok VC, Denver DR and Zasada IA** (2016) The mitochondrial genome of *Globodera ellingtonae* is composed of two circles with segregated gene content and differential copy numbers. *BMC Genomics* **17**, 706.
- Podsiadlowski L, Braband A, Struck TH, von Döhren J and Bartolomaeus T** (2009) Phylogeny and mitochondrial gene order variation in Lophotrochozoa in the light of new mitogenomic data from Nemertea. *BMC Genomics* **10**, 364.
- Robertson HE, Lapraz F, Egger B, Telford MJ and Schiffer PH** (2017) The mitochondrial genomes of the acoelomorph worms *Paratomella rubra*, *Isodiametra pulchra* and *Archaphanostoma ylvae*. *Scientific Reports* **7**, 1847.
- Schiffer P, Robertson H and Telford MJ** (2018) Orthonectids are highly degenerate annelid worms. *Current Biology* **28**, 1970–1974.
- Shao R, Campbell NJH and Barker SC** (2001) Numerous gene rearrangements in the mitochondrial genome of the Wallaby Louse, *Heterodoxus macropus* (Phthiraptera). *Molecular Biology and Evolution* **18**, 858–865.
- Shao R, Barker SC, Mitani H, Aoki Y and Fukunaga M** (2005) Evolution of duplicate control regions in the mitochondrial genomes of Metazoa: a case study with Australian Ixodes ticks. *Molecular Biology and Evolution* **22**, 620–629.
- Shao R, Kirkness EF and Barker SC** (2009) The single mitochondrial chromosome typical of animals has evolved into 18 minichromosomes in the human body louse, *Pediculus humanus*. *Genome Research* **19**, 904–912.
- Slyusarev GS and Starunov VV** (2016) The structure of the muscular and nervous systems of the female *Intoshia linei* (Orthonectida). *Organisms Diversity & Evolution* **16**, 65–71.
- Tobias ZJC, Yadav AK, Schmidt-Rhaesa A and Poulin R** (2017) Intra- and interspecific genetic diversity of New Zealand hairworms (Nematomorpha). *Parasitology* **144**, 1026–1040.
- Watanabe KI, Bessho Y, Kawasaki M and Hori H** (1999) Mitochondrial genes are found on minicircle DNA molecules in the mesozoan animal *Dicyema*. *Journal of Molecular Biology* **286**, 645–650.
- Webster BL and Littlewood DTJ** (2012) Mitochondrial gene order change in *Schistosoma* (Platyhelminthes: Digenea: Schistosomatidae). *International Journal for Parasitology* **42**, 313–321.
- Weigert A, Golombek A, Gerth M, Schwarz F, Struck TH and Bleidorn C** (2016) Evolution of mitochondrial gene order in Annelida. *Molecular Phylogenetics and Evolution* **94**, 196–206.