

The antiseptic and antipruritic principle in
WRIGHT'S COAL TAR SOAP

Somewhere about the middle of last century a determined effort was made to isolate the known antiseptic properties of Coal Tar from the inert residuum of substances lacking therapeutic value. This was achieved in 1862 in Wright's Liquor Carbonis Detergens and this preparation has been used and recommended for 80 years by skin specialists and specified repeatedly in authoritative reference works.



*** WRIGHT'S LIQUOR CARBONIS DETERGENS**

It followed that the inclusion of this Liquor Carbonis Detergens in the basic formula of Wright's Coal Tar Soap gave to this soap a character as individual as its own healthful aroma. It followed also that this powerful antiseptic principle could be—and has been—incorporated without the least risk of harshness to the skin. Wright's indeed has been recognised for generations as a supreme toilet and nursery soap.



Wright's
COAL TAR SOAP

CONTENTS

(All rights reserved)

	PAGE
KNOX, R., GELL, P. G. H. and POLLOCK, M. R. The selective action of tetrathionate in bacteriological media. A report to the Medical Research Council. (With Plate 2 and 1 Figure in the Text)	147
MURRAY, J. F. Diphtheria amongst the Bantu	159
MURRAY, J. F. Bantu immunity to scarlet-fever toxin	170
SHATTOCK, P. M. F. and MATTICK, A. T. R. The serological grouping of <i>Streptococcus lactis</i> (group N) and its relationship to <i>Streptococcus faecalis</i>	173
VAN DEN ENDE, M. An apparatus for the safe inoculation of animals with dangerous pathogens. With an Appendix by A. J. G. Hubbard. (With Plate 3 and 3 Figures in the Text)	189
EDWARD, D. G. ff. and LIDWELL, O. M. Studies on air-borne virus infections. III. The killing of aerial suspensions of influenza virus by hypochlorous acid. (With 1 Figure in the Text)	196
PIJPER, ADRIANUS and CROCKER, CLARICE G. Typhoid carriers and Vi agglutinins	201
HITCH, SURGEON CAPTAIN F. G., ASHCROFT, SURGEON CAPTAIN L. S. and GREEN, SURGEON COMMANDER C. A. Antigenicity of T.A.B.C. vaccine after admixture with tetanus toxoid for various periods. (With 1 Figure in the Text)	207
BARRITT, MAXWELL M. Notes on the α -naphthol modification of the Voges-Proskauer test and reference to the creatine modification	214
LOUREIRO, J. A. DE. A new principle for the production of thiamin-deficient diets and for the biological assay of thiamin in foodstuffs	217

The Journal of Hygiene is now issued irregularly. A volume is issued in paper covers at the following net prices:

Volumes I-III (1901-3). 17s. 6d. net per volume.

Volumes IV-XVI (1904-18), *excepting* Vol. VI, no. 4, Plague number. 25s. net per volume.

Volume XVII (1918). 31s. net.

Volumes XVIII-XXXI (1919-31). 42s. net per volume.

Volumes XXXII-XXXVII (1932-37). 47s. 6d. net per volume.

Volumes XXXVIII-XLII (1938-42). 75s. net per volume.

Quotations can be given for buckram binding cases, also for bound copies of back volumes.

Papers for publication should be sent to Dr G. S. GRAHAM-SMITH, Dept. of Pathology, Tennis Court Road, Cambridge. Other communications should be addressed to the University Press, Cambridge.

Papers forwarded to the Editor for publication are understood to be offered to *The Journal of Hygiene* alone, unless the contrary is stated.

Contributors receive twenty copies of their papers free. Twenty-five additional copies may be had at cost price: these should be ordered when the final proof is returned.

The subscription price is £3. 15s. 0d. per volume (post-free), payable in advance; single numbers 16s. 0d. net. Subscriptions may be sent to any Bookseller, or to The Cambridge University Press, Bentley House, 200 Euston Road, London, N.W. 1.

The Cambridge University Press has appointed the University of Chicago Press agents for the sale of *The Journal of Hygiene* in the United States of America.