



Review Article

Cite this article: O'Mahony S, Collins N, Gibney ER, and Doyle G (2025). Supermarkets, do they make the healthy choice the easy choice? A review of the healthfulness of the supermarket food environment. *Proceedings of the Nutrition Society*, page 1 of 11. doi: [10.1017/S0029665125000023](https://doi.org/10.1017/S0029665125000023)

Received: 14 June 2024
Revised: 29 October 2024
Accepted: 15 November 2024

Keywords:

Supermarkets; Retail food environment; Obesogenic environment; Diet-related health outcomes; Food-related ill-health

Corresponding author:

Sinead O'Mahony;
Emails: somahony@fsai.ie;
sinead.omahony@ucdconnect.ie

Supermarkets, do they make the healthy choice the easy choice? A review of the healthfulness of the supermarket food environment

Sinead O'Mahony^{1,2}, Nuala Collins¹, Eileen R. Gibney^{2,3} and Gerardine Doyle^{4,5}

¹Food Safety Authority of Ireland, Dublin 1, Ireland; ²Institute of Food and Health, School of Agriculture and Food Science, University College Dublin, Dublin 4, Ireland; ³School of Agriculture and Food Science, University College Dublin, Dublin 4, Ireland; ⁴College of Business, University College Dublin, Dublin 4, Ireland and ⁵UCD Geary Institute of Public Policy, University College Dublin, Dublin 4, Ireland

Abstract

The over consumption of high fat, sugar, and salt foods increases population risk of overweight, obesity and diet-related noncommunicable diseases. The food environment mediates consumer food choices and thus plays an important role in diet quality and related health outcomes. The built food environment, where most people in high-income countries access their food, has been found to be obesogenic. The aim of this review was to investigate the healthfulness of the supermarket food environment. Supermarkets are an important source of healthy foods in the built food environment. However, there are disparities in access to supermarkets, and in several countries, supermarkets located in areas of higher deprivation have an unhealthier consumer food environment. This double burden limits access to healthy foods amongst lower socio-economic groups, contributing to widening disparities in food-related ill health. There is a strong body of evidence supporting improved purchase of healthy foods by increasing the healthfulness of the supermarket consumer food environment. Voluntary measures co-designed with retailers to improve the healthfulness of the supermarket consumer food environment through restriction of product placement and private label reformulation have led to an increase in healthier food purchases. However, evidence also shows that mandatory, structural changes are most effective for improving disparities in the access to healthy food. Future research and policy related to the food environment should consider equitable access to healthy sustainable foods in built and online supermarkets.

A healthy and nutritious diet has long been recognised as an important component of protecting population health⁽¹⁾. Neufeld *et al.* define a healthy diet as 'health-promoting and disease-preventing. It provides adequacy, without excess, of nutrients and health-promoting substances from nutritious foods and avoids the consumption of health-harming substances'⁽²⁾. Food-based dietary guidelines (FBDGs) outline the composition of a healthy diet by grouping foods based on nutritional and health-promoting characteristics⁽³⁾. However, there is evidence that compliance with recommendations in FBDGs is low, and Leme *et al.* found almost 40 % of people living in low- and high-income countries do not adhere to FBDGs with many exceeding intakes of discretionary high fat, sugar and salt (HFSS) foods⁽⁴⁾.

The over consumption of HFSS foods increases the risk of overweight, obesity and diet-related noncommunicable diseases (NCDs), including cardiovascular disease (CVD), type 2 diabetes, and some cancers^(3,5–7). A review of the global burden of diet-related disease identified diets high in salt, trans fats, and sugar-sweetened beverages and low in wholegrains and fresh fruit and vegetables as major risk factors for the development of diet-related NCDs⁽⁸⁾. Diet-related NCDs are one of the leading causes of death in Europe and this is also true for Ireland where it is estimated 9,000 deaths per year are a result of CVD^(8,9). It is estimated that dietary improvement could prevent one in five deaths globally and a quarter of European deaths from NCDs^(8,10). With many countries globally, including Ireland, experiencing an ageing population, the prevention of diet-related NCDs and their associated risk factors, such as overweight and obesity, is of priority^(11,12). Reducing NCD risk, can enable people to live healthily for longer, increasing quality-adjusted life years⁽⁸⁾. Improvements in diet quality through increasing intakes of fresh fruit, vegetables and wholegrains whilst reducing consumption of HFSS foods, is an effective strategy to reduce population-wide NCD risk and promote healthy aging throughout the life course⁽⁸⁾. The aim of this review was to investigate the healthfulness of the supermarket food environment and its relationship with food choice, purchase and diet-related health outcomes.

Determinants of diet-related health outcomes

The determining factors of diet-related health outcomes are varied and complex, resulting from reciprocal relationships which influence food choice, dietary intake, and subsequent health

© The Author(s), 2025. Published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.






Fig. 1. Childhood overweight: a contextual model and recommendations for future research. Davison & Birch, *Obesity Reviews*, 2: 159-171 2001. Copyright © 2001 by Wiley Publications. Reprinted by permission of Wiley Publications⁽¹⁵⁾.

outcomes. Given this complexity, various models have been developed to describe and explain the determining factors of excess weight accumulation and/or development of diet-related NCDs⁽¹³⁾.

The Socio-Ecological Model (SEM) proposes that individual-level outcomes (such as over consumption of foods/nutrients, as described above) are affected by factors beyond the individual, including social, environmental, cultural and economic conditions⁽¹⁴⁾. Davison and Birch considered the context of food intake and energy balance, outlining a contextual model, focusing on child weight status, using ecological systems theory, outlined in Fig.1⁽¹⁵⁾. This model describes the micro-level factors including individual child characteristics and child risk factors, the meso-level factors including parenting styles and family characteristics, and the macro-level factors including community, demographic and societal characteristics contributing to child weight status, leading to childhood obesity⁽¹⁵⁾. A review of the contribution of each layer within the SEM to the accumulation of excess weight in childhood, found that meso and macro-level determinants such as parental demographic and neighbourhood characteristics made the most substantial contribution to predicting childhood weight accumulation in a population in New Jersey, United States of America (USA)⁽¹⁶⁾.

The importance of meso and macro-level factors in excess weight accumulation is also recognised in the *Foresight Obesity Map*⁽¹⁷⁾. The Obesity Map (Fig.2) was one of the first times the complex nature of factors underlying and driving obesity was explained as a system-wide issue⁽¹⁸⁾. This map is based on systems thinking, which includes well-established approaches in other disciplines, complemented by innovative methods to guide how to consider the bigger picture related to the topic of obesity⁽¹⁹⁾. Systems thinking is based on the idea that real-world phenomena exist within systems of contextually specific dynamic actors and factors. For example, in nutritional sciences, a person's nutritional status is determined by a wide range of factors, e.g. genetics, food

security, disease status and potential for absorption/increase requirements, food preferences, food environment and lifestyle factors⁽²⁰⁾. Together, these determinants and factors make up a complex system. In the *Foresight Obesity Map*, the reciprocal relationships between societal influences, individual psychology, individual activity, activity environment, biology, food product and food consumption are described⁽¹⁷⁾.

Both the SEM and *Foresight Obesity Map* demonstrate how factors, beyond the individual, play an important role in the development of overweight, obesity and diet-related NCDs in the population. Both also show how our living environment, including the food environment, are an important determinant in food choice, consumption and the development of overweight, obesity and diet-related NCDs^(21,22).

The food environment

Evidence supports a relationship between the availability and prominence of food in the food environment, food choice and health outcomes^(23,24). Swinburn *et al.* define the food environment as 'the collective physical, economic, policy, and sociocultural surroundings, opportunities, and conditions that affect people's food and beverage choices and nutritional status'⁽²⁵⁾. A health-enabling food environment has been described as one where the goods, beverages and meals which support dietary intakes in line with population FB DGs are available, affordable and accessible to all⁽²⁶⁾. However, today it is well documented that across the Western world our food environments are now considered 'obesogenic' meaning energy-dense and nutrient-poor foods are readily available whilst the environment is amenable to a sedentary lifestyle⁽²⁷⁻²⁹⁾.

Several frameworks have been developed to describe the relationship between the food environment, food choice and eating behaviours⁽³⁰⁻³³⁾. Glanz *et al.*, described the community nutrition

Foresight

Obesity System Map




Fig. 2. Foresight Obesity Systems Map. Figure sourced from Tackling Obesities: Future Choices Report (2007). Reprinted with permission of Foresight, Government Office for Science¹⁷. Contains public sector information licenced under the Open Government License v3-0 (www.nationalarchives.gov.uk/doc/opengovernment-licence/version/3/).




Fig. 3. Model of community nutrition environments; Glanz et al. Am J Health Promot. 2005 May-Jun;19(5):330-3. Copyright © 2005 by Sage Publications. Reprinted by Permission of Sage Publications³⁰.

environment using the sem to outline the relationship between policy variables, environmental variables, individual variables, and behaviour and how they interact to determine eating patterns⁽³⁰⁾. In this framework, shown in Fig.3, within the environmental variables, Glanz *et al.* describe the community food environment, referring to outlet type, location and availability; the consumer food environment, referring to in-store factors; and finally, the organisational food environment, referring to different settings where people spend significant amounts of time, such as work, school and care environments⁽³⁰⁾. Since the development of this model, much of the empirical food environment research has investigated these domains^(34,35).

In a review by Herforth and Ahmed, the bidirectional relationship between food environments and dietary intakes is described, with dietary intakes aligning with the foods available in the individuals food environment, rather than in line with FBDGs⁽³¹⁾. The authors outline how the availability of foods in the food environment precedes consumption, concluding that '*food cannot be consumed if it is not available at all*'⁽³¹⁾. This conclusion is supported by research across the world, examples of which are described here^(36–38). In the USA, Chen and Yang, using social media, examined the food environment experienced by individuals and how it affects food choice, concluding higher availability of fresh produce in the neighbourhood food environment was associated with increased likelihood of purchasing fresh and healthy foods⁽³⁶⁾. In Kenya, Chege *et al.* found that the diets of urban poor consumers could be improved by increasing the availability of fresh and healthy foods in traditional retail outlets such as mom-and-pop shops and informal restaurants⁽³⁷⁾. In Ireland, Kelly *et al.* found that high fast food outlet density around post-primary schools was associated with decreased odds of post-primary school-aged children consuming fruit (OR 0.79, 95 % CI 0.62, 0.99) and vegetable (OR 0.73, 95 % CI 0.60, 0.88) and increased odds of consuming soft drinks (OR 1.79, 95 % CI 1.30, 2.47) and chips (OR 1.82, 95 % CI 1.17, 2.83) daily⁽³⁸⁾. Furthermore, in a systematic review examining the impact of the consumer and neighbourhood food environment on dietary intake and obesity-related outcomes, Atanasova *et al.* found the availability of and distance to unhealthy food outlets was associated with increased likelihood of fast-food consumption and higher Body Mass Index (BMI) in children across all socio-economic groups and in some adult population groups⁽³⁹⁾. In addition, the availability of and distance to healthy food outlets was associated with improved children's dietary intake and BMI⁽³⁹⁾. In their systematic review and meta-analysis, Pineda *et al.* confirmed this finding, concluding fast-food outlet proximity was positively and significantly associated with obesity⁽²⁴⁾. However, despite this growing evidence, many food environments in high-income countries are not health enabling, meaning they do not align with FBDGs⁽²⁸⁾.

Supermarkets in the retail food environment

Whilst the food environment encompasses all access to food, the retail food environment (RFE) is specifically defined as '*the environment where all food and beverages are purchased by consumers, including food service operations such as restaurants*'⁽⁴⁰⁾. The RFE comprises of many different types of outlets such as restaurants, cafés, takeaways, street food vendors, pubs, convenience stores, grocery stores and supermarkets, and influences the population's food choices and dietary health outcomes^(40–42). In most high-income countries, there is a

dependence on the built food environment, where many retail outlets are located, and natural and cultivated food environments play a minimal role in access to foods, meaning populations are dependent on the RFE to access some if not all of their food⁽⁴³⁾. Globally, supermarkets are a significant actor in the RFE, with an increasing amount of food purchased from supermarkets^(44,45). Over the past three decades, the changing landscape of the RFE, such as a shift away from local supermarkets to larger grocery supermarkets and discount supermarkets, many with both a physical and online presence, as well as an increase in the availability and consumption of foods out of home (e.g. takeaways, cafes and restaurants) has changed food availability and eating habits⁽⁴⁰⁾. Evidence shows that different types of food outlets effect diet-related health outcomes in different ways. Pineda *et al.* found that RFE with high density of supermarkets had an inverse relationship with weight gain, whereas RFE with high density of fast-food outlets had a positive relationship with obesity⁽²⁴⁾. There is growing evidence that supermarkets influence dietary behaviours and because of this, they are increasingly becoming a focus of health-promoting interventions^(23,41). Indeed supermarkets have been identified as food environments with the potential to significantly influence consumer food choices and related health outcomes⁽⁴⁶⁾.

Supermarkets, the community food environment, diet quality and diet-related health outcomes

The community food environment describes the availability of food outlets in a person's surroundings⁽⁴⁷⁾. Food outlet location can be measured using Geographical Information Systems (GIS), which allows the visualisation of food outlet locations and the analysis of food outlet density and proximity using travel times to residential settings, schools and workplaces⁽⁴⁸⁾. Community food environments can be considered health-enabling or health-disabling; the latter is often called 'food swamps' or 'food deserts'^(28,49). Food swamps refer to areas with a high density of fast-food outlets serving energy-dense and nutrient-poor foods⁽⁴⁹⁾ and the tendency for fast-food outlets to be clustered in particular locations, even more so in areas of higher deprivation⁽²⁸⁾. Food deserts on the other hand refer to areas with limited access to food outlets offering fresh foods, such as supermarkets and green-grocers^(29,50). Food deserts have been identified in the United Kingdom (UK) and USA, but not to the same extent across Europe^(49,51,52). There is limited evidence on food swamps and deserts in the Irish context.

As previously discussed, supermarkets have been identified as an important source of healthy food in the built food environment⁽²⁴⁾. Increased availability of supermarkets has been associated with improved diet quality in some areas but decreased diet quality in others, with some concluding the relationship between supermarket density and proximity and diet-related health outcomes is cultural and dependant on the country or regional context⁽⁵³⁾.

Much of the scientific evidence around supermarket availability and diet-related health outcomes is from high-income countries. Overall, it would appear that increasing supermarket density and/or proximity has a positive effect on diet quality and is associated with reduced population obesity prevalence^(54–57). Laraia *et al.* found that pregnant women in the USA living more than four miles away from a supermarket were twice as likely to fall into the lowest diet quality score quartile, than pregnant women living within two miles of a supermarket⁽⁵⁴⁾. In addition, residents within

0.5 miles or less from a new supermarket, opened in the USA under an initiative to improve access to healthful food, had increased availability and consumption of healthy food following the opening of the supermarket⁽⁵⁵⁾. Lamichhane *et al.* found youths living with diabetes in the USA had improved diet quality following increased availability and accessibility to supermarkets⁽⁵⁶⁾. Increased supermarket proximity has also been shown to be associated with reduced frequency of fast-food consumption⁽⁵⁷⁾. The evidence on the relationship between supermarket proximity and obesity prevalence or high BMI is growing. Improved access to a supermarket has been found to be significantly associated with a decrease in BMI and lower population obesity prevalence^(24,58,59). However, some evidence from the USA is less positive, Elbel *et al.* found children aged 3–10 years living in close proximity to a new supermarket, specifically opened to tackle a food desert in New York, had no difference in the availability of healthy or unhealthy foods at home, or in children's dietary intake as a result of the supermarket opening⁽⁶⁰⁾. Furthermore, the clustering of fast-food outlets around supermarkets has been observed, which could diminish the effectiveness of supermarkets in increasing the purchase and consumption of healthy food options⁽⁶¹⁾.

Research outside of the USA is less conclusive^(62–64). In Canada, Stevenson *et al.* found no relationship between the availability of supermarkets and diet quality⁽⁶²⁾. In England, an inconsistent and complex relationship between supermarket availability and diet quality was found in children aged 9–10 years of age. In this cohort, density of supermarkets was associated with an increase in vegetable intake, but also an increase in the consumption of unhealthy foods⁽⁶⁴⁾. In Glasgow, Scotland, the introduction of a supermarket to improve diet and health in a low socio-economic community was associated with no effect on diet quality⁽⁶³⁾. In contrast, in Ireland an investigation into the relationship between availability of supermarkets and diet quality by Layte *et al.* found a significant difference in diet quality mediated by the distance to the closest supermarket and by socio-economic status (SES) groups (using a household measure of SES)⁽⁶⁵⁾.

Whilst proximity and density of supermarkets appears to be an important variable for diet quality and diet-related health outcomes, the effect of supermarket density and proximity is mediated by the healthfulness of the in-store supermarket environment, known as the consumer food environment⁽²⁴⁾.

Supermarkets, the consumer food environment and food purchase

Within outlet food availability has been described as the consumer food environment where the marketing mix: price, promotion, product, and placement, alongside food accessibility, convenience, and desirability, are variables influencing consumer food choice^(30,32). Studies investigating the in-store availability of unhealthy foods in supermarkets have found supermarkets have a high availability of unhealthy or discretionary foods. In Victoria, Australia, 63.2%–72.7% of shelf space was allocated to discretionary (unhealthy) foods, depending on the retailer⁽⁶⁶⁾. A nationwide study in New Zealand found that for every one metre (100 cm) of shelf space there was 58 cm of shelf space allocated to unhealthy food⁽⁶⁷⁾. In Flanders, Belgium, for every one metre (100 cm) of shelf space, 64 cm was allocated to unhealthy foods⁽⁶⁸⁾. A study examining the availability of healthy and unhealthy foods in supermarkets in Dublin, Ireland found that for every one metre (100 cm) of shelf space measured 68 cm was allocated to unhealthy food⁽⁶⁹⁾. This study also found high prominence areas in

supermarkets (locations highly visible to consumers such as check-outs and end of aisle) mostly contained unhealthy foods⁽⁶⁹⁾. These findings indicate that the current in-store supermarket food environment does not support consumers to make food choices in line with FBDGs recommendations. Given the high proportion of supermarket shelf space allocated to unhealthy foods, as well as the high visibility of this shelf space, there have been a number of studies examining interventions to improve the availability and prominence of healthy foods with the aim of increasing healthy food purchases and decreasing unhealthy food purchases.

Food purchase data is proposed as a reliable proxy indicator for food consumption in the adult population⁽⁷⁰⁾. Healthy in-store interventions, which aim to create health-enabling consumer retail environments utilise merchandising strategies targeting product, price, placement, and promotion to incentivise the purchase of healthy foods and disincentive the purchase of unhealthy foods^(71,72). Food pricing is a strategy used to increase sales and can be in the form of discounts, bulk buying offers or promotions⁽⁷³⁾. Adam and Jensen concluded that price promotions positively influenced the purchase of healthy food, and the effect was more significant the larger the price discount⁽⁷⁴⁾. Studies examining the effect of price promotion interventions on fresh fruit and vegetables found that discounts had a progressive and sustained impact in increasing the purchase of fresh fruit and vegetables^(75–77). Product placement is another strategy used by retailers to promote the purchase of healthy foods⁽⁷⁸⁾. Evidence suggests placing foods in more prominent locations increases product sales⁽⁷⁴⁾. A study investigating the placement of healthier food items at the point of sale increased their purchase⁽⁷⁹⁾. A third merchandising strategy to increase healthy food product sales is promotion, such as signage, shelf tags, floor stickers and special island displays. Promotions have been found to be an effective way to increase sales of healthy foods in New Zealand⁽⁸⁰⁾ and in Australia, shelf tags highlighting foods of lower nutrient quality effectively reduced their sale⁽⁸¹⁾. In fact, a combination of merchandising approaches is likely to be most effective in increasing the purchase of healthy foods, rather than any one approach alone. A randomised control trial investigating the effect of restricting unhealthy food promotion on food and beverage sales concluded restricted merchandising of unhealthy food alongside increased merchandising of healthy food can achieve public health gains⁽⁴⁶⁾. A systematic review by Adam and Jensen reported that n 36 of n 42 studies examining the effect of an in-store health-promoting intervention found a positive effect on the purchase of healthy foods⁽⁷⁴⁾. This review concluded that interventions combining merchandising techniques to improve the purchase of healthy foods are likely to be most effective, including interventions that combine price, information, and easy access to and availability of healthy food⁽⁷⁴⁾. Shaw *et al.* reached a similar conclusion, finding that in-store interventions which reduce the availability and prominence of unhealthy foods and increased the availability and prominence of healthy foods were associated with improved dietary health outcomes⁽⁴²⁾.

It is important to acknowledge that grocery retailers and supermarkets are businesses who are motivated by economic outcomes. Therefore, any intervention to improve the healthfulness of the supermarket food environment should also consider the feasibility and sustainability of healthy food retail strategies from the business outcome standpoint. In a review by, Blake *et al.* n 107 studies were considered to identify types of business outcomes that have been reported in healthy food retail strategy evaluations⁽⁸²⁾. The review found commercial viability and retailer and customer

perceptions were frequently measured as part of healthy food retail strategy evaluation. They concluded in most cases that retailer and customer perceptions, as well as product and promotional strategies were generally favourable⁽⁸²⁾. In a more recent review investigating the evidence on factors influencing the implementation, sustainability and scalability of healthy food retail interventions, Gupta *et al.* concluded that few studies reported on intervention sustainability and scalability and this should be considered in future studies⁽⁸³⁾.

Disparities in the healthfulness of the supermarket community and consumer food environment

In many high-income settings, obesity and diet-related NCD prevalence follow a social gradient, meaning overweight and obesity are more prevalent in lower socio-economic groups^(84,85). To address this health disparity, preventative policies targeting overweight, obesity and diet-related NCDs should be defined and implemented to ensure those with the greatest need benefit the most⁽⁸⁶⁾. Evidence suggests that unhealthy food environments also follow a social gradient in many contexts, meaning areas of higher deprivation have less access to healthy food outlets, including supermarkets⁽²⁹⁾. For example, in Australia, New Zealand and Belgium areas of higher deprivation had fewer supermarkets^(66–68). In the UK, using person-centred SES metrics, those with a higher education lived significantly closer to supermarkets than those with a lower education⁽⁸⁷⁾. Similarly in Ireland, distance to the nearest food store and diet quality was mediated by SES, with people of lower SES living further away from supermarkets and having lower diet quality⁽⁶⁵⁾. Evidence also suggests that the in-store food environment in supermarkets located in areas of high deprivation has significantly more unhealthy foods in some contexts such as Australia, New Zealand, and Belgium^(66–68). This means lower SES groups face a double burden in access to healthy food options, due to less access to supermarkets and less healthy foods available in supermarkets that are accessible. Given the important role supermarkets play in access to healthy foods, this situation could exacerbate disparities in diet-related health outcomes and requires policy action.

There is less evidence on the food environment, including the RFE, in low- and middle-income countries (LMIC)⁽⁸⁸⁾. Turner *et al.*, completed a systematic scoping review to investigate the relationship between the food environment, food choice and diet-related health outcomes in LMIC, concluding there were no published articles from low income countries and evidence regarding associations between the food environment and nutrition and health outcomes was inconclusive⁽⁸⁸⁾.

Improving the healthfulness of the supermarket food environment: policy options and opportunities

A variety of evidence-based policies can be used to improve the healthfulness of the supermarket food environment. Macro-level intervention is required as targeting individual behaviours to prevent obesity and food-related ill health will not have the desired impact without addressing the context in which these behaviours are practised⁽²³⁾. A Storey *et al.* concluded that to reduce food-related ill health and obesity, public health interventions are required that address environmental context and conditions in which people make health determining decisions⁽²³⁾. Macro-level structural policies, which change the food environment to one

which is health enabling are expected to have the highest and most equitable impact on overweight and obesity prevention⁽⁸⁹⁾.

As previously outlined, in the USA and Scotland, policies were introduced to increase the availability of supermarkets in food deserts, which had limited impact on improving healthy food purchase in some locations^(60,63) and positive impacts in others⁽⁵⁵⁾. The evidence also suggests that the availability of healthy food outlets does not buffer the effect of fast-food outlet proximity and higher BMI⁽⁹⁰⁾. Potentially the inverse could be true; where higher density and proximity of fast-food outlets could limit the impact of providing healthy food outlets, such as supermarkets, in food deserts; and this requires further investigation. Therefore, consideration should be given to the wider foodscape, and not policies which focus on increasing the availability of healthy food outlets, such as supermarkets, alone.

Both voluntary and mandatory policies have been implemented to create more health-enabling consumer food environments in supermarkets. In the creation of health enabling in-store food environments in supermarkets, co-design and co-creation approaches have been used to develop, implement, and evaluate interventions^(91,92). In designing and implementing these interventions, researchers describe some interventions as being more acceptable and easier to pursue, such as promotions on healthy and fresh foods, which lay the foundations for the design and implementation of interventions that can lead to more substantive change, such as price increases for HFSS foods⁽⁹²⁾. Co-design of healthy in-store interventions with retailers, which augment merchandising strategies to make healthier food choices the easy choice, has led to interventions which have increased purchases of healthy foods^(46,81,92,93). The Food (Promotion and Placement) (England) Regulation 2021, is an example of mandatory restriction on the placement and promotion of unhealthy foods in the grocery RFE in England⁽⁹⁴⁾. This legislation restricts the placement of HFSS foods in high-prominence locations in eligible grocery store outlets. Early investigations into the effectiveness of these mandatory measures highlight they are a good first step but require stronger enforcement and resourcing⁽⁹⁵⁾. The effectiveness of this intervention in reducing purchases of unhealthy foods will be of interest to many, following full evaluation of the measure.

Although not as widely covered in the scientific literature on the healthfulness of the supermarket food environment, retailers play an important role in the nutrient composition of commonly purchased and consumed prepacked foods. Private label, or own brand, food products make up a considerable part of prepacked food market share in Europe, where in 2014 between 17% and 44.5% of food product market share was private label^(44,96). Working with retailers in both the provision of and composition of these foods, is an opportunity to improve the healthfulness of the consumer RFE and make the healthier choice the easier choice⁽⁴⁴⁾. Food reformulation has been shown to be an effective strategy to reduce nutrients of public health concern, such as salt, in prepacked foods and consumer diets⁽¹⁰⁾. To reduce nutrients of public health concern in the food supply, food retailers have a responsibility to reduce the salt, sugar, and saturated fat in their private-label prepacked foods through food reformulation^(97,98). To address this, voluntary and mandatory reformulation policies have been established in many countries^(97–101). The reduction of salt by voluntary reformulation has led to reductions in prepacked foods⁽¹⁰²⁾. For example, a review of published literature found evidence of a consistent reduction in the sugar and salt content of breakfast cereals sold in the RFE⁽¹⁰³⁾. Despite this progress some researchers have proposed that reformulation under voluntary

policies has not gone far or fast enough and additional efforts, such as mandatory legislation, is needed⁽¹⁰⁴⁾.

Monitoring the healthfulness of the retail food environment

Measurement of the healthfulness of the RFE needs to consider several aspects, such as the availability, visibility and the nutrient composition of HFSS foods. To inform the development of contextually sensitive policies to improve the healthfulness of the supermarket food environment and to accurately measure their outcomes and impact, effective monitoring and evaluation strategies are required⁽¹⁰⁵⁾. The International Network for Food and Obesity/Non-communicable Diseases Research, Monitoring and Action Support (INFORMAS) has established a monitoring framework for the food environment⁽¹⁰⁵⁾. This framework outlines modules and protocols for monitoring the food environment, including the RFE and is an important source of standardised measures and tools to inform national food environment monitoring systems⁽¹⁰⁵⁾.

A combination of approaches is used to measure different aspects of the food environment. Internationally, the community food environment has been measured using GIS mapping which is critical for monitoring food outlet density by area-level deprivation and rurality⁽¹⁰⁶⁾. However, Wilkins *et al.* concluded that methods and reporting of GIS methods applied to the food environment are inconsistent and incomplete and proposed a reporting checklist, Geo-FERN, to support comprehensive measurement and reporting of the RFE⁽¹⁰⁷⁾. GIS mapping of the food environment will continue to play an important role in understanding the relationship between food outlet density, proximity, area-level deprivation and diet-related health outcomes.

The healthfulness of the consumer food environment is determined using methodological approaches that measure merchandising strategies based on the marketing mix, including product, price, placement, and promotion^(46,108). Several standardised approaches for measuring the healthfulness of the consumer food environment have been developed⁽¹⁰⁹⁾. A review by Brimblecombe *et al.* found *n* 47 tools have been applied to measure the healthiness of the consumer food environment⁽¹⁰⁹⁾. Tools that have been developed and applied in numerous settings to measure the healthiness of the in-store food environment include the Nutrition Environment Measures Survey (NEMS) and the INFORMAS Food Availability in Supermarkets tool^(35,110). Glanz *et al.* concluded that *n* 123 studies had used a modified version of the NEMS tool, meaning it has played a pivotal role in the development and increase of research on the food environment⁽³⁵⁾. The INFORMAS Food Availability in Supermarkets tool has been implemented in Australia, New Zealand, Argentina, Burkina Faso, Belgium, and Ireland^(66–68,111,112). More recently, Store Scout has been developed to rapidly measure the healthiness of the in-store food environment using an mHealth application^(109,113).

Changes in the composition of prepacked foods are measured using food label and composition information⁽¹¹⁴⁾. This means many data points (e.g. food product identifying information, description, nutrition composition, product weight and ingredients) for a large volume of prepacked foods are required to accurately monitor the evolution of the nutrient composition and ingredient content of the prepacked food supply over time⁽¹¹⁴⁾. Given the voluminous nature of data required to continually monitor the composition of the prepacked food supply, several

countries have piloted or established data repositories^(115–118). These repositories are known as branded food databases, which serve as valuable resources for scientific research, healthcare, public health monitoring, policy development, education, food innovation and others^(115,116). Branded food databases are compiled and maintained by different stakeholders in different countries; for example, foodDB was established and is maintained by a research group at Oxford University, the Deutch Branded Food Database, LEDA, is hosted and maintained by the Deutch Institute of Public Health and the Slovenia branded food database, CLAS, is hosted and maintained by their national nutrition institute^(115,117,119).

Branded food databases are only of value for policy development, monitoring and evaluation, if they are based on accurate and up-to-date nutrition declaration information⁽¹²⁰⁾. The European Commission guidelines on nutrition labelling tolerances sets out a tolerance for the acceptable variation in the declared and actual nutrient content of prepacked foods⁽¹²¹⁾. The range of what is allowable for legal labelling purposes, depends on the type of food, whether a claim is made on the food and the type and amount of the nutrient in the food⁽¹²¹⁾. The empirical evidence on the accuracy of nutrition declarations made on food labels varies with studies in Portugal and Italy finding 27 % and 35 % of products sampled were outside of the labelling tolerances, respectively^(120,122). In contrast, studies in Slovenia and Spain found high rates of declared sugar conformance with labelling tolerances, of 100 % and 98.4 % respectively^(123,124). Given the variability in the accuracy of nutrition declarations on food labels, food composition monitoring using food labels may not always be a reliable source of information and confirmatory laboratory analysis of nutrient content should be undertaken⁽¹¹⁴⁾. To reflect the potential for inaccuracies of nutrition declarations on food labels, guidance by research consortiums on methodologies to monitor the prepacked food supply recognise the role for laboratory analysis to determine the nutrient composition of foods^(114,125).

Commercial datasets which measure food product volume share can be used to complement, or sometimes instead of, the approaches described for measuring and monitoring the consumer food environment, and the nutrient composition of the prepacked food supply⁽¹²⁶⁾. In their systematic review, Bandy *et al.* found *n* 68 studies which used sales or purchase data from four commercial data providers (Euromonitor, GfK, Kantar and Nielsen) to measure and evaluate interventions to improve nutrient quality of foods, food purchasing including socio-demographic variations in purchasing⁽¹²⁶⁾.

Future perspectives

The supermarket food environment is extending into the digital space with the growth of online grocery stores and home delivery^(127,128). This evolution presents new considerations which will need to be captured in tools and frameworks for monitoring and evaluating the healthfulness of the supermarket food environment⁽¹²⁹⁾. A review by Maganja *et al.* found only one tool for determining the healthfulness of the online supermarket environment and concluded additional validated tools are needed to measure the new opportunities online grocery stores can use to influence consumers⁽¹²⁹⁾. The advancement in online grocery stores provides an opportunity to more efficiently collect food product information from retailer websites, which is already underway in some branded food databases such as FLIP in Canada and foodDB in the UK^(116,119). The potential for the use of online

data to support food composition monitoring should continue to be explored.

With the introduction of regulation to limit the placement and promotion of HFSS foods in high-prominence areas of supermarkets in England, the consumer food environment is adapting^(94,95). However, the impact of this regulation is yet to be evaluated. If this regulatory measure is effective in reducing the sales of HFSS foods and stimulating food product reformulation, similar strategies could be considered elsewhere. How retailers (and the food industry) respond to rules limiting the placement and promotion of foods, may require adaptations to the consumer food environment monitoring tools.

In high-income countries, overweight, obesity and diet-related NCDs follow a social gradient^(130–132). At the same time, lower socio-economic groups have less access to supermarkets in many contexts⁽⁶⁵⁾, and where they are accessible the consumer food environment is less healthy^(66,67). For healthy supermarket interventions to contribute to addressing this health disparity, policies addressing the healthfulness of the supermarket food environment will need to ensure the disparity in access to healthier food options is addressed. This can be done by prioritising areas of high deprivation for intervention first and evaluating the effectiveness of interventions across different socio-economic groups.

Finally, improving the sustainability of our food supply is a key component of addressing the climate crisis⁽¹³³⁾. Consumption has been described as an important pillar of sustainable food systems^(134,135). The supermarket food environment could play an important role in directing consumer food choice towards both healthy and sustainable foods⁽²³⁾. Currently, there are few tools which examine the availability of HFSS foods and food sustainability in tandem⁽¹³⁶⁾. Baker *et al.* reported only *n* 2 of *n* 58 studies investigating the availability of healthy foods also considered sustainability⁽¹³⁶⁾ and concluded there is a need to develop consumer nutrition environment measures that assess nutrient-dense food availability and food sustainability to inform improving consumer food environments for human and planetary health⁽¹³⁶⁾. Tools and frameworks developed to determine the healthfulness of the consumer food environment should consider food sustainability alongside healthiness⁽¹³⁶⁾.

Conclusion

The food environment mediates consumer food choices and thus plays an important role in diet quality and related health outcomes⁽²⁴⁾. Across high-income countries, the majority of populations are reliant on the built food environment to access food⁽⁴³⁾. In the Western world the built food environment has been described as obesogenic, meaning it does not support people to make food choices in line with FBDGs recommendations, whilst also being favourable to a sedentary lifestyle⁽²⁸⁾. Supermarkets are an important source of healthy foods in the built food environment⁽¹³⁷⁾. However, the extent to which supermarkets can influence access to healthy food depends on supermarket proximity, density, and the healthfulness of the supermarket consumer food environment^(24,65,74). The evidence supporting improved access to healthy foods and diet-related health outcomes by increasing supermarket density is mixed^(57,63). The evidence supports improved purchase of healthy foods by increasing the healthfulness of the supermarket consumer food environment^(42,74). Healthy in-store initiatives to improve the purchase of healthy foods using a combination of merchandising strategies

based on the marketing mix: product, price, promotion and placement, have been shown to be effective across a wide range of countries and supermarket types⁽⁷⁴⁾. Policies such as voluntary measures co-designed with retailers to improve the healthfulness of the supermarket consumer food environment through restriction of product placement and private label reformulation have also been shown to increase healthier food purchases^(46,93). However, evidence also shows that mandatory, structural changes are most effective in improving the disparities in access to healthy food and diet-related ill health⁽⁸⁹⁾. This is an important consideration for improving access to healthy foods for lower socio-economic groups, who carry a higher burden of food-related ill health⁽¹³¹⁾. The Food (Promotion and Placement) (England) Regulations 2021 is a legal measure to restrict placement and promotion of HFSS foods, and its impact on the purchase of healthy and unhealthy food could inform future policies⁽⁹⁴⁾. Future research and policy with a focus on the supermarket food environment should consider equitable access to healthy foods, sustainability alongside healthiness, and the growth of online supermarkets. Additional research is needed to understand how the density of fast-food outlets mediates the effectiveness of supermarkets in improving the purchase and consumption of healthy food, and the effectiveness of legal measures to restrict the in-store availability and purchase of unhealthy foods from supermarkets^(61,95).

Acknowledgements. The authors would like to thank the Nutrition Society for the opportunity to present this research at the Irish Section Postgraduate Meeting 2024.

Authors contributions. Conceptualisation SoM, ERG, NC and GD; literature review SoM; writing original draft SoM; writing review and edit SoM, ERG, GD and NC; supervision GD, ERG and NC; project administration SoM. All authors have read and agreed to the published version of the manuscript.

Financial support. The authors would like to thank the Food Safety Authority of Ireland for funding this research.

Competing interests. ERG has received research funding through the following funders and projects; Food for Health Ireland (www.fhi.ie) project, funded by Enterprise Ireland, co-funded with core partners Carbery, Kerry, Tirlan, Dairygold & Bord Bia; Science Foundation Ireland Insight Centre for Data Analytics, and Co-Centre for Sustainable Food Systems; Horizon Europe most recently in projects such as FNSSCloud, PLANEAT and MarieCurie CareerFIT; and PhD studentship funding from Société des Produits Nestlé, Switzerland. Eileen R Gibney has completed consultancy work for the following: Société des Produits Nestlé, Switzerland; Irish Advertising Standards Agency; and Food Safety Authority of Ireland. No personal payment was received, all payments were made into a research fund through Consult UCD.

SoM, NC and GD have no declarations of interest.

References

- Stafford N (2010) History: the changing notion of food. *Nat* **468**, S16–S7.
- Neufeld LM, Hendriks S & Hugas M (2023) Healthy Diet: A Definition for the United Nations Food Systems Summit 2021. In *Science and Innovations for Food Systems Transformation*, pp. 21–30 [J von Braun, K Afsana, LO Fresco and MHA Hassan, editors]. Cham: Springer International Publishing.
- Herforth A, Arimond M, Álvarez-Sánchez C, *et al.* (2019) A global review of food-based dietary guidelines. *Adv Nutr* **10**, 590–605.
- Leme ACB, Hou S, Fisberg RM, *et al.* (2021) Adherence to food-based dietary guidelines: a systemic review of high-income and low- and middle-income countries. *Nutrients* **13**, 1038.
- Nettleton JA, Brouwer IA, Geleijnse JM, *et al.* (2017) Saturated fat consumption and risk of coronary heart disease and ischemic stroke: a science update. *Ann Nutr Metab* **70**, 26–33.

6. Fiolet T, Srour B, Sellem L, *et al.* (2018) Consumption of ultra-processed foods and cancer risk: results from NutriNet-Santé prospective cohort. *BMJ* **360**, k322.
7. Wang Q, Afshin A, Yakoob MY, *et al.* (2016) Impact of nonoptimal intakes of saturated, polyunsaturated, and trans fat on global burdens of coronary heart disease. *J Am Heart Assoc* **5**, e002891.
8. Afshin A, Sur PJ, Fay KA, *et al.* (2019) Health effects of dietary risks in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* **393**, 1958–1972.
9. Central Statistics Office (2023) *Vital Statistics Yearly Summary 2022*. Cork: Central Statistics Office.
10. Gressier M, Swinburn B, Frost G, *et al.* (2021) What is the impact of food reformulation on individuals' behaviour, nutrient intakes and health status? A systematic review of empirical evidence. *Obesity Rev* **22**, e13139.
11. Milte CM & McNaughton SA (2016) Dietary patterns and successful ageing: a systematic review. *Eur J Nutr* **55**, 423–450.
12. Schoufour JD, Voortman T., Franco OH, *et al.* (2017) Dietary Patterns and Healthy Aging. In *Food for the Aging Population*, 2nd ed. Amsterdam: Elsevier Inc.
13. Bennett BJ, Hall KD, Hu FB, *et al.* (2015) Nutrition and the science of disease prevention: a systems approach to support metabolic health. *Ann N Y Acad Sci* **1352**, 1–12.
14. Bronfenbrenner U (1977) Toward an experimental ecology of human development. *Am Psychologist* **32**, 513–531.
15. Davison KK & Birch LL (2001) Childhood overweight: a contextual model and recommendations for future research. *Obesity Rev* **2**, 159–171.
16. Ohri-Vachaspati P, DeLia D, DeWeese RS, *et al.* (2015) The relative contribution of layers of the social ecological model to childhood obesity. *Public Health Nutr* **18**, 2055–2066.
17. Butland B, Jebb S, Kopelman P, *et al.* (2007) Foresight. Tackling obesity: future choices. *Obes Rev* **8**, 6–9.
18. Steele M, Mialon M, Browne S, *et al.* (2021) Obesity, public health ethics and the nanny state. *Ethics, Med Public Health* **19**, 100724.
19. World Health Organization (2022) *Systems Thinking for Noncommunicable Disease Prevention Policy; Guidance to Bring Systems Approaches into Practice*. Geneva: WHO.
20. Allender S, Brown AD, Bolton KA, *et al.* (2019) Translating systems thinking into practice for community action on childhood obesity. *Obesity Rev* **20**, 179–184.
21. Brug J (2008) Determinants of healthy eating: motivation, abilities and environmental opportunities. *Family Pract* **25**, i50–i5.
22. Hruby A & Hu FB (2015) The epidemiology of obesity: a big picture. *Pharmacoeconomics* **33**, 673–689.
23. Story M, Kaphingst KM, Robinson-O'Brien R, *et al.* (2008) Creating healthy food and eating environments: policy and environmental approaches. *Ann Rev Public Health* **29**, 253–272.
24. Pineda E, Stockton J, Scholes S, *et al.* (2024) Food environment and obesity: a systematic review and meta-analysis. *BMJ Nutr Prev Health* **7**, e000663.
25. Swinburn B, Sacks G, Vandevijvere S, *et al.* (2013) Informas overview. *Obes Rev* **14**, 1–12.
26. Swinburn B, Kraak V, Rutter H, *et al.* (2015) Strengthening of accountability systems to create healthy food environments and reduce global obesity. *Lancet* **385**, 2534–2545.
27. Lake A & Townshend T (2006) Obesogenic environments: exploring the built and food environments. *J Royal Soc for Promotion Health* **126**, 262–267.
28. Lake AA (2018) Neighbourhood food environments: food choice, foodscapes and planning for health. *Proc Nutr Soc* **77**, 239–246.
29. Cooksey-Stowers K, Schwartz MB & Brownell KD (2017) Food Swamps predict obesity rates better than food deserts in the United States. *Int J Environ Res Public Health* **14**, 1366.
30. Glanz K, Sallis JF, Saelens BE, *et al.* (2005) Healthy nutrition environments: concepts and measures. *Am J Health Promot* **19**, 330–333.
31. Herforth A & Ahmed S (2015) The food environment, its effects on dietary consumption, and potential for measurement within agriculture-nutrition interventions. *Food Secur* **7**, 505–520.
32. Turner C, Aggarwal A, Walls H, *et al.* (2018) Concepts and critical perspectives for food environment research: a global framework with implications for action in low- and middle-income countries. *Global Food Secur* **18**, 93–101.
33. Tufford AR, Diou C, Lucassen DA, *et al.* (2022) Toward systems models for obesity prevention: a big role for big data. *Curr Dev Nutr* **6**, nzac123.
34. Engler-Stringer R, Le H, Gerrard A, *et al.* (2014) The community and consumer food environment and children's diet: a systematic review. *BMC Public Health* **14**, 522.
35. Glanz K, Fultz AK, Sallis JF, *et al.* (2023) Use of the nutrition environment measures survey: a systematic review. *Am J Prev Med* **65**, 131–142.
36. Chen X & Yang X (2014) Does food environment influence food choices? A geographical analysis through "tweets". *Appl Geogr* **51**, 82–89.
37. Chege CGK, Wanyama R, Lundy M, *et al.* (2021) Does retail food diversity in urban food environments influence consumer diets?. *Sustainability* **13**, 7666.
38. Kelly C, Callaghan M, Molcho M, *et al.* (2019) Food environments in and around post-primary schools in Ireland: associations with youth dietary habits. *Appetite* **132**, 182–189.
39. Atanasova P, Kusuma D, Pineda E, *et al.* (2022) The impact of the consumer and neighbourhood food environment on dietary intake and obesity-related outcomes: a systematic review of causal impact studies. *Social Sci Med* **299**, 114879.
40. Winkler MR, Zenk SN, Baquero B, *et al.* (2020) A model depicting the retail food environment and customer interactions: components, outcomes, and future directions. *Int J Environ Res Public Health* **17**, 7591.
41. Glanz K, Bader MDM & Iyer S (2012) Retail grocery store marketing strategies and obesity: an integrative review. *Am J Preventative Med* **42**, 503–512.
42. Shaw SC, Ntani G, Baird J, *et al.* (2020) A systematic review of the influences of food store product placement on dietary-related outcomes. *Nutr Rev* **78**, 1030–1045.
43. Downs SM, Ahmed S, Fanzo J, *et al.* (2020) Food environment typology: advancing an expanded definition, framework, and methodological approach for improved characterization of wild, cultivated, and built food environments toward sustainable diets. *Foods* **9**, 532.
44. Pulker CE, Trapp GSA, Scott JA, *et al.* (2018) What are the position and power of supermarkets in the Australian food system, and the implications for public health? A systematic scoping review. *Obesity Rev* **19**, 198–218.
45. Hawkes C (2009) Sales promotions and food consumption. *Nutr Rev* **67**, 333–342.
46. Brimblecombe J, McMahon E, Ferguson M, *et al.* (2020) Effect of restricted retail merchandising of discretionary food and beverages on population diet: a pragmatic randomised controlled trial. *Lancet Planet Health* **4**, e463–e73.
47. Lake AA, Burgoine T, Greenhalgh F, *et al.* (2010) The foodscape: classification and field validation of secondary data sources. *Health Place* **16**, 666–673.
48. Charreire H, Casey R, Salze P, *et al.* (2010) Measuring the food environment using geographical information systems: a methodological review. *Public Health Nutr* **13**, 1773–1785.
49. Needham C, Strugnell C, Allender S, *et al.* (2022) Beyond food swamps and food deserts: exploring urban Australian food retail environment typologies. *Public Health Nutr* **25**, 1140–1152.
50. Vaughan CA, Cohen DA, Ghosh-Dastidar M, *et al.* (2017) Where do food desert residents buy most of their junk food? Supermarkets. *Public Health Nutr* **20**, 2608–2616.
51. Dubowitz T, Ghosh-Dastidar M, Cohen DA, *et al.* (2015) Diet and perceptions change with supermarket introduction in a food desert, but not because of supermarket use. *Health Aff (Millwood)* **34**, 1858–1868.
52. Macdonald L, Ellaway A & Macintyre S (2009) The food retail environment and area deprivation in Glasgow City, UK. *Int J Behav Nutr Physical Activity* **6**, 52.
53. Recchia D, Perignon M, Rollet P, *et al.* (2022) Associations between retail food environment and the nutritional quality of food purchases in French households: the Mont'Panier cross-sectional study. *PLOS ONE* **17**, e0267639.

54. Laraia BA, Siega-Riz AM, Kaufman JS, et al. (2004) Proximity of supermarkets is positively associated with diet quality index for pregnancy. *Prev Med* **39**, 869–875.
55. Rogus S, Athens J, Cantor J, et al. (2018) Measuring micro-level effects of a new supermarket: do residents within 0.5 mile have improved dietary behaviors?. *J Academy Nutr Diet* **118**, 1037–1046.
56. Lamichhane AP, Mayer-Davis EJ, Puett R, et al. (2012) Associations of built food environment with dietary intake among youth with diabetes. *J Nutr Educ Behavior* **44**, 217–224.
57. Athens JK, Duncan DT & Elbel B (2016) Proximity to Fast-Food Outlets and Supermarkets as Predictors of Fast-Food Dining Frequency. *J Academy Nutr Diet* **116**, 1266–1275.
58. Powell LM, Auld MC, Chaloupka FJ, et al. (2007) Associations between access to food stores and adolescent body mass index. *Am J Prev Med* **33**, S301–7.
59. Fiechtner L, Kleinman K, Melly SJ, et al. (2016) Effects of proximity to supermarkets on a randomized trial studying interventions for obesity. *Am J Public Health* **106**, 557–562.
60. Elbel B, Moran A, Dixon LB, et al. (2015) Assessment of a government-subsidized supermarket in a high-need area on household food availability and children's dietary intakes. *Public Health Nutr* **18**, 2881–2890.
61. Lamichhane AP, Warren JL, Puett RC, et al. (2013) Spatial patterning of supermarkets and fast food outlets with respect to neighborhood characteristics. *Health Place* **23**, 157–164.
62. Stevenson AC, Brazeau A-S, Dasgupta K, et al. (2019) Evidence synthesis-Neighbourhood retail food outlet access, diet and body mass index in Canada: a systematic review. *Health Promot Chronic Dis Prev Can* **39**, 261.
63. Cummins S, Petticrew M, Higgins C, et al. (2005) Large scale food retailing as an intervention for diet and health: quasi-experimental evaluation of a natural experiment. *J Epidemiol Community Health* **59**, 1035–1040.
64. Skidmore P, Welch A, van Sluijs E, et al. (2010) Impact of neighbourhood food environment on food consumption in children aged 9–10 years in the UK SPEEDY (sport, physical activity and eating behaviour: environmental determinants in young people) study. *Public Health Nutr* **13**, 1022–1030.
65. Layte R, Harrington J, Sexton E, et al. (2011) Irish exceptionalism? local food environments and dietary quality. *J Epidemiol Community Health* **65**, 881–888.
66. Schultz S, Cameron AJ, Grigsby-Duffy L, et al. (2021) Availability and placement of healthy and discretionary food in Australian supermarkets by chain and level of socio-economic disadvantage. *Public Health Nutr* **24**, 203–214.
67. Vandevijvere S, Waterlander W, Molloy J, et al. (2018) Towards healthier supermarkets: a national study of in-store food availability, prominence and promotions in New Zealand. *Eur J Clin Nutr* **72**, 971–978.
68. Vandevijvere S, Van Dam I, Inaç Y, et al. (2023) Unhealthy food availability, prominence and promotion in a representative sample of supermarkets in Flanders (Belgium): a detailed assessment. *Arch Public Health* **81**, 154.
69. O'Mahony S, Collins N, Doyle G, et al. (2024) A cross-sectional study of the relative availability and prominence of shelf space allocated to healthy and unhealthy foods in supermarkets in urban Ireland, by area-level deprivation. *BMC Public Health* **24**, 2689.
70. Vepsäläinen H, Nevalainen J, Kinnunen S, et al. (2022) Do we eat what we buy? Relative validity of grocery purchase data as an indicator of food consumption in the LoCard study. *Br J Nutr* **128**, 1780–1788.
71. Brimblecombe J, McMahon E, De Silva K, et al. (2020) Transforming food retail for better health: the healthy stores 2020 trial. *Eur J Public Health* **30**, ckaa165.072.
72. Cameron AJ, Charlton E, Ngan WW, et al. (2016) A systematic review of the effectiveness of supermarket-based interventions involving product, promotion, or place on the healthiness of consumer purchases. *Curr Nutr Rep* **5**, 129–138.
73. World Health Organization (2015) *Regional Office for E. Using price policies to promote healthier diets*. Copenhagen: World Health Organization.
74. Adam A & Jensen JD (2016) What is the effectiveness of obesity related interventions at retail grocery stores and supermarkets? —a systematic review. *BMC Public Health* **16**, 1247.
75. Geliebter A, Ang I, Bernales-Korins M, et al. (2013) Supermarket discounts of low-energy density foods: effects on purchasing, food intake, and body weight. *Obesity* **21**, E542–E8.
76. Glanz K & Yaroch AL (2004) Strategies for increasing fruit and vegetable intake in grocery stores and communities: policy, pricing, and environmental change. *Prev Med* **39**, S75–80.
77. Waterlander WE, de Boer MR, Schuit AJ, et al. (2013) Price discounts significantly enhance fruit and vegetable purchases when combined with nutrition education: a randomized controlled supermarket trial. *Am J Clin Nutr* **97**, 886–895.
78. Rose D, Hutchinson PL, Bodor JN, et al. (2009) Neighborhood food environments and Body Mass Index: the importance of in-store contents. *Am J Prev Med* **37**, 214–219.
79. Sigurdsson V, Larsen NM & Gunnarsson D (2014) Healthy food products at the point of purchase: an in-store experimental analysis. *J Appl Behavior Anal* **47**, 151–154.
80. Rosin M, Young L, Jiang Y, et al. (2023) Product promotional strategies in supermarkets and their effects on sales: a case study of breakfast cereals and drinks in New Zealand. *Nutr Diet* **80**, 463–471.
81. Cameron AJ, Brown A, Orellana L, et al. (2022) Change in the healthiness of foods sold in an Australian supermarket chain following implementation of a shelf tag intervention based on the health star rating system. *Nutrients* **14**, 2394.
82. Blake MR, Backholer K, Lancsar E, et al. (2019) Investigating business outcomes of healthy food retail strategies: a systematic scoping review. *Obesity Rev* **20**, 1384–1399.
83. Gupta A, Alston L, Needham C, et al. (2022) Factors influencing implementation, sustainability and scalability of healthy food retail interventions: a systematic review of reviews. *Nutrients* **14**, 294.
84. Cronin FM, Hurley SM, Buckley T, et al. (2022) Mediators of socioeconomic differences in overweight and obesity among youth in Ireland and the UK (2011–2021): a systematic review. *BMC Public Health* **22**, 1585.
85. Marmot M & Bell R (2019) Social determinants and non-communicable diseases: time for integrated action. *BMJ* **364**, l251.
86. Marmot MAJ, Boyce T, Goldblatt P, et al. (2020) Health equity in England: the marmot review 10 years on. *Inst Health Equity* **368**, m693.
87. Maguire ER, Burgoine T, Penney TL, et al. (2017) Does exposure to the food environment differ by socioeconomic position? Comparing area-based and person-centred metrics in the Fenland Study, UK. *Int J Health Geographics* **16**, 33.
88. Turner C, Kalamatianou S, Drewnowski A, et al. (2020) Food environment research in low- and middle-income countries: a systematic scoping review. *Adv Nutr* **11**, 387–397.
89. Backholer K, Beauchamp A, Ball K, et al. (2014) A framework for evaluating the impact of obesity prevention strategies on socioeconomic inequalities in weight. *Am J Public Health* **104**, e43–50.
90. van Erpecum CL, van Zon SKR, Bültmann U, et al. (2022) The association between the presence of fast-food outlets and BMI: the role of neighbourhood socio-economic status, healthy food outlets, and dietary factors. *BMC Public Health* **22**, 1432.
91. Vargas C, Whelan J, Brimblecombe J, et al. (2022) Co-creation of healthier food retail environments: a systematic review to explore the type of stakeholders and their motivations and stage of engagement. *Obesity Rev* **23**, e13482.
92. Middel CNH, Schuitmaker-Warnaar TJ, Mackenbach JD, et al. (2022) Designing a healthy food-store intervention; a co-creative process between interventionists and supermarket actors. *Int J Health Policy Manage* **11**, 2175–2188.
93. Vogel C, Crozier S, Penn-Newman D, et al. (2021) Altering product placement to create a healthier layout in supermarkets: outcomes on store sales, customer purchasing, and diet in a prospective matched controlled cluster study. *PLoS Med* **18**, e1003729.

94. Public Health, England (2021) *The Food (Promotion and Placement) (England) Regulations 2021*. London: Public Health, England.
95. Muir S, Dhuria P, Roe E, *et al.* (2023) UK government's new placement legislation is a 'good first step': a rapid qualitative analysis of consumer, business, enforcement and health stakeholder perspectives. *BMC Med* **21**, 33.
96. Olbrich R, Hundt M & Jansen HC (2016) Proliferation of private labels in food retailing: a literature overview. *Int J Marketing Stud* **8**, 63–76.
97. Public Health England (2017) *Sugar Reduction: Achieving the 20%*. London: Public Health, England.
98. Department of Health (2021) *A Roadmap for Food Product Reformulation in Ireland*. Dublin: Department of Health.
99. Ministerio de Sandid (2016) *Collaboration PLAN for the Improvement of the Composition of Food and Beverages and Other Measures 2020*. Madrid: Ministerio de Sandid.
100. Public Health England (2020) *Salt Reduction Targets for 2024*. London: Public Health, England.
101. Healthy Food Partnership (2018) *Health Food Partnership Reformulation Programme: Evidence Informing the Approach, Draft Targets and Modelling Outcomes*. Canberra ACT: Department of Health and Aged Care.
102. He FJ, Brown M, Tan M, *et al.* (2019) Reducing population salt intake—an update on latest evidence and global action. *J Clin Hypertens (Greenwich)* **21**, 1596–1601.
103. O'Mahony S, O'Donovan CB, Collins N, *et al.* (2023) Reformulation of processed yogurt and breakfast cereals over time: a scoping review. *Int J Environ Res Public Health* **20**, 3322.
104. Song J, Tan M, Wang C, *et al.* (2023) Salt intake, blood pressure and cardiovascular disease mortality in England, 2003–2018. *J Hypertens* **41**, 1713–1720.
105. Swinburn B, Vandevijvere S, Kraak V, *et al.* (2013) Monitoring and benchmarking government policies and actions to improve the healthiness of food environments: a proposed government healthy food environment policy index. *Obes Rev* **14**, 24–37.
106. Clelland D & Deprivation Hill C (2019) policy and rurality: the limitations and applications of area-based deprivation indices in Scotland. *Local Economy* **34**, 33–50.
107. Wilkins EL, Morris MA, Radley D, *et al.* (2017) Using geographic information systems to measure retail food environments: discussion of methodological considerations and a proposed reporting checklist (Geo-FERN). *Health Place* **44**, 110–117.
108. van Waterschoot W & van den Bulte C (1992) The 4P classification of the marketing mix revisited. *J Marketing* **56**, 83–93.
109. Brimblecombe J, Jaenke R, McMahon E, *et al.* (2021) Development and pilot of a tool to measure the healthiness of the in-store food environment. *Public Health Nutr* **24**, 243–252.
110. Ni Mhurchu C, Vandevijvere S, Waterlander W, *et al.* (2013) Monitoring the availability of healthy and unhealthy foods and non-alcoholic beverages in community and consumer retail food environments globally. *Obesity Rev* **14**, 108–119.
111. O'Mahony S, Collins NA, Doyle G, *et al.* (2022) Supermarket availability and socio-economic deprivation in urban Ireland. *Proc Nutr Soc* **81**, E100.
112. Elorriaga N, Moyano DL, López MV, *et al.* (2021) Urban retail food environments: relative availability and prominence of exhibition of healthy vs. unhealthy foods at supermarkets in Buenos Aires, Argentina. *Int J Environ Res Public Health* **18**, 944.
113. McMahon EJ, Jaenke R & Brimblecombe J (2020) A mobile app to rapidly appraise the in-store food environment: reliability, utility, and construct validity study. *JMIR Mhealth Uhealth* **8**, e16971.
114. Neal B, Sacks G, Swinburn B, *et al.* (2013) Monitoring the levels of important nutrients in the food supply. *Obes Rev* **14**, 49–58.
115. Pravst I, Hribar M, Žmitek K, *et al.* (2022) Branded foods databases as a tool to support nutrition research and monitoring of the food supply: insights from the Slovenian composition and labeling information system. *Front Nutr* **8**, 798576.
116. Ahmed M, Schermel A, Lee J, *et al.* (2022) Development of the food label information program: a comprehensive Canadian branded food composition database. *Front Nutr* **8**, 825050.
117. Westenbrink S, van der Vossen-Wijmenga W, Toxopeus I, *et al.* (2021) LEDA, the branded food database in the Netherlands: data challenges and opportunities. *J Food Compos Anal* **102**, 104044.
118. O'Mahony S, Pravst I, Hribar M, *et al.* (2023) Piloting a branded food database for reformulation monitoring in Ireland. *Eur J Public Health* **33**, ckad160.1090.
119. Harrington RA, Adhikari V, Rayner M, *et al.* (2019) Nutrient composition databases in the age of big data: foodDB, a comprehensive, real-time database infrastructure. *BMJ open* **9**, e026652.
120. Albuquerque TG, Nunes MA, Oliveira MBPP, *et al.* (2020) Compliance of declared vs. analysed values with EU tolerance limits for mandatory nutrients in prepacked foods. *Food Chem* **302**, 125330.
121. European Commission (2012) *Guidance with Regard to the Setting of Tolerances for Nutrient values Declared on a Label*. Luxembourg: European Commission.
122. Bragolusi M, Tata A, Massaro A, *et al.* (2023) Nutritional labelling of food products purchased from online retail outlets: screening of compliance with European Union tolerance limits by near infrared spectroscopy. *J Near Infrared Spectrosc* **31**, 89–99.
123. Hafner E, Lavriša Ž, Hribar M, *et al.* (2022) Verifying the use of food labeling data for compiling branded food databases: a case study of sugars in beverages. *Front Nutr* **9**, 794468.
124. Yusta-Boyo MJ, Bermejo LM, García-Solano M, *et al.* (2020) Sugar content in processed foods in Spain and a comparison of mandatory nutrition labelling and laboratory values. *Nutrients* **12**, 1078.
125. Dunford E, Webster J, Metzler AB, *et al.* (2012) International collaborative project to compare and monitor the nutritional composition of processed foods. *Eur J Prev Cardiol* **19**, 1326–1332.
126. Bandy L, Adhikari V, Jebb S, *et al.* (2019) The use of commercial food purchase data for public health nutrition research: a systematic review. *PLOS ONE* **14**, e0210192.
127. Kantar World Panel (2024) *Record-Breaking December for Grocery Sales in Ireland as Spend Surpasses €1.4 Billion*. London: Kantar World Panel.
128. Khandpur N, Zatz LY, Bleich SN, *et al.* (2020) Supermarkets in cyberspace: a conceptual framework to capture the influence of online food retail environments on consumer behavior. *Int J Environ Res Public Health* **17**, 8639.
129. Maganja D, Miller M, Trieu K, *et al.* (2022) Evidence gaps in assessments of the healthiness of online supermarkets highlight the need for new monitoring tools: a systematic review. *Curr Atheroscler Rep* **24**, 215–233.
130. Moore Heslin A, O'Donnell A, Kehoe L, *et al.* (2023) Adolescent overweight and obesity in Ireland—Trends and sociodemographic associations between 1990 and 2020. *Pediatr Obesity* **18**, e12988.
131. Andrade CAS, Mahrouseh N, Gabrani J, *et al.* (2023) Inequalities in the burden of non-communicable diseases across European countries: a systematic analysis of the Global Burden of Disease 2019 study. *Int J Equity Health* **22**, 140.
132. Abarca-Gómez L, Abdeen ZA, Hamid ZA, *et al.* (2017) Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128·9 million children, adolescents, and adults. *Lancet* **390**, 2627–2642.
133. Willett W, Rockström J, Loken B, *et al.* (2019) Food in the anthropocene: the EAT-lancet commission on healthy diets from sustainable food systems. *Lancet* **393**, 447–492.
134. Jarmul S, Dangour AD, Green R, *et al.* (2020) Climate change mitigation through dietary change: a systematic review of empirical and modelling studies on the environmental footprints and health effects of 'sustainable diets'. *Environ Res Lett* **15**, 123014.
135. Hallström E, Carlsson-Kanyama A & Börjesson P (2015) Environmental impact of dietary change: a systematic review. *J Cleaner Prod* **91**, 1–11.
136. Baker K, Burd L & Figueroa R (2024) Consumer nutrition environment measurements for nutrient-dense food availability and food sustainability: a scoping review. *Arch Public Health* **82**, 7.
137. Karpyn A, McCallops K, Wolgast H, *et al.* (2020) Improving consumption and purchases of healthier foods in retail environments: a systematic review. *Int J Environ Res Public Health* **17**, 7524.