

Original Article

Cite this article: Munsterman DK and Deckers J. Biostratigraphic ages and depositional environments of the upper Oligocene to lower Miocene Veldhoven Formation in the central Roer Valley Rift System (SE Netherlands-NE Belgium). *Netherlands Journal of Geosciences*, Volume 101, e6. <https://doi.org/10.1017/njg.2022.3>

Received: 3 October 2021

Revised: 11 February 2022

Accepted: 14 February 2022

Keywords:

dinoflagellate cysts; upper Palaeogene; lowermost Neogene; sedimentary facies; southern North Sea Basin; well-log correlation

Author for correspondence:

Dirk K. Munsterman,

Email: dirk.munsterman@tno.nl

Biostratigraphic ages and depositional environments of the upper Oligocene to lower Miocene Veldhoven Formation in the central Roer Valley Rift System (SE Netherlands-NE Belgium)

Dirk K. Munsterman¹  and Jef Deckers²

¹TNO, Geological Survey of the Netherlands, Utrecht, The Netherlands and ²VITO, Flemish Institute for Technological Research, Mol, Belgium

Abstract

Discussions on the age and the depositional environments of the Veldhoven Formation and its members are persistent in Belgium and the Netherlands. Uncertainties on stratigraphy and the constructive process of sediment accumulation continue today as a result of lack of data on this succession within the Roer Valley Rift System. The present study provides new information on the bio- and lithostratigraphy and facies from two boreholes based on dinoflagellate cyst taxa. The results were correlated by gamma-ray logs towards other key boreholes in the region and show a good consistency for stratigraphy and depositional environments for the members of the Veldhoven Formation.

After marginal to restricted marine conditions in the latest Rupelian (early Oligocene), the start of deposition of the Veldhoven Formation marked the transition towards a higher sea level, expressed by increased glauconite contents and gamma-ray values. The Voort Member in the lower part of the Veldhoven Formation has an early to late Chattian (Late Oligocene) age and comprises predominantly shallow marine (fluctuating restricted to open marine) conditions. The lithology in the lower part of this unit is often very clayey but is coarsening upward into sands. The superjacent Wintelre Member has a latest Chattian to early Aquitanian (early Miocene) age. This member is distinct by its clayey nature which is expressed by relatively high gamma-ray values. Earlier studies suggest a deeper marine facies for the Wintelre Member compared to the Someren and Voort members. However, the dinoflagellate cyst assemblages in this unit are mostly dominated by a single genus indicating a restricted marine setting, including salinities that deviate from normal marine conditions, most probably due to minor ventilation by narrow or lack of connection to the Atlantic Ocean. A glacio-eustatic sea-level fall around the Oligocene/Miocene boundary limited the sea coverage to the strongest subsiding areas, where deposition of the Wintelre Member is recorded, while non-deposition or erosion occurred in the surrounding highs, hence creating an isolated (sub)basin. The superjacent Someren Member was deposited during the late Aquitanian to middle Burdigalian and consists of shallow to open marine clayey fine sands. Increasing clay contents indicate a gradual development towards a higher sea level, which coincide with upward increasing gamma-ray values.

The biostratigraphic results of this study suggest that no major hiatuses are present in the differentially subsiding blocks of the Roer Valley Rift System during the late Oligocene to early Miocene.

Introduction

The Oligocene/Miocene boundary in the southern North Sea Basin often coincides with an important hiatus related to the Savian phase. The Savian tectonic pulse is believed to be caused by the regional stress regime induced by the Alpine orogeny, but also sea-level changes may have played a role (Wong et al., 2007). Most of the southern North Sea area emerged above sea level (Fig. 1A). In the city of Antwerp in the western Campine area (Fig. 1B), this hiatus spans the (late) Chattian (late Oligocene) to early Burdigalian (early Miocene; Louwye, 2005). A recent publication by Munsterman & Deckers (2020) showed that the range of this hiatus decreases from the western Campine area towards boreholes in the eastern Campine area, where it coincides with a gravel layer within shallow marine glauconitic sands of the Veldhoven Formation. Even further to the east, in boreholes in the northern Roer Valley Graben (RVG; Fig. 1B), Munsterman et al. (2019) did not find indications for a major Oligocene/Miocene hiatus within the Veldhoven Formation (Fig. 2). This is consistent with renewed activity of the RVG since the

© The Author(s), 2022. Published by Cambridge University Press on behalf of the Netherlands Journal of Geosciences Foundation. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.




Fig. 1. (A) The late Oligocene (middle Chattian) palaeogeography of the North Sea Basin and surrounding areas, modified from Gibbard & Lewin (2016). (B) The study area with the main structural elements of the Roer Valley Rift System. The location of the schematic cross-sections of Fig. 2A and 2B are shown with the two black dotted lines and the correlation panel of Fig. 3 is shown as a blue dotted line. CA, Campine area; GH, Groote Heide borehole; HB, Herkenbosch borehole; ML, Molenbeersel borehole; LBR, Limbricht borehole; KB, Krefelt Block; PB, Peel Block; RVG, Roer Valley Graben; VG, Venlo Graben; PBF, Peel Boundary Fault, TF, Tegelen Fault, VF, Viersen Fault. The fault lines were modified after Bense et al. (2003).

start of the late Oligocene along a well-pronounced fault (zone), which enabled continuous sedimentation across the Oligocene/Miocene boundary compared to more discontinuous sedimentation with hiatus(es) in the Campine area along its western flank.

In the Miocene, subsidence shifted towards the northern RVG (Michon et al., 2003) and sedimentation became more proximal in a south-eastern direction (Hager et al., 1998), which may give rise to larger sedimentary hiatuses in the central and southern RVG. One of the few boreholes that penetrates the Oligocene/Miocene boundary in the central RVG is the Molenbeersel borehole (DOV kb18d49w-B226, BGD 049W0226; Fig. 1B), which unfortunately lacks age information. Estimates of possible hiatuses in the Veldhoven Formation of this key borehole could therefore only be indirectly derived from log-interpretations and correlations with similar stratigraphic successions that were biostratigraphically analysed in nearby areas (Hager et al., 1998; Verbeek et al., 2002; Deckers, 2016). The uncertainty of these indirectly derived age interpretations is expressed by the different views that each of the above-mentioned authors has on the age of the Veldhoven Formation and its members in this borehole. This is unfortunate, as it not only causes major uncertainty on the late Oligocene to early Miocene depositional evolution but also the

tectonic evolution of the central RVG, as Deckers (2016) identified an important change in fault kinematics in this stratigraphic interval of the Molenbeersel borehole.

The Veldhoven Formation, originally defined by NAM & RGD (1980), was amended by Van Adrichem Boogaert & Kouwe (1993–1997) to include a coarsening upward sequence previously considered part of the former Breda Formation (now separated by the mid-Miocene Unconformity – MMU – into the formations of Groote Heide and Diessen by Munsterman et al., 2019). The Someren Member was introduced as a new member to accommodate the sands and sandy clays that constitute the upper part of the Veldhoven Formation in the south-eastern Netherlands (Van Adrichem Boogaert & Kouwe, 1993–1997). The Veldhoven Clay Member (middle unit of the Veldhoven Formation) originally also defined by NAM & RGD (1980) is recently renamed (to avoid confusion with the same name for the formation) in the Wintelre Member by the Dutch Stratigraphic Commission and officially published in the Dutch Stratigraphic Nomenclature online (Stratigraphic Nomenclature of the Netherlands online, 2021; June 2021).

The Chattian succession in Belgium was recorded in the past century by correlation of their mollusc faunas with the type Chattian in Germany (Schmitz & Stainier, 1909; Stainier, 1911).




Fig. 2. Schematic cross-section of the upper Oligocene to middle Miocene lithostratigraphy and chronostratigraphy. (A) from the Antwerp area (left side), across the Campine area towards the northern Roer Valley Graben (RVG) (right side). (B) Along the RVG, from north (left side) to south (right side). The lithostratigraphy is modified after Van Adrichem Boogaert & Kouwe (1993–1997), Deckers and Louwye (2019) and Munsterman et al. (2019). The location of these cross-sections is shown on Fig. 1B. The NSO dinocyst zones are those of Van Simaëys et al. (2005), while the M-dinocysts zones are those of Munsterman et al. (2019).

Consequently, the Voort Formation, for example in the Campine area, was established and assigned to a Chattian age. This succession becomes thicker in the RVG (borehole Molenbeersel), showing a more continuously development in clayey and sandy

units above the Voort Sands. Lithostratigraphically, this sequence is similar to the more extensively studied Veldhoven Formation in the Netherlands, where biostratigraphy revealed that the Veldhoven Formation grades into the early Miocene, crossing

the Chattian-Aquitainian boundary, and is separated from middle Miocene deposits by the Early Miocene Unconformity (EMU; Munsterman et al., 2019). Hence, it was proposed to harmonise Belgian and Dutch stratigraphic nomenclatures, making the more complete Veldhoven Formation applicable both in the Campine area and the Belgian and Dutch RVG (Dusar & Vandenberghe, 2020). After this new classification scheme, the Belgian Voort Formation became the Voort Member as the lower part of the Veldhoven Formation, of which the middle Wintelre clayey and upper Someren sandy members are only recognised in the RVG (Dusar & Vandenberghe, 2020).

Two boreholes located south and north the Molenbeersel borehole, respectively, that are provided with biostratigraphic age interpretations and log data of the Veldhoven Formation are the Limbricht (NLOG code: LBR-01) and Herkenbosch (Dinoloket code: B58G0192) boreholes (Fig. 1B). These boreholes were analysed for their content of dinoflagellate cysts by Munsterman (1997a, b) and Munsterman (1998), respectively. The aims of this study are (1) to update the results of Munsterman (1997a, b) and Munsterman (1998) to the biozonation currently in use (Munsterman et al., 2019), (2) to perform additional dinoflagellate cyst analyses for better insight in the late Oligocene to early Miocene depositional evolution of the Veldhoven Formation in the central RVG at the location of these boreholes, and (3) to correlate the studied boreholes with the previously biostratigraphically analysed Groote Heide borehole (Dinoloket code: B58F0064) in the Venlo Graben, located to the north of the Herkenbosch borehole, and with the key Molenbeersel borehole in the central RVG to provide a better understanding of the late Oligocene to early Miocene depositional environments in the area of the Roer Valley Rift System.

Geological background

The Roer Valley Rift System developed as a north-west trending branch of the Rhine-Graben-System (Ziegler, 1994) from the late Oligocene onwards, throughout the south-eastern part of the Netherlands, the northeastern part of Belgium and adjacent parts of Germany. In the Netherlands and Belgium, the Roer Valley Rift System can be subdivided into the Campine Block in the west, the RVG in the centre and the Peel High and Venlo Graben in the east (Fig. 1B). These blocks are bounded by major NW-SE striking fault systems. The Campine Block is separated from the differentially subsiding RVG by the Feldbiss Fault System (Dusar et al., 2001). The Peel High is a horst block (relatively high platform, Van den Berg, 1994) that is separated from the RVG by the Peel Boundary Fault and from the Venlo Graben by the Tegelen Fault (Van Adrichem Boogaert & Kouwe, 1993–1997). The Viersen Fault is the principal displacement zone that separates the Venlo Graben and the northeastern Krefeld Block (Munsterman & Brinkhuis, 2004). The RVG itself is about 30 km wide and 130 km long and is bounded by NW-SE striking border fault systems with vertical throws of several hundred metres. Subsidence strongly increased in the RVG during the Chattian initial rifting (Demyttenaere, 1989; Geluk, 1990). The start of the Chattian also marks an important decrease in sea level that resulted in a shift in lithology from the clayey silts of the Steensel Member in the late Rupelian to the glauconite- and fossil-rich sands of the Voort Member (Van Simaey, 2004; Van Simaey et al., 2004; De Man et al., 2010), which forms the lower part of the Veldhoven Formation (Fig. 2). The base of the Voort Member is characterised by marginal, brackish-marine

conditions, with water depths not exceeding 20 m, while higher up in the Chattian successions, conditions return to a normal marine environment (De Man & Van Simaey, 2004). In large parts of the Campine area, the (upper) Chattian and Aquitainian strata are represented by a hiatus (Louwye, 2005) that coincides with a gravel layer (Munsterman & Deckers, 2020; Fig. 2A). At the base of the Miocene in the Limburg Campine (Schmitz & Stainier, 1909), at the base of the Bolderberg (Halet, 1920), and at the base of the Kakert Member (Van Staaldunin & Zagwijn, 1975; Kuyll, 1980), this gravel bed is called the Elsloo Gravel Bed. This hiatus is known from large parts of the North Sea Basin and is related to the Savian tectonic phase (Utescher et al., 2000; Wong et al., 2001; Verbeek et al., 2002; Munsterman & Brinkhuis, 2004; Knox et al., 2010; Rasmussen et al., 2010; Eidvin et al., 2014; King et al., 2016; Dybkjær et al., 2020). At this boundary, an important change in the direction of maximum extension took place, which also coincides with an important shift in depocenter towards the northern RVG (Michon et al., 2003) and the stronger subsidence along the larger fault systems at the expense of the smaller ones in the RVG (Deckers, 2016). Around the Oligocene/Miocene (O/M) boundary, a phase of high $\delta^{18}\text{O}$ occurred, interpreted as a brief period of Antarctic ice sheet expansion, associated with global cooling (Steinthorsdottir et al., 2020). The $\delta^{18}\text{O}$ maximum is referred to as Mi-1 (Miller et al., 1991) or O/M (Climate) Transition (OMT; Beddow et al., 2018). Current studies show that the highest $\delta^{18}\text{O}$ values occur in the latest Oligocene (Liebrand et al., 2016). The event is recorded at several locations world-wide, indicating that cooling can be correlated on a global scale. A climatic deterioration is supported by investigations of successions around Antarctica, which point to ice sheets covering large parts of Antarctica around the O/M transition (e.g. Kennett, 1977; Naish et al., 2008; Barret, 2009). The storage of water into ice resulted in a pronounced eustatic sea-level fall of 50–70 m at this time (e.g. Haq et al., 1987; Miller et al., 2005). The North Sea Basin probably became a (semi-)enclosed basin during the latest Oligocene-earliest Miocene, with a narrow connection to the North Atlantic Ocean between the Shetland Isles and Norway (Rasmussen et al., 2008; Knox et al., 2010). In the latest Chattian, a shallowing occurred, probably reflecting the OMT glaciation event, as was also recorded in Denmark (Dybkjær & Rasmussen, 2007; Dybkjær et al., 2012). The last occurrences of the dinoflagellate cysts *Distatodinium biffii* and *Chiropteridium* spp. were used to identify the broader O/M boundary interval. This interval is referred to *Chiropteridium galea* dinocyst Zone in Denmark and Zone SNSM1 in the Netherlands. In very shallow marine settings, that is along the southern boundary of the North Sea Basin, this sea-level fall caused unconformities. The range of the Oligocene/Miocene hiatus decreases from west to east across the Campine area (Munsterman & Deckers, 2020; Fig. 2A). In the northern RVG, this hiatus is absent or of limited extent, as the Voort Member (c. 27.3–23.4 Ma) here is overlain by the uppermost Chattian to lower Aquitainian clays of the Wintelre Member (c. 23.4–21.5 Ma; Munsterman et al., 2019). During the late Aquitainian to middle Burdigalian, the Wintelre Member was transgressed and covered by the glauconitic sands of the Someren Member (c. 21.5–17.5 Ma), the uppermost member of the Veldhoven Formation. Contrary to the Wintelre Member, the age equivalent deposits to the Someren Member in the RVG, included in the lower part of the Berchem Formation (Fig. 2), covers most of the Campine area.

During the late Burdigalian, the Veldhoven Formation was overlain by the Groote Heide Formation (Munsterman et al., 2019; Fig. 2). In the distal northern RVG, the Groote Heide




Fig. 3. Correlation panel between the gamma-ray log of the analysed boreholes by this study in the central Roer Valley Rift System. The (NSM and NSO) dinocyst biozones as interpreted by Munsterman et al. (2019) for the Groote Heide borehole, by Munsterman (1997a, b) and this study for the Limbricht borehole and by Munsterman (1998) for the Herkenbosch borehole are shown next to the gamma- (and resistivity-) logs of these boreholes. The location of this panel is indicated on Fig. 1B.

Formation consists of homogeneous sequences of glauconite-rich sands that were deposited up to late in the Serravallian. In the more near-shore central RVG, it represents more heterogeneous sequences of the upper Burdigalian silty, glauconitic sands of the Kakert Member, followed by Langhian glauconite-poor and lignite-rich sands of the Heksenberg Member and finally the return of glauconite and decrease of lignite into the Serravallian Vrijherenberg Member (Munsterman et al., 2019; Fig. 2B).

Dataset and methodology

Dataset and correlations

The Limbricht borehole is located in the western flank of the intra-graben area of the central RVG (Fig. 1B). The Herkenbosch borehole is situated further north on the south-eastern part of the Peel High. Besides of lithological descriptions, both boreholes are provided with log data (Fig. 3). A schematic column of the lithological development of the successions (including photos of the sample material) in the boreholes, and interpreted lithostratigraphy, used for this study are publicly provided on internet by the Geological Survey of the Netherlands (DINOloket, n.d.; NLOG, n.d.). Detailed lithological descriptions of air-lifted (semi-cored) boreholes Groote Heide and Herkenbosch are also studied and available on request (DINOloket). For the purpose of this study, gamma-ray log patterns of the successions will be leading. The log data are cross-validated against lithological logs for consistency. Munsterman (1997a) and Munsterman (1998) performed dinoflagellate cyst analyses on some of the samples

from the Oligocene to middle Miocene succession in the Limbricht and Herkenbosch boreholes (palynological reports, including number and type of samples used are available on NLOG). For the interval around the Oligocene/Miocene boundary in the Limbricht borehole, Munsterman (1997b) made a revision of the results of Munsterman (1997a: see report on NLOG).

Based on these data, a dinoflagellate cyst zonation scheme (M zones) was defined for the Miocene in the southern North Sea Basin by Munsterman & Brinkhuis (2004). For the purpose of the present study, the results by Munsterman (1997a, b) and Munsterman (1998) were integrated in the updated version of this dinoflagellate cyst zonation (Munsterman et al., 2019). In addition, the interval between 244.5 and 271.5 m depth in the Limbricht borehole was resampled and the dinoflagellate cyst content was reanalysed for the purpose of the present study.

The gamma-ray logs of the Limbricht and Herkenbosch boreholes are correlated with those of the Groote Heide and Molenbeersel boreholes (Fig. 3). The Groote Heide borehole is located in the Venlo Graben and was selected because it was analysed biostratigraphically (based on dinoflagellate cysts) for the Oligocene and Miocene interval by Munsterman et al. (2019). The Molenbeersel borehole is located in the central RVG and was selected because it is provided with log data over the thick upper Oligocene and Miocene sedimentary successions, crossed by seismic lines and was used as a key borehole in a large number of studies on the sedimentary and tectonic evolution of the central RVG (Demyttenaere & Laga, 1988; Hager et al., 1998; Verbeek et al., 2002; Vandenberghe et al., 2004; Deckers, 2016; Deckers & Munsterman, 2020; Dusar & Vandenberghe, 2020).

Sedimentary Facies




Fig. 4. Schematically displayed depositional environments used as standard for interpretations in this paper (modified after Munsterman et al., 2012; Verreussel et al., 2018). At the base of this scheme, the extent of the depositional environments inferred from palynological analysis are compared with the detailed depositional setting, as can be inferred for instance from core description analysis.

Finally, well Veldhoven-1 (VEH-01) in northern part of the RVG is added, because it comprises the type section of the Veldhoven Formation [see Van Adrichem Boogaert & Kouwe (1993–1997, Annex I–6), for spontaneous potential (SP) and resistivity (RES) logs and lithological development; *DINOloket* and Munsterman (2001) for palynological report (NLOG)].

Dinoflagellate cyst analyses and zonation

Organic-walled dinoflagellate cysts analysis was utilised to support the interpretations of the Limbricht and Herkenbosch boreholes. This type of analysis led to significant improvements in dating and correlation of Neogene successions in NW Europe and beyond, and in understanding their palaeoenvironmental setting (e.g. Powell, 1986, 1992; Head et al., 1989; Brinkhuis et al., 1992; Zevenboom, 1995; De Verteuil & Norris, 1996; Head, 1998; Dybkjær & Rasmussen, 2000; Louwye, 2002; Louwye et al., 2004; Köthe, 2007; Köthe et al., 2008; De Schepper & Head, 2009; Dybkjær & Piasecki, 2010; Anthonissen, 2012; Quaijtaal et al., 2014; De Schepper et al., 2015, 2017; Dybkjær et al., 2019; Dybkjær et al., 2020). Figure 4 schematically displays the depositional environments used as standard for interpretations in this paper.

The potential of dinoflagellate cyst analyses in (bio)stratigraphic differentiation for upper Palaeogene and Neogene successions in the Netherlands has been proven (e.g. Munsterman & Brinkhuis, 2004; Van Simaëys et al., 2005; Munsterman et al., 2019). Standard palynological techniques, including HCL and HF digestion, and 15 µm sieving, were applied. No oxidation was used. The slides were mounted in glycerine jelly. Dinoflagellate

cyst taxonomy is according to that cited in Fensome et al. (2019). One microscope slide per sample was counted until a minimum of 200 palynomorphs (spores, pollen, and dinoflagellate cysts) had been identified. The remainder of the slides were scanned for rare taxa. Miscellaneous fossils (e.g. *Pediastrum*, *Botryococcus*) were also counted, but kept outside the total sum of 200 specimens.

The Miocene dinoflagellate cyst zonation is based on Munsterman & Brinkhuis (2004), recalibrated to the Geological Time Scale of Ogg et al. (2016) by Munsterman et al. (2019; Fig. 5). This zonation is based on consistent dinoflagellate cyst events (mainly on last occurrence datums) from available peer-reviewed palynological contributions from NW Europe and also uses a global compilation calibrated to palaeomagnetic, calcareous plankton and/or foraminifera/bolboforma stratigraphy (Zevenboom, 1995; De Verteuil and Norris, 1996; Van Leeuwen, 2000; and references therein). The age assessments have been cross-validated by correlation to NW European sea-level fluctuations (Hardenbol et al., 1998; Munsterman & Brinkhuis, 2004). For the Oligocene the dinoflagellate cyst zonation North Sea Oligocene (NSO) scheme of Van Simaëys et al. (2005) was used (Fig. 5).

Results

Borehole Limbricht-01 (RVG; Fig. 3)

Age assessment

The palynological (dinoflagellate cyst) analyses for the samples at different measured depths along the Limbricht borehole indicated the following results (Table 1):




Fig. 5. Composition of relevant parts of dinoflagellate cyst zonation schemes of Munsterman et al. (2019) and Van Simaëys et al. (2005).

The dating is based on:

- *1: Last occurrence datums (LOD's) of *Distatodinium paradoxum* and *Apteodinium spiridoides* both at 224.5 m (Munsterman et al., 2019).
- *2: The occurrences of several *Cordosphaeridium cantharellum* at 255 m (Munsterman & Brinkhuis, 2004; Dybkjær & Piasecki, 2010).
- *3: LOD of *Ectosphaeridium picenum* at 266 m (Zevenboom, 1995; Munsterman et al., 2019).
- *4: LOD of *Thalassiphora pelagica* at 271.5 m (Dybkjær & Piasecki, 2010).
- *5: The presence of dinoflagellate cyst *Homotryblum vallum* at 283 m (De Verteuil & Norris, 1996; Dybkjær & Piasecki, 2010; Munsterman et al., 2019).
- *6: LOD of *Chiropteridium galea* at 291.5 m (De Verteuil & Norris, 1996; Munsterman et al., 2019). The LOD of *Hystriochokolpoma pseudoceanica* at 291.5 m confirms the age assessment (Zevenboom, 1995). Reworking (*Deflandrea phosphorica*, *Enneadocysta pectiniformis*, *Wetzelella* spp.) from the Rupelian is recorded.
- *7: LOD of *Distatodinium biffii* at 322.5 m (Brinkhuis et al., 1992; Van Simaëys et al., 2005; Dybkjær & Piasecki, 2010).

- *8: LOD's of *Areoligera semicircularata* at 344.5 m, *Deflandrea phosphoritica* at 357.5 m, *Glaphyrocysta* cf. *semitecta* at 344.5 m and *Wetzelella symmetrica* at 344.5 m (Van Simaëys et al., 2005; Sliwinska, 2019). At the base of the current interval, 357.5–361.5 m, reworking from Late Jurassic – Early Cretaceous successions (*Chlamydophorella nyei*, *Dingodinium cerviculum*, *Gochteodinium mutabilis* and *Perisseiasphaeridium* spp.) is noticed.
- *9: LOD of *Achilleodinium biformoides* at 392 m (Stover and Hardenbol, 1994).

Stratigraphy, log patterns and depositional environments

The low gamma-ray values of the Steensel Member on top of the high gamma-ray values of the Boom Member are a good reference for the identification of these Rupelian strata. The Steensel Member shows a late Rupelian age based on the NSO5a dinoflagellate cyst biozone at 392–434.5 m. High numbers of the genus *Homotryblum* at the top of the Steensel Member indicate marginal to restricted marine conditions (Dybkjær, 2004). The Steensel Member is correlated to the Eigenbilzen Member in Belgium (Van Simaëys et al., 2005; Stratigraphic Nomenclature NL online, 2021).

Table 1. Limbricht-01 borehole.

Interval m (measured depth)	Age and dinocyst zonation	Lithostratigraphic interpretation
224.5–254	mid-Miocene, mid Langhian, Zone SNSM6, or older *1	Kakert Mbr (Groote Heide Fm)
255–257	early Miocene, mid-Burdigalian, Zone SNSM3 *2	Someren Mbr (Veldhoven Fm)
266–271.5	early Miocene, early Burdigalian, Zone SNSM2 *3	Someren Mbr (Veldhoven Fm)
271.5–274.5	early Miocene, base Burdigalian, Zone SNSM 2 *4	Someren Mbr (Veldhoven Fm)
283–291.5	early Miocene, late Aquitanian, base Zone SNSM 2 *5	Someren Mbr (Veldhoven Fm)
291.5–311	latest Chattian-early Aquitanian, Zone SNSM1 *6	Wintelre Mbr (Veldhoven Fm)
322.5–326	late Oligocene, Chattian, Zone NSO8 *7	Voort Mbr (Veldhoven Fm)
344.5–361.5	late Oligocene, Chattian, Zone NSO6 *8	Voort Mbr (Veldhoven Fm)
392–434.5	early Oligocene, late Rupelian, Zone NSO5a, or older *9	Steensel Mbr (Rupel Fm)

The transition from the Steensel Member towards the superjacent Voort Member of the Veldhoven Formation coincides with an increase in gamma-ray values as the result of an increase in clay content according to the borehole descriptions and resistivity log (the latter is not vectorially available and therefore not included in Fig. 3 of this study).

The increase in clay content may indicate a return of more open marine conditions. Indeed, an increase in relative abundance of *Spiniferites* and in the species diversity at 344.0–348.0 m suggests a development towards a more open marine depositional environment in the basal part of the Voort Member (Brinkhuis, 1994). The common occurrence of *Lingulodinium machaerophorum* and *Homotryblum floripes/plectilum* indicates, however, a near-shore depositional environment (Dybkjær, 2004; Dybkjær et al., 2019). This interval is dated as early Chattian based on the presence of the NSO6 dinoflagellate cyst biozone.

Within the lower part of the Voort Member, gamma-ray values show a gradual increase upward, which could be indicative for a continued deepening depositional environment. Maximum gamma-ray values are reached at 324–330 m, which may thus represent the maximum flooding surface. Indeed, the high numbers of *Spiniferites* and very rare occurrences of the genus *Homotryblum* at 322.5–326 m depth confirm a more open marine depositional environment (Brinkhuis, 1994). The latter interval was dated late Chattian based on the NSO 8 dinoflagellate cyst biozone.

From 324 m to 309 m, the gamma-ray values decrease upward. Decreasing gamma-ray values are indicative for a decrease in clay content, which – in view of the cross-correlation with lithofacies analysis – could be interpreted as a shallowing up. This shallowing is confirmed by the presence of P1 and P2 macerals (relatively dark coloured and large pieces of organic (continental) debris with low buoyancy properties) in the interval between 305 and 326 m (Whittaker, 1984). At 309 m, the gamma-ray values strongly increase towards a high plateau that continues up to 302 m, then drop towards lower values before rising again towards high gamma-ray values around 296 m (Fig. 3). The high gamma-ray values coincide with increase in clay content. In contrast to the interpretation of the results for the basal part of the Voort Member, that is more clay reflecting a phase of deepening of the depositional environment, here we interpret the results as indicating the establishment of lagoonal conditions for the clayey interval 305–296 m. The increase in clay content correlates with a distinct increase in abundance of *Homotryblum floripes/plectilum* (41% of the total sum of dinoflagellate cysts) and

Lingulodinium machaerophorum (6%). *Paralecaneia* is also represented (4%). This palaeoenvironmental interpretation is supported by the regional geological development indicating a sea-level fall (Munsterman & Deckers, 2020 and refs. herein). The Mi-1 glacial event resulted in a major global eustatic sea-level fall at the Oligocene-Miocene boundary (Miller et al., 2020). This interval with high gamma-ray values between 309 and 293 due to increased clay content in the central part of the Veldhoven Formation is typically identified as the Wintelre Member. The age of this interval is latest Chattian to earliest Aquitanian, based on the identification of the M1 dinoflagellate cyst biozone. Reworking (ca. 3% of the total sum of palynomorphs) with an origin from Carboniferous and Eocene to lower Oligocene deposits was recorded.

The boundary between the Wintelre and Someren Members is picked at 293 m, where gamma-ray values have returned to lower values (Fig. 3). From here, the gamma-ray values gradually increase upward towards 272 m. At sample locations at 291.5–283.0 m and 274.5–271.5 m depths, the abundant presence of *Homotryblum floripes/plectilum* (43% of the total sum of dinoflagellate cysts) remains indicating lowered salinity/restricted marine (possibly lagoonal) environment. This entire interval belongs to the late Aquitanian to early Burdigalian M2 dinoflagellate cyst biozone. The zone which has on average higher gamma-ray values between 272 and 257 m coincides with a more clayey-silty zone in the upper part of the Someren Member. The restricted marine influence has disappeared here and the genus *Spiniferites* dominates (56% of the total sum dinoflagellate cysts) the palynospectrum again, pointing to open marine conditions. In the uppermost part of the Someren Member at sample location 255.5–257 m, the middle Burdigalian M3 dinoflagellate cyst biozone was identified. The dominant palynomorphs in the associations hardly changed, showing for example 60% *Spiniferites* spp.

The top of the Someren Member is picked at the turn-around point from increasing to decreasing gamma-ray values in gamma-ray values at 257 m. Here, the Someren Member of the Veldhoven Formation changes into the Kakert Member of the Groote Heide Formation. In the Kakert Member, the gamma-ray values continue to decrease, with drops at 250 m and 228 m towards very low gamma-ray values of the superjacent Heksenberg Member. Our dinoflagellate cyst analyses show that the Kakert Member has an age not younger than dinoflagellate cyst zone M6 and no older than dinoflagellate cyst biozone M4. A dominance of terrestrial spores and pollen indicates that the terrestrial influence increases. Furthermore, the variety in marine species is reduced,

Table 2. Herkenbosch borehole.

Interval m (measured depth)	Age and dinocyst zonation	Lithostratigraphic interpretation
64.2–67.2	mid-Miocene, mid Langhian, Zone SNSM6, or older *1	Kakert Mbr (Groote Heide Fm)
70.2–72.2	early Miocene, mid-Burdigalian, Zone SNSM3 *2	Someren Mbr (Veldhoven Fm)
78.2–82.1	early Miocene, late Aquitanian, base Zone SNSM 2 *3	Someren Mbr (Veldhoven Fm)
88.7–91.6	latest Chattian-early Aquitanian, Zone SNSM1 *4	Wintelre Mbr (Veldhoven Fm)
109.75–112.75	late Oligocene, Chattian, Zone NSO7 *5	Voort Mbr (Veldhoven Fm)
121.75–144.25	late Oligocene, Chattian, Zone NSO6 *6	Voort Mbr (Veldhoven Fm)
148.75–151.25	late Oligocene, earliest Chattian, NSO5b, or older *7	Voort Mbr (Veldhoven Fm)
181.25–184.25	early Oligocene, late Rupelian, Zone NSO4a *8	Steensel Mbr (Rupel Fm)

while *Paralecaneilla*, a near-coastal marine indicator, increase in number. All these observations point towards a coastal marginal marine setting (Louwey et al., 2010; Dybkjær et al., 2019).

Borehole Herkenbosch (Peel High; Fig. 3)

Age assessment

The palynological analyses for the samples from the borehole Herkenbosch gave the following results (see Table 2):

The dating is based on:

- *1: LOD's of *Distatodinium paradoxum* and *Apteodinium tectatum* both at 64.2 m (Munsterman et al., 2019).
- *2: LOD of *Cordosphaeridium cantharellum* at 70.2 m (Munsterman & Brinkhuis, 2004; Dybkjær & Piasecki, 2010).
- *3: LOD's of *Ectosphaeridium picenum* (Zevenboom, 1995; Munsterman et al., 2019), *Thalassiphora pelagica* (Dybkjær & Piasecki, 2010), and the presence of dinoflagellate cyst *Homotryblum vallum* all at 78.2 m (De Verteuil & Norris, 1996; Munsterman et al., 2019).
- *4: LOD's of *Chiropteridium galea* and *Hystrichokolpoma cinctum* both at 88.7 m (De Verteuil & Norris, 1996; Munsterman et al., 2019).
- *5: LOD of *Glaphyrocysta cf. semitecta* at 109.75 m (Van Simaey et al., 2005).
- *6: LOD's of *Areoligera semicirculata* at 121.75 m, *Deflandrea phosphorica* at 141.75 m, and *Wetzeliella symmetrica* at 141.75 m (Van Simaey et al., 2005).
- *7: LOD of *Rhombodinium draco* at 148.75 m (Van Simaey et al., 2005).
- *8: LOD's of *Achilleodinium biformoides* and *Enneadocysta pectiniformis* both at 181.25 m (Stover & Hardenbol, 1994; Van Simaey et al., 2005).

Stratigraphy, log pattern and depositional environment

Low gamma-ray values of the sandy Steensel Member, interval 163.25–217.75 m, on top of the high gamma-ray values of the clayey Boom Member, interval 217.75–310.1 m, are once again good reference for the identification of the Rupelian strata. The Steensel Member is inferred as late Rupelian, based on the presence of the NSO 4a-5b dinoflagellate cyst interval biozones. The interpretation fits with Van Simaey et al. (2005), more detailed showing dinoflagellate zone NSO5b for the base of the Voort Member and NSO5a for the Eigenbilzen Formation. The abundance of the genus *Dapsilodinium* (up to 35% of the total sum of

dinoflagellate cysts) at the top of the Steensel Member (interval 163.25–178.25 m) indicates marginal to restricted marine conditions (Brinkhuis, 1994; Munsterman & Brinkhuis, 2004).

The transition from the Steensel Member towards the superjacent Voort Member of the Veldhoven Formation shows a sharp increase in gamma-ray values, unlike in borehole Limbricht, where a more gradual transition occurs. This increase in gamma-ray values is related to the upward increased clay content, which is expressed by an upward decrease in resistivity values. The increase in clay content coincides with a development towards more open marine conditions. Indeed, an increase in the relative abundances of the dinoflagellate cyst genera *Spiniferites*, *Systematophora* and in dinoflagellate cyst species diversity suggests an increased sea level in the Voort Member (Brinkhuis, 1994).

About 20 m above the base, the gamma-ray values increase and resistivity values decrease upward towards a 20 m thick clayey interval of early Chattian age (dinoflagellate cyst biozone NSO 6). From this clayey interval, the gamma-ray values decrease and the resistivity values increase upward again towards the upper part, more sandy part of the Voort Member, suggesting gradual shallowing upward. Despite the upward coarsening trend, shallow to open marine conditions still prevail at depth 103.75–100.75 m, mirrored by a relatively common occurrence of the genera *Operculodinium*, *Hystrichokolpoma* and 37% of *Spiniferites*. The interval between 148.75 and 109.75 m is dated as Chattian based on the presence of the NSO6 and NSO7 dinoflagellate cyst biozones.

From 101 m upward, the gamma-ray values increase again and the resistivity values flatten in the top of the Voort Member. At 92 m depth a sandstone is present. From this level and upward, an increase in gamma-ray values takes place, together with a sharp upward decrease in resistivity values. This boundary marks the transition from sand of the Voort Member towards the clay of the Wintelre Member. The interval of the Wintelre Member between 91.6 and 88.7 m is dated as latest Chattian to earliest Aquitanian (dinoflagellate cyst zone SNSM 1). As was the case in the Limbricht borehole, also here a lagoonal depositional setting is interpreted by palynological analysis of the clays of the Wintelre Member. Very high values of genus *Homotryblum* (>50% of the total sum dinoflagellate cysts) strongly indicate the presence of restricted marine conditions (Dybkjær, 2004).

From about 84 m upward, the resistivity values increase and the gamma-ray values decrease upward, which marks the transition from the silty clays of the Wintelre Member towards the sandy clays and sands of the Someren Member. Restricted marine conditions prevail at the base (82.1–78.2 m) of the Someren Member,

which was dated as dinoflagellate cyst biozone SNSM2 (late Aquitanian). Higher in the Someren Member, at depths between 72.2 and 70.2 m, the relative abundance of *Homotryblium* decreases and the diversity of dinoflagellate cyst species increases. The near-coastal marine dinoflagellate cyst genus *Apteodinium* comprises up to 15% of the total dinoflagellate cyst assemblage, which indicates a return to open marine conditions (Brinkhuis, 1994). This higher part of the Someren Member is dated as dinoflagellate cyst biozone SNSM3 (mid-Burdigalian in age).

The transition from the Someren Member (Veldhoven Formation) to the Kakert Member of the Groote Heide Formation is marked by the appearance of humic fragments. The boundary is expressed by lowering of gamma-ray values combined with the start of a further increase in values of the resistivity log. The boundary further coincides with a shift from dinoflagellate cyst biozone SNSM3 (middle Burdigalian) in the upper part of the Someren Member towards dinoflagellate cyst biozone SNSM4-6 (late Burdigalian-early Langhian) in the Kakert Member. Sea level is decreasing from the Someren Member towards the Kakert Member as shown by the occurrence of an acme of the marginal marine dinoflagellate cyst genus *Polysphaeridium* at depth 67.2–64.2 m (Brinkhuis, 1994).

Borehole correlations (Fig. 3)

Based on lithological descriptions and the gamma-ray log pattern, the Veldhoven Formation and its three members were identified and correlated between the Groote Heide, Herkenbosch, Limbricht and Molenbeersel boreholes, located in the Venlo Graben, on the Peel High and in the RVG, all part of the central Roer Valley Rift System (Fig. 1). For the Groote Heide, Herkenbosch and Limbricht boreholes, the three members of the Veldhoven Formation were identified for the first time in the present study. For the Molenbeersel borehole, the three-fold subdivision of the Veldhoven Formation was proposed by Broothaers et al. (2012).

Correlations with borehole Groote Heide (the Netherlands)

In the Groote Heide borehole, from which the dinoflagellate cyst content was studied previously by Munsterman & Brinkhuis (2004), the basal Veldhoven Formation is underlain by late Rupelian silty clay instead of sand as is the case for the Limbricht and Herkenbosch boreholes. Therefore, the lower boundary of the Veldhoven Formation in the Groote Heide borehole does not coincide with a major change in gamma-ray patterns. The gamma-ray values at the Steensel/Voort Member boundary are also influenced by a lithified shell bed at interval depth 642.25–643.7 m. The Steensel Member, below this bed, is once again interpreted as dinoflagellate cyst Zone NSO5a.

A three-fold subdivision of the Veldhoven Formation can be made, based on the lithological development in the Groote Heide borehole, albeit the gamma-ray pattern is much more subtle than within the Limbricht and Herkenbosch boreholes.

- Lower part or Voort Member consists of sandy clay in the lower part, which is reflected by relatively high gamma-ray values, gradually changing upward to very fine sand, as seen as lower gamma-ray values. The Voort Member is referred to the biozones NSO5b–NSO8. The palaeoenvironment is interpreted as fluctuating marginal to open marine. Shallow marine conditions are indicated in particular by the dinoflagellate cyst genus *Delflandrea* and less frequently also by the genus *Wetzeliella*

(Brinkhuis, 1994). These genera disappear at c. 590 m. Above this level, the more open marine dinoflagellate cyst genus *Spiniferites* increases occasionally in relative abundance, interspersed with increases in shallow and restricted marine species. Incidental assemblages show high concentrations of the genus *Homotryblium*, representing restricted marine conditions (silty clay sample interval at 564.75–566.65 m: 50% of the total sum dinoflagellate cysts). The relative sea level is finally decreasing towards the uppermost of the succession at c. 510 m. Shallow marine genera like *Chiropteridium* and *Apteodinium* relatively increase at the cost of *Spiniferites*. The sediments show an increase in mollusc content.

- Central part or Wintelre Member consists of very fine sand and sandy clay, reflected by high, but also very variable, gamma-ray values. The Wintelre Member is referred to the M1 dinoflagellate cyst zone. The palynofacies indicates restricted marine, possibly lagoonal conditions: the dinoflagellate cyst assemblage within the interval 470–446 m comprises 40–70% *Homotryblium floripes/pectilum*. Compared to the clays in the Voort Member, higher concentrations of shelves are evidently present.
- Upper part or Someren Member consists of fine sand with some silty intercalations and is referred to the dinoflagellate cyst zones M2–M3. It shows moderate and generally upward increasing gamma-ray values. In the basal part (interval 439–425 m) of this lithostratigraphic unit, marginal marine conditions are maintained, shown by relatively high values of *Homotryblium* and *Paralecaniella*, but in the overlying succession these taxa decrease in number and *Spiniferites* becomes the dominant genus, indicating a return to an open marine setting.

The upper boundary of the Veldhoven Formation is characterised by a decrease in gamma-ray values towards the Kakert Member of the Groote Heide Formation (Munsterman et al., 2019), which is very similar to the Limbricht and Herkenbosch boreholes. In the Groote Heide borehole, the upper boundary of the Veldhoven Formation coincides with the boundary between the dinoflagellate cyst zones M3 and M4. The Kakert Member shows the first continuous freshwater influence (continuous occurrence of *Pediastrum*) pointing to a change into an estuarine or an embayment setting.

Correlations with borehole Molenbeersel (Belgium)

The low gamma-ray values of the Steensel Formation on top of the high gamma-ray values of the Rupel Formation are a good reference for correlations of the Rupelian strata between the Limbricht, Herkenbosch and Molenbeersel boreholes. Based on correlations with surrounding boreholes, the low gamma-ray interval of the Molenbeersel borehole was referred to the Eigenbilzen Formation by Vandenberghe et al. (2003). The late Rupelian age for the Eigenbilzen Formation suggested by them and the late Rupelian NSO5a dinoflagellate cyst zone found in the Steensel Formation in the Limbricht and Herkenbosch boreholes in the present study support a correlation between the Belgian Eigenbilzen and the Dutch Steensel Formation.

The transition from the Steensel Member towards the overlying Veldhoven Formation coincides with an increase in gamma-ray values, reflecting an increase in glauconite content. The three-fold subdivision of the Veldhoven Formation in the Molenbeersel borehole into the Voort, Wintelre and Someren Members was introduced by Broothaers et al. (2012). The gamma-ray patterns on which this three-fold subdivision in the Molenbeersel borehole is based are similar to the patterns seen in the Limbricht and

Herkenbosch boreholes: the sandy Voort and Someren members show moderate gamma-ray values and are separated by the clayey, high gamma-ray interval referred to the Wintelre Member.

Correlation of the Wintelre Member in borehole Molenbeersel with the German lithostratigraphic subdivision, represented by Schneider and Thiele hydrostratigraphic codes (S&T) of the Lower Rhine Basin and Chattian age assessment based on correlation with type-sections in the Lower Rhine coal and salt districts, corresponds to S&T01 (Schneider and Thiele, 1965; Hager et al., 1998; Duser & Vandenberghe, 2020). These S&T units in the Köln Formation (Germany) are interpreted as marginal to lagoonal environments (see also Schäfer & Utescher, 2014).

The upper boundary of the Someren Member or Veldhoven Formation coincides with a gamma-ray spike in all three boreholes, and from there on, the gamma-ray values drop towards lower values in the lower part of the Kakert Member of the Groote Heide Formation. The transition from the Kakert Member towards the Heksenberg Member coincides with a further reduction towards very low gamma-ray values in all the above-mentioned boreholes as a result of a strong decrease in clay and glauconite content.

Correlation with holotype section well Veldhoven-1 (VEH-01) in the RVG (the Netherlands)

The holostratotype section of the Veldhoven Formation is established in well Veldhoven (NLOG code VEH-01) interval 860–1103 m (Van Adrichem Boogaert & Kouwe 1993–1997, Annex I-6). This well is drilled in the late 1950s. Unfortunately, the material of the well consists primarily of ditch cuttings, hampering lithological description and biostratigraphic interpretation by caving. Cores are only available at two intervals, viz. 907–913 m and 1041–1047 m. Furthermore, the type-well does not provide a gamma-ray log, but SP and resistivity (RES) logs, which make comparison with other wells challenging (Fig. 6).

In this well, the Veldhoven Formation overlies the Steensel Member of the Rupel Formation and its lower boundary is at the base of a sandy clay bed. Because of all limitations referred to caving, only several cuttings samples are palynologically analysed (Munsterman, 2001). At the top of the Steensel Member at 1105 m, the NSO5b Zone, or older is interpreted, which matches with wells Groote Heide, Herkenbosch and LBR-01. The upper boundary of the Veldhoven Formation (with Groote Heide Formation) at 860 m is interpreted at the shift of decreasing concentrations of silts and clays within the sandy succession. This corresponds to a lowering in the values on the SP log.

The SP log of the Veldhoven Formation in well VEH-01 shows a four-fold subdivision:

- Voort Member, interval 1047–1103 m: Low SP values, including very irregular high peaks in the lowest part. Best described as a stacking of coarsening upward sequences at the scale of tens of metres. The proportion of clays is relatively high in the lower part of the member. The basal clays are also shown in well Groote Heide, sandy clays are shown in the lower part of the other wells, although on the Peel High (well Herkenbosch) and in southern RVG (well LBR-01) the proportion of argillaceous sediments is very low. The middle part of VEH-01 stands out as a homogeneous clean sand, which is also represented on the SP log as low values. The interval is palynologically interpreted as Zone NSO6, like all basal successions of the Voort Member in the other wells. The sands are fine-grained,




Fig. 6. Holostratotype well Veldhoven-1 (VEH-01) sensu Van Adrichem Boogaert & Kouwe (1993–1997) and Stratigraphic Nomenclature of the Netherlands online.

grey-green and glauconitic. Locally, the sands may have a dark grey-green to black colour and comprise glauconites and marine molluscs. The uppermost part is fining upward into silts and becomes more clayey (in particular well-shown in well Molenbeersel).

- A transitional interval 995–1047 m: Serrated pattern of high SP values. The lithostratigraphy of this interval is doubtfully interpreted as Wintelre Member. The cuttings in this succession consist of changing proportions of sands, clays and glauconites. At 1009 m dinocyst Zone NSO8, or older is reached, which is based on the LOD of *Distatodinium biffii* at 1009 m (core

section). The upper part (995–1009 m) is interpreted as dinocyst Zone M1.

- Wintelre Member, interval 995–935 m: generally a very regular high signal is mirrored on the SP log. The sediments are grey to greenish-grey sandy clays, locally with brownish colours. The succession becomes more silty and sandy towards the top. Most typically developed in wells VEH-01 and Molenbeersel. This and most other trends are shown in a condensed manner in well LBR-01. The coarsening upward grain-size trend to the top is also shown in well Groote Heide (Venlo Graben). On the Peel High it is a relatively thin clay interval in between the Voort and Someren members. The dating is latest Chattian-earliest Aquitanian, dinocyst Zone M1.
- An irregular low pattern for the uppermost part corresponds to the Someren Member, interval 860–935 m. The succession consists of clayey-silty sands, like in all other studied wells here. The colour of the sand is greenish-grey, and the glauconite content is low. Marine shells occur locally. The palynological association of the cuttings samples indicates M2 and M3 dinocyst zones, which are in agreement with the recorded zones of the Someren Member in all wells of this study.

Discussion

Age interpretations

The existing biostratigraphic interpretations of the Groote Heide borehole by Munsterman et al. (2019) and of the Limbricht and Herkenbosch boreholes by Munsterman (1997a, b; 1998) combined with the new analyses of the present study provide consistent age information on the different members of the Veldhoven Formation: The Voort Member is of Chattian age (dinoflagellate cyst zones NSO5b-8), the Wintelre Member is of latest Chattian to earliest Aquitanian age (dinoflagellate cyst zone NSM1), and the Someren Member is of late Aquitanian to middle Burdigalian age (dinoflagellate cyst zones NSM2-3). This confirms the ages of the members of the Veldhoven Formation as shown in Figure 8 of Munsterman et al. (2019) for the northern part of the RVG (Fig. 2).

Nearly all the late Oligocene to early Miocene dinoflagellate cyst zones were identified in the Limbricht, Herkenbosch and Groote Heide boreholes excluding zone NSO7 in the Limbricht borehole and zone NSO8 in the Herkenbosch borehole. The missing zone is most probably due to the low sample resolution. It is therefore believed that no major hiatus (of longer duration than a dinoflagellate zone) occurred in the late Oligocene to early Miocene deposits in the central RVG, similar to what was found in the northern RVG by Munsterman et al. (2019).

The correlations of the biostratigraphically analysed Groote Heide, Herkenbosch and Limbricht boreholes with the Molenbeersel borehole allow age estimates for the upper Oligocene-lower Miocene succession in the latter key borehole. These results can then be compared with previously suggested ages for this succession by Hager et al. (1998), Verbeek et al. (2002) and Deckers (2016) as shown in Fig. 7. Similarly to the present study, the previous studies made age estimates for the stratigraphic successions in the Molenbeersel borehole based on correlation of log patterns with nearby boreholes that were provided with age information. The improvement and strength of this study compared to the previous ones is the proximity of the Limbricht borehole to the Molenbeersel borehole and its location in the same

AGE (Ma)	SERIES	STAGE	DINOCYST BIOZONES Munsterman & Brinkhuis (2004)	Hager et al. (1998)	Verbeek et al. (2002)	Deckers (2016)	This study
15	Middle Miocene	Langhian	M7 M6 M5				
	Early Miocene	Burdigalian	M4		Someren		
			M3				
20			M2			Someren Wintelre Clay	Someren Wintelre Clay
		Aquitanian	M1				
25	Lower Oligocene	Chattian			Wintelre Clay Voort	Voort	Voort
30	Early Oligocene	Rupelian					

Fig. 7. Comparison of the past and presently presumed ages of the Veldhoven Formation and its members in the Molenbeersel borehole.

structural entity, namely the RVG. Our datings are not in agreement with those of Hager et al. (1998) who referred to the whole interval here currently interpreted as the Veldhoven Formation to the Chattian. Our results also partly contradict those of Verbeek et al. (2002) since they suggested a Chattian age for the interval here referred to the Wintelre Member and a middle Burdigalian age for the Someren Member. Our results also contradict Deckers (2016) for his proposed Aquitanian age for the entire Someren Member, while his suggested Chattian and latest Chattian to Aquitanian ages for the Voort and Wintelre Members are confirmed by this study. Consequently, the change in fault kinematics in the RVG that Deckers (2016) observed around the stratigraphic level of the Wintelre Member is confirmed by this study to have an age spanning the Oligocene/Miocene boundary.

Depositional environment

The gamma-ray log patterns and relative position of biozones within the members of the Veldhoven Formation vary between the four studied boreholes. This shows that whereas the subdivision of the Veldhoven Formation in its three members was of a regional extent, the members themselves show lateral facies changes. This is in agreement with the results of Deckers (2016) who observed patterns of north-west progradation within both the Voort and Wintelre members in the area south and west of the Molenbeersel borehole. This progradation pattern confirms the idea of progressive infill of the RVG from a south-eastern source area as stated by Hager et al. (1998) whereby the type of sediment is dependent on the interaction between subsidence rates, eustatic sea-level and sediment input. The different subsidence rates and proximity to the sediment source area at the locations of the boreholes could explain the different gamma-ray log patterns for the members of the Veldhoven Formation in these boreholes. For each borehole, variation in lithology within the Veldhoven Formation seem, however, limited to in between the sandy clays and fine sands.

Despite these limited facies variations, the different members were deposited in similar environments, with predominantly shallow marine (fluctuating between open and restricted marine as shown in borehole Groote Heide) conditions in the Voort Member, restricted marine for the Wintelre Member that continue in the lower part of the Someren Member before returning again towards shallow marine conditions in the upper part of the Someren Member.

The marginal marine conditions during the earliest Chattian (NSO5b) followed by a short-term deepening in the early Chattian (NSO6) are in agreement with analyses of boreholes in northern Belgium by De Man & Van Simaey (2004). The latter authors stated that the early Chattian Voort Member was deposited in a shallower environment compared to the underlying Eigenbilzen Formation in boreholes in northern Belgium. In the Limbricht and Herkenbosch boreholes, further to the south-east and thus closer to the sediment source, on the other hand, the Steensel Member (Eigenbilzen Formation in Belgium) was deposited in a marginal to restricted marine depositional environment and the Voort Member represents a relative deepening of the depositional environment. The increase in water depths in the basal Chattian Voort Member coincides with an increase in clay content, expressed by an increase in gamma-ray values. For the overlying Wintelre Member, the increase in clay content also reflects a shallowing from shallow marine environments in the upper part of the Voort Member towards restricted marine environments. The clays of the Wintelre Member are therefore interpreted as lagoonal clay. These clays can be distinguished from most clays in the Voort Member by a higher concentration of shells as is obviously demonstrated in the lithological description of borehole Groote Heide. The dinoflagellate cyst assemblages are here dominated by a single genus indicating extreme depositional environments, including salinities that deviate from normal marine conditions, most probably due to minor ventilation by narrow or lack of connection to the Atlantic Ocean. The glacio-eustatic sea-level fall around the Oligocene/Miocene boundary (e.g. Miller et al., 2020; Steinthorsdottir et al., 2020) most likely limited the sea coverage to the strongest subsiding areas, where deposition of the Wintelre Member took place, while non-deposition or erosion occurred in the surrounding highs, hence creating an isolated (sub)basin. The lagoonal depositional environment of the Wintelre Member spanning the Oligocene/Miocene boundary, in the differentially subsiding Roer Valley Rift System, is in agreement with the missing late(st) Chattian-(early) Aquitanian sections at the relatively high flanks of the Rift System as observed by Louwye et al., (2020 and references therein; Fig. 2A).

The lateral persistence of the clayey Wintelre Member (unit S&T01) across all structural settings point to a eustasy-based model (Hager et al., 1998). This is in contrast to the sedimentological model of Schäfer et al. (2005), which proposes a lateral transition from marginal facies of the Köln Formation in the Lower Rhine Basin (Germany) across the shallow marine Dutch Voort Member in the Venlo/Peel Blocks to the deeper water Wintelre (Veldhoven Clay) Member in the RVG of the Netherlands. The latter is shared by the dynamic concept in De Mulder et al. (2003, fig. 172); Wong et al., (2007, fig. 9) suggesting that the Wintelre Member is a deeper marine facies compared to the Someren and Voort members (Dusar & VandenBerghe, 2020). This study disproves the latter vision by indicating restricted marine conditions for the Wintelre Member, interpreted as result of the associated glacio-eustatic sea-level fall around the Oligocene/Miocene boundary. The S&T06 unit (Hager et al., 1998) is not

systematically recorded as a clayey interval of consistent age in our studied wells.

Marginal marine conditions prevailed during the rest of the Aquitanian when the sands in the lower part of the Someren Member were deposited. As sea-level rose, the upper, more clayey part of the Someren Member was deposited in a shallow marine environment, and the flanking highs of the Roer Valley Rift System were transgressed during the Burdigalian (Fig. 2A). During a latest Burdigalian shallowing, the Veldhoven Formation became covered by the sands and clays of the Kakert Member (Groote Heide Formation). Input of freshwater is consistently recorded here.

Conclusions

New information on the bio- and lithostratigraphy and facies from Dutch boreholes Herkenbosch (Peel High) and Limbricht (RVG) and their lithological and gamma-ray log correlation to the boreholes Molenbeersel (RVG, Belgium) and Groote Heide (Venlo Graben, the Netherlands) provide good consistency on stratigraphy and depositional environments for the members of the Veldhoven Formation in the Roer Valley Rift System during the late Oligocene-early Miocene. Based on the new results of this study we were able to convincingly revise the previous views on, in particular, the Wintelre Member not as a deeper marine facies compared to the Someren and Voort members, but representing restricted marine conditions. Dinoflagellate cyst assemblages in the Wintelre Member are mostly dominated by a single genus, indicating a restricted marine depositional setting with salinities that deviate from normal marine conditions, most probably due to minor ventilation by narrow or lack of connection to the Atlantic Ocean. These circumstances would have resulted from the associated glacio-eustatic sea-level fall around the Oligocene/Miocene boundary in concert with the Savian tectonic phase. Both factors limited preserved marine successions to the strongest subsiding areas, where deposition of the Wintelre Member is recorded, while non-deposition or erosion occurred in the surrounding highs, hence creating an isolated (sub)basin.

Acknowledgements. The authors gratefully acknowledge financial support from the Land and Soil Protection, Subsoil, and Natural Resources Division of the Flemish Government. We would like to thank Katleen van Baelen for her excellent work on the figures. The three reviewers, Karin Dybkjær, Noël Vandenberghe and Frank Wesselingh are thanked for their very helpful suggestions and corrections improving the manuscript substantially. Finally we are much obliged to our editor Rick Donselaar (TU-Delft, the Netherlands) for his logistical support and corrections in the text.

References

- Anthoissen, E.**, 2012. A new Miocene biostratigraphy for the northeastern North Atlantic: an integrated foraminiferal, bolboformid, dinoflagellate and diatom zonation. *Newsletters on Stratigraphy* **45**(3): 281–307. doi: [10.1127/0078-0421/2012/0025](https://doi.org/10.1127/0078-0421/2012/0025).
- Barret, J.**, 2009. Cenozoic climate and sea level history from glacial marine strata off the Victoria Land coast, Cape Roberts Project, Antarctica. *In: Hambrey M.J., Christoffersen P., Glasser N.F. & Hubbard B.* (eds): *Glacial sedimentary processes and products*. Blackwell Publishing Ltd (Oxford, U.K): 259–287.
- Beddow, H.M., Liebrand, D., Wilson, D.S., Hilgen, F.J., Sluijs, A., Wade, B.S. & Lourens, L.J.**, 2018. Astronomical tunings of the Oligocene-Miocene transition from Pacific Ocean Site U1334 and implications for the carbon cycle. *Climate of the Past* **14**(3): 255–270. doi: [10.5194/cp-14-255-2018](https://doi.org/10.5194/cp-14-255-2018).
- Bense, V.F., Van Den Berg, E.H. & Van Balen, R.T.**, 2003. Deformation mechanisms and hydraulic properties of fault zones in unconsolidated sediments;

- the Roer Valley rift system, The Netherlands. *Hydrogeology Journal* **11**(3): 319–332.
- Brinkhuis, H.**, 1994. Late Eocene to Early Oligocene dinoflagellate cysts from the Priabonian type-area (Northeast Italy): biostratigraphy and paleoenvironmental interpretation. *Palaeogeography, Palaeoclimatology, Palaeoecology* **107**: 121–163.
- Brinkhuis, H., Powell, A.J. & Zevenboom, D.**, 1992. High-resolution dinoflagellate cyst stratigraphy of the Oligocene/Miocene transition interval in northwest and central Italy. *In*: Head M.J. & Wrenn J.H. (eds): Neogene and quaternary dinoflagellate cysts and acritarchs. American Association of Stratigraphic Palynologists Foundation (Dallas): 210–258.
- Broothaers, M., Deckers, J., Lagrou, D. & Matthijs, J.**, 2012. 3D-lagenmodel van de Tertiaire afzettingen in de Roerdalslenk in Vlaanderen. VITO-rapport 2012/SCT/R/191: 58 pp.
- De Man, E. & Van Simaey, S.**, 2004. Late Oligocene Warming Event in the southern North Sea Basin: benthic Foraminifera as paleotemperature proxies. *Netherlands Journal of Geosciences* **83**(3): 227–239. doi: [10.1017/S0016774600023520](https://doi.org/10.1017/S0016774600023520).
- De Man, E., Van Simaey, S., Vandenberghe, N., Harris, W.B. & Wampler, J.M.**, 2010. On the nature and chronostratigraphic position of the Rupelian and Chattian stratotypes in the southern North Sea Basin. *Episodes* **33**(1): 3–14. doi: [10.18814/epiugs/2010/v33i1/002](https://doi.org/10.18814/epiugs/2010/v33i1/002).
- De Mulder, E.F.J., Geluk, M.C., Ritsema, I.L., Westerhof, W.E. & Wong, T.E.**, 2003. De ondergrond van Nederland. Wolters Noordhoff (Groningen): 379 pp.
- De Schepper, S., Beck, K.M. & Mangerud, G.**, 2017. Late Neogene dinoflagellate cyst and acritarch biostratigraphy for Ocean Drilling Program Hole 642B, Norwegian Sea. *Review of Palaeobotany and Palynology* **236**: 12–32. doi: [10.1016/j.revpalbo.2016.08.005](https://doi.org/10.1016/j.revpalbo.2016.08.005).
- De Schepper, S. & Head, M.J.**, 2009. Pliocene and Pleistocene dinoflagellate cyst and acritarch zonation of DSDP Hole 610A, eastern North Atlantic. *Palynology* **33**(1): 179–218. doi: [10.2113/gspalynol.33.1.179](https://doi.org/10.2113/gspalynol.33.1.179).
- De Schepper, S., Schreck, M., Beck, K.M., Matthiessen, J., Fahl, K. & Mangerud, G.**, 2015. Early Pliocene onset of modern Nordic Seas circulation related to ocean gateway changes. *Nature Communications* **6**(1): 8659. doi: [10.1038/ncomms9659](https://doi.org/10.1038/ncomms9659).
- De Verteuil, L. & Norris, G.**, 1996. Miocene dinoflagellate stratigraphy and systematics of Maryland and Virginia. *Micropaleontology* **42**(Supplement): 1–172. doi: [10.2307/1485926](https://doi.org/10.2307/1485926).
- Deckers, J.**, 2016. The Late Oligocene to Early Miocene early evolution of rifting in the south-western part of the Roer Valley Graben. *International Journal of Earth Sciences* **105**(4): 1233–1243. <http://link.springer.com/article/10.1007/2Fs00531-015-1236-5>
- Deckers, J. & Louwye, S.**, 2019. Late Miocene increase in sediment accommodation rates in the southern North Sea Basin. *Geological Journal* **55**(1): 728–736. doi: [10.1002/gj.3438](https://doi.org/10.1002/gj.3438).
- Deckers, J. & Munsterman, D.**, 2020. Middle Miocene depositional evolution of the central Roer Valley Rift System. *Geological Journal* **55**(9): 6188–6197. doi: [10.1002/gj.3799](https://doi.org/10.1002/gj.3799).
- Demyttenaere, R.**, 1989. The post-Paleozoic geological history of north-eastern Belgium. *Mededelingen van de Koninklijke Academie voor Wetenschappen. Letteren en Schone Kunsten België* **51**: 51–81.
- Demyttenaere, R. & Laga, P.**, 1988. Breuken en isohypsenkaarten van het Belgische gedeelte van de Roerdal Slenk. *Belgische Geologische Dienst, Professional Paper* **234**: 20 pp.
- DINOLOket**, n.d. <https://www.dinoloket.nl>. <https://www.dinoloket.nl/sites/default/files/nomenclator/typelocaties/VEH-01.pdf>
- Dusar, M., Rijpens, J., Sintubin, M. & Wouters, L.**, 2001. Plio-Pleistocene fault pattern of the Feldbiss fault system (southern border of the Roer Valley Graben, Belgium) based on high resolution reflection seismic data. *Netherlands Journal of Geosciences* **80**: 79–93.
- Dusar, M. & Vandenberghe, N.**, 2020. Upper Oligocene lithostratigraphic units and the transition to the Miocene in North Belgium. *Geologica Belgica* **23**(3-4): 113–126. doi: [10.20341/gb.2020.009](https://doi.org/10.20341/gb.2020.009).
- Dybkjær, K.**, 2004. Morphological and abundance variations in Homotryblium-cyst assemblages related to depositional environments; uppermost Oligocene – Lower Miocene, Jylland. *Denmark Palaeogeography, Palaeoclimatology, Palaeoecology* **206**: 41–58.
- Dybkjær, K., King, C. & Sheldon, E.**, 2012. Identification and characterisation of the Oligocene-Miocene boundary (base Neogene) in the eastern North Sea Basin - based on dinocyst stratigraphy, micropalaeontology and $\delta^{13}\text{C}$ -isotope data. *Palaeogeography, Palaeoclimatology, Palaeoecology* **361-364**: 11–22. doi: [10.1016/j.palaeo.2012.08.007](https://doi.org/10.1016/j.palaeo.2012.08.007).
- Dybkjær, K. & Piasecki, S.**, 2010. Neogene dinoflagellate cyst zonation for the eastern North Sea Basin. *Denmark Review of Palaeobotany and Palynology* **161**: 1–29. doi: [10.1016/j.revpalbo.2010.02.005](https://doi.org/10.1016/j.revpalbo.2010.02.005).
- Dybkjær, K. & Rasmussen, E.S.**, 2000. Palynological dating of the Oligocene - Miocene successions in the Lille Bælt area. *Denmark Bulletin Geological Society Denmark* **47**: 87–103.
- Dybkjær, K. & Rasmussen, E.S.**, 2007. Dinocyst stratigraphy in an expanded Oligocene-Miocene boundary section in the eastern North Sea Basin (the Frida-1 well, Denmark) and correlation from basinal to marginal areas. *Journal of Micropaleontology* **26**: 1–17.
- Dybkjær, K., Rasmussen, E.S., Eidvin, T., Grøsfjeld, K., Riis, F., Piasecki, S. & Śliwińska, K.K.**, 2020. A new stratigraphic framework for the Miocene – Lower Pliocene deposits offshore Scandinavia: a multiscale approach. *Geological Journal* **2020**: 1–27. doi: [10.1002/gj.3982](https://doi.org/10.1002/gj.3982).
- Dybkjær, K., Rasmussen, E.S., Śliwińska, K.K., Esbensen, K.H. & Mathiesen, A.**, 2019. A palynofacies study of past fluvio-deltaic and shelf environments, the Oligocene-Miocene succession, North Sea Basin: a reference data set for similar Cenozoic systems. *Marine and Petroleum Geology* **100**(5): 111–147. doi: [10.1016/j.marpetgeo.2018.08.012](https://doi.org/10.1016/j.marpetgeo.2018.08.012).
- Eidvin, T., Fridtjof, R. & Rasmussen, E.S.**, 2014. Oligocene to Lower Pliocene deposits of the Norwegian continental shelf, Norwegian Sea, Svalbard, Denmark and their relation to the uplift of Fennoscandia: a synthesis. *Marine and Petroleum Geology* **56**: 184–221. doi: [10.1016/j.marpetgeo.2014.04.006](https://doi.org/10.1016/j.marpetgeo.2014.04.006).
- Fensome, R.A., Williams, G.L. & MacRae, R.A.**, 2019. Lentin and Williams index of fossil dinoflagellates. *AASP Contribution Series* **50**: 1173 pp.
- Geluk, M.C.**, 1990. The Cenozoic Roer Valley Graben, southern Netherlands. *Mededelingen Rijks Geologische Dienst* **44**(4): 63–72.
- Gibbard, PhL & Lewin, J.**, 2016. Filling the North Sea Basin: Cenozoic sediment sources and river styles. *Andre Dumont medallist lecture 2014. Geologica Belgica* **19**(3-4): 201–217. doi: [10.20341/gb.2015.017](https://doi.org/10.20341/gb.2015.017).
- Hager, H., Vandenberghe, N., Van den Bosch, M., Abraham, M., Von der Hocht, F., Rescher, K., Laga, P., Nickel, E., Verstrealen, A., Leroi, S., Van Leeuwen, R.J.W.**, 1998. The geometry of the Rupelian and Chattian depositional bodies in the Lower Rhine district and its border area: implications for Oligocene lithostratigraphy. *Geological Society Denmark* **45**: 53–62.
- Halet, F.**, 1920. La géologie tertiaire de la Campine anversoise et limbourgeoise. La falaise d'Elslloo et son gravier fossilifère. *Bulletin de la Société Belge de Géologie* **30**: 84–100.
- Haq, B.U., Hardenbol, J. & Vail, P.R.**, 1987. Chronology of fluctuating sea levels since the Triassic (250 million years ago to present). *Science* **235**: 1156–1167.
- Hardenbol, J., Thierry, J., Farley, M.B., Jacquin, Th, De Graciansky, P.C.H. & Vail, P.R.**, 1998. Mesozoic and Cenozoic sequence chronostratigraphic framework of European Basins, Mesozoic and Cenozoic sequence chronostratigraphic chart 1. *In*: Graciansky, P.C.H., Hardenbol, J., Jacquin, T. & Vail, P.R. (eds): Mesozoic and Cenozoic sequence stratigraphy of European Basins. SEPM (Tulsa, OK).
- Head, M.J.**, 1998. Marine environmental change in the Pliocene and early Pleistocene of eastern England: the dinoflagellate evidence reviewed. *In*: Van Kolfshoten, T. & Gibbard, P.L. (eds): The dawn of the Quaternary. *Mededelingen NITG-TNO* **60**: 199–225.
- Head, M.J., Norris, G. & Mudie, P.J.**, 1989. Palynology and dinoflagellate cyst stratigraphy of the Upper Miocene and lowermost Pliocene, ODP Leg 105, Site 646, Labrador Sea. *Proceedings of the Ocean Drilling Program, Scientific Results* **105**: 423–451. doi: [10.2973/odp.proc.sr.105.135.1989](https://doi.org/10.2973/odp.proc.sr.105.135.1989).
- Kennett, J.P.**, 1977. Cenozoic evolution of Antarctic glaciation, the circum Antarctic Ocean, and their impact on global paleoceanography. *Journal of Geophysical Research* **82**: 3843–3860.
- King, C., Gale, A.S. & Barry, T.L.**, 2016. A revised correlation of Tertiary rocks in British Isles and adjacent areas of NW Europe. *Geological Society Special Reports*, vol. 27. Geological Society (London): 719 pp. doi: [10.1144/SR27](https://doi.org/10.1144/SR27).

- Knox, R.W.O.'B., Bosch, J.H.A., Rasmussen, E.S., Heilmann-Clausen, C., Hiss, M., De Lugt, I.R., Kasiński, J., King, C., Köthe, A., Slodkowska, B., Standke, G., Vandenbergh, N.**, 2010. Cenozoic. In: Doornenbal J.C. & Stevenson A.G. (eds): Petroleum geological atlas of the Southern Permian Basin area. EAGE Publications B.V. (Houten): 211–223.
- Köthe, A.**, 2007. Cenozoic biostratigraphy from the German North Sea sector (G-11-1 borehole, dinoflagellate cysts, calcareous nannoplankton). *Zeitschrift der Deutschen Gesellschaft für Geowissenschaften* **158**(2): 287–327. doi: [10.1127/1860-1804/2007/0158-0287](https://doi.org/10.1127/1860-1804/2007/0158-0287).
- Köthe, A., Gaedicke, C. & Lutz, R.**, 2008. Erratum: The age of the Mid-Miocene Unconformity (MMU) in the G-11-1 borehole, German North Sea sector. *Zeitschrift der Deutschen Gesellschaft für Geowissenschaften* **159**(4): 687–689. doi: [10.1127/1860-1804/2008/0159-0687](https://doi.org/10.1127/1860-1804/2008/0159-0687).
- Kuyl, O.S.**, 1980. Toelichtingen bij de Geologische Kaart van Nederland 1:50.000. Blad Heerlen, 62 W oostelijke helft, 62 O westelijke helft. Rijks Geologische Dienst (Haarlem): 206 pp.
- Liebrand, D., Beddow, H.M., Lourens, L.J., Pälike, H., Raffi, I., Bohaty, S.M. & et al.**, 2016. Cyclostratigraphy and eccentricity tuning of the early Oligocene through early Miocene (30.1–17.1 Ma): Cibicides mundulus stable oxygen and carbon isotope records from Walvis Ridge Site 1264. *Earth and Planetary Science Letters* **450**: 392–405. doi: [10.1016/j.epsl.2016.06.007](https://doi.org/10.1016/j.epsl.2016.06.007).
- Louwe, S.**, 2002. Dinoflagellate cyst biostratigraphy of the Upper Miocene Deurne Sands (Diest Formation) of northern Belgium, southern North Sea Basin. *Geological Journal* **37**(1): 55–67. doi: [10.1002/gj.900](https://doi.org/10.1002/gj.900).
- Louwe, S.**, 2005. The Early and Middle Miocene transgression at the southern border of the North Sea Basin (northern Belgium). *Geological Journal* **40**(4): 441–456. doi: [10.1002/gj.1021](https://doi.org/10.1002/gj.1021).
- Louwe, S., Deckers, J., Verhaegen, J., Adriaens, J. & Vandenbergh, N.**, 2020. A review of the lower and middle Miocene of northern Belgium. *Geologica Belgica* **23**(3-4): 137–156. doi: [10.20341/gb.2020.010](https://doi.org/10.20341/gb.2020.010).
- Louwe, S., Head, M.J. & De Schepper, S.**, 2004. Dinoflagellate cyst stratigraphy and palaeoecology of the Pliocene in northern Belgium, southern North Sea Basin. *Geological Magazine* **141**(3): 353–378. doi: [10.1017/S0016756804009136](https://doi.org/10.1017/S0016756804009136).
- Louwe, S., Marquet, R., Bosselaers, M. & Lambert, O.**, 2010. Stratigraphy of an Early-Middle Miocene sequence near Antwerp in northern Belgium (southern North Sea Basin). *Geologica Belgica* **13**: 269–284.
- Michon, L., Van Balen, R.T., Merle, O. & Pagnier, H.**, 2003. The Cenozoic evolution of the Roer Valley rift system integrated at a European scale. *Tectonophysics* **367**: 101–126.
- Miller, K.G., Browning, J.V., Kopp, R.E., Montain, G.S. & Wright, J.D.**, 2020. Cenozoic sea-level and cryospheric evolution from deep-sea geochemical and continental margin records. *Science Advances* **6**(20): 15 pp. doi: [10.1126/sciadv.aaz1346](https://doi.org/10.1126/sciadv.aaz1346).
- Miller, K.G., Kominz, M.A., Browning, J.V., Wright, J.D., Mountain, G.S., Katz, M.E., Sugarman, P.J., Cramer, B.S., Christie-Blick, N., Pekar, S.F.**, 2005. The Phanerozoic record of global sea-level change. *Science* **310**: 1293–1298.
- Miller, K.G., Wright, J.D. & Fairbanks, R.G.**, 1991. Unlocking the Ice House: Oligocene-Miocene oxygen isotopes, eustasy and margin erosion. *Journal of Geophysical Research* **96**(B4): 6829–6848.
- Munsterman, D. & Deckers, J.**, 2020. The Oligocene/Miocene boundary in the ON-Mol-1 and Weelde boreholes along the southern margin of the North Sea Basin, Belgium. *Geologica Belgica* **23**(3-4). doi: [10.20341/gb.2020.007](https://doi.org/10.20341/gb.2020.007).
- Munsterman, D.K.**, 1997a. De resultaten van het dinoflagellaten onderzoek naar de ouderdom en het afzettingmilieu van boring Limbricht-01, traject: 205,5-516,5 m. TNO rapport NITG 97-109-B. Available at <https://www.nlog.nl/>
- Munsterman, D.K.**, 1997b. De resultaten van het aanvullend en herzien dinoflagellatenonderzoek naar de overgang van de Veldhoven-/Breda Formatie in boring Limbricht-01 (LBR-01), traject 224,5-326,0 m. TNO rapport NITG 97-193-B. Available at <https://www.nlog.nl/>
- Munsterman, D.K.**, 1998. I De resultaten van het palynologisch onderzoek naar de ouderdom en het afzettingmilieu van boring 58G192 (Herkenbosch), traject: 43.8-260.25 m. TNO rapport NITG 98-224-B. Available at <https://www.dinoloket.nl/ondergrondmodellenforlithologicalandlithostratigraphicinterpretation>
- Munsterman, D.K.**, 2001. Het dinoflagellatenonderzoek aan boring VEH-01 (Veldhoven-01), traject: 830- 1130 m. TNO rapport NITG 01-146-B.
- Munsterman, D.K. & Brinkhuis, H.**, 2004. A southern North Sea Miocene dinoflagellate cyst zonation. *Netherlands Journal of Geosciences* **83**(4): 267–285. doi: [10.1017/S0016774600020369](https://doi.org/10.1017/S0016774600020369).
- Munsterman, D.K., ten Veen, J.H., Menkovic, A., Deckers, J., Witmans, N., Verhaegen, J., Kerstholt-Boegehold, S.J., van de Ven, T. & Busschers, F.S.**, 2019. An updated and revised stratigraphic framework for the Miocene and earliest Pliocene strata of the Roer Valley Graben and adjacent blocks. *Netherlands Journal of Geosciences* **98**: 239. doi: [10.1017/njg.2019.10](https://doi.org/10.1017/njg.2019.10).
- Munsterman, D.K., Verreussel, R.M.C.H., Mijnlief, H.F., Witmans, N., Kerstholt-Boegehold, S.J. & Abbink, O.A.**, 2012. Revision and update of the Callovian-Ryazanian Stratigraphic Nomenclature in the northern Dutch offshore, i.e. Central Graben Subgroup and Scruff Group. *Netherlands Journal of Geosciences* **91**(4): 555–590.
- Naish, T.R., Wilson, G.S., Dunbar, G.B. & Barret, P.J.**, 2008. Constraining the amplitude of Late Oligocene bathymetric changes in western Ross Sea during orbitally induced oscillations in the East Antarctic Ice Sheet: (2) Implications for global sea-level changes. *Palaeogeography, Palaeoclimatology, Palaeoecology* **260**: 66–76.
- NAM & RGD**, 1980. Stratigraphic nomenclature of The Netherlands. *Verhandelingen van het Koninklijk Nederlands Geologisch Mijnbouwkundig Genootschap* **32**: 77 pp.
- NLOG**, n.d. Available at <https://www.nlog.nl/zoeken-boringen>
- Ogg, J.G., Ogg, G.M. & Gradstein, F.M.**, 2016. A concise geologic time scale. Elsevier (Amsterdam): 234 pp. doi: [10.1016/C2009-0-64442-1](https://doi.org/10.1016/C2009-0-64442-1).
- Powell, A.J.**, 1986. Latest Palaeogene and earliest Neogene dinoflagellate cysts from the Lemme section, northwest Italy. *AASP Contribution Series* **17**: 83–104.
- Powell, A.J.**, 1992. Dinoflagellate cysts of the Tertiary System. In: Powell A.J. (eds): A stratigraphic index of dinoflagellate cysts. Springer (Dordrecht): 155–272.
- Quaijtaal, W., Donders, T.H., Persicoc, D. & Louwe, S.**, 2014. Characterising the middle Miocene Mi-events in the Eastern North Atlantic realm: a first high-resolution marine palynological record from the Porcupine Basin. *Palaeogeography, Palaeoclimatology, Palaeoecology* **399**(4): 140–159. doi: [10.1016/j.palaeo.2014.02.017](https://doi.org/10.1016/j.palaeo.2014.02.017).
- Rasmussen, E.S., Dybkjaer, K. & Piasecki, S.**, 2010. Lithostratigraphy of the Upper Oligocene – Miocene Succession of Denmark. *Geological Survey of Denmark and Greenland Bulletin* **22**: 92.
- Rasmussen, E.S., Heilmann-Clausen, C., Waagstein, R. & Eidvin, T.**, 2008. Tertiary of the Norden. *Episodes* **31**: 66–72.
- Schäfer, A. & Utescher, T.**, 2014. Origin, sediment fill, and sequence stratigraphy of the Cenozoic Lower Rhine Basin (Germany) interpreted from well logs. *German Journal of Geoscience* **165**(2): 287–314. doi: [10.1127/1860-1804/2014/0062](https://doi.org/10.1127/1860-1804/2014/0062).
- Schäfer, A., Utescher, T., Klett, M. & Valdivia-Manchego, M.**, 2005. The Cenozoic Lower Rhine Basin – rifting, sediment input, and cyclic stratigraphy. *International Journal of Earth Sciences (Geologische Rundschau)* **94**(4): 621–639. doi: [10.1007/s00531-005-0499-7](https://doi.org/10.1007/s00531-005-0499-7).
- Schmitz, G. & Stainier, W.**, 1909. Découverte en campine de l'Oligocène supérieure marin : la question de l'âge du Boldérien de Dumont. *Annales de la Société géologique de Belgique* **36**: 253–367.
- Schneider, H. & Thiele, S.**, 1965. Geohydrologie des Erftgebietes. Ministerium für Ernährung, Landwirtschaft und Forsten Land Nordrhein-Westfalen (Düsseldorf): 185 pp.
- Sliwinska, K.K.**, 2019. Early Oligocene dinoflagellate cysts as a tool for palaeoenvironment reconstruction and stratigraphical framework – a case study from a North Sea well. *J. Micropalaeontology* **38**(2): 143–176. doi: [10.5194/jm-38-143-2019](https://doi.org/10.5194/jm-38-143-2019).
- Stainier, X.**, 1911. Sur les recherches de sel en Campagne. *Annales des Mines de Belgique* **16**: 117–170.
- Steinthorsdottir, M., Coxall, H.K., De Boer, A.M., Huber, M., Barbolini, N., Bradshaw, C.D., Burls, N.J., Feakins, S.J., Gasson, E., Henderiks, J., Holbourn, A.E., Kiel, S., Kohn, M.J., Knorr, G., Kürschner, W.M., Lear, C.H., Liebrand, D., Lunt, D.J., Mörs, T., Pearson, P.N., Pound, M.J., Stoll, H. & Strömberg, C.A.E.**, 2020. The Miocene: the future of the past.

- Available at https://www.researchgate.net/publication/347960864_The_Miocene_The_Future_of_the_Past
- Stover, L.E. & Hardenbol, J.**, 1994. Dinoflagellates and depositional sequences in the Lower Oligocene (Rupelian) Boom Clay Formation. Belgium. *Bulletin Society Belgium Geology* **102**(1-2): 5–77.
- Stratigraphic Nomenclature of the Netherlands online**, 2021. Available at <https://www.dinoloket.nl/stratigrafische-nomenclator>
- Utescher, T., Mosbrugger, V. & Ashraf, A.R.**, 2000. Terrestrial climate evolution in Northwest Germany over the last 25 million years. *Palaios* **15**: 430–449.
- Van Adrichem Boogaert, A.H. & Kouwe, W.F.P.**, 1993–1997. Stratigraphic Nomenclature of the Netherlands, revision and update by RGD and NOGEP. Mededelingen Rijks Geologische Dienst **50**(section I): 39 pp.
- Van den Berg, M.W.**, 1994. Neotectonics of the Roer Valley rift system. Style and rate of crustal deformation inferred from syntectonic sedimentation. *Geologie en Mijnbouw* **73**: 143–156.
- Van Leeuwen, R.J.**, 2000. Foraminiferen- & bolboformastratigrafievian boring Groote Heide B58F0064. TNO report NITG 00-15-B.
- Van Simaëys, S.**, 2004. The Rupelian-Chatthian boundary in the North Sea Basin and its calibration to the international time scale. *Netherlands Journal of Geosciences* **83**(3): 241–248. doi: [10.1017/S0016774600023532](https://doi.org/10.1017/S0016774600023532).
- Van Simaëys, S., De Man, E., Vandenberghe, N., Brinkhuis, H. & Steurbaut, E.**, 2004. Stratigraphic and palaeoenvironmental analysis of the Rupelian-Chatthian transition in the type region: evidence from dinoflagellate cysts, foraminifera and calcareous nannofossils. *Palaeogeography, Palaeoclimatology, Palaeoecology* **208**: 31–58. doi: [10.1016/j.palaeo.2004.02.029](https://doi.org/10.1016/j.palaeo.2004.02.029).
- Van Simaëys, S., Munsterman, D.K. & Brinkhuis, H.**, 2005. Oligocene dinoflagellate cyst biostratigraphy of the southern North Sea Basin. *Review of Palaeobotany and Palynology* **134**: 105–128. doi: [10.1016/j.revpalbo.2004.12.003](https://doi.org/10.1016/j.revpalbo.2004.12.003).
- Vandenberghe, N., Brinkhuis, H. & Steurbaut, E.**, 2003. The Eocene/Oligocene boundary in the North Sea Area: sequence stratigraphic approach. In: Prothero, D.D., Ivany, L.C. & Nesbitt, E.A. (eds): *From greenhouse to icehouse: The marine Eocene-Oligocene transition*. Columbia University Press (New York): 419–437.
- Vandenberghe, N., Van Simaëys, S., Steurbaut, E., Jagt, J.W.M. & Felder, P.J.**, 2004. Stratigraphic architecture of the Upper Cretaceous and Cenozoic along the southern border of the North Sea Basin in Belgium. *Netherlands Journal of Geosciences* **83**: 155–171. doi: [10.1017/S0016774600020229](https://doi.org/10.1017/S0016774600020229).
- Verbeek, J.W., De Leeuw, C.S., Parker, N. & Wong, Th E.**, 2002. Characterization and correlation of Tertiary seismostratigraphic units in the Roer Valley Graben. *Netherlands Journal of Geosciences* **81**(2): 159–166. doi: [10.1017/S0016774600022393](https://doi.org/10.1017/S0016774600022393).
- Verreussel, R.M.C.H., Bouroullac, R., Munsterman, D.K., Dybkjær, K., Geel, C.R., Houben, A.J.P. & Kerstholt-Boegehold, S.J.**, 2018. Stepwise basin evolution of the Middle Jurassic to Early Cretaceous rift phase in the Central Graben area of Denmark, Germany and the Netherlands. Mesozoic resource potential in the Southern Permian Basin. *Geological Society, London, Special Publications* **469**(1): 305–340. doi: [10.1144/SP469.23](https://doi.org/10.1144/SP469.23).
- Whittaker, M.F.**. The usage of palynostratigraphy and palynofacies in definition of Troll field geology. In: *Norwegian Petroleum Society Offshore North Seas conference*, paper 66, 1984: 1–50.
- Wong, Th E., Batjes, A.J. & De Jager, J.**, 2007. *Geology of the Netherlands*. Royal Netherlands Academy of Arts and Sciences (Amsterdam): 354.
- Wong, Th E., Parker, N. & Horst, P.**, 2001. Tertiary sedimentary development of the Broad Fourteens area, the Netherlands. *Netherlands Journal of Geosciences* **80**(1): 85–94. doi: [10.1017/S0016774600022186](https://doi.org/10.1017/S0016774600022186).
- Zagwijn, W.H. & Van Staaldunin, C.J.**, 1975. *Toelichtingen bij Geologische overzichtskaarten van Nederland*. Rijks Geologische Dienst (Haarlem): 134 pp.
- Zevenboom, D.**, 1995. *Dinoflagellate cysts from the Mediterranean Late Oligocene and Miocene*. Unpublished Ph.D. Thesis, University of Utrecht, Utrecht: 221 pp.
- Ziegler, P.A.**, 1994. Cenozoic rift system of western and central Europe: an overview. *Geologie en Mijnbouw* **73**: 99–127.