


Erwinia Chrysanthemi:
A plant pathogen causing Stem
Rot in Chrysanthemums and
Soft Rot in fleshy vegetables



a

cure

for

chrysanthemums

When your business is selling Chrysanthemums, Stem Rot can be disastrous. Faced with this menace a certain large firm of growers turned to us. The problem was to isolate infected cuttings early enough to prevent cross infection in the Nursery.

Now, in this one business alone upwards of a million cuttings a year are being laboratory tested in Oxoid Culture Media before rooting, to eradicate infected stock.

FOR PROGRESSIVE LABORATORIES




CULTURE MEDIA

The use of Oxoid Culture Media ensures constant quality and uniformity in your laboratory technique. Absolutely reliable, quick, convenient and economical, Oxoid Culture Media are available in tablet or granular form. Full details from:

**OXOID DIVISION OF OXO LTD., Thames House,
Queen Street Place, London, E.C.4 (CENTRAL 9781)**

Hygiene, 58, 1

(i)



A MICROBE COULD DESTROY A MISSILE

GUIDED MISSILES depend for their accurate functioning on incredibly sensitive and delicate instruments. Their mechanisms are machined to within one millionth of an inch. Therefore, if even the most minute particles get into the hydraulic control system, a failure could occur when the missile is fired. Consequently, micro-filtration of the hydraulic fluid is essential and, as a check on its efficiency, increasing use is now being made of Oxoid Membrane Filters

These filters, made of Cellulose Acetate, have a pore size of 0.5 to 1.0 micron and will permit rapid filtration of fluids and gases.


Oxoid Membrane Filters were primarily developed for the bacteriologist and are widely used for the bacteriological examination of fluids and in sterility testing. A range of special Oxoid culture media for the membrane filtration technique will shortly be available.

FOR PROGRESSIVE LABORATORIES

OXOID

MEMBRANE FILTERS

For information about OXOID Membrane Filters and the new OXOID range of membrane filtration media in tablet form, write to: The Oxoid Division of Oxo Limited, Thames House, Queen Street Place, London, E.C.4. (Central 9781)



“THE BOTTOM DROPPED OUT OF MY BARGE!”

This was the problem someone recently faced. When we heard about it we were sympathetic, for the barge in question was a houseboat, and most of the owner's worldly goods had vanished in the catastrophe.

Many people would not think of looking to bacteriology for the answer. But if they did they would learn that members of the *Desulphovibrio* group, which are commonly present in soil and water, can corrode steel.

These anaerobic organisms, first described in 1895 by Beijerinck, are capable of reducing sulphates to sulphides. Their activities, often accompanied by the production of hydrogen sulphide, may result in the corrosion of buried iron pipes and the steel hulls of barges.

By the use of Baar's or Starkey's media they are comparatively easy to isolate as crude cultures.

*Electron micrograph:
desulphovibrio
desulphuricans*




FOR PROGRESSIVE LABORATORIES

OXOID

CULTURE MEDIA

The use of Oxoid Culture Media ensures constant uniformity and quality in your laboratory technique. Absolutely reliable, quick, convenient and economical, Oxoid Culture Media are available in tablet or granular form. If you would like the manual of all Oxoid preparations, write to: The OXOID Division of OXO Ltd., Thames House, Queen Street Place, London, E.C.4 (Central 9781).



DISPOSABLE PETRI DISHES

eliminate time-wasting washing, sterilizing

The Oxoid Disposable Petri Dish is ready for instant use and can afterwards be discarded. Made of clear, moulded polystyrene, it is sterile and biologically inert as well as light in weight and virtually unbreakable.

Because of its low cost and the elimination of washing and sterilizing, Oxoid Disposable Petri Dishes offer considerable economy in use. The dishes are

supplied in cartons of 160 or 500. Full details, prices and quantity rates will gladly be sent on application.

.....
 : *NEW Oxoid Disposable Plastic Precipitin* :
 : *Tubes are now available—made from* :
 : *crystal-clear polystyrene, they are chemi-* :
 : *cally clean and ready for immediate use.* :
 : *Further details supplied on request.* :
 :
 :

FOR PROGRESSIVE LABORATORIES



BACTERIOLOGICAL SERVICES

Made for the Oxoid Division of Oxo Ltd.

Thames House, Queen Street Place, London E.C.4. Telephone: CENtral 9781

The Clonal Selection Theory of Acquired Immunity

SIR MACFARLANE BURNET

A discussion of the population dynamics of clones—with applications to the phenomena of acquired immunity, antibody production, and the problems of neoplastic growth. This book will be of interest to microbiologists, immunologists, and to all who are engaged in cancer research. Based on the Abraham Flexner Lectures for 1958. 22s. 6d. net

CAMBRIDGE UNIVERSITY PRESS

THE WORLD HEALTH ORGANIZATION

invites applications, which should be made within six weeks of the date of issue of this advertisement, for two posts of

TECHNICAL ASSISTANT

QUALIFICATIONS: University degree in some branch of biological sciences, preferably bacteriology or immunology. Original research in biology or in communicable diseases and some laboratory experience desirable. Good knowledge of English and French. Knowledge of other European languages, particularly Russian, desirable.

SALARY: U.S. \$3600 per annum, net of income tax, rising by annual increments to \$5000 plus cost of living and dependants allowances and education grant for children under certain conditions. Initial contract for two years.

Applications giving brief biographical data should be addressed to Chief, Personnel, World Health Organization, Palais des Nations, Geneva, marking the outside of the envelope "VN. 570."

Further information and application forms will be sent only to those candidates who seem suitable for serious consideration.

Full particulars of the

JOURNALS

published by the

CAMBRIDGE
UNIVERSITY PRESS

may be had from

*The Manager
Cambridge University Press
Bentley House, 200 Euston Road
London, N.W. 1*

The Journal of General Microbiology

Editors: B. C. J. G. KNIGHT and A. F. B. STANDFAST

Vol. 23, No. 3

35s.

December 1960

- PREB, L., HOWELL, A. JR. and WATSON, S. J. Studies of the morphological, physiological, and biochemical characters of *Actinomyces bovis*.
- TRENTHAM, H. The adaptive degradation of L-histidine by *Paracolobatum aerogenoides*.
- KRIS, A. E. and MITSKEVICH, I. N. The distribution of filamentous-cluster organisms (Kraussnikovise) in sea and ocean depths.
- COETZEE, J. N. and SACKS, T. G. Transduction of streptomycin resistance in *Proteus mirabilis*.
- BAUCHOP, T. and ELSDEN, S. R. The growth of microorganisms in relation to their energy supply.
- BROWN, A. D. Some properties and a proposed classification of a Gram-negative heterotrophic marine bacterium.
- LASCÉLLES, J. The synthesis of enzymes concerned in bacteriochlorophyll formation in growing cultures of *Rhodospseudomonas spheroides*.
- LASCÉLLES, J. The formation of ribulose 1:5-diphosphate carboxylase by growing cultures of Athiorhodaceae.
- KORNBERG, H. L. and LASCÉLLES, J. The formation of isocitrate by the Athiorhodaceae.
- KERRIDGE, D. The effect of inhibitors on the formation of flagella by *Salmonella typhimurium*.
- MUKAI, F. Inter-relationship between colicin sensitivity and phage resistance.
- BRADLEY, D. E. and KAY, D. The fine structure of bacteriophages.
- KARSKA, B., KOZAK, W., NARBUROWICZ, B. and STOLARSKA, A. Study on the lysogenicity of staphylococci isolated in Poland.
- TAYLOR, J., LEE, M. M., EDWARDS, P. R. and RAMSEY, C. H. A new type of flagellar variation associated with new antigens in the Salmonella group.
- WEITZ, B. The properties of some antigens of *Trypanosoma brucei*.
- COWAN, S. T., STEEL, K. J., SHAW, C. and DUGUID, J. P. A classification of the Klebsiella group.
- FURNESS, G., GRAHAM, D. M. and REEVE, P. The titration of Trachoma and Inclusion Blepharorrhoea Viruses in cell cultures.
- SHARPE, M. E. and FEWINS, B. G. Serological typing of strains of *Streptococcus faecium* and unclassified group D streptococci isolated from canned hams and pig intestines.

Orders should sent to

CAMBRIDGE UNIVERSITY PRESS
BENTLEY HOUSE, 200 BUSTON ROAD, LONDON, N.W. 1

The Quarterly Journal of Medicine

Edited by D. H. BRINTON, W. D. W. BROOKS, A. M. COOKE,
R. B. HUNTER, H. L. MARRIOTT, J. D. N. NABARRO, M. L. ROSENHEIM

CONTENTS

NEW SERIES. Vol. XXIX. No. 115. JULY 1960

- Staphylococcal Septicaemia and Pyaemia. By D. R. HAY. With Plate 27.
- The Absorption of Folic Acid and Labelled Cyanocobalamin in Intestinal Malabsorption. By A. DOIG and R. H. GIRDWOOD.
- Fulminant Ulcerative Colitis. By J. G. RANKIN, S. J. M. GOULSTON, R. W. BODEN and A. W. MORROW. With Plates 28 to 32.
- Recurrent Focal Nephritis. By J. H. ROSS. With Plates 33 and 34.
- The Metabolic Disorder in Hartnup Disease. By M. D. MILNE, M. A. CRAWFORD, C. B. GIRAŐ and LAVINIA W. LOUGHRIDGE.
- Some Investigations into Endocrine Exophthalmos. By D. A. MCGILL. With Plate 35.
- Gout and the Serum Uric Acid in Diabetes Mellitus. By A. GORDON BECKETT and J. G. LEWIS.
- Embolic Occlusion of the Aorta in Patients with Mitral Stenosis. By A. E. A. READ, K. P. BALL and C. G. ROB. With Plates 36 and 37.

Single Numbers 20s. net each Subscription 60s. per annum

OXFORD UNIVERSITY PRESS, LONDON E.C. 4

CONTENTS

(All rights reserved)

	PAGE
COLEBROOK, LEONARD, LOWBURY, E. J. L. and HURST, THE LATE L. The growth and death of wound bacteria in serum, exudate and slough	357
MARSHALL, J. H. and KELSEY, J. C. A standard culture medium for general bacteriology	367
DIMMICK, ROBERT L. Measurement of the physical decay of aerosols by a light scatter method compared to a radioactive tracer method	373
SMITH, H. WILLIAMS. The effect of feeding pigs on food naturally contaminated with salmonellae	381
KAPLAN, COLIN. The antigenicity of γ -irradiated vaccinia virus	391
KENDALL, E. J. C., VOLLMUM, B. L., BESWICK, T. S. L., MILLER, ANN and TOBIN J. O'H. The antigenic activity of some poliomyelitis vaccines used in the United Kingdom in 1958	399
THORNE, H. V. and BURROWS, T. M. Aerosol sampling methods for the virus of foot-and-mouth disease and the measurement of virus penetration through aerosol filters	409
POLSON, A., HAMPTON, J. W. F. and DEEKS, D. Zone electrophoresis of poliovirus partially inactivated by formaldehyde. (With Plates 1 and 2)	419
BLOWERS, ROBERT and CREW, BERYL. Ventilation of operating-theatres. (With Plates 1 and 2)	427
LIDWELL, O. M. and WILLIAMS, R. E. O. The ventilation of operating-theatres	449
OLIVER, R. and TOMLINSON, A. H. The sterilization of surgical rubber gloves and plastic tubing by means of ionizing radiation	465
FERGUSON, G. S. and ROBERTSON, A. Brucellosis in dairy herds—some applications of the milk ring test	473
MARSHALL, I. D. and FENNER, FRANK. Studies in the epidemiology of infectious myxomatosis of rabbits. VII. The virulence of strains of myxoma virus recovered from Australian wild rabbits between 1951 and 1959	485

The Journal of Hygiene is issued in four parts to a volume.

Quotations can be given for back volumes on application.

Papers for publication should be sent to Lt.-Col. H. J. BENSTED, c/o Lister Institute of Preventive Medicine, Chelsea Bridge Road, London, S.W. 1. Other communications should be addressed to the University Press, Cambridge.

Papers forwarded to the Editor for publication are understood to be offered to *The Journal of Hygiene* alone, unless the contrary is stated.

Contributors receive twenty-five copies of their paper free. Additional copies may be had at cost price: these should be ordered when the final proof is returned.

The subscription price is £4. 10s. per volume (post-free), payable in advance; single numbers 30s. net. Subscriptions may be sent to any Bookseller, or to The Cambridge University Press, Bentley House, 200 Euston Road, London, N.W. 1.

The subscription price in the U.S.A., is \$17.50. Single parts \$5.00. Enquiries should be addressed to the American Branch of the Press at 32 East 57th Street, New York 22, N.Y.