

**PUBLISHED BY  
THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE**

**The Pitt Building, Trumpington Street, Cambridge CB2 1RP  
32 East 57th Street, New York, NY 10022, USA  
10 Stamford Road, Oakleigh, Melbourne 3166, Australia**

**© Cambridge University Press 1984**

***Printed in Great Britain by J. W. Arrowsmith Ltd, Bristol***

# Contents

## PART 1 MARCH 1984

Fat baker's transformation	<i>J. C. Alexander and J. A. Yorke</i>	1
On orbits of unipotent flows on homogeneous spaces	<i>S. G. Dani</i>	25
Dynamics of $\exp(z)$	<i>R. L. Devaney and M. Krych</i>	35
Flow equivalence of subshifts of finite type	<i>J. Franks</i>	53
Flots d'Anosov sur les 3-variétés en cercles	<i>E. Ghys</i>	67
Combined structural and topological stability are equivalent to Axiom A and the strong transversality condition	<i>M. Hurley</i>	81
Kakutani equivalence of ergodic $Z^n$ actions	<i>A. del Junco and D. J. Rudolph</i>	89
Any flow is an orbit factor of any flow	<i>D. Ornstein and B. Weiss</i>	105
Bowen's equidistribution theory and the Dirichlet density theorem	<i>W. Parry</i>	117
A complex Ruelle-Perron-Frobenius theorem and two counter-examples	<i>M. Pollicott</i>	135
Lorenz knots are prime	<i>R. F. Williams</i>	147

## PART 2 JUNE 1984

Rational ergodicity of geodesic flows	<i>J. Aaronson and D. Sullivan</i>	165
Simple permutations with order a power of two	<i>C. Bernhardt</i>	179
One-parameter automorphism groups of operator algebras allowing spectral projections	<i>G. A. Elliott and L. Zsidó</i>	187
Skew products and minimal dynamical systems on separable Hilbert manifolds	<i>A. Fathi</i>	213
A return time invariant for finitary isomorphisms	<i>U. R. Fiebig</i>	225
Uncountably many topological models for ergodic transformations	<i>S. Glasner and D. Rudolph</i>	233
Twofold mixing implies threefold mixing for rank one transformations	<i>S. A. Kalikow</i>	237
The Lyapunov dimension of a nowhere differentiable attracting torus	<i>J. L. Kaplan, J. Mallet-Paret and J. A. Yorke</i>	261
The entropies of topological Markov shifts and a related class of algebraic integers	<i>D. A. Lind</i>	283
Non-existence of invariant circles	<i>J. N. Mather</i>	301
Ergodic rational maps with dense critical point forward orbit	<i>M. Rees</i>	311

PART 3 SEPTEMBER 1984

Weak disjointness and the equicontinuous structure relation <i>J. Auslander, D. C. McMahon, J. C. S. P. van der Woude and T.-S. Wu</i>	323
Ergodic and mixing sequences of transformations <i>D. Berend and V. Bergelson</i>	353
Shift equivalence and the Jordan form away from zero <i>M. Boyle</i>	367
Stability of the weak Pinsker property for flows <i>A. Fieldsteel</i>	381
Twist sets for maps of the circle <i>M. Misiurewicz</i>	391
On the notion of the dimension with respect to a dynamical system <i>Ya. B. Pesin</i>	405
Unfoldings of discrete dynamical systems <i>J. W. Robbin</i>	421

PART 4 DECEMBER 1984

The backward continued fraction map and geodesic flow <i>R. L. Adler and L. Flatto</i>	487
Rotation intervals of endomorphisms of the circle <i>R. Bamon, I. P. Malta, M. J. Pacifico and F. Takens</i>	493
Minimal sets on tori <i>D. Berend</i>	499
Subordinate Šil'nikov bifurcations near some singularities of vector fields having low codimension <i>H. W. Broer and G. Vegter</i>	509
A criterion for the positivity of the Liapunov characteristic exponent <i>E. Cornelis and M. Wojtkowski</i>	527
Approximation by Brownian motion for Gibbs measures and flows under a function <i>M. Denker and W. Philipp</i>	541
Non-integrability of the 1:1:2 resonance <i>J. J. Duistermaat</i>	553
Billiards on almost integrable polyhedral surfaces <i>E. Gutkin</i>	569
A topological version of a theorem of Mather on twist maps <i>G. R. Hall</i>	585
Transitivity and invariant measures for the geometric model of the Lorenz equations <i>C. Robinson</i>	605
The Perron-Frobenius operator in spaces of smooth functions on an interval <i>B. Szewc</i>	613
Index to Volume 4	645

## THE PREPARATION OF TYPESCRIPTS

The attention of authors is particularly directed to the following requests:

### 1 *Typescript*

Papers should be typed, double-spaced, on one side of white paper (of which A4, 210 by 297 mm, is a suitable size). The pages must be numbered. Generous margins should be left at the side, top and bottom of each page. The copy sent must be clear.

A cover page should give the title, the author's name and institution, with the address to which mail is to be sent.

The title, while brief, must be informative (e.g. *A new proof of the ergodic theorem*, whereas *Some applications of a theorem of Birkhoff* would be useless).

The first paragraph or two should form a summary of the main theme of the paper, providing an abstract intelligible to mathematicians.

For a typescript to be accepted for publication, it must accord with the standard requirements of publishers, and be presented in a form in which the author's intentions regarding symbols etc. are clear to a printer (who is not a mathematician).

The following notes are intended to help the author in preparing the typescript. New authors may well enlist the help of senior colleagues, both as to the substance of their work and the details of setting it out correctly and attractively.

### 2 *Notation*

Notation should be chosen carefully so that mathematical operations are expressed with all possible neatness, to lighten the task of the compositor and reduce the chance of error.

For instance  $n_k$  ( $n$  sub  $k$ ) is common usage, but avoid if possible using  $c$  sub  $n$  sub  $k$ . Fractions are generally best expressed by a solidus. Complicated exponentials like

$$\exp \{z^2 \sin \theta / (1 + y^2)\}$$

should be shown in this and no other way.

In the typescript, italics, small capitals and capitals are specified by single, double and triple underlining. Bold-faced type is shown by wavy underlining; wavy will be printed **wavy**.

It helps if displayed equations or statements which will be quoted later are numbered in order on the right of their line. They can then be referred to by, for example, 'from (7)'.  
The author must enable the printer (if necessary by pencilled notes in the margin) to distinguish between similar symbols such as  $o$ ,  $O$ ,  $o$ ,  $O$ ,  $0$ ;  $x$ ,  $X$ ,  $\times$ ;  $\phi$ ,  $\Phi$ ,  $\emptyset$ ;  $l$ ,  $1$ ;  $\epsilon$ ,  $\in$ ;  $\kappa$ ,  $k$ .

Greek letters can be denoted by Gk in the margin.

If an author wishes to mark the end of the proof of a theorem, the sign  $\square$  may be used.

Footnotes should be avoided.

### 3 *Diagrams*

It is extremely helpful if diagrams are drawn in Indian ink on white card, faintly blue or green-lined graph paper, or tracing cloth or paper. *Symbols, legends and captions should be given on a transparent overlay*. Each text figure must be numbered as Figure 1, Figure 2, ... and its intended position clearly indicated in the typescript:

Figure 1 here

The author's name in pencil must be on all separate sheets of diagrams.

A figure is expensive to reproduce and should be included only when the subject matter demands it, or when it greatly clarifies the exposition.

The publisher recognizes that some authors do not have the facilities for producing drawings of a sufficiently high standard to be reproduced directly and is therefore willing to have such diagrams re-drawn, provided that they are clear.

### 4 *Tables*

Tables should be numbered (above the table) and set out on separate sheets. Indicate the position of each in the text as for figures:

Table 3 here

### 5 *References*

References should be collected at the end of the paper numbered in alphabetical order of the authors' names. A reference to a book should give the title, in italics, and then in roman type the publisher's name and the place and year of publication:

[4] N. Dunford & J. T. Schwartz *Linear Operators* Part I. Wiley: New York, 1958.

A reference to a paper should give in italics the title of the periodical, the number of the volume and year, and the beginning and end pages of the paper. Titles should be abbreviated as in *Mathematical Reviews*:

[6] J. E. Littlewood. The 'pits effect' for functions in the unit circle. *J. Analyse Math.* **23** (1970), 236–268.

# *Ergodic theory and dynamical systems*

VOLUME 4 PART 4 DECEMBER 1984

## CONTENTS

<i>Adler, R. L. and Flatto, L.</i> The backward continued fraction map and geodesic flow	487
<i>Bamon, R., Malta, I. P., Pacifico, M. J. and Takens, F.</i> Rotation intervals of endomorphisms of the circle	493
<i>Berend, D.</i> Minimal sets on tori	499
<i>Btoer, H. W. and Vegter, G.</i> Subordinate Šil'nikov bifurcations near some singularities of vector fields having low codimension	509
<i>Cornelis, E. and Wojtkowski, M.</i> A criterion for the positivity of the Liapunov characteristic exponent	527
<i>Denker, M. and Philipp, W.</i> Approximation by Brownian motion for Gibbs measures and flows under a function	541
<i>Duistermaat, J. J.</i> Non-integrability of the 1:1:2-resonance	553
<i>Gutkin, E.</i> Billiards on almost integrable polyhedral surfaces	569
<i>Hall, G. R.</i> A topological version of a theorem of Mather on twist maps	585
<i>Robinson, C.</i> Transitivity and invariant measures for the geometric model of the Lorenz equations	605
<i>Szewc, B.</i> The Perron-Frobenius operator in spaces of smooth functions on an interval	613
Index to Volume 4	645

© Cambridge University Press 1984

CAMBRIDGE UNIVERSITY PRESS

The Pitt Building, Trumpington Street, Cambridge CB2 1RP

32 East 57th Street, New York, NY 10022, USA

10 Stamford Road, Oakleigh, Melbourne 3166, Australia

*Printed in Great Britain by J. W. Arrowsmith Ltd, Bristol*