

ADDENDUM TO
"ON A THEOREM OF NIVEN"

Kenneth S. Williams

Can. Math. Bull., Vol. 10, no.4, pp. 573-578

Gordon Pall has kindly pointed out that the result of this paper was obtained by him in

Sums of two squares in a quadratic field, Duke Math. Jour.,
18 (1951), 399-409.

He gives the result in a slightly different form on page 405. We note that the two formulae are indeed the same. In my paper write (3) as

$$z = \epsilon(1+i)^\alpha \pi_1^{\alpha_1} \dots \pi_s^{\alpha_s} \bar{\pi}_1^{\alpha'_1} \dots \bar{\pi}_s^{\alpha'_s} q_1^{\beta_1} \dots q_l^{\beta_l}$$

so that

$$z\bar{z} = 2^\alpha p_1^{\alpha_1 + \alpha'_1} \dots p_s^{\alpha_s + \alpha'_s} q_1^{2\beta_1} \dots q_l^{2\beta_l} .$$

If $p_i^{\gamma_i} \mid\mid (x, y)$, then in Pall's notation $\gamma_i = \min(\alpha_i, \alpha'_i)$

and $\alpha_i + \alpha'_i = 2\gamma_i + \delta_i$,

giving $\gamma_i + \delta_i = \alpha_i + \alpha'_i - \min(\alpha_i, \alpha'_i) = \max(\alpha_i, \alpha'_i)$.

Hence $(1 + \gamma_i)(1 + \gamma_i + \delta_i)$

$$= (1 + \min(\alpha_i, \alpha'_i))(1 + \max(\alpha_i, \alpha'_i))$$

$$= (1 + \alpha_i)(1 + \alpha'_i) ,$$

showing that Pall's formula (22) is the same as my formula (19).