

THE SUBMODULE STRUCTURE OF THE  
QUANTUM SYMMETRIC POWERS

LARS THAMS

In this paper we study the symmetric powers of quantum  $GL_n$ . These are comodules and we give a complete description of the submodule structure.

In [3] Doty gives a complete description of the submodule structure for the symmetric powers of the natural module for  $GL_n$ . We want to do the same thing for quantum  $GL_n$ , that is, determine the complete submodule structure of the  $q$ -symmetric powers (defined in [2] by Dipper and Donkin).

Throughout the paper we shall look at quantum  $GL_n$  defined over some field  $k$ . From [2, Theorem 2.1.9] we get a  $k$  basis of the  $q$ -symmetric powers. Following Doty's idea [3] we shall determine the submodule generated by a basis element by relating it to the carry pattern which we shall define, and hence get a complete description of the submodule structure. As one would expect, the quantum parameter  $q \in k$  will play an important role in this description.

1. PRELIMINARIES

We shall first recall the set-up from [2].

1.1 Let  $R$  be a commutative ring,  $q \in R$  and  $2 \leq n \in \mathbf{Z}$ . Let  $A_q(n)$  be defined as the  $R$ -algebra generated by  $\{c_{i,j} \mid 1 \leq i, j \leq n\}$  with the following relations:

$$\begin{aligned} c_{i,k}c_{j,l} &= qc_{j,l}c_{i,k} \quad \text{for } i > j \text{ and } k \leq l \\ c_{i,k}c_{j,l} &= c_{j,l}c_{i,k} + (q - 1)c_{j,k}c_{i,l} \quad \text{for } i > j \text{ and } k > l \\ c_{i,k}c_{i,l} &= c_{i,l}c_{i,k} \quad \text{for all } i, k \text{ and } l. \end{aligned}$$

1.2 For some positive integer  $r$ , we denote by  $I(n, r)$  the set of  $\mathbf{i} = (i_1, \dots, i_r)$  where  $i_j \in \{1, \dots, n\}$  ( $1 \leq j \leq r$ ). For  $\mathbf{i}, \mathbf{s} \in I(n, r)$  let  $c_{\mathbf{i}, \mathbf{s}} = c_{i_1, s_1} c_{i_2, s_2} \dots c_{i_r, s_r} \in A_q(n)$ . Let  $A_q(n, r)$  denote the subspace of  $A_q(n)$  spanned by  $\{c_{\mathbf{i}, \mathbf{s}} \mid \mathbf{i}, \mathbf{s} \in I(n, r)\}$ .

---

Received 13th September, 1993.

I would like to thank S. Donkin for his help and for making me interested in this problem.

---

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9729/94 \$A2.00+0.00.

Then  $A_q(n)$  is a graded  $R$ -algebra  $A_q(n) = \bigoplus_{0 \leq r} A_q(n, r)$  and it is free as an  $R$ -module. The graded part  $A_q(n, r)$  has  $R$ -basis  $\{c_{i,s} \mid (i, s) \in I^2(n, r)^+\}$ . (See [2] for a definition of  $I^2(n, r)^+$ . It means that a basis element will be of the form  $c_{1,1}^{a_{11}} c_{1,2}^{a_{12}} \dots c_{1,n}^{a_{1n}} c_{2,1}^{a_{21}} \dots c_{n,n}^{a_{nn}}$  where  $a_{ij} \geq 0$  for all  $i, j$  and  $\sum_{i,j} a_{ij} = r$ .)

1.3  $A_q(n)$  is made into a bialgebra with comultiplication  $\Delta(c_{i,s}) = \sum_{j=1}^n c_{i,j} \otimes c_{j,s}$  and counit  $\varepsilon(c_{i,s}) = \delta_{i,s}$ . For every  $0 \leq r \in \mathbb{Z}$ ,  $A_q(n, r)$  is a subcoalgebra of  $A_q(n)$

1.4 Let  $E$  be the free  $R$ -module of dimension  $n$  with basis  $\{e_1, \dots, e_n\}$ . This means  $E = Re_1 \oplus \dots \oplus Re_n$ . Let  $E^{\otimes r}$  denote the  $r$ -fold tensor product of  $E$  with itself. For  $\mathbf{i} = (i_1, i_2, \dots, i_r) \in I(n, r)$  let  $e_{\mathbf{i}} = e_{i_1} \otimes e_{i_2} \otimes \dots \otimes e_{i_r} \in E^{\otimes r}$ . Now  $E^{\otimes r}$  becomes a right  $A_q(n, r)$ -comodule with structure map  $\tau_r : E^{\otimes r} \rightarrow E^{\otimes r} \otimes A_q(n, r)$  defined by  $\tau_r(e_{\mathbf{s}}) = \sum_{\mathbf{i} \in I(n, r)} e_{\mathbf{i}} \otimes c_{\mathbf{i}, \mathbf{s}}$ . Let  $T(E)$  denote the tensor algebra  $T(E) = \bigoplus_{0 \leq r} E^{\otimes r}$ , which is also a right  $A_q(n)$ -comodule.

1.5 There is also a left comodule structure on  $E^{\otimes r}$ , but to distinguish this from the right comodule it is given the name  $V^{\otimes r}$ , that is,  $V$  is the free  $R$ -module with basis  $\{v_1, \dots, v_n\}$  and  $V^{\otimes r}$  becomes a left  $A_q(n, r)$ -comodule with structure map  $\sigma_r : V^{\otimes r} \rightarrow A_q(n, r) \otimes V^{\otimes r}$  defined by  $\sigma_r(v_{\mathbf{s}}) = \sum_{\mathbf{i} \in I(n, r)} c_{\mathbf{i}, \mathbf{s}} \otimes v_{\mathbf{i}}$  (again  $v_{\mathbf{i}} = v_{i_1} \otimes \dots \otimes v_{i_r} \in V^{\otimes r}$  for  $\mathbf{i} = (i_1, \dots, i_r) \in I(n, r)$ ). The tensor algebra  $T(V)$  is a left  $A_q(n)$ -comodule.

1.6 The ideal  $\langle e_i \otimes e_j - e_j \otimes e_i \mid 1 \leq i, j \leq n \rangle$  is a subcomodule of  $T(E)$ . By  $S_q(E)$  we denote the factor comodule. The ideal  $\langle q(v_i \otimes v_j) - v_j \otimes v_i \mid 1 \leq i < j \leq n \rangle$  is a subcomodule of  $T(V)$ . By  $S_q(V)$  we denote the factor comodule.

**THEOREM.** [2, 2.1.9]

- (i)  $S_q(E)$  is a graded algebra  $S_q(E) = \bigoplus_{0 \leq r} S_q^r(E)$ . The graded part  $S_q^r(E)$  is  $R$ -free with basis  $\{e_{i_1} \dots e_{i_r} \mid i_1 \leq i_2 \leq \dots \leq i_r\}$ .
- (ii)  $S_q(V)$  is a graded algebra  $S_q(V) = \bigoplus_{0 \leq r} S_q^r(V)$ . The graded part  $S_q^r(V)$  is  $R$ -free with basis  $\{v_{i_1} \dots v_{i_r} \mid i_1 \leq i_2 \leq \dots \leq i_r\}$ .

(Here  $e_{i_1} \dots e_{i_r}$  (respectively  $v_{i_1} \dots v_{i_r}$ ) denotes the image of  $e_{i_1} \otimes \dots \otimes e_{i_r}$  (respectively  $v_{i_1} \otimes \dots \otimes v_{i_r}$ ) in  $S_q^r(E)$  (respectively  $S_q^r(V)$ ).

We shall call  $S_q^r(E)$  and  $S_q^r(V)$  the  $q$ -symmetric powers of  $E$  and  $V$ .

2. CALCULATIONS

We want to give a complete description of the subcomodule structure of  $S_q^r(E)$  and  $S_q^r(V)$ . To do so we shall, following Doty [3], determine the subcomodule generated by some basis element  $e_{i_1} \dots e_{i_r} \in S_q^r(E)$  (respectively  $v_{i_1} \dots v_{i_r} \in S_q^r(V)$ ). From now on we shall assume that  $R$  is some field  $k$ .

**2.1** For  $s \in \mathbf{Z}$  let  $[s] = (q^s - 1)/(q - 1) \in k$ . We have that  $[s] = 1 + q + \dots + q^{s-1}$ . For  $s, t \in \mathbf{N}$  let  $\begin{bmatrix} s \\ t \end{bmatrix} = ([s][s-1] \dots [s-t+1])/([1][2] \dots [t])$  (called the Gaussian polynomial). Now we get:

**LEMMA.** Let  $s, t \in \mathbf{N}$ .

- (i) If  $q$  is a non root of unity then  $\begin{bmatrix} s \\ t \end{bmatrix} \neq 0$  for all  $t \leq s$ .
- (ii) If  $q$  is an  $l^{\text{th}}$  root of unity, then write  $s = s_1l + s_0, t = t_1l + t_0$  where  $0 \leq t_0, s_0 < l$ . Then we have

$$\begin{bmatrix} s \\ t \end{bmatrix} = \binom{s_1}{t_1} \begin{bmatrix} s_0 \\ t_0 \end{bmatrix}$$

where  $\binom{s_1}{t_1}$  is the ordinary binomial coefficient. Furthermore we have that

$$\begin{bmatrix} s_0 \\ t_0 \end{bmatrix} \neq 0 \Leftrightarrow s_0 \geq t_0.$$

**PROOF:**

- (i) This is clear.
- (ii) This is proved by induction on  $t$  using that

$$\begin{bmatrix} s \\ t \end{bmatrix} \begin{bmatrix} t \\ 1 \end{bmatrix} = \begin{bmatrix} s \\ t-1 \end{bmatrix} \begin{bmatrix} s-t+1 \\ 1 \end{bmatrix}.$$

□

For more information about Gaussian polynomials see [1].

**2.2** We shall first consider the right comodule. As  $\tau_1(e_j) = \sum_{i=1}^n e_i \otimes c_{i,j}$  ( $1 \leq j \leq n$ )

we have that  $e_1^{a_1} \dots e_n^{a_n} \in S_q^r(E)$  is sent to

$$(*) (e_1 \otimes c_{1,1} + \dots + e_n \otimes c_{n,1})^{a_1} (e_1 \otimes c_{1,2} + \dots + e_n \otimes c_{n,2})^{a_2} \dots (e_1 \otimes c_{1,n} + \dots + e_n \otimes c_{n,n})^{a_n}$$

by  $\tau_r$ . So we shall determine the term in (\*) involving some arbitrary basis element  $e_1^{b_1} \dots e_n^{b_n} \in S_q^r(E)$ .

**LEMMA 2.3.** Let  $\sum_{i=1}^n b_i = a_j$  ( $b_i \in \mathbf{N}$  for all  $i$ ). The term involving  $e_1^{b_1} e_2^{b_2} \dots e_n^{b_n}$  in  $(e_1 \otimes c_{1,j} + e_2 \otimes c_{2,j} + \dots + e_n \otimes c_{n,j})^{a_j}$  is equal to

$$e_1^{b_1} e_2^{b_2} \dots e_n^{b_n} \otimes \begin{bmatrix} a_j \\ b_1 \end{bmatrix} \begin{bmatrix} a_j - b_1 \\ b_2 \end{bmatrix} \begin{bmatrix} a_j - b_1 - b_2 \\ b_3 \end{bmatrix} \dots \begin{bmatrix} a_j - b_1 - b_2 \dots - b_{n-1} \\ b_n \end{bmatrix} c_{1,j}^{b_1} c_{2,j}^{b_2} \dots c_{n,j}^{b_n}$$

PROOF: Assume first  $n = 2$ . So we shall look at  $(e_1 \otimes c_{1,j} + e_2 \otimes c_{2,j})^{a_j}$  for  $1 \leq j \leq 2$ . If  $a_j = 1$  the lemma is true, so by induction we shall assume that the lemma is true for  $a_j - 1$ . Hence the term involving  $e_1^{b_1} e_2^{b_2}$  is equal to

$$\begin{aligned} & \left( e_1^{b_1} e_2^{b_2-1} \otimes \begin{bmatrix} a_j - 1 \\ b_1 \end{bmatrix} \begin{bmatrix} b_2 - 1 \\ b_2 - 1 \end{bmatrix} c_{1,j}^{b_1} c_{2,j}^{b_2-1} \right) (e_2 \otimes c_{2,j}) \\ & + \left( e_1^{b_1-1} e_2^{b_2} \otimes \begin{bmatrix} a_j - 1 \\ b_1 - 1 \end{bmatrix} \begin{bmatrix} b_2 \\ b_2 \end{bmatrix} c_{1,j}^{b_1-1} c_{2,j}^{b_2} \right) (e_1 \otimes c_{1,j}) \\ & = e_1^{b_1} e_2^{b_2} \otimes \left( \begin{bmatrix} a_j - 1 \\ b_1 \end{bmatrix} \begin{bmatrix} b_2 - 1 \\ b_2 - 1 \end{bmatrix} + q^{b_2} \begin{bmatrix} a_j - 1 \\ b_1 - 1 \end{bmatrix} \begin{bmatrix} b_2 \\ b_2 \end{bmatrix} \right) c_{1,j}^{b_1} c_{2,j}^{b_2} \\ & = e_1^{b_1} e_2^{b_2} \otimes \begin{bmatrix} a_j \\ b_1 \end{bmatrix} \begin{bmatrix} b_2 \\ b_2 \end{bmatrix} c_{1,j}^{b_1} c_{2,j}^{b_2} \end{aligned}$$

as  $b_2 = a_j - b_1$ .

Now for the general case. Using the same argument as above we can first move  $c_{1,j}$  to the left, that is, the term involving  $e_1^{b_1} \dots e_n^{b_n}$  will be of the form

$$e_1^{b_1} \dots e_n^{b_n} \otimes \begin{bmatrix} a_j \\ b_1 \end{bmatrix} c_{1,j}^{b_1} (\dots)$$

where  $(\dots)$  does not include  $c_{1,j}$ . Now using the same argument on  $(\dots)$  we can move  $c_{2,j}$  to the left in  $(\dots)$  and we have to multiply by  $\begin{bmatrix} a_j - b_1 \\ b_2 \end{bmatrix}$ . By repeating this we get the lemma. □

**COROLLARY 2.4.** Let  $e^a = e_1^{a_1} \dots e_n^{a_n}$ ,  $e^b = e_1^{b_1} \dots e_n^{b_n} \in S_q^r(E)$ . The term involving  $e^b$  in

$$(e_1 \otimes c_{1,1} + \dots + e_n \otimes c_{n,1})^{a_1} (e_1 \otimes c_{1,2} + \dots + e_n \otimes c_{n,2})^{a_2} \dots (e_1 \otimes c_{1,n} + \dots + e_n \otimes c_{n,n})^{a_n}$$

is a sum of elements of the form

$$e_1^{b_1} e_2^{b_2} \dots e_n^{b_n} \otimes d q^s c_{1,1}^{d_{1,1}} c_{1,2}^{d_{1,2}} \dots c_{1,n}^{d_{1,n}} c_{2,1}^{d_{2,1}} \dots c_{n,n}^{d_{n,n}}$$

where  $\sum_i d_{i,j} = a_j$ ,  $\sum_j d_{i,j} = b_i$ ,  $s$  is some integer and

$$d = \begin{bmatrix} a_1 \\ d_{1,1} \end{bmatrix} \begin{bmatrix} a_1 - d_{1,1} \\ d_{2,1} \end{bmatrix} \dots \begin{bmatrix} a_1 - d_{1,1} - \dots - d_{n-1,1} \\ d_{n,1} \end{bmatrix} \begin{bmatrix} a_2 \\ d_{1,2} \end{bmatrix} \dots \begin{bmatrix} a_n - d_{1,n} - \dots - d_{n-1,n} \\ d_{n,n} \end{bmatrix}.$$

PROOF: From the lemma it follows that we have a sum of elements of the form

$$e_1^{b_1} e_2^{b_2} \dots e_n^{b_n} \otimes d_{c_{1,1}^{d_{1,1}} c_{2,1}^{d_{2,1}} \dots c_{n,1}^{d_{n,1}} c_{1,2}^{d_{1,2}} \dots c_{n,n}^{d_{n,n}}}.$$

And as  $c_{i,j}c_{s,t} = qc_{s,t}c_{i,j}$  for  $i > s$  and  $j < t$  the corollary follows. □

2.5 Now consider the left comodule case. As  $\sigma_1(v_j) = \sum_{i=1}^n c_{j,i} \otimes v_i$  ( $1 \leq j \leq n$ ) we

have that  $v_1^{a_1} \dots v_n^{a_n} \in S_q^r(V)$  is sent to

(\*)

$$(c_{1,1} \otimes v_1 + \dots + c_{1,n} \otimes v_n)^{a_1} (c_{2,1} \otimes v_1 + \dots + c_{2,n} \otimes v_n)^{a_2} \dots (c_{n,1} \otimes v_1 + \dots + c_{n,n} \otimes v_n)^{a_n}$$

by  $\sigma_r$ . So we shall describe the term in (\*) involving some arbitrary basis element  $v_1^{b_1} \dots v_n^{b_n} \in S_q^r(V)$ .

LEMMA 2.6. Let  $\sum_{i=1}^n b_i = a_j$  ( $b_i \in \mathbb{N}$  for all  $i$ ). Then the term involving  $v_1^{b_1} v_2^{b_2} \dots v_n^{b_n}$  in  $(c_{j,1} \otimes v_1 + c_{j,2} \otimes v_2 + \dots + c_{j,n} \otimes v_n)^{a_j}$  is equal to

$$c_{j,1}^{b_1} c_{j,2}^{b_2} \dots c_{j,n}^{b_n} \otimes \begin{bmatrix} a_j \\ b_1 \end{bmatrix} \begin{bmatrix} a_j - b_1 \\ b_2 \end{bmatrix} \begin{bmatrix} a_j - b_1 - b_2 \\ b_3 \end{bmatrix} \dots \begin{bmatrix} a_j - b_1 - b_2 \dots - b_{n-1} \\ b_n \end{bmatrix} v_1^{b_1} v_2^{b_2} \dots v_n^{b_n}.$$

PROOF: Assume first  $n = 2$ . If  $a_j = 1$  the lemma is true, so by induction we shall assume that the lemma is true for  $a_j - 1$ . Hence the term involving  $v_1^{b_1} v_2^{b_2}$  is equal to

$$\begin{aligned} & \left( c_{j,1}^{b_1-1} c_{j,2}^{b_2} \otimes \begin{bmatrix} a_j - 1 \\ b_1 - 1 \end{bmatrix} \begin{bmatrix} b_2 \\ b_2 \end{bmatrix} v_1^{b_1-1} v_2^{b_2} \right) (c_{j,1} \otimes v_1) \\ & + \left( c_{j,1}^{b_1} c_{j,2}^{b_2-1} \otimes \begin{bmatrix} a_j - 1 \\ b_1 \end{bmatrix} \begin{bmatrix} b_2 - 1 \\ b_2 - 1 \end{bmatrix} v_1^{b_1} v_2^{b_2-1} \right) (c_{j,2} \otimes v_2) \\ & = c_{j,1}^{b_1} c_{j,2}^{b_2} \otimes \left( q^{b_2} \begin{bmatrix} a_j - 1 \\ b_1 - 1 \end{bmatrix} \begin{bmatrix} b_2 \\ b_2 \end{bmatrix} + \begin{bmatrix} a_j - 1 \\ b_1 \end{bmatrix} \begin{bmatrix} b_2 - 1 \\ b_2 - 1 \end{bmatrix} \right) v_1^{b_1} v_2^{b_2} \\ & = c_{j,1}^{b_1} c_{j,2}^{b_2} \otimes \begin{bmatrix} a_j \\ b_1 \end{bmatrix} \begin{bmatrix} b_2 \\ b_2 \end{bmatrix} v_1^{b_1} v_2^{b_2} \end{aligned}$$

as  $b_2 = a_j - b_1$ .

Now for the general case the lemma follows by the same argument as in 2.3. □

COROLLARY 2.7. Let  $v^a = v_1^{a_1} \dots v_n^{a_n}, v^b = v_1^{b_1} \dots v_n^{b_n} \in S_q^r(V)$ . The term involving  $v^b$  in

$$(c_{1,1} \otimes v_1 + \dots + c_{1,n} \otimes v_n)^{a_1} (c_{2,1} \otimes v_1 + \dots + c_{2,n} \otimes v_n)^{a_2} \dots (c_{n,1} \otimes v_1 + \dots + c_{n,n} \otimes v_n)^{a_n}$$

is a sum of elements of the form

$$c_{1,1}^{d_{1,1}} c_{1,2}^{d_{1,2}} \dots c_{1,n}^{d_{1,n}} c_{2,1}^{d_{2,1}} \dots c_{n,n}^{d_{n,n}} \otimes d q^s v_1^{b_1} v_2^{b_2} \dots v_n^{b_n}$$

where  $\sum_j d_{i,j} = a_i, \sum_i d_{i,j} = b_j, s$  is some integer and

$$d = \begin{bmatrix} a_1 \\ d_{1,1} \end{bmatrix} \begin{bmatrix} a_1 - d_{1,1} \\ d_{1,2} \end{bmatrix} \dots \begin{bmatrix} a_1 - d_{1,1} - \dots - d_{1,n-1} \\ d_{1,n} \end{bmatrix} \begin{bmatrix} a_2 \\ d_{2,1} \end{bmatrix} \dots \begin{bmatrix} a_n - d_{n,1} - \dots - d_{n,n-1} \\ d_{n,n} \end{bmatrix}.$$

PROOF: From the lemma it follows that we have a sum of elements of the form

$$c_{1,1}^{d_{1,1}} c_{1,2}^{d_{1,2}} \dots c_{1,n}^{d_{1,n}} c_{2,1}^{d_{2,1}} \dots c_{n,n}^{d_{n,n}} \otimes d v_1^{d_{1,1}} v_2^{d_{1,2}} \dots v_n^{d_{1,n}} v_1^{d_{2,1}} \dots v_n^{d_{n,n}}.$$

And as  $v_i v_j = q v_j v_i$  for  $1 \leq j < i \leq n$  the corollary follows. □

**2.8** Let  $e^a, e^b$  (respectively  $v^a, v^b$ ) be basis elements of  $S_q^r(E)$  (respectively  $S_q^r(V)$ ). From 2.4 and 2.7 it follows that

$$e^b \in \langle e^a \rangle \Leftrightarrow v^b \in \langle v^a \rangle$$

where  $\langle e^a \rangle$  (respectively  $\langle v^a \rangle$ ) is the subcomodule of  $S_q^r(E)$  (respectively  $S_q^r(V)$ ) generated by  $e^a$  (respectively  $v^a$ ). Hence the subcomodule structure of  $S_q^r(E)$  and  $S_q^r(V)$  is the same. From now on we shall only consider the right comodule case.

### 3. THE MAIN RESULT

In this section we shall give a complete description of the subcomodule structure of the  $q$ -symmetric powers. In 2.4 we have described the subcomodule of  $S_q^r(E)$  generated by some basis element. Now we want to know when the coefficient  $d$  in 2.4 is non zero. We shall do so by introducing the carry pattern. If the quantum parameter  $q \in k$  is a non root of unity, then there is no work to be done (see 3.1). If, however,  $q$  is a root of unity, then we shall consider two cases: when  $k$  is of characteristic 0 (see 3.2–3.6) and when  $k$  is of prime characteristic (see 3.7–3.10).

**THEOREM 3.1.** *Let  $q$  be a non root of unity. Then  $S_q^r(E)$  is irreducible for each  $r \geq 0$ .*

PROOF: By 2.1 and 2.4 it follows that the subcomodule generated by some arbitrary basis element  $e^a \in S_q^r(E)$  is equal to  $S_q^r(E)$ . □

From now on we shall assume that  $q$  is an  $l$ 'th root of unity ( $l \geq 1$ ).

**3.2** Assume first that  $k$  is of characteristic 0. Let  $e^a = e_1^{a_1} e_2^{a_2} \dots e_n^{a_n} \in S_q^r(E)$ . For each  $1 \leq i \leq n$  write  $a_i = a_i^1 l + a_i^0$  where  $0 \leq a_i^0 < l$ . Furthermore let  $r = r_1 l + r_0$  where  $0 \leq r_0 < l$ .

DEFINITION:  $C(e^a)$  is defined by

$$\sum_{i=1}^n a_i^0 = C(e^a)l + r_0$$

Following [3], we call  $C(e^a)$  the carry pattern of  $e^a$ .

**PROPOSITION 3.3.** Let  $e^a = e_1^{a_1} e_2^{a_2} \dots e_n^{a_n}$ ,  $e^b = e_1^{b_1} e_2^{b_2} \dots e_n^{b_n} \in S_q^r(E)$ . Then  $C(e^a) \geq C(e^b)$  if and only if there exists  $d_{i,j} \in \mathbb{N}$  ( $1 \leq i, j \leq n$ ) such that  $\sum_{i=1}^n d_{i,j} = a_j$ ,  $\sum_{j=1}^n d_{i,j} = b_i$  and

$$\begin{aligned} \begin{bmatrix} a_1 \\ d_{1,1} \end{bmatrix} \begin{bmatrix} a_1 - d_{1,1} \\ d_{2,1} \end{bmatrix} \dots \begin{bmatrix} a_1 - d_{1,1} - \dots - d_{n-1,1} \\ d_{n,1} \end{bmatrix} \begin{bmatrix} a_2 \\ d_{1,2} \end{bmatrix} \\ \dots \begin{bmatrix} a_n - d_{1,n} - \dots - d_{n-1,n} \\ d_{n,n} \end{bmatrix} \neq 0. \end{aligned}$$

PROOF: Assume  $C(e^a) \geq C(e^b)$ . This means that  $\sum_{i=1}^n a_i^0 \geq \sum_{i=1}^n b_i^0$ . For  $1 \leq i, j \leq n$  we shall define  $d_{i,j}$ . If  $b_i \leq a_i$  set  $d_{i,i} = b_i^1 l + b_i^0$  if  $b_i^0 \leq a_i^0$  and set  $d_{i,i} = b_i^1 l + a_i^0$  otherwise. If  $b_i > a_i$  set  $d_{i,i} = a_i^1 l + a_i^0$  if  $b_i^0 > a_i^0$  and set  $d_{i,i} = a_i^1 l + b_i^0$  otherwise. As  $\sum_{i=1}^n a_i^0 \geq \sum_{i=1}^n b_i^0$  we see by 2.1 (ii) that we can choose  $d_{i,j}$  ( $i \neq j$ ) such that we get the statement in the proposition.

Assume now that there exist  $d_{i,j} \in \mathbb{N}$  ( $1 \leq i, j \leq n$ ) as required in the proposition. For each  $d_{i,j}$  we write  $d_{i,j} = d_{i,j}^1 l + d_{i,j}^0$  where  $0 \leq d_{i,j}^0 < l$ . For each  $1 \leq j \leq n$

$$\begin{bmatrix} a_j \\ d_{1,j} \end{bmatrix} \begin{bmatrix} a_j - d_{1,j} \\ d_{2,j} \end{bmatrix} \dots \begin{bmatrix} a_j - d_{1,j} - \dots - d_{n-1,j} \\ d_{n,j} \end{bmatrix} \neq 0.$$

Hence by 2.1 (ii) we have  $a_j^0 \geq d_{1,j}^0 + \dots + d_{n,j}^0$ . But as  $\sum_{i=1}^n d_{i,j} = a_j$  we must then have that

$$a_j^0 = d_{1,j}^0 + \dots + d_{n,j}^0$$

for each  $1 \leq j \leq n$ . Now as  $\sum_{j=1}^n d_{i,j} = b_i$  we have (for  $1 \leq i \leq n$ ) that  $\sum_{j=1}^n d_{i,j}^0 = n_i l + b_i^0$  where  $n_i \geq 0$ . Hence

$$\sum_{j=1}^n a_j^0 = \sum_{j=1}^n \sum_{i=1}^n d_{i,j}^0 = \sum_{i=1}^n \sum_{j=1}^n d_{i,j}^0 = \sum_{i=1}^n (n_i l + b_i^0) = \sum_{i=1}^n n_i l + \sum_{i=1}^n b_i^0,$$





**3.7** Let now  $k$  be of prime characteristic  $p$ . Let  $e^a = e^{a_1} \dots e^{a_n} \in S_q^r(E)$ , and for each  $1 \leq i \leq n$  write  $a_i = a_i^1 l + a_i^0$  where  $0 \leq a_i^0 < l$ . Furthermore let  $a_i^1 = \sum_j c_i^{1,j} p^j$  where  $0 \leq a_i^{1,j} < p$  for all  $i, j$ . Let  $r = r_1 l + r_0$ ,  $0 \leq r_0 < l$  and  $r_1 = \sum_j r_1^j p^j$ ,  $0 \leq r_1^j < p$  for all  $j$ .

Let  $C_0(e^a)$  be defined by

$$\sum_{i=1}^n a_i^0 = C_0(e^a)l + r_0,$$

and for  $t \geq 1$  define  $C_t(e^a)$  by

$$C_{t-1}(e^a) + \sum_{i=1}^n a_i^{1,t-1} = C_t(e^a)p + r_1^{t-1}.$$

**DEFINITION:** The carry pattern of  $e^a$  is defined to be

$$C(e^a) = (C_0(e^a), C_1(e^a), \dots, C_m(e^a))$$

where  $m$  is the biggest  $j$  such that  $r_1^j > 0$ .

The carry pattern is defined as a mixture of the carry pattern in characteristic 0, and the carry pattern defined in [3].

**EXAMPLE.** Let  $n = 3$ ,  $r = 99$ ,  $p = 3$  and  $l = 5$ . Let  $e^a = e_1^{29} e_2^{37} e_3^{33}$  and  $e^b = e_1^{26} e_2^{52} e_3^{21}$ . Then we have that

$$C(e^a) = (1, 1, 2) \text{ and } C(e^b) = (0, 1, 1).$$

**3.8** Let  $C(r)$  be the set of carry patterns which occur as carry pattern for some  $e^a \in S_q^r(E)$ . We let  $C(r)$  be partially ordered by  $c \leq c'$  if and only if  $c_i \leq c'_i$  for all  $i$  (where  $c = (c_0, c_1, \dots, c_m)$  and  $c' = (c'_0, c'_1, \dots, c'_m)$ ).

**PROPOSITION.** Let  $e^a, e^b$  be basis elements of  $S_q^r(E)$ . Then

$$e^b \in \langle e^a \rangle \Leftrightarrow C(e^b) \leq C(e^a).$$

**PROOF:** The proposition can be proved as in 3.3. If  $s, t \in \mathbb{N}$  then by 2.1

$$\begin{bmatrix} s \\ t \end{bmatrix} = \begin{pmatrix} s_1 \\ t_1 \end{pmatrix} \begin{bmatrix} s_0 \\ t_0 \end{bmatrix}$$

where  $s = s_1 l + s_0$ ,  $t = t_1 l + t_0$ ,  $0 \leq s_0, t_0 < l$ .

$C_0(e^a)$  plays exactly the same role as the carry pattern does in the characteristic 0 case, and can be handled as in the proof of 3.3. But we have to be a bit more careful with the binomial coefficient. We have that

$$\binom{s_1}{t_1} \equiv \binom{s_1^m}{t_1^m} \binom{s_1^{m-1}}{t_1^{m-1}} \cdots \binom{s_1^0}{t_1^0} \pmod p$$

where  $s_1 = \sum_j s_1^j p^j$ ,  $t_1 = \sum_j t_1^j p^j$ ,  $0 \leq s_1^j, t_1^j < p$  for all  $j$ .

The proof of 3.3 uses the fact that  $\begin{bmatrix} s_0 \\ t_0 \end{bmatrix} \neq 0 \Leftrightarrow s_0 \geq t_0$ . As  $\binom{s_1^j}{t_1^j} \neq 0 \Leftrightarrow t_1^j \leq s_1^j$  the present proposition can be proved using the proof of 3.3 on each level of the  $p$ -adic expansion. □

**3.9** For  $c \in C(r)$  let  $L(c)$  denote the span of all  $e^a \in S_q^r(E)$ , where  $C(e^a) = c$ . For a subset  $B \subset C(r)$  let  $T(B) = \sum_{c \in B} L(c)$ . We say that the subset  $B$  is order closed if it contains the predecessors of all its elements under the given order relation. As a consequence of 3.8 we now have a complete description of the subcomodule structure of  $S_q^r(E)$ .

**THEOREM.** *We have a 1-1 correspondence*


$$B \leftrightarrow T(B)$$

*between the set of order closed subsets of  $C(r)$  and the set of subcomodules of  $S_q^r(E)$ .*

In particular, the irreducible composition factors of  $S_q^r(E)$  are in one-to-one correspondence with  $L(c)$  for  $c \in C(r)$ , and all composition factors have multiplicity one.

**3.10** We can determine  $C(r)$  and hence the order closed subsets of  $C(r)$  by combining 3.6 with Lemma 3 in [3].

**EXAMPLE.** Let  $n = 3$ ,  $l = 5$ ,  $p = 7$  and  $r = 51 = (1 p + 3)l + 1$ . Then  $C(r) = \{(0, 0), (0, 1), (1, 0), (1, 1), (2, 0), (2, 1)\}$ . Hence by 3.9 the subcomodule structure of  $S_q^{51}(E)$  can be described by



REFERENCES

[1] G.E. Andrews, *The theory of partitions*, Encyclopedia Math. **2** (Addison-Wesley, 1976).  
 [2] R. Dipper, S. Donkin, 'Quantum  $GL_n$ ', *Proc. London Math. Soc.* **63** (1991), 165–211.  
 [3] S. Doty, 'Submodules of symmetric powers of the natural module for  $GL_n$ ', in *Contemp. Math.* **88** (American Mathematical Society, Providence, R.I., 1989), pp. 185-191.

School of Mathematics and Statistics  
 The University of Sydney  
 New South Wales 2006  
 Australia