

CORRIGENDUM

The utilization of large savanna trees by elephant in southern Kruger National Park

Graeme Shannon, Dave J. Druce, Bruce R. Page, Holger C. Eckhardt, Rina Grant and Rob Slotow
doi:10.1017/S0266467408004951, Published by Cambridge University Press, 13 May 2008.

On page 284, Table 1; the decimal point had been omitted from the second column “Density (trees ha⁻¹). The correct table is featured below.

Table 1. Patterns of elephant utilization for the five most common species (*Acacia nigrescens*, *Spirostachys africana*, *Combretum apiculatum*, *Sclerocarya birrea* and *Terminalia sericea*) and six height classes. Utilization, pushing and debarking are recorded as proportions of the total within each category.

Species	Density (trees ha ⁻¹)	Utilized	Pushed (% of those which are dead)	Debarked	Mean intensity of utilization (%)
<i>Acacia nigrescens</i>	12.4	0.56	0.14 (54)	0.16	18
<i>Combretum apiculatum</i>	8.5	0.57	0.24 (14)	0.03	18
<i>Sclerocarya birrea</i>	5.7	0.75	0.13 (26)	0.05	21
<i>Spirostachys africana</i>	5.3	0.44	0.03 (41)	0.08	7
<i>Terminalia sericea</i>	2.0	0.50	0.01 (0)	0.05	7
Height class (m)					
5–6.5	22.7	0.57	0.09 (27)	0.09	14
6.5–8.0	12.6	0.59	0.09 (37)	0.10	14
8.0–9.5	6.9	0.58	0.15 (35)	0.10	17
9.5–11.0	2.4	0.55	0.15 (42)	0.17	16
11.0–12.5	1.1	0.66	0.14 (63)	0.21	20
>12.5	0.4	0.67	0.0 (0)	0.19	12

REFERENCE

SHANNON, G., DRUCE, D. J., PAGE, B. R., ECKHARDT, H. C., GRANT, R. & SLOTOW, R. The utilization of large savanna trees by elephant in southern Kruger National Park. *Journal of Tropical Ecology* 24:281–289.
Published by Cambridge University Press, 13 May 2008. doi:10.1017/S0266467408004951