

Evidence on the prevalence and geographic distribution of major cardiovascular risk factors in Italy

Roberta Laccetti^{1,*}, Andrea Pota², Saverio Stranges³, Claudio Falconi¹, Bruno Memoli², Leopoldo Bardaro⁴ and Bruna Guida¹

¹Department of Neuroscience, Physiology-Nutrition Section, University 'Federico II' of Naples, Via S. Pansini 5, 80131 Naples, Italy; ²Department of Nephrology, University 'Federico II' of Naples, Naples, Italy; ³Health Sciences Research Institute, University of Warwick Medical School, Coventry, UK; ⁴University Claude Bernard-Lyon1, Lyon, France

Submitted 22 March 2011: Final revision received 22 January 2012: Accepted 30 March 2012: First published online 30 May 2012

Abstract

Objective: To assess the prevalence and geographic distribution of major cardiovascular risk factors in a large community-wide sample of the Italian population.

Design: A cross-sectional survey. Standardized methods were used to collect and measure cardiovascular risk factors. Data were adjusted for survey weightings. Qualitative and quantitative variables were compared with parametric and non-parametric tests, as appropriate.

Setting: Towns (n 193) across different Italian regions.

Subjects: Unselected adults (n 24 213; 12 626 men; 11 587 women) aged 18–98 years (mean age 56.9 (sd 15.3) years), who volunteered to participate in a community-wide screening programme over a 2 d period in 2007.

Results: Overall, the prevalence of major cardiovascular risk factors was: obesity, 22.7% (women 18.9%, men 26.1%); overweight, 44.7% (women 31.6%, men 56.7%); hypertension, 59.6% (women 48.3%, men 70.0%); dyslipidaemia, 59.1% (women 57.7%, men 60.3%); diabetes, 15.3% (women 11.2%, men 19.0%) and smoking, 19.8% (women 14.0%, men 25.2%). We found a high prevalence of unhealthy eating habits; fruit and vegetable consumption was below the recommended range in 60% of the study population. Ninety per cent of the study population had more than one cardiovascular risk factor and 84% had between two and five cardiovascular risk factors. There were differences among Italian macro-areas mainly for obesity, hypertension, dyslipidaemia and diabetes.

Conclusions: The study provides alarming evidence on current prevalence data for major cardiovascular risk factors in a large sample of the Italian population. Particularly, obesity and hypertension represent a relevant public health problem. There is a pressing need for effective preventive health measures which must also take into account the differences among Italian macro-areas.

Keywords

Cardiovascular risk factors
Italy
Nutrition
Population
Epidemiology

Mortality rates from CVD have halved in Western Europe in the last 20 years⁽¹⁾. Many factors that predispose to the development of CVD have been thoroughly investigated⁽²⁾. Some of these cardiovascular risk factors (CRF) cannot be modified (advanced age, male gender, heredity), while some can be modified (smoking, hypertension, dyslipidaemia, diabetes mellitus, physical inactivity, overweight and obesity). Additional modifiable CRF include specific dietary patterns such as wine, fruit and vegetable consumption as well as physiological factors such as stress.

Europe has been experiencing a marked increase in obesity rates, which have doubled over just a few years⁽³⁾.

This continuing trend in obesity is a critical public health threat because of its potential transition into higher prevalences of diabetes and hypercholesterolaemia⁽⁴⁾. Furthermore, recent national survey data show that hypertension prevalence is much higher in European countries than in North America⁽⁵⁾.

Few published data are available to characterize current CRF prevalence and to monitor trends in CRF in the Italian population^(6–9). The aims of the present study were therefore to assess the prevalence of major CRF in a large community-wide sample of the Italian population and to examine potential differences in the distribution of CRF across Italian macro-areas.

*Corresponding author: Email robertalaccetti@hotmail.com

Methods

Study population

The present cross-sectional survey included 24 213 unselected adults (12 626 men and 11 587 women) aged 18 years and older (mean age 56.9 (SD 15.3) years; median (25th, 75th percentile) 59 (46, 68) years), who volunteered to participate in a community-wide screening programme to promote healthy lifestyle ('Misuriamoci'), organized by the Italian Red Cross and supported by the Italian Ministry of Health. Participants were recruited in 193 Italian towns over a 2 d period in March 2007 in response to an extensive media campaign. Advertising was conducted by radio and television announcements, newspapers, magazines, billboards and bus posters.

Measurements

Standardized methods were used to collect and measure risk factors.

Weight and height were measured without shoes and heavy clothing; a standard electronic scale and a wall height ruler were used respectively for weight and height. BMI (kg/m^2) was calculated as weight (in kilograms) divided by the square of height (in metres). The classification of overweight and obesity was based on the following categories⁽¹⁰⁾: overweight as $\text{BMI} = 25.0\text{--}29.9 \text{ kg}/\text{m}^2$ and obesity as $\text{BMI} \geq 30.0 \text{ kg}/\text{m}^2$.

Waist circumference was measured with an anelastic tape with the participant in standing position, at the level of the umbilicus. According to the National Cholesterol Education Program Adult Treatment Panel III criteria⁽¹¹⁾, abdominal obesity was defined as waist circumference ≥ 102 cm in men and ≥ 88 cm in women.

Blood pressure was recorded as the average of two measurements after participants had been sitting for 5 min, applying an appropriately sized cuff on the right arm. The average of the two consecutive readings was used in the analysis. Hypertensive participants were defined as those with systolic/diastolic blood pressure $\geq 140/\geq 90$ mmHg or on regular antihypertensive treatment⁽¹²⁾.

Fasting cholesterol, TAG and glucose levels were determined from finger-prick capillary blood samples using a Multicare[®] diagnostic device (<http://www.multicare.it>). High blood total cholesterol was defined as $\geq 200 \text{ mg}/\text{dl}$ ⁽¹³⁾. Hypertriglycerolaemia was considered as serum TAG $\geq 150 \text{ mg}/\text{dl}$ ⁽¹¹⁾. Lipid-lowering therapy was defined if participants reported taking lipid-lowering drugs. Dyslipidaemia was considered as participants reporting high blood total cholesterol and/or high serum TAG and/or taking lipid-lowering medication^(11,13). Impaired fasting glucose (hyperglycaemia) was defined as a fasting plasma glucose level in the range 100.0–125.9 mg/dl and diabetes was considered as fasting plasma glucose $\geq 126 \text{ mg}/\text{dl}$ or if participants reported taking antidiabetic drugs⁽¹⁴⁾.

For wine, fruit and vegetable consumption, a commonly used serving size was specified and participants

were asked 'if and how often they had consumed that unit'. Current wine consumption was recorded as glasses/week (one glass is defined as 4 fl oz of wine or a drink equivalent). An average daily intake of 1 or 2 drinks for men and 0.5 to 1 drink for women was considered as recommended⁽¹⁵⁾. Daily and/or weekly consumption was asked for fruit and vegetables. An average of 8–10 servings/d for fruit and vegetables was considered as recommended⁽¹⁶⁾.

Current smokers were defined as participants who smoked one or more cigarettes daily.

As regards geographic distribution, data were analysed as a whole and by the following macro-areas (i.e. areas comprising more regions, generally contiguous): North-west (Valle d'Aosta, Piemonte, Liguria and Lombardia); North-east (Trentino-Alto Adige, Friuli-Venezia Giulia, Veneto and Emilia Romagna); Centre (Toscana, Marche, Umbria and Lazio); South (Campania, Abruzzo, Molise, Calabria, Puglia and Basilicata); and Islands (Sicily and Sardinia).

Statistical analysis

Data are expressed as means and standard deviations or as frequencies in presenting continuous or categorical variables, and are adjusted by age, averaging the 5-year age groups and stratifying the data for men and women. All the data have been weighted using the Italy population pyramid for 2005 (US Census Bureau, international database); in particular, we stratified by age and sex and then we performed a weighted average.

A survey regression analysis was used for averaged age and gender adjustment for each town unit and then we adjusted for age and gender at the individual level. Separate prevalence results are presented for men and women in single macro-areas and linear and logistic regression models were performed to obtain comparisons and to take into account the role of potential confounders (age and sex).

A *post hoc* power calculation indicates that the available sample allows analysis of variance comparison tests to detect a difference of 5% between groups, with power $>94\%$ and $\alpha < 0.05$.

Analyses were performed using the statistical software package STATA 9.2 SE (StataCorp., College Station, TX, USA). A *P* value < 0.05 was considered significant.

Results

The main demographic, clinical and biochemical characteristics of the study population according to gender and macro-area are shown in Table 1. Overall, average BMI was 26.9 (SD 4.5) kg/m^2 and men had a higher mean BMI than women (27.6 (SD 4.1) *v.* 26.3 (SD 4.8) kg/m^2). Men and women on average exceeded the sex-specific waist circumference cut-off (mean: 101 (SD 11) cm in men;




Fig. 1 Maps showing (a) the number of surveyed towns and macro-areas (■, Islands; □, South; ■, Centre; ■, North-west; ■, North-east) and (b) geographical differences in the prevalence (■, very high; ■, high; □, low) of cardiovascular risk factors according to macro-area; Italy, 2007

(iii) dyslipidaemia; (iv) diabetes and hyperglycaemia; (v) smoking; (vi) fruit and vegetable consumption lower than the recommended intake; and (vii) wine consumption higher than the recommended intake. The prevalence of having between two and five CRF was 84%.

Table 4 describes the prevalence of having zero to seven CRF (obesity/overweight/high waist circumference; hypertension; dyslipidaemia; diabetes/hyperglycaemia; smoking; fruit and vegetable consumption lower than the recommended intake; wine consumption higher than the recommended intake) among Italian macro-areas and the effect of macro-area on the distribution of CRF in the study population.

Consistent with previous studies^(17,18), when stratifying our data into three age groups, i.e. 18–39, 40–65 and >65 years, we observed an increased prevalence of some CFR with age. Particularly, the prevalence of hypertension, diabetes, wine consumption and fruit and vegetable consumption increased with age, while smoking decreased with age and dyslipidaemia reached a peak in the 40–65-year age group (see Appendices 1 to 3).

Discussion

The present study provides novel evidence about the current prevalence and distribution of major CRF in a

Table 2 Prevalence of cardiovascular risk factors in the study participants stratified by gender and macro-area; Italy, 2007

	Study population			North-west			North-east			Centre			South			Islands			P‡
	Men	Women	P†	Men	Women	P†	Men	Women	P†	Men	Women	P†	Men	Women	P†	Men	Women	P†	
Obesity (%)	26.1	18.9	0.003	20.5	15.2	0.041	19.9	12.9	0.001	24.2	18.1	0.010	31.5	23.5	0.020	24.6	16.8	0.004	0.001
Overweight (%)	56.7	31.6	0.001	55.3	29.2	0.001	56.0	32.2	0.001	57.5	32.5	0.016	52.6	32.3	0.017	57.5	30.8	0.010	0.001
High waist circumference (%)	57.0	56.4	0.001	53.8	53.0	0.001	52.9	51.8	0.001	53.6	58.1	0.028	60.3	60.6	0.022	61.5	53.9	0.044	0.001
Hypertension (%)	70.0	48.3	0.001	73.4	52.1	0.001	73.4	51.0	0.001	72.5	47.0	0.022	69.3	48.2	0.009	61.0	40.6	0.010	0.001
Hypercholesterolaemia (%)	26.8	32.7	0.001	29.0	35.2	0.001	29.7	34.9	0.001	26.5	33.2	0.028	25.9	29.3	0.001	24.6	32.9	0.001	0.001
Hypertriacylglycerolaemia (%)	47.8	43.1	0.002	48.7	48.1	0.022	52.0	45.8	0.002	49.6	45.8	0.010	46.7	38.5	0.512	43.1	36.5	0.248	0.001
Lipid-lowering therapy (%)	9.5	9.0	0.007	9.5	10.5	0.001	6.5	7.8	0.001	10.0	8.5	0.372	9.7	9.2	0.030	10.6	9.6	0.488	0.001
Dyslipidaemia (%)	60.3	57.7	0.001	63.5	63.3	0.001	65.2	60.1	0.001	60.9	59.9	0.001	58.2	52.2	0.010	56.2	53.7	0.003	0.001
Diabetes (%)	19.0	11.2	0.001	18.1	11.7	0.004	15.3	10.4	0.036	16.8	9.6	0.001	21.5	12.6	0.021	20.6	11.3	0.014	0.001
Hyperglycaemia (%)	15.5	10.8	0.019	16.7	10.7	0.006	14.7	8.6	0.030	21.7	15.2	0.003	12.2	9.0	0.098	12.7	7.6	0.040	0.001
Smoking (%)	25.2	14.0	0.001	18.8	11.6	0.010	20.6	13.7	0.011	23.2	16.8	0.022	28.4	11.9	0.001	30.5	17.9	0.001	0.001

†Survey regression analysis adjusted for each town unit for average age and gender.

‡Linear and logistic regression models adjusted for age and gender.

Table 3 Prevalence of fruit and vegetable consumption within the recommended range and wine consumption higher than the recommended intake in the study participants stratified by gender and macro-area; Italy, 2007

	Study population			North-west			North-east			Centre			South			Islands			P‡
	Men	Women	P†	Men	Women	P†	Men	Women	P†	Men	Women	P†	Men	Women	P†	Men	Women	P†	
Fruit and vegetables (%)§	38.4	41.0	0.028	51.7	57.3	0.033	49.9	51.4	0.018	43.6	46.8	0.029	24.3	26.0	0.033	29.1	32.3	0.040	0.021
Wine (%)																			
Abstainer	74.7	78.4		64.6	76.2		70.1	75.9		73.9	74.3		78.1	84.7		80.7	84.8		
Moderate	23.2	15.5	0.020	32.6	12.2	0.001	26.2	12.7	0.029	22.4	13.4	0.044	18.7	11.2	0.017	18.1	9.9	0.023	0.042
Habitual	2.1	6.1		2.8	11.6		3.7	11.4		3.7	12.3		3.2	4.1		1.2	5.3		

†Survey regression analysis adjusted for each town unit for average age and gender.

‡Linear and logistic regression models adjusted for age and gender.

§Fruit and vegetable consumption within the recommended range of 8–10 servings/d.

||Wine consumption higher than the recommended intake: moderate, 1–2 drinks/d for men and 0.5–1 drink/d for women; habitual, >2 drinks/d for men and >1 drink/d for women.




Fig. 2 Prevalence of (from zero to seven) cardiovascular risk factors (CRF; obesity/overweight/high waist circumference, hypertension, dyslipidaemia, diabetes/hyperglycaemia, smoking, fruit and vegetable consumption lower than the recommended intake, wine consumption higher than the recommended intake) in the overall population (■) and by gender (□, men; ▒, women); Italy, 2007

Table 4 Prevalence (%) of (from zero to seven) of cardiovascular risk factors (CRF; obesity/overweight/high waist circumference, hypertension, dyslipidaemia, diabetes/hyperglycaemia, smoking, fruit and vegetable consumption lower than the recommended intake, wine consumption higher than the recommended intake) in the study participants stratified by macro-area; Italy, 2007

Number of CRF	North-west	North-east	Centre	South	Islands
0	1.77	1.84	2.02	1.06	1.57
1	9.45	10.05	8.51	8.00	8.85
2	20.56	20.41	18.88	18.29	21.05
3	27.04	28.73	27.10	27.26	28.95
4	23.09	22.82	24.43	26.56	24.85
5	13.13	11.51	13.71	14.26	11.29
6	4.18	3.53	4.54	4.04	2.63
7	0.78	1.12	0.82	0.53	0.81

Effect of macro-area on the distribution of CRF in the Italian study population: $P < 0.001$ among macro-areas, adjusted for age and gender.

large community-wide sample of the Italian adult population, including 24 213 participants. Altogether the findings are worrisome and clearly indicate that obesity and hypertension represent a relevant public health problem in Italy. However, results should be interpreted with caution because of self-selection bias; this may be a possible source of discrepancy with other national results.

There was a consistent north-to-south increasing gradient in the prevalence of obesity, diabetes and smoking, while a consistent south-to-north increasing gradient was found in the prevalence of hypertension and dyslipidaemia (Table 2, Fig. 1). Wine and fruit and vegetable consumption was significantly higher in the northern macro-areas than in the south (Table 3).

Comparing our results with data from the National Institute for Statistics (ISTAT) survey in 2007⁽⁷⁾ reveals that the prevalence of obesity has increased considerably over the last few years (overall 22.7% *v.* 9.8%), while the prevalence of overweight has increased by a smaller degree (overall 44.7% *v.* 34.2%). This discrepancy in obesity prevalence may be due to self-reported data

collected by ISTAT, compared with direct measurements used in our study. In fact, obese individuals tend to overestimate their height and to underestimate their weight, leading to an underestimation of obesity prevalence in studies based on self-report measurements⁽¹⁹⁾. Moreover, comparing our data with previous studies^(8,9,20) using direct measurements, almost the same prevalence of obesity and overweight was found. According to previous reports the prevalence of obesity was quite similar between men and women, while the prevalence of overweight was nearly double in men compared with women^(7,9). Recent changes in lifestyle in Western and developing countries, such as physical inactivity and unhealthy eating habits, are likely to account for the increasing prevalence of obesity. Nowadays, women have more stringent weight standards and engage in more efforts to avoid being overweight⁽²¹⁾.

Consistent with the ISTAT data⁽⁷⁾, the Southern population showed the highest prevalence of obesity in the present study while the lowest prevalence of overweight was found in the North-east population. We believe that excessive total energy intake and a low degree of physical activity may account for the high prevalence of obesity in the Southern population^(17,22,23).

By comparing our results with data from the Italian Epidemiologic Cardiovascular Observatory (OEC) as part of the Progetto CUORE⁽⁸⁾, we found an increase in the mean waist circumference value both in men (101 *v.* 95 cm) and women (91 *v.* 85 cm). This agrees with a positive trend of the waist circumference gradient observed over the last few years⁽²⁴⁾. The prevalence of abdominal obesity was higher in the South than in other macro-areas and in men than women in the general population; however, according to previous studies the prevalence of abdominal obesity was higher in women than men in the Centre macro-area^(9,17).

A striking finding of the present study is the high prevalence of hypertension (59.6%) in our population, which is much higher than in previous reports of both

ISTAT (12.9%; 13.6%)^(6,7) and OEC (48.5%)⁽⁸⁾. However, ISTAT surveys analysed a wide age range (0 to >80 years) and used self-reported data which may underestimate the prevalence of hypertension, failing to detect hypertensive people who are not aware of their condition. Moreover, according to more recent guidelines⁽¹²⁾, we used cut-off points for hypertension lower than those used in previous studies⁽⁶⁻⁸⁾. Interestingly, when analysing our data using the cut-off of $\geq 160/95$ mmHg similar to OEC⁽⁸⁾, we found comparable prevalence figures for hypertension prevalences in both genders (men: 34% *v.* 33%; women: 28% *v.* 28%). Based on current guidelines⁽¹²⁾, the prevalence of hypertension has raised in our population.

We found differences in the prevalence of dyslipidaemia between genders, with a higher prevalence of men on pharmacological treatment than women (Table 2). In particular, high TAG levels are commonly observed in individuals with abdominal obesity⁽²⁵⁾. In parallel to the rise in prevalence of overweight/obesity and abdominal obesity, the prevalence of hypertriglycerolaemia was almost doubled in our study compared with a previous study⁽⁸⁾. This could be the result of unhealthy eating habits in the Italian population and increased intakes of fast foods and other foods with high fat content, over the last years. On the contrary, in our study the prevalence of hypercholesterolaemia was lower than in a previous survey⁽⁸⁾, but almost comparable to that in a more recent study⁽⁹⁾.

Also the prevalence of diabetes and hyperglycaemia was almost double compared with previous surveys^(7,8), but the prevalence of diabetes was similar to that in a more recent study⁽⁹⁾. The growing prevalence of diabetes in our study population and especially in the Southern population is likely due to the increased prevalence of obesity^(23,26). In addition, consistent with previous studies^(8,17,27), men showed a higher prevalence of diabetes and hyperglycaemia than women, while women showed a higher prevalence of dyslipidaemia than men⁽²⁸⁾.

Over the last 50 years, many studies have evaluated the associations between food groups and chronic diseases or weight gain, with an increasing recognition of the role of nutritional factors in the aetiology of these diseases^(29,30). A low incidence of CVD in the Mediterranean countries, mainly due to the typical nutritional model, has been demonstrated⁽³¹⁾. Unfortunately, in Italy there has been a gradual abandonment of this nutritional model during the past few decades^(32,33). Actually, nutritional recommendations for fruit and vegetable consumption have been raised from 5 servings/d to 7 (or 9) servings/d⁽¹⁶⁾ in order to prevent the major chronic diseases. In our study population, many participants (60%) were far from reaching the optimal intake for these foods⁽¹⁶⁾. Even by considering the recommended range as between 3 and 5 servings/d⁽³⁴⁻³⁷⁾, only 69% of the study population reached the optimal intake for fruit and vegetables⁽²⁸⁾. Unexpectedly, in South and Island macro-areas, the cradle of the Mediterranean diet concept, we found the lowest

prevalence of people reporting fruit and vegetable consumption in the recommended range. Although we did not have data concerning the participants' socio-economic status, we believe that these findings could be related to low household income in the South and Islands macro-areas⁽³⁸⁾.

No recent surveys have examined wine consumption according to the recommended amount⁽¹⁵⁾. In the last few years there has been a steady decline in wine consumption, whereas the consumption of other alcoholic beverage types (e.g. beer) has increased⁽³⁹⁾. Interestingly, a very low prevalence of habitual wine consumption (6%) was found in the general population. However, although moderate wine consumption is considered part of healthy lifestyle, the prevalence of current wine abstainers was very high (76%) in our population. Moreover, consistent with ISTAT data⁽⁶⁾, we found a south-to-north increasing gradient in the prevalence of people showing high wine consumption.

Trends of smoking prevalence are declining as supported by our data (19.8%) and findings from previous studies such as OEC (25.5%) and ISTAT (23.9%; 21.7%)⁽⁶⁻⁸⁾. In particular, the proportion of people who have quit smoking has increased over the last few years⁽⁷⁾, after the banning of smoking in public places with the introduction of the Italian anti-smoking legislation in January 2005⁽⁴⁰⁾. However, consistent with previous studies⁽⁶⁻⁸⁾, men smoked more than women in the present survey; the South and Islands macro-areas showed the highest prevalence of smokers⁽⁸⁾.

As hypertension, obesity/overweight/high waist circumference, dyslipidaemia and diabetes/hyperglycaemia occurred frequently in the study population, it is not surprising that we found 84% of people having between two and five CRF and 90% of people showing more than one CRF (Fig. 2). These findings are deeply alarming. Even if the co-prevalences of CRF were quite similar among the Italian macro-areas, the South and Islands seemed to be the highest-risk macro-areas because of showing the highest prevalence of participants having between two and five CRF (Table 4).

Study strengths and limitations

The major strength of our study is the wide community sample across several Italian macro-areas. Furthermore, anthropometric measurements were made by direct examination according to standard protocols. In addition, we analysed several classical and new CRF.

However, there are some limitations to our study, which should be considered for the interpretation of the present results.

First, surveys based on volunteers may be affected by selection bias. Volunteers with unhealthy lifestyle habits may be more motivated to perform the screening programme, resulting in the screening detecting a high prevalence of CRF in this population. However, selection bias may also operate in the opposite direction, whereby

the more educated and health-conscious (and likely healthier) individuals of the population may volunteer for screening, resulting in the screening detecting a low prevalence of CRF in this population⁽⁴¹⁾. Moreover, because participants were recruited in 193 Italian towns in response to an extensive media campaign, the rural population may be not adequately represented.

Second, a further limitation of the study is the absence of data on previous personal history of CVD and of data concerning socio-economic status, physical activity levels, insulin resistance, HDL cholesterol, LDL cholesterol and minor CRF.

Third, fruit and vegetable consumption, wine intake and cigarette smoking are based on self-reported data and thus the observed prevalence may have been affected by recall bias. In particular, it is known that the intake of health promoting foods such as fruits and vegetables is over-reported by individuals and their consumption is known to remain seasonal in many countries in spite of the spread of the global economy. Self-reported wine consumption is likely to underestimate the real level of consumption. Estimates for cigarette smoking were not validated by biochemical tests.

Conclusions

The present study provided novel evidence on current prevalences and geographic variations for major CRF in Italy. Data are worrisome and suggest a high prevalence of obesity/overweight and hypertension in the Italian population.

Although the decreased cardiovascular mortality in Italy⁽⁴²⁾ has been attributed to improved medical therapies and changes in major CRF, CVD prevalence remains high. Therefore, there is a pressing need for targeted prevention programmes aimed at modifying the lifestyle and eating habits of the Italian population to reduce the risk of CVD. Our results will provide the evidence base to inform effective prevention programmes, which must also take into account the differences in the prevalence of CRF among Italian macro-areas.

Acknowledgements

The work was supported by 'la Federazione nazionale dei titolari di farmacia italiani' (FEDERFARMA), Takeda Italia Farmaceutici, Multicare, FederFARMACO, Pacini editore and the Italian Ministry of Health. The authors declare that there are no conflicts of interest. Authors' contributions were as follows: study design, B.G., R.L., B.M.; statistical analysis, A.P.; collecting data, C.F.; data processing, R.L., B.G., S.S.; writing, R.L., B.G., L.B., S.S. The authors thank Dr Giuliano da Villa and Brunella Maruotto for their assistance in the collection of the data presented in this paper.

References

1. Rayner M & Petersen S (2008) *European Cardiovascular Disease Statistics*. London: British Heart Foundation; available at <http://www.heartstats.org/homepage.asp>
2. Yusuf S, Hawken S, Ounpuu S *et al.*; INTERHEART Study Investigators (2004) Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* **364**, 937–952.
3. Finucane MM, Stevens GA & Cowan MJ (2011) National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants. *Lancet* **377**, 557–567.
4. Mokdad AH, Ford ES, Bowman BA *et al.* (2003) Prevalence of obesity, diabetes, and obesity-related health risk factors, 2001. *JAMA* **289**, 76–79.
5. Wolf-Maier K, Cooper RS, Banegas JR *et al.* (2003) Hypertension prevalence and blood pressure levels in 6 European countries, Canada, and the United States. *JAMA* **289**, 2363–2369.
6. The National Institute for Statistics (2005) Stili di vita e condizioni di salute, Indagine multiscopo sulle famiglie "Aspetti della vita quotidiana", 2003. http://www.istat.it/dati/catalogo/20051118_00/ (accessed January 2007).
7. The National Institute for Statistics (2007) Condizioni di salute, fattori di rischio e ricorso ai servizi sanitari. http://www.istat.it/salastampa/comunicati/non_calendario/20070302_00/ (accessed June 2007).
8. Atlante Italiano delle Malattie Cardiovascolari. II edizione. Osservatorio Epidemiologico Cardiovascolare Italiano (2004) *Ital Heart J* **5**, Suppl. 3, S49–S92.
9. Donfrancesco C, Lo Noce C, Brignoli O *et al.* (2008) Italian network for obesity and cardiovascular disease surveillance: a pilot project. *BMC Fam Pract* **9**, 53.
10. World Health Organization (1998) *Obesity: Preventing and Managing the Global Epidemic. Report of a WHO Consultation on Obesity. WHO Technical Report Series no. 894*. Geneva: WHO.
11. Grundy SM, Cleeman JJ, Daniels SR *et al.*; American Heart Association; National Heart, Lung, and Blood Institute (2005) Diagnosis and management of the metabolic syndrome. An American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* **112**, 2735–2752.
12. US Department of Health and Human Services, National Institutes of Health, National Heart, Lung, and Blood Institute (2004) *The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. NIH Publication no. 04-5230*. Washington, DC: DHSS; available at <http://www.nhlbi.nih.gov/guidelines/hypertension/jnc7full.pdf>
13. National Heart, Lung, and Blood Institute (2007) High blood cholesterol: what you need to know. <http://www.nhlbi.nih.gov/health/public/heart/chol/wyntk.htm> (accessed November 2007).
14. American Diabetes Association (2007) Standards of medical care in diabetes. *Diabetes Care* **30**, Suppl. 1, S4–S41.
15. American Heart Association Nutrition Committee, Lichtenstein AH, Appel LJ, Brands M *et al.* (2006) Diet and lifestyle recommendations revision 2006: a scientific statement from the American Heart Association Nutrition Committee. *Circulation* **114**, 82–96.
16. US Department of Health and Human Services & US Department of Agriculture (2005) Dietary Guidelines for Americans. <http://www.healthierus.gov/dietaryguidelines> (accessed March 2007).
17. Miccoli R, Bianchi C, Odoguardi L *et al.* (2005) Prevalence of the metabolic syndrome among Italian adults according to ATP III definition. *Nutr Metab Cardiovasc Dis* **15**, 250–254.

18. Hu G, Qiao Q, Tuomilehto J *et al.* (2004) Prevalence of the metabolic syndrome and its relation to all-cause and cardiovascular mortality in nondiabetic European men and women. *Arch Intern Med* **164**, 1066–1076.
19. Visscher TL, Viet AL, Kroesbergen IH *et al.* (2006) Under-reporting of BMI in adults and its effect on obesity prevalence estimations in the period 1998 to 2001. *Obesity (Silver Spring)* **14**, 2054–2063.
20. Berghöfer A, Pischon T, Reinhold T *et al.* (2008) Obesity prevalence from a European perspective: a systematic review. *BMC Public Health* **8**, 20.
21. Hedley AA, Ogden CL, Johnson CL *et al.* (2004) Prevalence of overweight and obesity among US children, adolescents, and adults, 1999–2002. *JAMA* **291**, 2847–2850.
22. Barbagallo CM, Cavera G, Sapienza M *et al.* (2002) Nutritional characteristics of a rural Southern Italy population: The Ventimiglia di Sicilia Project. *J Am Coll Nutr* **21**, 523–529.
23. Giampaoli S & Vanuzzo D (2004) II Progetto CUORE – Studi Longitudinali. In Atlante italiano delle malattie cardiovascolari. II Edizione 2004. *Ital Heart J* **5**, Suppl. 3, S94–S101.
24. Ike SO, Chandra DKM, Boev A *et al.* (2004) Abdominal adiposity in US adults: prevalence and trends, 1960–2000. *Prev Med* **39**, 197–206.
25. Bard JM, Charles MA, Juhan-Vague I *et al.*; BIGPRO Study Group (2001) Accumulation of triglyceride-rich lipoprotein in subjects with abdominal obesity: the biguanides and the prevention of the risk of obesity (BIGPRO) 1 study. *Arterioscler Thromb Vasc Biol* **21**, 407–414.
26. Wild S, Roglic G, Green A *et al.* (2004) Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care* **27**, 1047–1053.
27. Verhave JC, Hillege HL, Burgerhof JG *et al.*; PREVEND Study Group (2003) Cardiovascular risk factors are differently associated with urinary albumin excretion in men and women. *J Am Soc Nephrol* **14**, 1330–1335.
28. Sofi F, Innocenti G, Dini C *et al.* (2006) Low adherence of a clinically healthy Italian population to nutritional recommendations for primary prevention of chronic diseases. *Nutr Metab Cardiovasc Dis* **16**, 436–444.
29. Joshipura KJ, Hu FB, Manson JE *et al.* (2001) The effect of fruit and vegetable intake on risk for coronary heart disease. *Ann Intern Med* **134**, 1106–1114.
30. Buijsse B, Feskens EJ, Schulze MB *et al.* (2009) Fruit and vegetable intakes and subsequent changes in body weight in European populations: results from the project on Diet, Obesity, and Genes (DiOGenes). *Am J Clin Nutr* **90**, 202–209.
31. Keys A, Menotti A, Karvonen MJ *et al.* (1986) The diet and 15-year death rate in the Seven Countries Study. *Am J Epidemiol* **124**, 903–915.
32. Trichopoulos D & Lagiou P (2004) Mediterranean diet and overall mortality differences in the European Union. *Public Health Nutr* **7**, 949–951.
33. Centritto F, Iacoviello L, di Giuseppe R *et al.* on behalf of Molisani Investigators (2009) Dietary patterns, cardiovascular risk factors and C-reactive protein in a healthy Italian population. *Nutr Metab Cardiovasc Dis* **19**, 697–706.
34. World Health Organization (2003) *Diet, Nutrition and the Prevention of Chronic Diseases. Joint WHO/FAO Expert Consultation. WHO Technical Report Series* no. 916. Geneva: WHO.
35. Istituto Nazionale di Ricerca per gli Alimenti e la Nutrizione (2003) Linee Guida per una Sana Alimentazione Italiana. Revisione. <http://www.inran.it> (accessed October 2007).
36. Società Italiana di Nutrizione Umana (1996) Livelli di Assunzione Giornalieri Raccomandati di Nutrienti per la Popolazione Italiana (LARN). http://www.sinu.it/larn/tab_rias.asp (accessed November 2007).
37. World Health Organization (2004) *Fruit and Vegetables for Health. Report of a Joint FAO/WHO Workshop, 1–3 September 2004, Kobe, Japan*. Geneva: WHO.
38. Giskes K, Turrell G, Patterson C *et al.* (2002) Socio-economic differences in fruit and vegetable consumption among Australian adolescents and adults. *Public Health Nutr* **5**, 663–669.
39. World Health Organization (2004) Global status report: Alcohol policy. http://www.who.int/substance_abuse/publications/alcohol/en/ (accessed March 2007).
40. Cesaroni G, Forastiere F, Agabiti N *et al.* (2008) Effect of the Italian smoking ban on population rates of acute coronary events. *Circulation* **117**, 1183–1188.
41. Froom P, Melamed S, Kristal-Boneh E *et al.* (1999) Healthy volunteer effect in industrial workers. *J Clin Epidemiol* **52**, 731–735.
42. Palmieri L, Bennett K, Giampaoli S *et al.* (2010) Explaining the decrease in coronary heart disease mortality in Italy between 1980 and 2000. *Am J Public Health* **100**, 684–692.

Appendix 1

Prevalence of cardiovascular risk factors in the study participants aged 18–39 years stratified by gender and macro-area; Italy, 2007

	Study population			North-west			North-east			Centre			South			Islands			P
	Men	Women	P	Men	Women	P	Men	Women	P	Men	Women	P	Men	Women	P	Men	Women	P	
Obesity (%)	14.8	13.1	0.090	10.5	12.6	0.461	11.8	10.9	0.710	13.9	12.8	0.610	17.1	14.9	0.220	16.2	11.4	0.007	0.004
Overweight (%)	57.0	39.9	0.070	55.4	35.4	0.001	49.5	37.1	0.002	54.4	39.0	0.001	61.5	43.5	0.001	57.5	39.5	0.001	0.001
High waist circumference (%)	30.6	44.7	0.001	31.4	43.2	0.006	22.4	40.0	0.001	28.7	47.2	0.001	32.0	45.0	0.001	36.5	45.4	0.023	0.044
Hypertension (%)	29.2	25.3	0.003	30.3	25.9	0.263	26.1	26.1	0.999	31.3	23.5	0.003	30.4	25.7	0.003	25.0	26.3	0.690	0.782
Hypercholesterolaemia (%)	17.9	28.0	0.001	21.9	29.2	0.064	20.0	31.0	0.004	17.7	27.7	0.001	16.7	26.7	0.001	16.4	27.7	0.001	0.071
Hypertriglyceridaemia (%)	32.7	33.0	0.787	32.0	36.7	0.256	36.0	34.4	0.687	28.5	36.7	0.012	34.4	30.1	0.056	31.0	27.5	0.342	0.020
Lipid-lowering therapy (%)	1.3	1.4	0.826	0.6	1.9	0.222	0	0.6	0.266	0.3	1.3	0.074	2.6	1.4	0.080	1.6	1.9	0.789	0.032
Dyslipidaemia (%)	40.8	49.1	0.001	44.6	53.2	0.047	43.5	50.4	0.088	36.9	51.1	0.001	41.5	46.5	0.042	40.1	46.1	0.126	0.026
Diabetes (%)	8.2	8.2	0.940	6.3	8.5	0.344	9.8	9.8	0.998	7.9	8.0	0.979	8.4	7.6	0.550	7.9	8.5	0.207	0.614
Hyperglycaemia (%)	8.2	8.5	0.748	8.6	8.3	0.922	10.7	7.7	0.179	10.3	11.7	0.452	6.6	7.5	0.468	6.7	5.4	0.477	0.001
Smoking (%)	34.7	22.6	0.001	26.9	17.8	0.010	37.4	22.3	0.001	32.6	25.6	0.001	37.6	20.1	0.001	34.1	30.1	0.278	0.001
Fruit and vegetables (%)*	20.7	32.9	0.001	33.7	44.1	0.015	27.1	37.7	0.006	24.1	40.4	0.001	13.7	23.1	0.001	18.2	26.1	0.019	0.001
Wine (%) [†]																			
Abstainer	91.8	87.3		85.7	86.4		93.0	84.4		90.8	84.9		92.9	90.2		94.1	88.4		
Moderate	7.5	8.4	0.001	13.1	7.2	0.001	6.5	10.2	0.002	8.4	9.5	0.001	6.3	7.5	0.037	5.9	8.5	0.008	0.001
Habitual	0.7	4.3		1.1	6.4		0.5	5.4		0.8	5.6		0.8	2.4		0.0	3.1		

*Fruit and vegetable consumption within the recommended range of 8–10 servings/d.

[†]Wine consumption higher than the recommended intake: moderate, 1–2 drinks/d for men and 0.5–1 drink/d for women; habitual, >2 drinks/d for men, and >1 drink/d for women.

Appendix 2

Prevalence of cardiovascular risk factors in the study participants aged 40–65 years stratified by gender and macro-area; Italy, 2007

	Study population			North-west			North-east			Centre			South			Islands			P
	Men	Women	P	Men	Women	P	Men	Women	P	Men	Women	P	Men	Women	P	Men	Women	P	
Obesity (%)	24.4	21.8	0.002	20.0	17.9	0.386	20.1	15.9	0.042	23.2	20.0	0.052	29.0	29.0	0.972	22.9	23.9	0.725	0.001
Overweight (%)	51.7	37.3	0.001	49.2	35.5	0.001	50.2	35.9	0.001	54.5	36.7	0.001	50.2	40.7	0.001	54.7	37.7	0.001	0.001
High waist circumference (%)	49.6	65.1	0.001	45.7	58.7	0.001	46.2	57.9	0.001	47.5	65.3	0.001	51.6	73.5	0.001	54.8	65.0	0.001	0.001
Hypertension (%)	59.8	57.8	0.047	60.1	59.1	0.679	63.6	59.5	0.111	61.3	55.2	0.002	60.1	61.2	0.539	52.1	48.5	0.251	0.001
Hypercholesterolaemia (%)	26.6	41.5	0.001	28.3	42.7	0.001	27.3	41.5	0.001	27.1	41.6	0.001	26.3	40.0	0.001	23.8	43.5	0.001	0.004
Hypertriglyceridaemia (%)	44.9	51.4	0.001	46.6	53.4	0.005	49.2	52.5	0.203	46.5	52.5	0.002	43.0	49.8	0.001	41.7	46.5	0.127	0.001
Lipid-lowering therapy (%)	7.9	10.0	0.001	8.5	10.3	0.221	5.6	6.6	0.415	7.5	9.5	0.071	8.1	11.5	0.001	10.0	12.9	0.144	0.001
Dyslipidaemia (%)	56.9	69.4	0.001	60.0	73.8	0.001	61.0	69.5	0.001	57.6	70.2	0.001	54.9	66.3	0.001	54.1	67.5	0.001	0.001
Diabetes (%)	13.5	11.4	0.001	15.2	12.7	0.138	9.5	9.5	0.959	11.3	8.5	0.018	15.2	14.5	0.579	14.6	10.6	0.080	0.001
Hyperglycaemia (%)	13.8	12.5	0.057	15.0	11.9	0.061	12.4	10.5	0.269	20.3	18.1	0.177	10.7	9.5	0.284	11.0	9.5	0.454	0.001
Smoking (%)	26.1	15.8	0.001	19.4	12.6	0.001	19.8	16.7	0.134	25.0	20.6	0.009	29.3	12.5	0.001	32.2	17.1	0.001	0.001
Fruit and vegetables (%)*	33.4	47.2	0.001	47.2	58.1	0.001	45.4	56.7	0.001	39.8	51.3	0.001	22.2	31.5	0.001	27.3	42.0	0.001	0.001
Wine (%) [†]																			
Abstainer	76.9	73.4		68.5	67.8		74.0	68.5		75.4	70.7		80.1	80.1		82.8	82.6		
Moderate	21.6	15.9	0.001	29.9	15.6	0.001	24.0	16.4	0.001	23.2	17.0	0.001	18.4	15.6	0.001	16.8	13.2	0.001	0.001
Habitual	1.5	10.7		1.6	16.6		2.0	15.1		1.4	12.3		1.4	4.3		0.4	4.2		

*Fruit and vegetable consumption within the recommended range of 8–10 servings/d.

[†]Wine consumption higher than the recommended intake: moderate, 1–2 drinks/d for men and 0.5–1 drink/d for women; habitual, >2 drinks/d for men, and >1 drink/d for women.

Appendix 3

Prevalence of cardiovascular risk factors in the study participants aged 65 years and over stratified by gender and macro-area; Italy, 2007

	Study population			North-west			North-east			Centre			South			Islands			<i>P</i>
	Men	Women	<i>P</i>	Men	Women	<i>P</i>	Men	Women	<i>P</i>	Men	Women	<i>P</i>	Men	Women	<i>P</i>	Men	Women	<i>P</i>	
Obesity (%)	24.7	24.4	0.750	21.1	18.3	0.175	16.3	14.8	0.470	21.9	23.7	0.315	30.6	37.4	0.001	25.8	23.6	0.504	0.001
Overweight (%)	41.8	39.7	0.001	51.6	34.8	0.001	56.3	41.3	0.001	53.5	42.2	0.001	49.2	40.1	0.171	51.1	37.9	0.001	0.001
High waist circumference (%)	59.2	77.0	0.001	55.3	68.1	0.001	56.4	69.1	0.001	54.2	79.0	0.001	64.0	88.5	0.001	62.1	73.8	0.001	0.001
Hypertension (%)	76.7	76.9	0.803	81.9	77.4	0.036	80.4	75.6	0.036	78.9	75.4	0.051	74.7	81.2	0.001	68.8	69.2	0.910	0.001
Hypercholesterolaemia (%)	22.7	37.5	0.001	23.9	40.9	0.001	27.6	40.7	0.001	21.6	39.2	0.001	21.5	29.9	0.001	22.0	40.5	0.001	0.001
Hypertriacylglycerolaemia (%)	43.5	57.9	0.001	42.5	63.7	0.001	46.9	60.3	0.001	46.4	61.1	0.001	42.6	49.7	0.001	37.7	51.9	0.001	0.001
Lipid-lowering therapy (%)	11.4	18.8	0.001	10.4	19.7	0.001	8.0	17.1	0.001	13.1	17.1	0.009	11.6	20.4	0.001	12.1	21.1	0.001	0.001
Dyslipidaemia (%)	54.3	71.3	0.001	55.8	76.6	0.001	60.0	74.5	0.001	56.4	75.2	0.001	52.0	61.3	0.001	49.5	68.5	0.001	0.001
Diabetes (%)	24.0	17.6	0.001	20.2	16.3	0.059	19.6	14.5	0.015	21.2	15.7	0.001	27.7	22.0	0.001	27.1	21.1	0.063	0.001
Hyperglycaemia (%)	15.6	14.6	0.203	16.6	14.3	0.239	14.5	9.7	0.007	21.4	20.0	0.433	12.5	13.6	0.431	13.5	11.9	0.511	0.001
Smoking (%)	14.2	7.2	0.001	10.9	9.4	0.354	10.7	7.7	0.061	12.3	8.7	0.006	16.6	3.8	0.001	18.4	6.1	0.001	0.001
Fruit and vegetables (%)*	40.6	47.8	0.001	57.2	60.1	0.263	57.6	59.2	0.541	48.9	53.4	0.029	26.5	24.1	0.178	30.2	46.1	0.001	0.001
Wine (%)†																			
Abstainer	71.4	70.0		62.0	62.8		63.4	65.6		68.6	69.6		76.0	77.1		81.1	79.2		
Moderate	26.4	17.5	0.001	36.2	19.4	0.001	34.0	16.7	0.001	29.4	17.9	0.001	20.9	17.5	0.001	18.1	13.1	0.001	0.001
Habitual	2.2	12.5		1.8	17.8		2.6	17.7		2.0	12.5		3.1	5.4		0.8	7.7		

*Fruit and vegetable consumption within the recommended range of 8–10 servings/d.

†Wine consumption higher than the recommended intake: moderate, 1–2 drinks/d for men and 0.5–1 drink/d for women; habitual, >2 drinks/d for men, and >1 drink/d for women.