

Platelet ice and the land-fast sea ice of McMurdo Sound, Antarctica

INGA J. SMITH,¹ PATRICIA J. LANGHORNE,¹ TIMOTHY G. HASKELL,² H. JOE TRODAHL,³
RUSSELL FREW,⁴ M. ROSS VENNEL⁵

¹*Department of Physics, University of Otago, Dunedin, New Zealand*

²*Industrial Research Limited, P.O. Box 31-310, Lower Hutt, New Zealand*

³*School of Physics and Chemistry, Victoria University, Wellington, New Zealand*

⁴*Department of Chemistry, University of Otago, Dunedin, New Zealand*

⁵*Department of Marine Science, University of Otago, Dunedin, New Zealand*

ABSTRACT. Dendritic crystals of platelet ice appear beneath the columnar land-fast sea ice of McMurdo Sound, Antarctica. These leaf-like crystals are frozen into place by the advancing columnar growth. The platelets most probably begin to appear during July, although in some parts of the Sound they may not appear at all. In addition, the amount and extent of platelet ice within the Sound varies from year to year. Previous authors have suggested that the formation of platelet ice is linked to the presence of the nearby ice shelf. It is a matter of debate whether these platelets form at depth and then float upwards or whether they grow in slightly supercooled water at the ice/water interface. The phenomenon is similar to that observed in the Weddell Sea region, but previous authors have suggested the two regions may experience different processes. This paper presents the results of field-work conducted in McMurdo Sound in 1999. Ice-structure analysis, isotopic analysis and salinity and temperature measurements near the ice/water interface are presented. Freezing points are calculated, and the possible existence of supercooling is discussed in relation to existing conjectures about the origin of platelets.

INTRODUCTION

The term *platelet ice* is used by different authors to mean either the loose platelets underlying a sea-ice sheet, but not yet frozen into place (e.g. Dayton, 1989), or the incorporated layer consisting of platelets and columnar ice formed between the platelets (e.g. Jeffries and others, 1993; Gow and others, 1998). In this paper we will discuss both loose and incorporated platelet ice, and have attempted to make this distinction clear in the terminology used if it is not clear from the context.

There is strong evidence that platelet-ice formation is linked to the presence of ice shelves. Moretskiy (1965) found that loose billows of platelet ice beneath the fast ice off Mirny station, Antarctica, in late November 1958 were thickest adjacent to the ice shelf. Gow and others (1998) saw “no systematic variation of [incorporated] platelet thicknesses with location in McMurdo Sound”. Crocker and Wadhams (1989) noted the thickest billows of loose platelets beneath the fast ice of McMurdo Sound in 1986 occurred in regions where Lewis and Perkin (1985) had observed maximal supercooling in 1982. The generally accepted explanation for the supercooling observed is that sea water near its surface freezing point has been transported by currents beneath the ice shelf and then has been cooled by contact with the ice shelf (Lewis and Perkin, 1985). When this water rises to the surface along the underside of the ice shelf, it becomes supercooled due to the change in freezing point with depth (Foldvik and Kvinge, 1974; Robin, 1979; Crocker and Wadhams, 1989). Of some debate is whether the platelets then form at considerable depths or at

the underside of the sea-ice cover. There is evidence for both hypotheses: Dieckmann and others (1986) observed a large cloud of platelets at 250 m depth in the Weddell Sea, while Gow and others (1998) state that the orientation of the platelet crystals beneath McMurdo Sound fast ice indicates that the platelets grow attached to the ice above, rather than forming at depth. It has been acknowledged (Eicken and Lange, 1989) that different processes may cause platelet-ice formation in different regions of Antarctica. Smetacek and others (1992) suggested that platelet ice observed under pack ice could be caused by different processes than those that cause platelet ice observed under fast ice. This possibility was also alluded to by Moretskiy (1965).

In this paper the results of field measurements made in McMurdo Sound in 1999 are presented and discussed in relation to consistency with the conjecture that platelet ice in McMurdo Sound forms *in situ*.

LOCATIONS

Three sites in McMurdo Sound were studied (see Fig. 1).

- (1) Site AS (array site) was on the sea ice off Arrival Heights. This site was the location of a thermistor (temperature sensor) array, part of an investigation of the thermal conductivity of sea ice by one of us (H.J.T.). The thermistor-array global positioning system (GPS) coordinates were 77°50.197' S, 166°36.764' E. Temperatures were automatically logged simultaneously at each depth every 30 min. Thermistors were spaced 100 mm apart along the 2 m length of the probe. Water salinity and temperature meas-




Fig. 1. Map of locations in McMurdo Sound.

Measurements were made and ice cores taken 150 m north of the array. The site was examined twice: once on 23 August 1999, and again on 9 October 1999.

- (2) Site CH (Camp Haskell) was located between Big Razorback Island and Inaccessible Island on the sea ice between the Dellbridge Islands. The GPS coordinates of the CH site were 77°40.627' S, 166°27.038' E.
- (3) Site CE (Cape Evans) was located on the sea ice approximately 150 m from Scott's Hut at Cape Evans. (Coordinates of Scott's Hut at Cape Evans are roughly 77°38'10" S, 166°25'10" E.)

Table 1. Summary of ice structure at the sites

	Loose snow	Columnar ice	Incorporated platelet ice	Total ice thickness
	mm	mm	mm	mm
AS 24 August 1999	60	1450	300	1750
AS 10 October 1999	65	1460	690	2150
CH 21 October 1999	Minimal	1200–1400	540–740	1940
CE 17 October 1999	20	1850	None present	1850

Measurements at CH and CE were conducted on 17 October 1999.

ICE-STRUCTURE ANALYSIS

Ice cores were taken at CE, CH and AS sites for structural analysis. In addition, a slab approximately 1 m by 1 m by 1.93 m was extracted at CH, and vertical thick sections were made from this. When this slab was removed, large platelets approximately 2 mm thick and up to 200 mm in diameter were found attached to the underside of the ice. These were discoloured by brown algae. The base of such a slab is shown in Figure 2. The results of thin- and thick-section analysis of ice cores and slabs from the locations are shown schematically in Figure 3 and summarized in Table 1.

From Figure 3 and Table 1, it should be noted that although the CE and CH sites were only approximately 3 km apart, there was no platelet ice observed at CE. A possible explanation for this difference is the existence of a submarine ridge between Cape Evans and Inaccessible Island (Hydrographer of the Navy, 1974; U.S. National Imagery and Mapping Agency, 1997). This ridge could block



Fig. 2. Base of sea-ice slab taken at CH site, showing attached platelet ice.




Fig. 3. Ice-structure diagram for the sites.

the flow of ice-shelf water (ISW) to the area off Cape Evans, resulting in an absence of platelet ice in the area. Alternatively, the ridge could block the advection of loose platelet crystals from further south in the Sound.

The *c*-axis plot for the horizontal thin section prepared from 2.12 m depth in the AS core from 10 October 1999 (Fig. 4) shows crystals at large angles to each other, which is characteristic of incorporated platelet ice. This is in contrast to the aligned or girdle distributions that would have been obtained for the columnar ice in the study. We have not distinguished between the platelet crystals and the columnar ice between them. Our core was not oriented with respect to a fixed azimuth.

The two AS cores were taken just over a month apart. The onset of incorporated platelet ice beneath the columnar ice occurs at the same depth in each core, as would be expected. The increase in the thickness of the incorporated

platelet-layer thickness suggests active platelet growth was still occurring between the two core extractions. The layer of loose platelet ice observed on 24 August 1999 was approximately 50 mm thick. This is insufficient to account for the incorporated platelet-ice thickness observed on 10 October 1999 unless platelets were continually advected upwards or were continuously nucleating attached to the ice undersurface. It is possible these platelets were formed at depth and were advected upwards. Gow and others (1987) observed *banded frazil* in Weddell Sea pack-ice floes, where plates of ice seemed to have floated up from depth and the resulting *c*-axis orientations were vertical. In contrast, the observations presented here are more consistent with those of Gow and others (1998). In that paper, attached platelets below the incorporated layer of McMurdo Sound fast ice had a near-horizontal *c*-axis orientation, and the authors hence concluded that those platelets nucleated directly onto the bottom of the sea-ice cover. We did not observe any purely columnar bands within the incorporated platelet-ice layer, as Jeffries and others (1993) did in their core RS-29. In the absence of such bands, advection theory would require continuous advection of platelets from depth to the site. Otherwise it might be expected that one could observe bands of just columnar ice where all the advected platelets present had been frozen into place and then columnar growth continued.

ISOTOPE ANALYSIS

Oxygen isotope analysis was carried out on sea-ice samples from each site. At each site a core was divided into 100 mm sections for analysis. The samples from AS were taken on 24 August 1999. These were melted and bottled at Scott Base, then transported to New Zealand. Analysis was carried out in February and March 2000. The analysis was referenced to Vienna Standard Mean Ocean Water (VSMOW), with a precision of ±0.04‰. Duplicate subsamples were taken for the analysis. The samples from CH were taken on 19 October 1999 and were melted and bottled at the camp in 100 mL glass bottles with aluminium caps fitted with rubber seals. A duplicate core was sectioned and transported to New Zealand frozen for later alternative preparation and




Fig. 4. *c*-axis plot for horizontal thin section from 2.12 m depth in the AS core from 10 October 1999.




Fig. 5. Oxygen isotope measurements through the ice cores from CE and CH.

analysis. The samples from CE were also transported frozen back to New Zealand. These samples had the centre of each section removed at room temperature on 10 February 2000, and these were then melted and bottled.

The preliminary isotope results for CH and CE are shown in Figure 5. Although an obvious change from columnar to platelet ice is observed in vertical thin sections of each core, there is no such obvious change in the isotope values at that level. In other words, the isotope values seem to be independent of ice type. This is consistent with the observations of Lange and others (1990) and Tison and others (1998), although in both those cases the transition in ice types was not columnar to platelet. However, the loose platelets do exhibit a lower $\delta^{18}\text{O}$ ($\approx 1.4\text{‰}$) than the mixture of platelet and columnar ice above ($\delta^{18}\text{O} \approx 1.7\text{‰}$). This could be due to the presence of sea water in the loose platelets, or it could indicate that the platelets form from fresher water (e.g. from ISW). It is also possible that the platelets and interstitial columnar ice grow at different rates, and the $\delta^{18}\text{O}$ values are therefore an average of the results of the two different growth rates. Below 0.6 m, the CE $\delta^{18}\text{O}$ values are consistently higher (+2.0 to +2.3‰) than the CH values (+1.5 to +1.8‰) at the same depths. The salinity profiles (Fig. 6) show the CE ice salinities lower than the CH ice salinities by at least 0.5‰ below 0.6 m. Since no surface water samples were taken for




Fig. 6. Salinity results from the ice cores from CE and CH.

isotopic analysis at these two sites, it cannot be determined whether the same water types were present. If the ice formed from the same water type and desalination was minimal, then Tison and others (1998) state that the isotopic profiles will be determined by the changes in freezing rate at each depth. Tison and others (1998) then state that a lower freezing rate will cause lower bulk salinity and increased $\delta^{18}\text{O}$ values in the ice. Hence, if the water at CH and CE were identical, then the $\delta^{18}\text{O}$ and salinity values reported here could be explained by lower freezing rates at CE. As discussed below, however, our salinity and temperature measurements indicate that the water was slightly more saline at CE than at CH, which may indicate an absence of ISW at CE. CTD casts at depth are planned for the CE and CH areas in future work, and water samples will be taken for isotope analysis. This will allow determination of the presence or absence of ISW in these areas.

SALINITY AND TEMPERATURE MEASUREMENTS

Many previous characterizations of the oceanography of the Ross Sea region have been based on ship-based measurements over open water during the austral summer (Jacobs and others, 1985). However, Lewis and Perkin (1985) measured the presence of supercooling beneath the McMurdo Sound fast ice using a Guildline Type 8706 CTD (conductivity- (salinity-), temperature- and depth-measuring device). Due to the presence of the ice cover and the influence of the hole, they relied solely on measurements below 4 m. Their results show supercoolings of up to approximately 0.025°C at the 4 m level.

Our experiments were designed to make similar measurements, but closer to the ice/water interface and away from the influence of the hole.

At the CH, CE and AS sites, a Sea-Bird Electronics (SBE) system was used. The instrument was configured for making measurements beneath the ice and away from the hole. SBE 3plus (temperature) and SBE 4C (conductivity (salinity)) sensors were connected to an SBE 5T pump, and the cables were then connected to a desktop multi-channel counter (SBE 3l), driven by a laptop computer running SeaSoft software. This allowed monitoring of the conditions during the experiment, which we found invaluable for detecting any possible icing of the C cell. No depth sensor was used, since all measurements were to be near the ice/water interface. We therefore refer to our equipment as a CT rather than the usual CTD. The cell and pump were mounted on a hinged, alloy arm (Fig. 7) that could be lowered through a hole vertically and then winched into position so that an L-shaped configuration resulted. All experiments were conducted from inside a heated dive hut with a removable floor. This allowed the CT to be kept warmer than the water into which it was being lowered, thus preventing freezing of the cells. The hut exhibited strong temperature gradients (e.g. in the CH experiments the temperature at head height was $+13^\circ\text{C}$, while the floor was -7°C). Because of this, the arm was stored standing upright, with the cells near the roof, until deployment. The CT was set running in the vertical position to remove any air bubbles in the system, and salinity and temperature were recorded while the cells were put into position. These data were removed from the final analysis. The final positioning of the CT sensors with respect to the underside of the ice was aided by a small black-and-white video camera mounted at the end of the arm (Fig. 8). The depth of the sensors was determined later by measuring along the arm from a mark made to show the ice surface position. The




Fig. 7. Photo of CT mounted on arm.

CT was in the water for at least 10 min, with data logged 24 times a second. The arm was designed so that the cells were 2 m away from the hole, which ensures the measurements are away from the convective effects of the hole when measuring close to the ice/water interface (Scorer, 1957).

Salinities were automatically calculated from the conductivities by the SeaSoft software using the Unesco Practical Salinity Scale 1978 (Unesco, 1981). Following the recommendation of that work, salinities are often given without units (Lewis and Perkin (1985) followed this convention). However, we have given our results with psu (practical salinity units) following salinity values. This is an alternative convention that has developed in the oceanographic literature (e.g. Jenkins and Bombosch, 1995). It clarifies the distinction between sea-water salinities which are given in psu, and ice salinities, which we have given in parts per thousand.

The freezing points were calculated using the Millero (1978) formula, with pressure taken to be atmospheric. The errors quoted are based on manufacturers' calibration before and after the fieldwork.

Near-surface values at the three sites are shown in Table 2.

The results of the measurements from AS on 9 October 1999 are shown in Figure 9. The upper three lines are the freezing-point temperatures (centre) and error limits, calculated from salinity values. The bottom three lines are the temperature measurements (centre) with error limits. The time-scale is data points, with data recorded 24 times a second.

It can be seen from Figure 9 and Table 2 that at the AS




Fig. 8. Diagram of CT arm in deployed position.

Table 2. Mean measured S and T values at AS, CH and CE sites, and supercoolings calculated from these values

Site	Date	Salinity psu	Temperature °C	$T_{f(\text{mean})} - T_{(\text{mean})}$	$\Delta T_{(\text{min})}$ °C	z m
AS	23 Aug. 1999	34.60 ± 0.01	-1.910 ± 0.001	0.011 ± 0.004	0.006	0.41
AS	9 Oct. 1999	34.65 ± 0.01	-1.913 ± 0.001	0.011 ± 0.004	0.006	0.38
CH	17 Oct. 1999	34.67 ± 0.01	-1.911 ± 0.001	0.008 ± 0.004	0.003	0.57
CE	17 Oct. 1999	34.73 ± 0.01	-1.910 ± 0.001	0.004 ± 0.004	Not discernible	0.35

Notes: $T_{f(\text{mean})} - T_{(\text{mean})}$ = mean freezing-point temperature – mean temperature; $\Delta T_{(\text{min})}$ = minimum mean supercooling; z = distance of cells beneath ice/water interface.

site the water was supercooled by at least 0.006°C on 9 October 1999. At the same site on 23 August 1999, we also measured supercooling of at least 0.006°C (Table 2). However (Table 2), the water was also colder on average by 0.003°C for the October measurements and saltier on average by 0.05 psu. This can be explained as the result of ongoing rejection of salt as the ice continued to thicken in the late winter/early spring.

The supercooling measured at CH on 17 October 1999 is of a similar magnitude to that observed at the AS site (Table 2). The average salinity of 34.67 psu is higher by 0.02 psu than that observed at AS. In contrast, there is negligible discernible supercooling from the CE site on 17 October 1999 (Table 2). This may be indicative of the absence of ISW in the CE area, but this needs further investigation.

ICE GROWTH IN SUPERCOOLED WATER

The analysis of measurements from the thermistor array at the AS site yields a result similar to that described in Trodahl and others (2000) for a probe at the same site in 1997, namely, that not all the latent heat associated with incorporated platelet-ice formation was conducted to the atmosphere. The experiments described in that paper could not determine whether this was due to platelet formation at the ice/water




Fig. 9. Graph showing water temperatures and calculated freezing points from CT measurements at AS on 9 October 1999.

interface in a supercooled layer or due to advection of platelet crystals that had nucleated at depth, or whether a combination of these processes was occurring. Based on observations of the base of a slab of sea ice taken at the CH site, 10% of the volume of the incorporated platelet ice was estimated to be pure platelet ice. The other 90% of that ice type was interstitial congelation ice that formed between the individual platelets by heat conduction to the air through the ice. The observed thicknesses of incorporated platelet ice at the AS and CH sites were of the order of 0.7 m by mid-October. The pure platelet ice is composed of dendritic crystals, implying growth under supercooled conditions.

Currents in McMurdo Sound have been measured in the Cape Armitage area to be 0.043 m s^{-1} (Barry and Dayton, 1988). If a horizontal current transported supercooled water from beneath an ice shelf and this then interacted with the skeletal interface beneath a growing sea-ice cover, the skeletal layer could experience enhanced growth conditions. This mechanism could explain the predominantly near-horizontal c -axis orientation noted in the platelet ice of McMurdo Sound. The following calculation shows how this forced convection may explain the amounts of platelet ice observed in this study.

The equation for heat flux in the buffer zone, where molecular and eddy transfer are occurring (Kay and Nedderman, 1974), is

$$q = \rho c_p (\alpha + \epsilon) \frac{dT}{dz},$$

where q is the heat flux, ρ and c_p are the density and specific heat capacity of sea water, respectively, α is the thermal diffusivity of sea water (conduction term), ϵ is the kinematic eddy viscosity (forced convection term) and dT/dz is the temperature gradient in the buffer zone.

The formation of 0.07 m^3 of pure platelet ice (10% of a

0.7 m thick section of incorporated platelet ice in a 1 m^2 area) requires the removal of approximately 21000 kJ of latent heat, Q . Assuming the platelet ice began forming in early July (Crocker and Wadhams, 1989), this gives approximately 100 days for the observed thickness of incorporated platelet ice to form. Thus we can approximate the rate of heat transfer over an area of 1 m^2 as;

$$q = \frac{1}{A} \frac{dQ}{dt} \approx \frac{1}{A} \frac{\Delta Q}{\Delta t} \\ \approx \frac{2.1 \times 10^4 \text{ kJ}}{1 \text{ m}^2 \times 8.64 \times 10^6 \text{ s}} \approx 2 \times 10^{-3} \text{ kJ m}^{-2} \text{ s}^{-1}.$$

Assuming that the water directly in contact with the ice/water interface is not supercooled, and since the CT cells were 0.5 m below the ice/water interface, we can approximate the temperature gradient as

$$\frac{dT}{dz} \approx \frac{\Delta T}{\Delta z} = \frac{0.01 \text{ K}}{0.5 \text{ m}} = 0.02 \text{ K m}^{-1}.$$

Using values for sea water of $\rho = 1028 \text{ kg m}^{-3}$ (density), $c_p = 4.2177 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (specific heat), $\alpha = 1.3 \times 10^{-7} \text{ m}^2 \text{ s}^{-1}$ (thermal diffusivity), and solving for the kinematic eddy viscosity gives

$$\epsilon \approx 2 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}.$$

Since eddy viscosity tends to zero with decreasing current velocity and decreasing depth below the ice, this value is not inconsistent with values of kinematic eddy viscosity reported beneath first-year land-fast sea ice in the Arctic (Crawford and others, 1996), where for current velocities of $5\text{--}10 \text{ cm s}^{-1}$ and at a depth of 2 m below the ice, the eddy viscosity was reported to be $0.005 \text{ m}^2 \text{ s}^{-1}$.

Therefore the amount of incorporated platelet ice observed at the AS and CH sites could be explained by in situ nucleation of pure platelets due to current-enhanced convection of supercooled water adding to the conduction

process, and the interstitial congelation ice being the result of conduction of heat to the air due to the temperature gradient between the air and water.

CONCLUSIONS

This study confirms the presence of supercooling in sea water immediately below actively growing platelet ice, and the absence of pronounced supercooling in an area where no platelet ice is present. Combined with the ice-growth calculations in a horizontal current, this result adds weight to the argument that McMurdo Sound fast-ice platelets grow attached to the fast-ice above, rather than being advected from depth. The near-horizontal *c*-axis orientation of the attached platelets beneath the McMurdo Sound fast ice in August and October 1999 is consistent with the platelets growing at the ice/water interface. If such platelets were advected from depth, it would be expected that the *c* axes would be more vertically oriented, and that supercooling would already have been relieved, so that no supercooling would be observed next to the ice/water interface. However, it is possible that in other areas of McMurdo Sound, or beneath the Ross Sea pack ice, platelets advecting from depth may be observed, and future work will examine whether different platelet formation processes are present in the wider McMurdo Sound and Ross Sea areas. The local bathymetry may play a significant role in the presence or absence of platelet ice, with a submarine ridge off Cape Evans either blocking the flow of ISW or blocking the advection of platelets from depth to that area. In future work, it is intended to conduct deeper CTD casts in the CH (Dellbridge Islands) and CE (Cape Evans) areas. The results of these casts will be combined with water-sample isotopic analysis to compare the water masses present in the two areas, and determine if ISW is present in the areas. Isotope and salinity profiles from CH and CE exhibit characteristics that indicate the ice below 600 mm at CE was subject to a lower freezing rate than the ice at similar depths at CH. The detailed analysis of these profiles is ongoing, and a comparison of isotope and salinity profiles with ice-growth rates from AS (sea ice off Arrival Heights) is intended in the future.

ACKNOWLEDGEMENTS

We wish to thank the Foundation for Research, Science and Technology for their ongoing support of Event K131 research. I.J.S is grateful to Antarctica New Zealand for awarding her the 1999/2000 Antarctica New Zealand Sir Robin Irvine Scholarship and for the research opportunities that resulted from it. Thanks also to Antarctica NZ and the Scott Base staff of the 1998/99 and 1999/2000 seasons for providing essential field and logistical support. We are grateful to the staff of McMurdo station for their assistance, especially W. Haals, R. Score of the Crary Laboratory and Jeff and Dennis of the Survey Team.

REFERENCES

- Barry, J. P. and P. K. Dayton. 1988. Current patterns in McMurdo Sound, Antarctic and their relationship to local biotic communities. *Polar Biol.*, **8**(5), 367–376.
- Crawford, G. B., L. Padman and M. G. McPhee. 1996. Turbulent mixing in Barrow Strait. *Continental Shelf Res.*, **19**(2), 205–245.
- Crocker, G. B. and P. Wadhams. 1989. Modelling Antarctic fast-ice growth. *J. Glaciol.*, **35**(119), 3–8.
- Dayton, P. K. 1989. Interdecadal variation in an Antarctic sponge and its predators from oceanographic climate shifts. *Science*, **245**(4925), 1484–1486.
- Dieckmann, G. S., G. Rohardt, H. Hellmer and J. Kipfstuhl. 1986. The occurrence of ice platelets at 250 m depth near the Filchner Ice Shelf and its significance for sea ice biology. *Deep-Sea Res.*, **33**(2), Part A, 141–148.
- Eicken, H. 1998. Deriving modes and rates of ice growth in the Weddell Sea from microstructural, salinity and stable-isotope data. In Jeffries, M. O., ed. *Antarctic sea ice: physical processes, interactions and variability*. Washington, DC, American Geophysical Union, 89–122. (Antarctic Research Series 74)
- Eicken, H. and M. A. Lange. 1989. Development and properties of sea ice in the coastal regime of the southeastern Weddell Sea. *J. Geophys. Res.*, **94**(C6), 8193–8206.
- Foldvik, A. and T. Kvinge. 1974. Conditional instability of sea water at the freezing point. *Deep-Sea Res.*, **21**(3), 169–174.
- Gow, A. J., S. F. Ackley, K. R. Buck and K. M. Golden. 1987. Physical and structural characteristics of Weddell Sea pack ice. *CRREL Rep.* 87-14.
- Gow, A. J., S. F. Ackley, J. W. Govoni and W. F. Weeks. 1998. Physical and structural properties of land-fast sea ice in McMurdo Sound, Antarctica. In Jeffries, M. O., ed. *Antarctic sea ice: physical processes, interactions and variability*. Washington, DC, American Geophysical Union, 355–374. (Antarctic Research Series 74)
- Hydrographer of the Navy. 1974. *The Antarctic pilot (comprising the coasts of Antarctica and all islands southward of the usual routes of vessels)*. Fourth edition. Taunton, Somerset, Ministry of Defence. Hydrographer of the Navy.
- Jacobs, S. S., R. G. Fairbanks and Y. Horibe. 1985. Origin and evolution of water masses near the Antarctic continental margin: evidence from H₂¹⁸O/H₂¹⁶O ratios in seawater. In Jacobs, S. S., ed. *Oceanology of the Antarctic continental shelf*. Washington, DC, American Geophysical Union, 59–85. (Antarctic Research Series 43)
- Jeffries, M. O., W. F. Weeks, R. Shaw and K. Morris. 1993. Structural characteristics of congelation and platelet ice and their role in the development of Antarctic land-fast sea ice. *J. Glaciol.*, **39**(132), 223–238.
- Jenkins, A. and A. Bombosch. 1995. Modeling the effects of frazil ice crystals on the dynamics and thermodynamics of ice shelf water plumes. *J. Geophys. Res.*, **100**(C4), 6967–6981.
- Kay, J. M. and R. M. Nedderman. 1974. *An introduction to fluid mechanics and heat transfer*. Third edition. Cambridge, Cambridge University Press.
- Lange, M. A., P. Schlosser, S. F. Ackley, P. Wadhams and G. S. Dieckmann. 1990. ¹⁸O concentrations in sea ice of the Weddell Sea, Antarctica. *J. Glaciol.*, **36**(124), 315–323.
- Lewis, E. L. and R. G. Perkin. 1985. The winter oceanography of McMurdo Sound, Antarctica. In Jacobs, S. S., ed. *Oceanology of the Antarctic continental shelf*. Washington, DC, American Geophysical Union, 145–165. (Antarctic Research Series 43)
- Millero, F. J. 1978. Appendix 6. Freezing point of sea water. In *Eighth report of the Joint Panel on Oceanographic Tables and Standards*. Paris, Unesco, 29–31. (Technical Papers in Marine Science 28.)
- Moretskiy, V. N. 1965. Morskoy vnutrivodnyy led [Underwater sea ice]. *Probl. Arkt. Antarkt.* **19**, 32–38. (Translation by E. R. Hope, DRB Canada Report T-497-R, 1968.)
- Robin, G. de Q. 1979. Formation, flow, and disintegration of ice shelves. *J. Glaciol.*, **24**(90), 259–271.
- Scorer, R. S. 1957. Experiments on convection of isolated masses of buoyant fluid. *J. Fluid Mech.*, **2**, 583–594.
- Smetacek, V. and 6 others. 1992. Early spring phytoplankton blooms in ice platelet layers of the southern Weddell Sea, Antarctica. *Deep-Sea Res.*, **39**(2), 153–168.
- Tison, J.-L., R. D. Lorrain, A. Bouzette, M. Dini, A. Bondesan and M. Stiévenard. 1998. Linking landfast sea ice variability to marine ice accretion at Hells Gate Ice Shelf, Ross Sea. In Jeffries, M. O., ed. *Antarctic sea ice: physical processes, interactions and variability*. Washington, DC, American Geophysical Union, 375–407. (Antarctic Research Series 74)
- Trodahl, H. J. and 6 others. 2000. Heat transport in McMurdo Sound first-year fast ice. *J. Geophys. Res.*, **105**(C5), 11,347–11,358.
- Unesco. 1981. Annex 1. The practical salinity scale 1978 and the international equation of state of seawater 1980. In *Tenth report of the Joint Panel on Oceanographic Tables and Standards*. Paris, Unesco, 13–17. (Technical Papers in Marine Science 36.)
- United States National Imagery and Mapping Agency. 1997. *Pub. 200 – sailing directions (planning guide and enroute) Antarctica*. Third edition. Bethesda, MD, U.S. National Imagery and Mapping Agency.