

The role of microRNAs in the regulation of critical genes and signalling pathways that determine endometrial receptivity

Review Article

Cite this article: Wang Y *et al.* (2024). The role of microRNAs in the regulation of critical genes and signalling pathways that determine endometrial receptivity. *Zygote*, page 1 of 10. doi: [10.1017/S0967199424000297](https://doi.org/10.1017/S0967199424000297)


Received: 22 May 2024
Revised: 30 July 2024
Accepted: 7 August 2024

Keywords:

miRNA; endometrial receptivity; embryo implantation; endometrial epithelial cells

Corresponding authors:

Xingping Wang;
Email: wxp@nxu.edu.cn;
Zhuoma Luoreng;
Email: luorenzhuoma@nxu.edu.cn

Yumei Wang , Zhongxiang Ji, Ni Yao, Ximin Hu, Ran Zhou, Xingping Wang and Zhuoma Luoreng

College of Animal Science and Technology, Ningxia University, Yin Chuan, NingXia, 750021, China

Summary

Endometrial receptivity is the ability of the endometrium to accept embryos. Thus, endometrial receptivity dysfunction is an important factor leading to embryo implantation failure. A good endometrial receptivity provides a suitable environment for embryo implantation, improving the embryo implantation rate. The “implantation window” stage, or the receptive stage of the endometrium, is regulated by various hormones, genes, proteins and cytokines, among which microRNAs (miRNAs) and their target genes have a regulatory effect on endometrial receptivity. This review outlines the relationship between endometrial receptivity and pregnancy, the mRNAs and related signalling pathways that regulate endometrial receptivity, and the regulatory role of miRNA in endometrial receptivity, providing a deeper understanding of the regulatory mechanisms of miRNA on endometrial receptivity in humans and animals and reference for the endometrial receptivity-related research.

Introduction

Endometrial receptivity is the ability of the endometrium to allow normal embryo implantation. It is the intricate process undertaken by the endometrial tissue to prepare for blastocyst implantation and pregnancy initiation. A good endometrial receptivity provides opportunities for embryo attachment, invasion and development, culminating in a new individual and species continuation (Neykova *et al.*, 2022; Lessey and Young, 2019). Embryo implantation occurs during the implantation window, a mid-secretory phase of the menstrual cycle. Many molecular pathways involving hormones, adhesion molecules, cytokines and growth factors synchronize the implantation window during this phase. Thus, a loss of synchronisation or failure to achieve endometrial receptivity can lead to a miscarriage or infertility (Governini *et al.*, 2021; Blanco-Breindel *et al.*, 2023).

Mammals successfully implant embryos only when the endometrium is in a receptive state, during which the endometrial epithelial cells (EECs) are reshaped to accept embryo implantation. Generally, embryo implantation requires a two-way communication between the blastocyst and the receptive endometrium, with poor quality blastocysts or defective endometrial receptivity leading to an implantation failure (Makrigiannakis *et al.*, 2021; Paulson and Comizzoli, 2021; Bui *et al.*, 2022). During the early implantation stage, the EECs undergo epithelial-mesenchymal transition (EMT) to prepare for subsequent embryo implantation. Subsequently, the endometrial stromal cells (ESCs) undergo decidualization to create a favourable developmental environment for the embryo. Based on this, the EMT and decidualization are crucial for maintaining a good endometrial receptivity. A good endometrial receptivity is the biological basis for embryo implantation and normal development (Crha *et al.*, 2019; Cai *et al.*, 2022). Therefore, improving endometrial receptivity is a key factor in increasing pregnancy rates.

This article provides an overview of mRNA and its related signalling pathways that regulate endometrial receptivity, as well as the regulatory role of miRNA in endometrial receptivity. It provides an important reference for a deeper understanding of the molecular regulatory mechanisms of miRNA on human and animal endometrial receptivity.

mRNAs regulating endometrial receptivity

Homeobox (HOX)

HOX is important in cell proliferation and differentiation and is the main regulatory factor for cell identity and fate during embryonic development (Steens and Klein, 2022; Smith *et al.*, 2019). For example, *HOXA10* can bind to the E-cadherin promoter region and directly regulate its expression, thereby improving endometrial receptivity, which subsequently increases embryo

adhesion and implantation (Zhang *et al.*, 2017). Besides, the mechanism of metformin treatment, which enhances endometrial receptivity, is associated with the upregulation of the *HOXA10* gene (Cheng *et al.*, 2022). Notably, *HOXA9*, *HOXA11* and *HOXD10* are involved and highly expressed during endometrial receptivity. The embryo implantation rate decreases when the *HOXA9*, *HOXA11* and *HOXA10* genes are silenced in the mouse uterus (Kara *et al.*, 2019; Xu *et al.*, 2014). In addition, the expression levels of *HOXA10* and *HOXA11* are significantly reduced in women with endometriosis compared to the non-endometriotic controls, which potentially significantly affects endometrial remodelling and the expression of endometrial receptivity markers (Jana *et al.*, 2013). Therefore, a high expression of *HOX* genes, including *HOXA10*, *HOXA9*, *HOXA11* and *HOXD10*, can serve as markers for achieving endometrial receptivity.

Interleukin (IL)

ILs are lymphokines excreted by the white blood cell and other immune cell (Khan *et al.*, 2023). Among them, *IL-1* is positively correlated with endometrial receptivity in patients with polycystic ovary syndrome (PCOS); thus, it is a therapeutic target for PCOS to improve endometrial receptivity (Zhao *et al.*, 2019). In addition, the production of *IL-1 β* promotes oocyte maturation, fertilisation, and endometrial receptivity, thereby promoting embryo implantation (Rehman *et al.*, 2015). Specifically, *IL-1 β* is highly expressed in the mid-luteal phase of implantation and positively correlated with endometrial thickness, serum estradiol and progesterone levels, implying that *IL-1 β* improves endometrial receptivity, which promotes embryonic development (Wang *et al.*, 2018). Additionally, *IL-10* promotes the attachment of BeWo spheroids to Ishikawa cells and increases *HOXA10* expression through phosphorylation of signal transducer and activator of transcription 3 (STAT3), thereby promoting endometrial receptivity (Wang *et al.*, 2018). *IL-33* promotes endometrial receptivity by increasing *HOXA10* expression through the phosphorylation of STAT3. Decreased intracellular *IL-33* impairs endometrial receptivity in women with adenomyosis (He *et al.*, 2022). In summary, the positive correlation between *IL* members, such as *IL-1*, *IL-1 β* , *IL-10* and *IL-33*, and endometrial receptivity suggests that *IL* promotes endometrial receptivity and embryo implantation.

Leukaemia inhibitory factor (LIF)

LIF is a cytokine in the *IL-6* family, which regulates the adhesion characteristics of EECs, enhances endometrial receptivity, and promotes embryo implantation and pregnancy (Camargo-Diaz *et al.*, 2017). Based on in vitro and in vivo studies on the effect of the *Cnidium officinale* Makino (CoM) roots on endometrial receptivity using embryo implantation models, CoM enhances the adhesion of JAr cells to Ishikawa cells by activating the expression of *LIF* and integrin, thereby enhancing endometrial receptivity (Chung *et al.*, 2019). Hajipour *et al.* (2021) also revealed that the upregulation of *LIF* is one of the mechanisms by which sildenafil citrate enhances endometrial receptivity. In addition, various bioactive substances such as paeoniflorin increase the adhesion of JAr cells derived from the trophoblastic ectoderm to Ishikawa cells in the endometrium by activating *LIF* expression, which enhances endometrial receptivity (Eun-Yeong *et al.*, 2019; Park *et al.*, 2021; Choi *et al.*, 2017). Therefore, activating *LIF* expression enhances the adhesion between JAr cells and Ishikawa cells, promoting endometrial receptivity.

Vascular endothelial growth factor (VEGF)

Vascular endothelial growth factor (*VEGF*), or vascular permeability factor, is a highly specific endothelial cell mitogen and the most important factor regulating angiogenesis (Dvorak, 2000). *VEGF* stimulates embryo implantation by promoting embryonic development, improving endometrial receptivity, and enhancing the interaction between developing embryos and the endometrium (Guo *et al.*, 2021; Zarei *et al.*, 2019). For example, Bu Shen Zhu Yun Decoction activates the mitogen-activated protein kinase (MAPK) signalling pathway by upregulating *VEGF* and *VEGF* receptor-2, thereby improving endometrial receptivity and embryo implantation rate (Li *et al.*, 2019). In addition, Tiaojing Zhu Yun Formula improves endometrium receptivity and facilitates embryo implantation of controlled ovarian hyperstimulation (COH) rats by upregulating *VEGF* and enhancing the phosphatidylinositol 3 kinase/protein kinase B (PI3K/Akt) signalling pathway (You *et al.*, 2022a). Endometrial macrophages modulate uterine receptivity by regulating the expression of *VEGFA*, thereby affecting embryo implantation (Wang *et al.*, 2016). In summary, the upregulation of *VEGF* improves endometrial receptivity and promotes embryo implantation and development.

Other mRNAs that regulate endometrial receptivity

Other mRNAs are also closely related to endometrial receptivity in various species such as mouse, rat, human and goat (Table 1). Their expression trends are also clear; thus, they can serve as potential biomarkers of endometrial receptivity.

miRNA-mediated regulation of the key signalling pathways in endometrial receptivity

MicroRNAs (miRNAs) are a class of non-coding single-stranded ribonucleic acid (RNA) molecules, typically 20-24 nt in length encoded by endogenous genes. MiRNA-encoding genes are transcribed by RNA polymerase II to generate primary transcripts, which are then processed into small RNAs of approximately 21 nucleotides by RNase III endonucleases DROSHA and DICER. All miRNAs are loaded into Argonaute protein in RNA-induced silencing complex and act as post-transcriptional regulators by binding to the 3'-untranslated region of mRNA (Stavast and Erkeland, 2019; Michlewski and Caceres, 2019). MiRNAs play an important regulatory role in endometrial receptivity and the formation of related factors during embryo implantation (Liang *et al.*, 2017; Shekibi *et al.*, 2022; Goharitaban *et al.*, 2022). Notably, miRNAs regulate endometrial receptivity-related key genes to promote or suppress endometrial receptivity by activating or inhibiting signalling pathways, such as PI3K/Akt, extracellular signal-regulated kinase/mechanistic target of rapamycin (ERK/mTOR), NF- κ B, Wnt/ β -Catenin and LIF/STAT3 signalling pathways.

Regulation of gene expression by miRNA via the PI3K/Akt signalling pathway

The PI3K/Akt signalling pathway plays a crucial role in cell functions such as proliferation, adhesion, invasion, migration, metabolism and survival through phosphorylation or dephosphorylation. It also plays a role in angiogenesis, endometrial receptivity and successful embryo implantation (Shen *et al.*, 2023). In addition, acupuncture and moxibustion improve endometrial angiogenesis by activating the PI3K/Akt signalling pathway, and

Table 1. Other mRNAs regulating endometrial receptivity

Gene name	Species	The role of mRNA	Change in expression	References
<i>GRIM-19</i>	mouse	Regulating endometrial receptivity and embryo implantation by regulating adhesion, cell apoptosis, and immune tolerance	down	(Yang <i>et al.</i> , 2017)
<i>TLR2/6</i>	mouse	Inhibiting endometrial receptivity by reducing the in vitro adhesion between trophoblast spheres and endometrial cells	down	(Sanchez-Lopez <i>et al.</i> , 2014)
<i>MUC-1</i>	mouse	<i>MUC1</i> expression is negatively correlated with fertility	down	(Shokrzadeh <i>et al.</i> , 2019)
$\alpha V\beta 3$ and $\alpha V\beta 5$	mouse	Promoting the process of embryo endometrial adhesion	up	(Choi <i>et al.</i> , 2017)
<i>MMP9</i>	rat	Promoting endometrial invasion and lesions to suppress endometrial receptivity	down	(Cheng <i>et al.</i> , 2022)
<i>ILK</i>	human	By activating Wnt/ β -Catenin signalling improves the formation of endometrial receptivity	up	(Chen <i>et al.</i> , 2020)
<i>HB-EGF</i>	mouse/human	Positive correlation with endometrial thickness	up	(Shokrzadeh <i>et al.</i> , 2018; Sutaji <i>et al.</i> , 2023)
<i>OPN</i>	mouse/rat	Promoting blastocyst formation and embryo implantation	up	(Liu <i>et al.</i> , 2013; Zarrin <i>et al.</i> , 2020)
<i>COX-2</i>	mouse	Expression is negatively correlated with endometrial receptivity	down	(Yu <i>et al.</i> , 2021)
<i>ITG$\beta 3$</i>	human	Participate in cell adhesion and cell surface-mediated signalling, and play an important role in regulating endometrial receptivity	up	(Luo <i>et al.</i> , 2021)
<i>EMX2</i>	human	High expression in the endometrium often leads to abnormal development of the endometrium	down	(Luo <i>et al.</i> , 2021)
<i>FOXA2</i>	human	Potential regulatory genes affecting uterine receptivity, blastocyst implantation, and stromal cell decidualization	up	(Kelleher <i>et al.</i> , 2019)
<i>PODXL (PCX)</i>	human	<i>PODXL (PCX)</i> is a key negative regulator of human endometrial epithelial receptivity for embryo implantation	down	(Paule <i>et al.</i> , 2021)
<i>PGR</i>	mouse	Control epithelial and stromal progesterone reactivity and epithelial cell differentiation to provide normal uterine receptivity and subsequent embryo attachment	up	(Gebriel <i>et al.</i> , 2020)
<i>BCL2L15</i>	goat	<i>BCL2L15</i> is the key regulator of endometrial receptivity in goats, regulating the endometrial receptivity via the STAT1 pathway	up	(Yang <i>et al.</i> , 2020)
<i>HIF-1α</i>	human	<i>HIF-1α</i> is positively associated with endometrial receptivity by regulating PKM ₂	up	(Liang <i>et al.</i> , 2023)
<i>IGF1R</i>	mouse	Promote embryo attachment	up	(Kang <i>et al.</i> , 2015)




Figure 1. The mechanism of miRNA in regulating endometrial receptivity via the PI3K/AKT signalling pathway. Some miRNAs promote the activation of the PI3K/Akt signalling pathway by regulating the expression of target mRNA, thereby altering the expression of key genes involved in endometrial receptivity, consequently improving endometrial receptivity. The figure is made by WPS, which is a Chinese office software.

increase endometrial receptivity and implantation sites in PCOS rats (Xing *et al.*, 2022). Liu *et al.* (2018) also revealed that circ-8073 regulates *CEP55* by sponging miR-449a in dairy goat EECs,

promoting the proliferation of the EECs via the PI3K/Akt/mTOR pathway, and regulating endometrial receptivity in dairy goat (Figure 1). In addition, Wang *et al.* (2023) confirmed that human

chorionic gonadotropin activates the PI3K/Akt/eNOS pathway via the miR-126-3p/*PIK3R2* axis, promoting EEC proliferation and improving endometrial receptivity (Figure 1). Moreover, Talin1 and phosphatidylinositol 4,5-phosphate kinase (*PIP1K1*) competitively bind miR-1285-3p to activate the PI3K/Akt signalling pathway. The activation of the PI3K/Akt signalling pathway further promotes the expression of *ERK2*, *FAK* and Vinculin, $\alpha\beta3$ molecules crucial in the embryo implantation process, promoting the adhesion ability of EECs to the embryo (Li, 2019) (Figure 1). In summary, miRNA promotes the activation of the PI3K/Akt signalling pathway by regulating the expression of target mRNA, which alters the expression of endometrial receptivity key genes, thereby improving endometrial receptivity.

Regulation of gene expression by miRNA via the ERK/mTOR signalling pathway

Various studies have revealed that the mTOR pathway is important in endometrial receptivity (Qi *et al.*, 2022; Niknafs *et al.*, 2022). For example, dexamethasone disrupts endometrial receptivity by altering the expression of miR-223, miR-200a, LIF, Muc1, SGK1 and ENaC via the ERK1/2-mTOR pathway (Shariati *et al.*, 2019). On the contrary, flhydrocortisone (Hesam *et al.*, 2019) and calcitonin (Shokrzadeh *et al.*, 2018) improve endometrial receptivity by regulating the expression of endometrial receptivity-related genes (*HB-EGF*, *MSX.1* and miRNA *Let-7a*) and activating the ERK1/2-mTOR pathway (Figure 2). Additionally, Cui *et al.* (2020) revealed that miR-184 targets and inhibits the expression of *STC2* via the RAS/RAF/MEK/ERK signalling pathway, thereby promoting the expression of *FOXM1* and *VEGF* and apoptosis of EECs, consequently promoting endometrial receptivity in dairy goats (Figure 2). In summary, miRNA promotes endometrial receptivity by activating the ERK/mTOR signalling pathway to regulate endometrial receptivity-related genes.

Regulation of gene expression by miRNA via the NF- κ B signalling pathway

Progesterone and melatonin improve endometrial receptivity in mice and cows, increasing their pregnancy rates, which is closely related to the NF- κ B signalling pathway (Feng *et al.*, 2022; Guan *et al.*, 2022; Zheng *et al.*, 2022). Notably, high progesterone levels during early pregnancy in cows induce high expression of miR-20b. MiR-20b blocks the NF- κ B signalling pathway by negatively regulating *NCOA3* and inhibits the immune response of bovine EECs, crucial in uterine tissue remodelling, changes in uterine immune microenvironment and placental formation (Yang, 2021) (Figure 3). In addition, miR-182-5p activates the NF- κ B signalling pathway by negatively regulating the target gene *NDRG1*, which reduces the E-cadherin expression, inhibits endometrial receptivity and causes embryo implantation failure (Yu *et al.*, 2022) (Figure 3).

The NF- κ B signalling pathway is also an important pathway regulating inflammation (Capece *et al.*, 2022). Interestingly, enhancing the endometrium inflammatory process reduces endometrial receptivity (Zhang *et al.*, 2021; Pirtea *et al.*, 2021). Therefore, some of the miRNAs in the NF- κ B signalling pathway could be regulating the expression of inflammatory factors, interfering with the embryo implantation process, and reducing the ability of the endometrium to receive embryos. In summary, miRNAs often activate the NF- κ B signalling pathway, suppressing endometrial receptivity. Therefore, future studies should explore




Figure 2. The mechanism of miRNA in regulating endometrial receptivity via the ERK/mTOR signalling pathway. MiR-184 regulates genes associated with endometrial receptivity by activating the ERK/mTOR signalling pathway, thus promoting endometrial receptivity. Some drug treatments regulate the expression of endometrial receptivity-related genes and miRNAs via the ERK/mTOR signalling pathway, consequently altering endometrial receptivity. The figure is made by WPS, which is a Chinese office software.




Figure 3. The mechanism of miRNA in regulating endometrial receptivity via the NF- κ B signalling pathway. MiRNA typically inhibits target mRNA to activate NF- κ B signalling pathway inhibits endometrial receptivity. MiRNA typically inhibits target mRNA to block NF- κ B signalling pathway promotes endometrial receptivity. The figure is made by WPS, which is a Chinese office software.

the miRNAs that inhibit the NF- κ B signalling pathway to improve endometrial receptivity.

Regulation of gene expression by miRNA via the Wnt/ β -catenin signalling pathway

The Wnt/ β -catenin signalling pathway is a highly conserved pathway involved in various processes such as cell development, proliferation, differentiation, apoptosis and autophagy. It plays a crucial role in embryonic development and maintaining dynamic homeostasis in the body (Ma *et al.*, 2023; Liu *et al.*, 2023). *Integrin-*




Figure 4. The mechanism of miRNA in regulating endometrial receptivity via the Wnt/ β -catenin signalling pathway. MiRNA typically inhibits target mRNA, thereby inactivating the Wnt/ β -catenin signalling pathway, leading to the inhibition of endometrial receptivity. The figure is made by WPS, which is a Chinese office software.

linked kinase (*ILK*) (Chen *et al.*, 2020) and baicalin (Zhang *et al.*, 2015) promote endometrial receptivity and embryo implantation via the Wnt/ β -catenin signalling pathway (Figure 4). Zheng *et al.* (2017) also revealed the miR-200c targeted inhibition of fucosyltransferase IV (*FUT4*) expression while reducing the glycoprotein CD44 α 1,3-fucosylation inactivates the Wnt/ β -catenin signalling pathway, leading to endometrial receptivity dysfunction (Figure 4). Besides, miR-543 inhibits *MAPK* expression and inactivates the Wnt/ β -catenin pathway, inducing the inhibition of the EMT of ESCs by transforming growth factors β (Wang *et al.*, 2021) (Figure 4). In summary, the Wnt/ β -catenin signalling pathway plays an important regulatory role in initiating endometrial receptivity. The miRNAs often activate the Wnt/ β -Catenin signalling pathway, promoting endometrial receptivity, or suppress the Wnt/ β -catenin signalling pathway, inhibiting endometrial receptivity.

Regulation of gene expression by miRNA via the LIF/STAT3 signalling pathway

The LIF/STAT3 signalling pathway also affects endometrial receptivity (Liang *et al.*, 2018, 2021; Hu *et al.*, 2019). For example, implantation failure in subclinical hypothyroidism rats is related to abnormal LIF/STAT3 signalling (Shan *et al.*, 2019). Generally, COH induces endometrial receptivity dysfunction during the implantation window. However, electroacupuncture therapy reduces cell adhesion molecules such as E-Cadherin, β -catenin and *CLDN1*, which activates the LIF/STAT3 signalling pathway, improves endometrial receptivity, and promotes blastocyst implantation in COH rats (You *et al.*, 2021) (Figure 5). In addition, miR-223-3p expression inhibits the LIF/STAT3 signalling pathway in the rat endometrium. However, electroacupuncture therapy enhances endometrial receptivity by inhibiting miR-223-3p expression in the LIF/STAT3 signalling pathway (You *et al.*, 2022b) (Figure 5). In summary, miRNA promotes endometrial receptivity by regulating the LIF/STAT3 signalling pathway. However, there are few reports on the specific regulatory effects of miRNA on endometrial receptivity. Therefore, the




Figure 5. The mechanism of miRNA in regulating endometrial receptivity via the Leukaemia inhibitory factor (LIF)/STAT3 signalling pathway. Electroacupuncture therapy reduces cell adhesion molecules, thereby activating the LIF/STAT3 signalling pathway, which improves endometrial receptivity. Electroacupuncture therapy activates the LIF/STAT3 signalling pathway by inhibiting the expression of miR-223-3p, thereby enhancing endometrial receptivity. The figure is made by WPS, which is a Chinese office software.

LIF/STAT3 signalling pathway is a potential signalling pathway for miRNA regulation of endometrial receptivity, though more research is needed.

The regulatory role of miRNA in endometrial receptivity

A good endometrial receptivity is a prerequisite for successful embryo implantation. Therefore, scholars are constantly exploring methods to improve endometrial receptivity. Many miRNAs regulate endometrial receptivity in mouse and other animals and can serve as useful diagnostic and therapeutic targets for successful embryo implantation, providing reference for relevant scientific research and clinical application (Table 2).

The regulatory effect of miRNA on endometrial receptivity in mouse

Cullin 3 (*CUL3*), an E3 ubiquitin ligase, interacts with beclin 1 (*BECN1*), thereby promoting K48-linked ubiquitination and degradation of *BECN1*. In mice, miR-23a-3p inhibits target gene *CUL3* expression by negatively regulating β -Catenin ubiquitination to promote endometrial receptivity and embryo implantation (Huang *et al.*, 2020). According to Akbar *et al.* (2020), inhibiting miR-183-5p also significantly reduces the embryo implantation rate and increases the expression of target gene catenin alpha 2 (*CTNNA2*) in vivo using a mouse pregnancy model, implying that miR-183-5p regulates endometrial receptivity and enhances embryo implantation. Another study revealed that miR-let-7a/g enhances mouse endometrial receptivity by inhibiting the Wnt signalling pathway by negatively regulating oestrogen and progesterone, improving embryo attachment and growth related to implantation (Li *et al.*, 2020).

Poly (ADP-ribose) polymerase 2 (*PARP-2*) is a gene involved in endometrial receptivity for trophoblast implantation. Moreover, miR-149 inhibits endometrial receptivity by downregulating the

Table 2. The regulatory role of miRNA in endometrial receptivity

miRNA	Species	Target Gene	Improved/Impaired Receptivity
miR-23a-3p	mouse	<i>CUL3</i>	Improved
miR-183-5p	mouse	<i>CTNNA2</i>	Improved
miR-let-7a/g	mouse	–	Improved
miR-149	mouse	<i>PARP-2</i>	Impaired
miR-223-3p	mouse	<i>LIF</i>	Impaired
miR-181	mouse	<i>LIF</i>	Impaired
miR-494-3p	mouse	<i>LIF</i>	Impaired
miR-183	mouse	<i>HB-EGF</i> and <i>LAMC1</i>	Impaired
miR-205-5p	mouse	<i>ZEB1</i>	Improved
miR-145 and miR-199	human	–	Improved
miR-30d	human	<i>Snai1</i>	Improved
miR-133b	human	<i>SGK1</i>	Improved
miR-1290	human	<i>LHX6</i>	Improved
miR-192-5p	human	<i>ARHGAP19</i>	Impaired
miR-125b	human	<i>MMP26</i>	Impaired
miR-146a-3p	human	<i>ZNF23</i>	Impaired
miR-29c	human	<i>COL4A1</i>	Impaired
miR-23b-3p	human	<i>SFRP4</i>	Improved
miR-543	human	–	Improved
miR-31	human	<i>FOXP3</i> and <i>CXCL12</i>	Improved
miR-124-3p	human	<i>LIF</i> , <i>MUC1</i> and <i>BCL2</i>	Impaired
miR-449a	goat	<i>LGR4</i>	Improved
miR-182	dairy goat	<i>HOXA10</i>	Improved
miR-26a	dairy goat	<i>PTEN</i>	Improved
miR-15b	dairy goat	<i>LIF</i>	Improved
miR-34a/c	caprine	<i>CEP55</i>	Improved
bta-miR-200b	bovine	<i>MYB</i>	Improved
ssc-miR-21-5p	pig	<i>PDCD4</i>	Impaired
miR-1246	buffalo	<i>MUC1</i> and β - <i>catenin</i>	Improved
miR-134-5p	sheep	<i>CREB1</i>	Impaired

target gene *PARP-2* and upregulating caspase-8 (Verma et al., 2022; Soni et al., 2021). At the same time, miR-223-3p and miR-181 inhibit endometrial receptivity and embryo implantation in mice by targeting and downregulating the *LIF*, a key regulatory factor for endometrial receptivity (Dong et al., 2016; Chu et al., 2015). On the contrary, electroacupuncture therapy enhances endometrial receptivity by inhibiting the expression of miR-223-3p on the *LIF/STAT3* signalling pathway (You et al., 2022b). Yuan et al. (2021) also revealed that miR-494-3p targeting *LIF* regulates endometrial receptivity via the *PI3K/Akt/mTOR* pathway, and inhibiting miR-494-3p increases

the *LIF* and *VEGF* expression levels, thereby increasing endometrial receptivity in mice. Besides, miR-183 inhibits embryo implantation by regulating *HB-EGF* and *LAMC1* in the mouse uterus, suggesting that miR-183 inhibits endometrial receptivity (Cao et al., 2020). Exosomes miR-205-5p upregulate E-cadherin by targeting *ZEB1*, thereby improving endometrial receptivity (Yu et al., 2023).

The regulatory effect of miRNA on endometrial receptivity in human

miR-145 and miR-199 transfection into Ishikawa cells significantly reduces the *podocalyxin (PODXL)* expression, which significantly promotes the endometrial receptivity of the Ishikawa monolayer (Shekibi et al., 2022). Polychlorinated biphenyls induce endometrial receptivity dysfunction by disrupting the interaction between the key gene miR-30d regulating endometrial receptivity and the EMT process. *Snai1* is targeted by miR-30d and induced by polychlorinated biphenyls (Cai et al., 2016). Lack of miR-30d in the mother or embryo results in a significant downregulation of the endometrial receptivity markers, which reduces the implantation rate and damages embryonic growth (Balaguer et al., 2019). In addition, Kong et al. (2016) revealed that miR-133b targets *SGK1* to increase *HOXA10* expression and promote the attachment of BeWo spheres to Ishikawa cells, which induce endometrial receptivity and embryo attachment. Shi et al. (2021) also revealed that miR-1290 is an extracellular vesicle derived from placental trophoblasts, which stimulates the interaction between the endometrium and embryo by targeting *LHX6*, thereby promoting endometrial receptivity.

Moreover, miR-192-5p is an upstream regulatory factor of *ARHGAP19*, which promotes the transition of EECs from non-receptive to receptive states by regulating the remodelling of connexin and membrane-related cytoskeleton (Liang et al., 2021). Another study revealed that miR-192-5p inhibits endometrial receptivity by hindering epithelial transformation during embryo implantation (Liang et al., 2020). Therefore, sustained high levels of miR-192-5p are harmful to embryo implantation. At the same time, miR-125b inhibits EEC migration and invasion by downregulating *MMP26*, which weakens the embryo attachment and subsequent endometrial invasion, leading to decreased endometrial receptivity (Chen et al., 2016). The overexpression of miR-146a-3p also significantly inhibits the expression of *LIF*, integrin3, claudin4 and *DKK1* in EECs, which inhibits endometrial receptivity (Wei et al., 2021). In the endometrium of infertile women, the overexpression of miR-29c and downregulation of *COL4A1* reduces the EECs adhesion ability in vitro, suggesting that miR-29c plays a role in the secretion phase of infertile women, decreasing receptivity and implantation success rate (Griffiths et al., 2019). MiR-23b-3p also regulates the adhesion of human EECs, suggesting that it may be a key regulator of endometrial epithelial adhesion and endometrial receptivity (Barton et al., 2023).

Furthermore, research has also revealed that the downregulation of miR-543 affects embryo implantation, leading to endometriosis-related infertility (Yang et al., 2019). Another study revealed that miR-31 is the best potential biomarker for endometrial receptivity and may function through immunosuppressive mechanisms (Kresowik et al., 2014). MiR-124-3p negatively impacts embryo implantation by suppressing endometrial receptivity formation and embryo development through targeted inhibition of *LIF*, *MUC1* and *BCL2* expression (Yao et al., 2024).

The regulatory effect of miRNA on endometrial receptivity in livestock

The miR-449a/c expression in the uterine cycle is associated with endometrial development (Naydenov *et al.*, 2022). MiR-449a promotes endometrial receptivity by regulating the apoptosis of goat ESCs and negatively regulating *LGR4* (An *et al.*, 2017). Similarly, miR-182 promotes endometrial receptivity by down-regulating pleiotrophin and regulating apoptosis-related genes and endometrial receptivity marker genes such as *osteopontin* (*OPN*), *cyclooxygenase-2* (*COX-2*) and *prolactin receptor* (*PRLR*) in dairy goat EECs (Zhang *et al.*, 2017). miR-26a also regulates the expression of *OPN*, *VEGF*, *COX-2* and *prolactin* (*PRL*) in endometrial cells, thereby promoting endometrial receptivity (Zhang *et al.*, 2018). At the same time, miR-26a promotes the proliferation of EECs and induces ESC apoptosis via the *PTEN*-PI3K/Akt pathway in dairy goats (Zhang *et al.*, 2018).

Furthermore, miR-15b is competitively bound by lncRNA882 and LIF in EECs, which upregulates *LIF* and induces endometrial receptivity in dairy goats (Zhang *et al.*, 2019). Liu *et al.* (2020) also revealed that miR-34a and miR-34c are highly expressed in goats and induce apoptosis of EECs by binding to circ-8073 and *CEP55* via the RAS/RAF/MEK/ERK and PI3K/Akt/mTOR pathways, which induce endometrial receptivity in dairy goats. The apoptosis of EECs is an important process for endometrial receptivity induction and embryo implantation. However, bta-miR-200b affects the apoptosis of bovine EECs by targeting *MYB* genes, thereby promoting endometrial receptivity (Lyu *et al.*, 2023). In pigs, knocking down ssc-miR-21-5p inhibits Akt phosphorylation by targeting *PDCD4*, which hinders endometrial receptivity, decreasing the embryo implantation rate (Hua *et al.*, 2020). Besides, in buffalo, miR-1246 secreted by the trophoblast reduces the expression of endometrial receptivity gene (*MUC1* and *β-catenin*), implying that miR-1246 is a potential marker for buffalo endometrial receptivity (Dubey *et al.*, 2023). In the same line, miR-134-5p inhibits the proliferation of sheep ESCs by inhibiting the target gene *CREB1*, thereby inhibiting the establishment of endometrial receptivity (Li *et al.*, 2023).

Conclusions and future perspectives

endometrial receptivity is crucial for successful embryo implantation and pregnancy. endometrial receptivity dysfunction induces EEC attachment failure or poor embryo reception, which decreases the pregnancy rate. Various factors influence endometrial receptivity via multiple signalling pathways, including mRNA and miRNAs, which play a crucial role in endometrial receptivity induction by participating in the signalling pathways, such as the PI3K/Akt signalling pathway. Although the regulatory mechanisms of some miRNAs regulating endometrial receptivity have been studied, the regulatory mechanism of miRNAs on endometrial receptivity is still unclear. Therefore, further research on the molecular regulatory network of miRNA on endometrial receptivity is needed to improve endometrial receptivity and pregnancy rates. In addition, miRNAs should be explored as biomarkers or therapeutic targets for detecting or improving endometrial receptivity in maternal therapy.

Acknowledgements. We would like to thank the editor of MogoEdit professional English editing company (Mogo Internet Technology Co., LTD, Shaanxi, China) for help in the preparation of this manuscript.

Author contributions. The idea of the manuscript was created by Yumei Wang. All authors contributed to the initial conception and creation structure of the manuscript. Yumei Wang wrote the first draft. All authors contributed to the improvement and final edition of the manuscript.

Competing interests. The authors declare no conflict of interest.

Funding. This research was supported by Project of Ningxia Hui Autonomous Region Key Research and Development Program (No. 2021BEF02029); Project of Ningxia Hui Autonomous Region Key Research and Development Program (Special Talent Introduction) (No. 2020BEB04006); and the earmarked fund for CARS (No. CARS-36).

References

- Akbar, R., Ullah, K., Rahman, T. U., Cheng, Y., Pang, H. Y., Jin, L. Y., Wang, Q. J., Huang, H. F. and Sheng, J. Z. (2020) miR-183-5p regulates uterine receptivity and enhances embryo implantation. *Journal of Molecular Endocrinology*, **64**, 43–52.
- An, X., Liu, X., Zhang, L., Liu, J., Zhao, X., Chen, K., Ma, H., Li, G., Cao, B. and Song, Y. (2017) MiR-449a regulates caprine endometrial stromal cell apoptosis and endometrial receptivity. *Scientific Reports*, **7**, 12248.
- Balaguer, N., Moreno, I., Herrero, M., Gonzalez-Monfort, M., Vilella, F. and Simon, C. (2019) MicroRNA-30d deficiency during preconception affects endometrial receptivity by decreasing implantation rates and impairing fetal growth. *American Journal of Obstetrics and Gynecology*, **221**, 41–46.
- Barton, S., Zhou, W., Santos, L. L., Menkhorst, E., Yang, G., Tinn, T. W., Ang, C., Lucky, T. and Dimitriadis, E. (2023) miR-23b-3p regulates human endometrial epithelial cell adhesion implying a role in implantation. *Reproduction*, **165**, 407–416.
- Blanco-Breindel, M. F., Singh, M. and Kahn, J. (2023) *Endometrial Receptivity*. StatPearls Publishing.
- Bui, A. H., Timmons, D. B. and Young, S. L. (2022) Evaluation of endometrial receptivity and implantation failure. *Current Opinion in Obstetrics and Gynecology*, **34**, 107–113.
- Cai, J. L., Liu, L. L., Hu, Y., Jiang, X. M., Qiu, H. L., Sha, A. G., Wang, C. G., Zuo, Z. H. and Ren, J. Z. (2016) Polychlorinated biphenyls impair endometrial receptivity in vitro via regulating miR-30d expression and epithelial mesenchymal transition. *Toxicology*, **365**, 25–34.
- Cai, X., Xu, M., Zhang, H., Zhang, M., Wang, J., Mei, J., Zhang, Y., Zhou, J., Zhen, X., Kang, N., Yue, Q., Sun, H., Jiang, R. and Yan, G. (2022) Endometrial stromal PRMT5 plays a crucial role in decidualization by regulating NF-kappaB signaling in endometriosis. *Cell Death Discovery*, **8**, 408.
- Camargo-Diaz, F., Garcia, V., Ocampo-Barcenas, A., Gonzalez-Marquez, H. and Lopez-Bayghen, E. (2017) Colony stimulating factor-1 and leukemia inhibitor factor expression from current-cycle cannula isolated endometrial cells are associated with increased endometrial receptivity and pregnancy. *BMC Women's Health*, **17**, 63.
- Cao, D., Liang, J., Feng, F., Shi, S., Tan, Q. and Wang, Z. (2020) MiR-183 impeded embryo implantation by regulating Hbegf and Lamc1 in mouse uterus. *Theriogenology*, **158**, 218–226.
- Capece, D., Verzella, D., Flati, I., Arboretto, P., Cornice, J. and Franzoso, G. (2022) NF-kappaB: blending metabolism, immunity, and inflammation. *Trends in Immunology*, **43**, 757–775.
- Chen, C., Zhao, Y., Yu, Y., Li, R. and Qiao, J. (2016) MiR-125b regulates endometrial receptivity by targeting MMP26 in women undergoing IVF-ET with elevated progesterone on HCG priming day. *Scientific Reports*, **6**, 25302.
- Chen, Q., Ni, Y., Han, M., Zhou, W. J., Zhu, X. B. and Zhang, A. J. (2020) Integrin-linked kinase improves uterine receptivity formation by activating Wnt/beta-catenin signaling and up-regulating MMP-3/9 expression. *American Journal of Translational Research*, **12**, 3011–3022.
- Cheng, J., Li, C., Ying, Y., Lv, J., Qu, X., McGowan, E., Lin, Y. and Zhu, X. (2022) Metformin alleviates endometriosis and potentiates endometrial receptivity via decreasing VEGF and MMP9 and increasing leukemia inhibitor factor and HOXA10. *Frontiers in Pharmacology*, **13**, 750208.

- Choi, H. J., Chung, T. W., Park, M. J., Jung, Y. S., Lee, S. O., Kim, K. J. and Ha, K. T. (2017) Water-extracted tubers of *Cyperus rotundus* L. enhance endometrial receptivity through leukemia inhibitory factor-mediated expression of integrin α Vbeta3 and α Vbeta5. *Journal of Ethnopharmacology*, **208**, 16–23.
- Chu, B., Zhong, L., Dou, S., Wang, J., Li, J., Wang, M., Shi, Q., Mei, Y. and Wu, M. (2015) miRNA-181 regulates embryo implantation in mice through targeting leukemia inhibitory factor. *Journal of Molecular Cell Biology*, **7**, 12–22.
- Chung, T. W., Park, M. J., Lee, H., Kim, K. J., Kim, C. H., Choi, H. J. and Ha, K. T. (2019) Enhancement of endometrial receptivity by *cnidium officinale* through expressing LIF and integrins. *Evidence-Based Complementary and Alternative Medicine*, **2019**, 7560631.
- Crha, I., Ventruba, P., Zakova, J., Jeseta, M., Pilka, R., Lousova, E. and Papikova, Z. (2019) Uterine microbiome and endometrial receptivity. *Ceska Gynekologie*, **84**, 49–54.
- Cui, J., Liu, X., Yang, L., Che, S., Guo, H., Han, J., Zhu, Z., Cao, B., An, X., Zhang, L. and Song, Y. (2020) MiR-184 combined with STC2 promotes endometrial epithelial cell apoptosis in dairy goats via RAS/RAF/MEK/ERK pathway. *Genes*, **11**, 1052.
- Dong, X., Sui, C., Huang, K., Wang, L., Hu, D., Xiong, T., Wang, R. and Zhang, H. (2016) MicroRNA-223-3p suppresses leukemia inhibitory factor expression and pinopodes formation during embryo implantation in mice. *American Journal of Translational Research*, **8**, 1155–1163.
- Dubey, P., Batra, V., Sarwalia, P., Nayak, S., Baithalu, R., Kumar, R. and Datta, T. K. (2023) miR-1246 is implicated as a possible candidate for endometrium remodelling facilitating implantation in buffalo (*Bubalus bubalis*). *Veterinary Medicine and Science*, **9**, 443–456.
- Dvorak, H. F. (2000) VPF/VEGF and the angiogenic response. *Seminars in Perinatology*, **24**, 75–78.
- Eun-Yeong, K., Tae-Wook, C., Hee-Jung, C., Ki-Tae, H., Yeon-Seop, J., Syng-Ook, L., Jun-Yong, C., Hyung, S. K., Sooseong, Y. and Myeong, S. L. (2019) Extracts from *Paeonia lactiflora* Pallas, *Rehmannia Glutinosa* var. *Purpurea* Makino, *Perilla Frutescens* var. *Acuta* Kudo may increase the endometrial receptivity through expression of leukemia inhibitory factor and adhesion molecules. *Journal of Traditional Chinese Medicine*, **39**, 15–25.
- Feng, R., Qin, X., Li, Q., Olugbenga, A. S., Huang, F., Li, Y., Zhao, Q. and Zheng, P. (2022) Progesterone regulates inflammation and receptivity of cells via the NF- κ B and LIF/STAT3 pathways. *Theriogenology*, **186**, 50–59.
- Gebril, M., Hirota, Y., Aikawa, S., Fukui, Y., Kaku, T., Matsuo, M., Hirata, T., Akaeda, S., Hiraoka, T., Shimizu-Hirota, R., Takeda, N., Taha, T., Balah, O. A., Elnoury, M., Fujii, T. and Osuga, Y. (2020) Uterine Epithelial Progesterone Receptor Governs Uterine Receptivity Through Epithelial Cell Differentiation. *Endocrinology*, **161**, bqaa195.
- Goharitaban, S., Abedelahi, A., Hamdi, K., Khazaei, M., Esmacilvand, M. and Niknafs, B. (2022) Role of endometrial microRNAs in repeated implantation failure (mini-review). *Frontiers in Cell and Developmental Biology*, **10**, 936173.
- Governini, L., Luongo, F. P., Haxhiu, A., Piomboni, P. and Luddi, A. (2021) Main actors behind the endometrial receptivity and successful implantation. *Tissue and Cell*, **73**, 101656.
- Griffiths, M., Van Sinderen, M., Rainczuk, K. and Dimitriadis, E. (2019) miR-29c overexpression and COL4A1 downregulation in infertile human endometrium reduces endometrial epithelial cell adhesive capacity in vitro implying roles in receptivity. *Scientific Reports*, **9**, 8644.
- Guan, X., Liu, D., Zhou, H., Dai, C., Wang, T., Fang, Y., Jia, Y. and Li, K. (2022) Melatonin improves pregnancy outcomes in adenomyosis mice by restoring endometrial receptivity via NF- κ B/apoptosis signaling. *Annals of Translational Medicine*, **10**, 1317.
- Guo, X., Yi, H., Li, T. C., Wang, Y., Wang, H. and Chen, X. (2021) Role of vascular endothelial growth factor (VEGF) in human embryo implantation: clinical implications. *Biomolecules*, **11**, 253.
- Hajipour, H., Sambrani, R., Ghorbani, M., Mirzamohammadi, Z. and Nouri, M. (2021) Sildenafil citrate-loaded targeted nanostructured lipid carrier enhances receptivity potential of endometrial cells via LIF and VEGF upregulation. *Naunyn-Schmiedeberg's Archives of Pharmacology*, **394**, 2323–2331.
- He, B., Teng, X. M., Hao, F., Zhao, M., Chen, Z. Q., Li, K. M. and Yan, Q. (2022) Decreased intracellular IL-33 impairs endometrial receptivity in women with adenomyosis. *Frontiers in Endocrinology*, **13**, 928024.
- Hesam, S. M., Seghinsara, A. M., Shokrzadeh, N. and Niknafs, B. (2019) The effect of fludrocortisone on the uterine receptivity partially mediated by ERK1/2-mTOR pathway. *Journal of Cellular Physiology*, **234**, 20098–20110.
- Hu, W., Liang, Y. X., Luo, J. M., Gu, X. W., Chen, Z. C., Fu, T., Zhu, Y. Y., Lin, S., Diao, H. L., Jia, B. and Yang, Z. M. (2019) Nucleolar stress regulation of endometrial receptivity in mouse models and human cell lines. *Cell Death & Disease*, **10**, 831.
- Hua, R., Zhang, X., Li, W., Lian, W., Liu, Q., Gao, D., Wang, Y. and Lei, M. (2020) Ssc-miR-21-5p regulates endometrial epithelial cells proliferation, apoptosis and migration via the PDCD4/AKT pathway. *Journal of Cell Science*, **133**, jcs248898.
- Huang, K., Chen, G., Fan, W. and Hu, L. (2020) miR-23a-3p increases endometrial receptivity via CUL3 during embryo implantation. *Journal of Molecular Endocrinology*, **65**, 35–44.
- Jana, S. K., Banerjee, P., Mukherjee, R., Chakravarty, B. and Chaudhury, K. (2013) HOXA-11 mediated dysregulation of matrix remodeling during implantation window in women with endometriosis. *Journal of Assisted Reproduction and Genetics*, **30**, 1505–1512.
- Kang, Y. J., Lees, M., Matthews, L. C., Kimber, S. J., Forbes, K. and Aplin, J. D. (2015) MiR-145 suppresses embryo-epithelial juxtacrine communication at implantation by modulating maternal IGF1R. *Journal of Cell Science*, **128**, 804–814.
- Kara, M., Ozcan, S. S., Aran, T., Kara, O. and Yilmaz, N. (2019) Evaluation of endometrial receptivity by measuring HOXA-10, HOXA-11, and leukemia inhibitory factor expression in patients with polycystic ovary syndrome. *Gynecology and Minimally Invasive Therapy*, **8**, 118–122.
- Kelleher, A. M., Behura, S. K., Burns, G. W., Young, S. L., DeMayo, F. J. and Spencer, T. E. (2019) Integrative analysis of the forkhead box A2 (FOXA2) cistrome for the human endometrium. *The FASEB Journal*, **33**, 8543–8554.
- Khan, A. W., Farooq, M., Hwang, M. J., Haseeb, M. and Choi, S. (2023) Autoimmune neuroinflammatory diseases: Role of interleukins. *International Journal of Molecular Sciences*, **24**, 7960.
- Kong, C., Sun, L., Zhang, M., Ding, L., Zhang, Q., Cheng, X., Diao, Z., Yan, Q., Zhang, H., Fang, T., Zhen, X., Hu, Y., Sun, H. and Yan, G. (2016) miR-133b reverses the hydrosalpinx-induced impairment of embryo attachment through down-regulation of SGK1. *The Journal of Clinical Endocrinology & Metabolism*, **101**, 1478–1489.
- Kresowik, J. D., Devor, E. J., Van Voorhis, B. J. and Leslie, K. K. (2014) MicroRNA-31 is significantly elevated in both human endometrium and serum during the window of implantation: a potential biomarker for optimum receptivity. *Biology of Reproduction*, **91**, 17.
- Lessey, B. A. and Young, S. L. (2019) What exactly is endometrial receptivity? *Fertility and Sterility*, **111**, 611–617.
- Li J. (2019) Talin1/miR-1285-3p/PIPK1 Axis Regulating Endometrial Adhesion via Influencing PI3K/AKT Signaling Pathway. Doctoral thesis. Guangxi Medical University (In Chinese).
- Li, L., Jiang, H., Wei, X., Geng, D., He, M. and Du, H. (2019) Bu Shen Zhu Yun decoction improves endometrial receptivity via VEGFR-2-mediated angiogenesis. *Evidence-Based Complementary and Alternative Medicine*, **2019**, 1–14.
- Li, Q., Liu, W., Chiu, P. and Yeung, W. (2020) MiR-let-7a/g enhances uterine receptivity via suppressing Wnt/beta-catenin under the modulation of ovarian hormones. *Reproductive Sciences*, **27**, 1164–1174.
- Li, X., Yao, X., Li, K., Guo, J., Deng, K., Liu, Z., Yang, F., Fan, Y., Yang, Y., Zhu, H. and Wang, F. (2023) CREB1 is involved in miR-134-5p-mediated endometrial stromal cell proliferation, apoptosis, and autophagy. *Cells*, **12**, 2554.
- Liang, J., Cao, D., Zhang, X., Liu, L., Tan, Q., Shi, S., Chen, K., Liang, J. and Wang, Z. (2020) miR-192-5p suppresses uterine receptivity formation through impeding epithelial transformation during embryo implantation. *Theriogenology*, **157**, 360–371.
- Liang, J., Li, K., Chen, K., Liang, J., Qin, T., He, J., Shi, S., Tan, Q. and Wang, Z. (2021) Regulation of ARHGAP19 in the endometrial epithelium: a

- possible role in the establishment of uterine receptivity. *Reproductive Biology and Endocrinology*, 19, 2.
- Liang, J., Wang, S. and Wang, Z. (2017) Role of microRNAs in embryo implantation. *Reproductive Biology and Endocrinology*, 15, 90.
- Liang, L., Yang, Y., Yang, L., Zhang, X., Xu, S., Liu, Y., Wu, X. and Chao, L. (2023) HIF-1 α is positively associated with endometrial receptivity by regulating PKM2. *Journal of Obstetrics and Gynaecology Research*, 49, 2734–2745.
- Liang, Y. X., Liu, L., Jin, Z. Y., Liang, X. H., Fu, Y. S., Gu, X. W. and Yang, Z. M. (2018) The high concentration of progesterone is harmful for endometrial receptivity and decidualization. *Scientific Reports*, 8, 712.
- Liang, Y., Shuai, Q., Wang, Y., Jin, S., Feng, Z., Chen, B., Liang, T., Liu, Z., Zhao, H., Chen, Z., Wang, C. and Xie, J. (2021) 1-Nitropyrene exposure impairs embryo implantation through disrupting endometrial receptivity genes expression and producing excessive ROS. *Ecotoxicology and Environmental Safety*, 227, 112939.
- Liu, D. X., Hao, S. L. and Yang, W. X. (2023) Crosstalk between beta-CATENIN-mediated cell adhesion and the WNT signaling pathway. *DNA and Cell Biology*, 42, 1–13.
- Liu, N., Zhou, C., Chen, Y. and Zhao, J. (2013) The involvement of osteopontin and beta3 integrin in implantation and endometrial receptivity in an early mouse pregnancy model. *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 170, 171–176.
- Liu, X., Zhang, L., Liu, Y., Cui, J., Che, S., An, X., Song, Y. and Cao, B. (2018) Circ-8073 regulates CEP55 by sponging miR-449a to promote caprine endometrial epithelial cells proliferation via the PI3K/AKT/mTOR pathway. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research*, 1865, 1130–1147.
- Liu, X., Zhang, L., Yang, L., Cui, J., Che, S., Liu, Y., Han, J., An, X., Cao, B. and Song, Y. (2020) miR-34a/c induce caprine endometrial epithelial cell apoptosis by regulating circ-8073/CEP55 via the RAS/RAF/MEK/ERK and PI3K/AKT/mTOR pathways. *Journal of Cellular Physiology*, 235, 10051–10067.
- Luo, X., Yang, R., Bai, Y., Li, L., Lin, N., Sun, L., Liu, J. and Wu, Z. (2021) Binding of microRNA-135a (miR-135a) to homeobox protein A10 (HOXA10) mRNA in a high-progesterone environment modulates the embryonic implantation factors beta3-integrin (ITGbeta3) and empty spiracles homeobox-2 (EMX2). *Annals of Translational Medicine*, 9, 662.
- Lyu, S., Zhai, Y., Zhu, X., Shi, Q., Chen, F., Zhang, G., Zhang, Z. and Wang, E. (2023) Bta-miR-200b promotes endometrial epithelial cell apoptosis by targeting MYB in cattle. *Theriogenology*, 195, 77–84.
- Ma, Q., Yu, J., Zhang, X., Wu, X. and Deng, G. (2023) Wnt/beta-catenin signaling pathway—a versatile player in apoptosis and autophagy. *Biochimie*, 211, 57–67.
- Makrygiannakis, A., Makrygiannakis, F. and Vrekoussis, T. (2021) Approaches to improve endometrial receptivity in case of repeated implantation failures. *Frontiers in Cell and Developmental Biology*, 9, 613277.
- Michlewski, G. and Caceres, J. F. (2019) Post-transcriptional control of miRNA biogenesis. *RNA*, 25, 1–16.
- Naydenov, M., Nikolova, M., Apostolov, A., Glogovitis, I., Salumets, A., Baev, V. and Yahubyan, G. (2022) The dynamics of miR-449a/c expression during uterine cycles are associated with endometrial development. *Biology*, 12, 55.
- Neykova, K., Tosto, V., Giardina, I., Tsbizova, V. and Vakilov, G. (2022) Endometrial receptivity and pregnancy outcome. *The Journal of Maternal-Fetal & Neonatal Medicine*, 35, 2591–2605.
- Niknafs, B., Shokrzadeh, N., Reza, A. M. and Bakhtiar, H. S. M. (2022) The effect of dexamethasone on uterine receptivity, mediated by the ERK1/2-mTOR pathway, and the implantation window: An experimental study. *International Journal of Reproductive Biomedicine*, 20, 47–58.
- Park, H. R., Choi, H. J., Kim, B. S., Chung, T. W., Kim, K. J., Joo, J. K., Ryu, D., Bae, S. J. and Ha, K. T. (2021) Paeoniflorin enhances endometrial receptivity through leukemia inhibitory factor. *Biomolecules*, 11, 439.
- Paule, S. G., Heng, S., Samarajeewa, N., Li, Y., Mansilla, M., Webb, A. I., Nebl, T., Young, S. L., Lessey, B. A., Hull, M. L., Scelwyn, M., Lim, R., Vollenhoven, B., Rombauts, L. J. and Nie, G. (2021). Podocalyxin is a key negative regulator of human endometrial epithelial receptivity for embryo implantation. *Human Reproduction*, 36, 1353–1366.
- Paulson, E. E. and Comizzoli, P. (2021) Endometrial receptivity and embryo implantation in carnivores—commonalities and differences with other mammalian species. *Biology of Reproduction*, 104, 771–783.
- Pirtea, P., Cicinelli, E., De Nola, R., de Ziegler, D. and Ayoubi, J. M. (2021) Endometrial causes of recurrent pregnancy losses: Endometriosis, adenomyosis, and chronic endometritis. *Fertility and Sterility*, 115, 546–560.
- Qi, Y., Wang, X., Hou, S., Wu, Z., Xu, X. and Pang, C. (2022) Intracavitary physiotherapy combined with acupuncture mediated AMPK/mTOR signaling to improve endometrial receptivity in patients with thin endometrium. *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 277, 32–41.
- Rehman, R., Jawed, S., Zaidi, S. F., Baig, M. and Ahmeds, K. (2015) Role of interleukin-13 in conception after intracytoplasmic sperm injection. *Journal of the Pakistan Medical Association*, 65, 49–53.
- Sanchez-Lopez, J. A., Caballero, I., Montazeri, M., Maslehat, N., Elliott, S., Fernandez-Gonzalez, R., Calle, A., Gutierrez-Adan, A. and Fazeli, A. (2014) Local activation of uterine Toll-like receptor 2 and 2/6 decreases embryo implantation and affects uterine receptivity in mice. *Biology of Reproduction*, 90, 87.
- Shan, L., Zhou, Y., Peng, S., Wang, X., Shan, Z. and Teng, W. (2019) Implantation failure in rats with subclinical hypothyroidism is associated with LIF/STAT3 signaling. *Endocrine Connections*, 8, 718–727.
- Shariati, M., Niknafs, B., Seghinsara, A. M., Shokrzadeh, N. and Alivand, M. R. (2019) Administration of dexamethasone disrupts endometrial receptivity by alteration of expression of miRNA-223, 200a, LIF, Muc1, SGK1, and ENaC via the ERK1/2-mTOR pathway. *Journal of Cellular Physiology*, 234, 19629–19639.
- Shekibi, M., Heng, S. and Nie, G. (2022) MicroRNAs in the regulation of endometrial receptivity for embryo implantation. *International Journal of Molecular Sciences*, 23, 6210.
- Shekibi, M., Heng, S., Wang, Y., Samarajeewa, N., Rombauts, L. and Nie, G. (2022) Progesterone suppresses podocalyxin partly by up-regulating miR-145 and miR-199 in human endometrial epithelial cells to enhance receptivity in in vitro models. *Molecular Human Reproduction*, 28, gaac034.
- Shen, M., Liu, Y., Ma, X. and Zhu, Q. (2023) Erbu Zhuyu decoction improves endometrial angiogenesis via uterine natural killer cells and the PI3K/Akt/eNOS pathway a mouse model of embryo implantation dysfunction. *American Journal of Reproductive Immunology*, 89, e13634.
- Shi, S., Tan, Q., Liang, J., Cao, D., Wang, S., Liang, J., Chen, K. and Wang, Z. (2021) Placental trophoblast cell-derived exosomal microRNA-1290 promotes the interaction between endometrium and embryo by targeting LHX6. *Molecular Therapy-Nucleic Acids*, 26, 760–772.
- Shokrzadeh, N., Alivand, M. R., Abedelahi, A., Hessam, S. M. and Niknafs, B. (2018) Upregulation of HB-EGF, Msx.1, and miRNA Let-7a by administration of calcitonin through mTOR and ERK1/2 pathways during a window of implantation in mice. *Molecular Reproduction and Development*, 85, 790–801.
- Shokrzadeh, N., Alivand, M. R., Abedelahi, A., Hessam, S. M. and Niknafs, B. (2019) Calcitonin administration improves endometrial receptivity via regulation of LIF, Muc-1 and microRNA Let-7a in mice. *Journal of Cellular Physiology*, 234, 12989–13000.
- Smith, J., Zyou, A. and Allegrucci, C. (2019) A case of identity: HOX genes in normal and cancer stem cells. *Cancers*, 11, 512.
- Soni, U. K., Chadchan, S. B., Gupta, R. K., Kumar, V. and Kumar, J. R. (2021) miRNA-149 targets PARP-2 in endometrial epithelial and stromal cells to regulate the trophoblast attachment process. *Molecular Human Reproduction*, 27, gaab039.
- Stavast, C. J. and Erkeland, S. J. (2019) The non-canonical aspects of microRNAs: many roads to gene regulation. *Cells*, 8, 1465.
- Steens, J. and Klein, D. (2022) HOX genes in stem cells: Maintaining cellular identity and regulation of differentiation. *Frontiers in Cell and Developmental Biology*, 10, 1002909.
- Sutaji, Z., Abu, M. A., Sayutti, N., Elias, M. H., Ahmad, M. F., Nur, A. A., Chew, K. T., Abdul, K. A., Aziz, N., Mokhtar, M. H., Zin, R. and Hussein, T. (2023) Endometrial heparin-binding epidermal growth factor gene expression and hormone level changes in implantation window of obese women with polycystic ovarian syndrome. *Biomedicine*, 11, 276.

- Verma, R. K., Soni, U. K., Chadchan, S. B., Maurya, V. K., Soni, M., Sarkar, S., Pratap, J. V. and Jha, R. K. (2022) miR-149-PARP-2 signaling regulates E-cadherin and N-cadherin expression in the murine model of endometrium receptivity. *Reproductive Sciences*, **29**, 975–992.
- Wang, H., Shi, G., Li, M., Fan, H., Ma, H. and Sheng, L. (2018) Correlation of IL-1 and HB-EGF with endometrial receptivity. *Experimental and Therapeutic Medicine*, **16**, 5130–5136.
- Wang, J., Huang, C., Jiang, R., Du Y., Zhou, J., Jiang, Y., Yan, Q., Xing, J., Hou, X., Zhou, J., Sun, H. and Yan, G. (2018) Decreased endometrial IL-10 impairs endometrial receptivity by downregulating HOXA10 expression in women with adenomyosis. *BioMed Research International*, **2018**, 2549789.
- Wang, J., Xie, D., Liu, M., Gong, Y., Shi, X., Wei, J. Y. and Quan, S. (2016) Uterine macrophages affect embryo implantation via regulating vascular endothelial growth factor A in mice. *Journal of Southern Medical University*, **36**, 909–914.
- Wang, L., Liu, D., Wei, J., Yuan, L., Zhao, S., Huang, Y., Ma, J. and Yang, Z. (2021) MiR-543 inhibits the migration and epithelial-to-mesenchymal transition of TGF- β -treated endometrial stromal cells via the MAPK and Wnt/ β -catenin signaling pathways. *Pathology and Oncology Research*, **27**, 1609761.
- Wang, W., Ge, L., Zhang, L. L., Wang, L. R., Lu, Y. Y., Gou, L., Gou, R. Q., Xu, T. Y., Ma, X. L. and Zhang, X. H. (2023) Mechanism of human chorionic gonadotropin in endometrial receptivity via the miR-126-3p/PI3K/Akt/eNOS axis. *The Kaohsiung Journal of Medical Sciences*, **39**, 468–477.
- Wei, P., Wang, H., Li, Y. and Guo, R. (2021) Nucleolar small molecule RNA SNORA75 promotes endometrial receptivity by regulating the function of miR-146a-3p and ZNF23. *Aging*, **13**, 14924–14939.
- Xing, L., Chen, Y., He, Z., He, M., Sun, Y., Xu, J., Wang, J., Zhuang, H., Ren, Z., Chen, Y., Yang, J., Cheng, S. and Zhao, R. (2022) Acupuncture improves endometrial angiogenesis by activating PI3K/AKT pathway in a rat model with PCOS. *Evidence-Based Complementary and Alternative Medicine*, **2022**, 1–15.
- Xu, B., Geerts, D., Bu, Z., Ai, J., Jin, L., Li, Y., Zhang, H. and Zhu, G. (2014) Regulation of endometrial receptivity by the highly expressed HOXA9, HOXA11 and HOXD10 HOX-class homeobox genes. *Human Reproduction*, **29**, 781–790.
- Yang C. (2021) The Study on miR-20b Targeting NCOA3 to Regulate Progesterone-Induced Immune Response in Bovine Endometrium. Master's thesis. Huazhong Agricultural University (In Chinese).
- Yang, D., Liu, A., Wu, Y., Li, B., Nan, S., Yin, R., Zhu, H., Chen, J., Ding, Y. and Ding, M. (2020) BCL2L15 depletion inhibits endometrial receptivity via the STAT1 signaling pathway. *Genes*, **11**, 816.
- Yang, P., Wu, Z., Ma, C., Pan, N., Wang, Y. and Yan, L. (2019) Endometrial miR-543 is downregulated during the implantation window in women with endometriosis-related infertility. *Reproductive Sciences*, **26**, 900–908.
- Yang, Y., Sun, Y., Cheng, L., Li, A., Shen, Y., Jiang, L., Deng, X. and Chao, L. (2017) GRIM-19, a gene associated with retinoid-interferon-induced mortality, affects endometrial receptivity and embryo implantation. *Reproduction, Fertility and Development*, **29**, 1447–1455.
- Yao, K., Kang, Q., Chen, K., Shi, B. and Jin, X. (2024) MiR-124-3p negatively impacts embryo implantation via suppressing uterine receptivity formation and embryo development. *Reproductive Biology and Endocrinology*, **22**, 16.
- You, F., Du X., Zhang, T., Wang, Y., Lv, Y. and Zeng, L. (2021) High-frequency electroacupuncture improves endometrial receptivity via regulating cell adhesion molecules and leukemia inhibitory factor/signal transducer and activator of transcription signaling pathway. *Bioengineered*, **12**, 10470–10479.
- You, F., Du X., Zhang, T., Wang, Y., Lv, Y. and Zeng, L. (2022a) TJZYF improves endometrial receptivity through regulating VEGF and PI3K/AKT signaling pathway. *BioMed Research International*, **2022**, 9212561.
- You, F., Du X., Zhang, T., Wang, Y., Lv, Y. and Zeng, L. (2022b) Electroacupuncture improves endometrial receptivity through miRNA-223-3p-mediated regulation of leukemia inhibitory factor/signal transducer and activator of transcription 3 signaling pathway. *Bioengineered*, **13**, 10298–10312.
- Yu, S. L., Jeong, D. U., Noh, E. J., Jeon, H. J., Lee, D. C., Kang, M., Kim, T. H., Lee, S. K., Han, A. R., Kang, J. and Park, S. R. (2023) Exosomal miR-205-5p improves endometrial receptivity by upregulating E-cadherin expression through ZEB1 inhibition. *International Journal of Molecular Sciences*, **24**, 15149.
- Yu, S., Kang, Y., Jeong, D., Lee, D. C., Jeon, H. J., Kim, T., Lee, S. K., Han, A. R., Kang, J. and Park, S. (2022) The miR-182-5p/NDRG1 axis controls endometrial receptivity through the NF- κ B/ZEB1/E-cadherin pathway. *International Journal of Molecular Sciences*, **23**, 12303.
- Yu, Y., Cao, Y., Huang, W., Liu, Y., Lu, Y. and Zhao, J. (2021) β -Sitosterol ameliorates endometrium receptivity in PCOS-like mice: The mediation of gut microbiota. *Frontiers in Nutrition*, **8**, 667130.
- Yuan, L., Feng, F., Mao, Z., Huang, J. Z., Liu, Y., Li, Y. L. and Jiang, R. X. (2021) Regulation mechanism of miR-494-3p on endometrial receptivity in mice via PI3K/AKT/mTOR pathway. *General Physiology & Biophysics*, **40**, 351–363.
- Zarei, R., Nikpour, P., Rashidi, B., Eskandari, N. and Aboutorabi, R. (2019) Evaluation of diabetes effects on the expression of leukemia inhibitory factor and vascular endothelial growth factor A genes and proteins at the time of endometrial receptivity after superovulation in rat model. *Advanced Biomedical Research*, **8**, 66.
- Zarrin, Y., Bakhteyari, A., Nikpour, P., Mostafavi, F. S., Eskandari, N., Matinfar, M. and Aboutorabi, R. (2020) A study on the presence of osteopontin and alpha3beta1 integrin in the endometrium of diabetic rats at the time of embryo implantation. *Journal of Reproduction & Infertility*, **21**, 87–93.
- Zhang, L., Liu, X. R., Liu, J. Z., Song, Y. X., Zhou, Z. Q. and Cao, B. Y. (2017) miR-182 selectively targets HOXA10 in goat endometrial epithelium cells in vitro. *Reproduction in Domestic Animals*, **52**, 1081–1092.
- Zhang, L., Liu, X., Cui, J., Che, S., Liu, Y., An, X., Cao, B. and Song, Y. (2019) LncRNA882 regulates leukemia inhibitory factor (LIF) by sponging miR-15b in the endometrial epithelium cells of dairy goat. *Journal of Cellular Physiology*, **234**, 4754–4767.
- Zhang, L., Liu, X., Liu, J., Ma, X., Zhou, Z., Song, Y. and Cao, B. (2018) miR-26a promoted endometrial epithelium cells (EECs) proliferation and induced stromal cells (ESCs) apoptosis via the PTEN-PI3K/AKT pathway in dairy goats. *Journal of Cellular Physiology*, **233**, 4688–4706.
- Zhang, L., Liu, X., Liu, J., Zhou, Z., Song, Y., Cao, B. and An, X. (2017) miR-182 aids in receptive endometrium development in dairy goats by down-regulating PTN expression. *Plos One*, **12**, e179783.
- Zhang, Y. M., Zhang, Y. Y., Bulbul, A., Shan, X., Wang, X. Q. and Yan, Q. (2015) Baicalin promotes embryo adhesion and implantation by upregulating fucosyltransferase IV (FUT4) via Wnt/beta-catenin signaling pathway. *FEBS Letters*, **589**, 1225–1233.
- Zhang, Z., Li, T., Xu, L., Wang, Q., Li, H. and Wang, X. (2021) Extracellular superoxide produced by *Enterococcus faecalis* reduces endometrial receptivity via inflammatory injury. *American Journal of Reproductive Immunology*, **86**, e13453.
- Zhao, D. M., Shan, Y. H., Li, F. H., Jiang, L. and Qu, Q. L. (2019) Correlation between endometrial receptivity with expressions of IL-1 and VEGF in rats with polycystic ovary syndrome. *European Review for Medical & Pharmacological Sciences*, **23**, 5575–5580.
- Zheng, P., Qin, X., Feng, R., Li, Q., Huang, F., Li, Y., Zhao, Q. and Huang, H. (2022) Alleviative effect of melatonin on the decrease of uterine receptivity caused by blood ammonia through ROS/NF- κ B pathway in dairy cow. *Ecotoxicology and Environmental Safety*, **231**, 113166.
- Zheng, Q., Zhang, D., Yang, Y. U., Cui, X., Sun, J., Liang, C., Qin, H., Yang, X., Liu, S. and Yan, Q. (2017) MicroRNA-200c impairs uterine receptivity formation by targeting FUT4 and alpha1,3-fucosylation. *Cell Death & Differentiation*, **24**, 2161–2172.