



# *Weeds*

VOLUME 7  
JANUARY 1959  
NUMBER 1

*Journal of the Weed Society of America*

# Weeds

*Issued Quarterly by the Weed Society of America*

K. P. BUCHHOLTZ, *Editor*, Dept. of Agronomy, Univ. of Wisconsin,  
Madison, Wisconsin.

W. C. JACOB, *Business Manager*, Dept. of Agronomy, Univ. of  
Illinois, Urbana, Illinois.

## EDITORIAL COMMITTEE

O. C. LEE, Dept. of Botany and Plant Pathology, Purdue Univ.,  
Lafayette, Indiana.

R. A. PETERS, Dept. of Plant Science, Univ. of Connecticut, Storrs,  
Connecticut.

W. C. ROBOCKER, Crops Research Division, ARS, USDA, State  
College of Washington, Pullman, Washington.

E. G. RODGERS, Dept. of Agronomy, Univ. of Florida, Gainesville,  
Florida.

WEEDS is a quarterly journal published by the Weed Society of America. Editorial offices are located at the University of Wisconsin, Madison, Wisconsin. Printing is by the W. F. Humphrey Press Inc., Geneva, New York. Subscription price is \$6.00 yearly for four issues: single copies \$1.50. Address all communications regarding subscriptions, advertising and reprints to W. C. Jacob, Department of Agronomy, University of Illinois, Urbana, Illinois. Inquiries concerning information on manuscripts and other material for publication should be addressed to the Editorial Offices. All checks, money orders and other remittances should be made payable to the Weed Society of America.

Entered as second-class matter at the post office at Urbana,  
Illinois with additional entry at Geneva, New York.

## Table of Contents

|                                                                                                                                                                                                                   | <i>Page</i> |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|
| The Effects of Quackgrass on Germination and Seedling Development of Certain Crop Plants. Thor Kommedahl, J. B. Kotheimer and J. V. Bernardini. . . . .                                                           | 1           |
| The Seasonal Development of Johnson Grass Plants. Edwin B. Oyer, G. A. Gries, and B. J. Rogers. . . . .                                                                                                           | 13          |
| A Study of the Absorption and Translocation of Several Chemicals in Johnson Grass, and an Evaluation of Their Effectiveness for its Control under Field Conditions. Ellis W. Hauser and Jack T. Thompson. . . . . | 20          |
| A Method for Inducing Dormancy in Wild Oats ( <i>Avena fatua</i> L.). J. R. Hay and B. G. Cumming. . . . .                                                                                                        | 34          |
| Methods of Obtaining Field and Laboratory Germination of Seeds of Bindweeds, Lady's Thumb and Velvet Leaf. G. P. Steinbauer and Buford Grigsby. . . . .                                                           | 41          |
| Neburon as a Selective Herbicide for Tartary Buckwheat and Wild Buckwheat. H. A. Friesen and D. E. Forsberg. . . . .                                                                                              | 47          |
| Effects of Light Quality on Herbicide Toxicity to Plants. Subhash C. Datta and Stuart Dunn. . . . .                                                                                                               | 55          |
| A Method of Comparing Herbicides and Assessing Herbicide Mixtures at the Screening Level. Donald P. Gowing. . . . .                                                                                               | 66          |
| Some Effects of Various Esters of 2,4-Dichlorophenoxyacetic Acid on the Morphological Development of the Strawberry. Ronald W. Campbell. . . . .                                                                  | 77          |
| Chemical Weed Control in Garden Chrysanthemums. R. J. Stadtherr and R. E. Widmer. . . . .                                                                                                                         | 82          |
| The Relationship of Soil Adsorption of EPTC to Oats Injury in Various Soil Types. Floyd M. Ashton and Thomas J. Sheets. . . . .                                                                                   | 88          |
| A Logarithmic-Concentration Sprayer for Small Plot Use. J. K. Leasure. . . . .                                                                                                                                    | 91          |
| A Visual Aid for Illustrating Herbicidal Chemical Structures and Teaching Chemical Nomenclature. W. A. Gentner, W. C. Shaw, and L. L. Jansen. . . . .                                                             | 98          |
| Keeping Extension Agents and Dealers Informed on Weed Control. Paul W. Santelmann and Andrew A. Duncan. . . . .                                                                                                   | 104         |
| Regulating the Use of 2,4-D in the Southern States. Ernest A. Epps, Jr. . . . .                                                                                                                                   | 108         |
| News and Notes. . . . .                                                                                                                                                                                           | 111         |
| Sustaining Members. . . . .                                                                                                                                                                                       | 112         |
| Bibliography of Weed Investigations, April to June, 1958. . . . .                                                                                                                                                 | 113         |


## Advertisers Index

|                                     |     |
|-------------------------------------|-----|
| Spraying Systems Co. . . . .        | ii  |
| Union Carbide Chemicals Co. . . . . | iii |
| duPont de Nemours & Co. . . . .     | iv  |
| U. S. Borax & Chemical Co. . . . .  | v   |
| Stauffer Chemical Co. . . . .       | vi  |
| Chipman Chemical Co. . . . .        | vii |




the precision nozzle for effective spraying


Supplied in a full range of interchangeable orifice tip and strainer sizes to meet every capacity requirement. TeeJet Spray Nozzles for Weed Control by spraying make it possible to take maximum advantage of the chemical and sprayer unit. TeeJet nozzles are precision built and provide a flat spray with uniform distribution. Atomization is properly controlled to give coverage with an absolute minimum of driftage. Patented tip design, with set-back orifice opening protects precision orifice from accidental damage. TeeJet spray nozzles are built for use on spray booms and portable sprayers.



TEEJET SPRAY NOZZLE male pipe connection



TEEJET SPRAY NOZZLE female pipe connection



INTERCHANGEABLE ORIFICE TIPS flat and cone spray types

**OFF-CENTER SPRAY NOZZLES**

Spraying Systems Spray Nozzles with TeeJet tips are supplied in a variety of special body types to meet any unusual spraying requirement. For example, one type of off-center spray nozzle with swivel body provides a flat spray up to 35 feet wide for spraying areas with a single nozzle, that are not accessible with a boom.

**SUPPLEMENTARY EQUIPMENT**

Complete accessories relating to nozzle use are supplied. These include strainers, special nozzle fittings, and hand valve equipment.

TeeJet Spray Nozzles are supplied for Weed Control... as well as all other types of agricultural spraying. For complete information and reference data write for Catalog 30.

**SPRAYING SYSTEMS CO.**  
*Engineers and Manufacturers*

3296 RANDOLPH STREET

BELLWOOD, ILLINOIS

**Crag** Mylone  
BRAND SOIL FUMIGANT

HELPS

FOREST TREE SEEDLINGS

GET OFF TO A  
BETTER START



In numerous tests throughout the country, formulations with Crag Mylone soil fumigant have given excellent control of weeds, nematodes, and soil fungi in forest tree seed beds. Seedlings have a better chance to grow without interference from these pests.

When formulated, Mylone is an easy-to-handle powder that can be applied to the soil with a fertilizer spreader or as a drench. Application should be made at least three weeks before seeds are planted, except tobacco seed. Soil mixing is not necessary; no plastic cover is needed over the bed.

Formulations of Crag Mylone are now commercially available for use in certain ornamental propagating beds, tobacco seed beds, and in tomato, pepper, egg plant, lettuce, and cabbage seed beds. It is also sold for weed and dry rot control in gladiolus in Florida. Experimental work is continuing with other crops.

Write to the address below for formulations in test quantities or more information. Names and addresses of commercial formulators are also available.

"Crag", "Mylone", and "Union Carbide" are trade marks of Union Carbide Corporation.



**FORMULATIONS CONTAINING  
MYLONE NOW AVAILABLE  
FOR VEGETABLE SEED BEDS**

A commercial label has been accepted for Mylone pre-planting use in seed beds of certain vegetables. Growers can now use it for the control of weeds, nematodes, and soil fungi in tomato, pepper, cabbage, egg plant, and lettuce seed beds.



CRAG Agricultural Chemicals

**Union Carbide Chemicals Company**

Division of Union Carbide Corporation

180 South Broadway, White Plains, New York

**For Agriculture and Industry . . .**

# **Du Pont WEED KILLERS**

*offer new economies  
and efficiency in killing  
weeds, grass and brush*

**KARMEX®** diuron for weed control in many crops; irrigation and drainage ditches; and industrial areas.

**KARMEX® DL** for pre-emergence weed control in cotton.

**TELVAR®** monuron for weed control in many crops; irrigation and drainage ditches; and industrial areas.

**KLOBEN** neburon for weed and grass control in nursery plantings and tomatoes.

**DYBAR** fenuron—a pelleted product for dry application to the soil for control of brush in fence rows, drainage ditches, utility and railroad right-of-ways, and other non-cultivated areas.

**TRYBEN 200**, a new weed killer based on trichlorobenzoic acid, for control of bindweed and other noxious weeds; mixed broad-leaf weeds; and certain woody vines and brush.

**AMMATE® X** for long-term, low-cost brush control.



Better Things for Better Living . . . . *Through Chemistry*

Four  
easy  
ways to  
Destroy  
Weeds

## 1. UREABOR®

A nonselective, granular complex of sodium borate and substituted urea. Low application rates are a feature. Apply with the special new PCB Spreader for best results.

## 2. DB® Granular

A combination of 2,4-D and sodium borates. Kills deep-rooted, noxious weeds. Low application rates for maximum control with the utmost economy; use the PCB Spreader.

*(Not intended for control of grass.)*

## 3. POLYBOR-CHLORATE®

Highly soluble; for spray or dry application. It gives a quick knock-down; destroys top growth and roots. A general nonselective herbicide.

## 4. Concentrated BORASCU®

A nonselective, granular material. Apply by hand or with a mechanical spreader. Long residual action.

When you want Nonselective Herbicides

for Dependable Action

look to United States Borax  
& Chemical Corporation

PACIFIC COAST BORAX COMPANY DIVISION



630 Shatto Place, Los Angeles 5, Calif.



## Now Eptam® offers the farmer a truly selective herbicide


Eptam (EPTC) not only gives excellent control of most grassy weeds, but it is also effective against nutgrass. This highly selective herbicide controls many broadleaf weeds as well, including pigweed and purslane. Eptam is now registered for use on field and sweet corn, beets, flax, snap beans, forage legumes, strawberries and ornamentals. It can be applied at any time before the weeds emerge. For additional information, write us at 380 Madison Avenue, New York 17.



New York • San Francisco • Houston • Omaha • Los Angeles • Tampa  
North Portland • Weslaco • Lubbock • Harvey • North Little Rock

©Eptam is Stauffer Chemical Company's trade-mark (registered in principal countries) for *ethyl-di-n propylthiocarbamate*, a selective herbicide.





**ATLACIDE:** Safer chlorate weed killer...widely used for non-selective eradication of bindweed, Canada thistle, quack grass, Johnson grass and other tough perennials. Kills roots...discourages regrowth. Applied dry or dissolved in water for use as a spray.

**ATLACIDE-2,4-D:** A combination of Atlacide and 2,4-D acid. Offers dual killing action of chlorate and 2,4-D.

**CHLOREA:** A non-separating combination of sodium chlorate, borate and monuron in powder form. Kills weeds and grasses. Combines the proven effectiveness of chlorate on deep-rooted weeds with the soil-surface action of monuron on shallow-rooted grasses and annual seedling growth. Lasting residual effect inhibits new growth. Does not create a fire hazard when used as directed. Applied dry or as a water-mixed spray. For industrial, railroad and certain agricultural uses.

**CHLOREA GRANULAR** Similar to Chlorea, but a granular material. No mixing or diluting...“pellets” are easy to apply by hand or with mechanical spreader.

**CHLORAX “40”:** A composition

of sodium chlorate and borate...for weed and grass control. Has lasting residual effect. Does not create a fire hazard. Applied dry or as a spray.

**CHLORAX LIQUID:** Similar to Chlorax “40”...in liquid form.

**ATLAS “A”:** A 40% sodium arsenite solution (4 lbs. arsenic trioxide per gal.). Destroys submersed vegetation and algae in ponds and lakes. Controls crabgrass, chickweed and clover in turf. Used as general weed killer and to kill trees and stumps. Also used to kill potato vines prior to harvesting.

**SODIUM ARSENITE:** A powder containing 75% arsenic trioxide. Used for the same purposes as Atlas “A”. Applied dry or as a spray.

**2,4-D & 2,4,5-T WEED KILLERS:** A complete line...available as 2,4-D Amine and 2,4-D Ester liquids; 2,4-D Ester dusts; Low Volatile 2,4,5-T and Brush Killer.

**METHOXONE:** Contains 2 pounds of MCP sodium salt per gallon. Used for weed control in small grains, flax, rice and grass. Controls same weeds as 2,4-D; considered safer for selective spraying.

*Write for Weed Control Booklets*

**CHIPMAN CHEMICAL COMPANY, INC.**

Chicago, Ill.

Bound Brook, N. J.

Portland, Ore.

Palo Alto, Calif.

Pasadena, Tex.

Bessemer, Ala.

*Manufacturers of Weed Killers Since 1912*