

The 14 μm Band of Carbon Stars¹

I. Yamamura, T. de Jong², L.B.F.M. Waters³, J. Cami³

*Astronomical Institute 'Anton Pannekoek', University of Amsterdam,
Kruislaan 403, 1098 SJ, The Netherlands*

K. Justtanont

Stockholm Observatory, 13336 Saltsjöbaden, Sweden

Abstract. We have studied the absorption bands around 14 μm in the spectra of 11 carbon stars with mass-loss rates ranging from 10^{-8} to $10^{-4} M_{\odot} \text{yr}^{-1}$, based on data obtained with the Short Wavelength Spectrometer (SWS) on board the Infrared Space Observatory (ISO). All stars clearly show a C_2H_2 absorption band peaking at 13.7 μm , while the contribution from HCN molecules is small in this wavelength region. A simple plane-parallel LTE model with two layers at different temperatures is used to derive the C_2H_2 abundances in the outer photosphere and in the circumstellar envelope. We find that (1) the column density of the hot-layer, placed at about $3R_{\star}$ with a temperature of 1400 K is roughly the same for all stars regardless of the mass-loss rate, and (2) the contribution of cool molecules in the circumstellar envelope increases with the dust mass-loss rate, (3) the abundance of C_2H_2 in the two layers is about the same, i.e. no obvious depletion of C_2H_2 molecules seems to occur in the circumstellar envelope.

1. Introduction

Acetylene (C_2H_2) is the most abundant molecule in the atmosphere of carbon stars after CO and HCN (Tsuji 1964). The molecule probably plays an essential role in the atmosphere of these stars, not only in the energy budget through their ro-vibrational transitions but also as a seed molecule for dust formation. C_2H_2 has been little studied, however, since the linear and symmetric structure of C_2H_2 prevents us to observe the molecule directly at radio frequencies. The abundance of the molecule has been estimated from its photodissociation products such as C_2H (Keady & Hinkle 1988; Fuente et al. 1998). The ro-vibrational

¹Based on observations with ISO, an ESA project with instruments funded by ESA Member States (especially the PI countries: France, Germany, the Netherlands and the United Kingdom) with the participation of ISAS and NASA. The SWS is a joint project of SRON and MPE.

²also SRON-Utrecht, Sorbonnelaan 2, 3584 CA, Utrecht, The Netherlands

³also SRON-Groningen, Landleven 12, 9700 AV, Groningen, The Netherlands

transitions in the infrared are also not easy to measure. The $3.05\ \mu\text{m}$ absorption band is accessible from the ground, but very high resolution is required in order to avoid confusion with other molecules such as HCN. The band centers of other major transitions at 7.5 and $13.7\ \mu\text{m}$ are inaccessible from the ground, and have only been partially observed (Keady & Ridgway 1993).

The ISO/SWS (Kessler et al. 1996; de Graauw et al. 1996) provides infrared spectra in a broad wavelength range with moderately high resolution. As a part of the ISO guaranteed time program AGBSTAR, we observed 11 carbon stars with various mass-loss rates ranging from 10^{-8} and $10^{-4}\ M_{\odot}\ \text{yr}^{-1}$. Based on the infrared spectra of these stars we concluded that not only the energy distribution and the broad dust features but also several narrow molecular features drastically change with the mass-loss rate (Yamamura et al. 1998). In this article, we focus on the C_2H_2 band at $13.7\ \mu\text{m}$, and discuss the C_2H_2 distribution around the stars. This band has a sharp Q-branch, which makes it easy to identify and distinguish from other molecular features.

2. Observations and results

The observations were carried out using the SWS AOT01 full-grating scan mode. All stars were observed with a resolution of about 300, except IRC +10216 which was observed at a resolution of about 1000. The data were reduced by the SWS Interactive Analysis package developed by the SWS instrument dedicated team, using the most recent set of calibration parameters (as of April 1998).

The carbon star sample is divided into four groups (II–V) based on their near-infrared color temperature, T_{NIR} , which is supposed to be a good indicator of mass-loss rate (Groenewegen et al. 1992). The spectra of our carbon stars between 13 and $15\ \mu\text{m}$ are shown in Fig. 1, labeled with their name and group number. The spectra are normalized to the *local continuum*. The spectra are also shifted slightly with respect to the model spectra in order to correct for the ambiguity in the absolute continuum level. The uncertainty in the continuum determination will be discussed in Sect. 4.1.

The C_2H_2 band at $13.7\ \mu\text{m}$ is clearly detected in *absorption* in all stars, regardless of the mass-loss rate. The band is actually a mixture of many Q-branches including the ν_5 fundamental transition. The band is broader in stars with smaller mass-loss rate, indicating that the absorption is due to hot gas near the photosphere. On the other hand, the HCN band at $14.0\ \mu\text{m}$ is hardly recognized in any of the spectra in Fig. 1. Indeed, the feature is much weaker than expected from the far-infrared spectrum of IRC +10216 (Cernicharo et al. 1996) or from the $3.6\ \mu\text{m}$ feature in the same SWS spectra. This may be explained in terms of the distribution of the molecule as well as the excitation mechanism, but it is a subject for future studies.

The spectra are in contrast with those of oxygen-rich stars, which exhibit CO_2 emission bands in the same wavelength range (Justtanont et al. 1998; Cami et al. 1997). This indicates that there is always a continuum source behind the C_2H_2 molecule.




Figure 1. The 14 μm band of carbon stars (*solid line*) is shown with the model spectra (*gray line*). The spectra are ordered by decreasing near-infrared color temperature (T_{NIR}) from bottom to top. The group number of each object is indicated after the object name. The spectra are normalized to the *local continuum*. In addition, they are shifted by at most 7% when compared with the models, in order to correct for the uncertainty in the absolute continuum levels. The fit is not so good for group II stars especially between 13.8 and 14.0 μm due to insufficient molecular line data.

3. Modeling

Quantitative analysis of the spectra is attempted with a simple plane-parallel model. In order to simplify the problem as much as possible, we try to explain the observed spectra by a combination of two components: a *hot-component* representing the photosphere and a *cool-component* corresponding to the circumstellar envelope. Each layer is characterized by three parameters, the column density (N), the excitation temperature (T), and the turbulent velocity (v_{turb}). In the present calculation, v_{turb} is fixed to 3 km s^{-1} in both layers. The results are insensitive to the value of v_{turb} . For instance, for $v_{\text{turb}} = 10 \text{ km s}^{-1}$ the column density would only decrease by 0.1 dex. Only C_2H_2 is considered. The line data are based on Kabbadj et al. (1991), Rinsland et al. (1982), Palmer et al. (1972), and Weber et al. (1994). About 20 000 lines are included in the wavelength region in Fig. 1. From a comparison with laboratory spectra (Kawaguchi, private communication), it is known that the line data are incomplete, especially between 13.8 and 14.0 μm . We ignored that part in the fitting.

First, the temperatures of the two layers are determined. We assume that group II and group V stars represent the hot- and cool-component, respectively, and we fit their spectra by a single layer model. For every combination in the parameter space of $10^{15} \leq N \leq 10^{19} \text{ cm}^{-2}$ with 0.1 dex step and $300 \leq T \leq 1800 \text{ K}$ with 50 K interval, the model spectrum is characterized by the *rms* of the residual from the observed spectrum, and the best fit is the one which minimizes the *rms*. We obtain $T_{\text{hot}} = 1400 \text{ K}$ and $T_{\text{cool}} = 450 \text{ K}$. In the second step, we fix the temperatures at these values and fit all stars with a two-layers model by the same procedure in ($N_{\text{hot}}, N_{\text{cool}}$) space. The best fit spectra are shown in Fig. 1 by gray lines, and the derived column densities for the spectra are plotted in Fig. 2 as a function of T_{NIR} .

The emission from the dust envelope fills in and weakens the absorption in the photosphere. The correction of this effect is attempted as follows: A model spectrum with N_{hot} and T_{hot} is calculated. The spectrum is then inserted in a spherical dust envelope model with a certain mass-loss rate, and a synthesized spectrum is calculated. A star with a blackbody temperature of 2500 K and radius $3 \times 10^{13} \text{ cm}$ is used when the dust temperature distribution is calculated. The envelope is assumed to extend from $10 R_{\star}$ to $10^4 R_{\star}$. The weakened molecular feature in the output spectrum may have an intensity similar to that with a lower column density and no dust effect. The difference between the original and weakened spectrum in terms of the column density is the correction factor corresponding to the dust mass-loss rate. The dust mass-loss rate of each star is determined by adopting the same model to their SWS spectra. The corrected values of N_{hot} are also plotted in Fig. 2.

4. Discussion

4.1. Uncertainty in the result

C_2H_2 molecules can be as abundant as 10^{-4} in the photosphere when the temperature is below about 2000 K (Tsuji 1964). We find that when the column density exceeds about 10^{19} cm^{-2} and the temperature is high, the sharp feature at 13.7 μm disappears and the C_2H_2 band becomes a rather smooth and deep




Figure 2. The column densities of C_2H_2 molecules in hot (*squares*) and cool (*circles*) layers are plotted against the near-infrared temperature. The hot-layer column densities after correction for dust emission effect are also plotted (*triangle*). The correction may be proper only for the low mass-loss rate stars (*filled triangles*). The error bars given in some data indicate the approximate uncertainty of the results in general. Symbols with arrows are upperlimits.

absorption from 10 μm to over 20 μm due to heavy saturation effects. Presumably, the *continuum* we draw in the present analysis is actually this continuous absorption, and the features we call *hot-component* arise from a layer just above this hottest, innermost part of the atmosphere.

The dust envelope also affects the results of the cool layer. Especially when the envelope has a moderate optical depth, the present plane parallel model potentially underestimates the column density of the cool layer, since it does not consider the size difference between two layers. This effect is small even in the most optically thick envelopes (group V), in which the circumstellar dust emission is the only continuum source.

4.2. Column densities of two layers

The column density in the hot-layer is almost constant at $10^{18.7} \text{ cm}^{-2}$ in group II and III stars. The value increases considerably in groups IV and V, but this may be an artifact because the dust mass-loss rate of these stars is too high for this correction method to be applicable.

N_{cool} increases as T_{NIR} decreases. Since T_{NIR} refers to the dust mass-loss rate of the stars, the good correlation between the two quantities indicates that the C_2H_2 molecule is associated with the dust envelope. One interesting exception is R Scl, which shows a significant N_{cool} although its T_{NIR} is the highest in the sample. This star is known to have a rather thick, probably detached circumstellar shell (Olofsson et al. 1990) and a relation between the cool C_2H_2 and such an envelope is probable.

4.3. C₂H₂ abundance

Despite the uncertainties described above, it is still worth to make estimates of the abundance of C₂H₂ in the two layers. According to the dynamic model atmosphere by Höfner & Dorfi (1997), the temperature of the hot layer, 1400 K, can be achieved at a distance of about 3 R_* from the central star. The density of the region is about 3×10^9 H₂ cm⁻³. Assuming that the molecule is distributed between the surface (1 R_*) and 3 R_* , the column density $10^{18.7}$ cm⁻² is converted to an abundance of 2×10^{-5} . In the circumstellar envelope, we assume that the density decreases proportionally to the square of radius. We take AFGL 3068 as an example. The expansion velocity and the mass-loss rate measured by CO observations is 14 km s⁻¹ and 2×10^{-5} M_⊙ yr⁻¹ (Loup et al. 1993), and the column density derived by the model analysis is $10^{17.7}$ cm⁻². Assuming the inner radius of the envelope as 3×10^{14} cm, we obtain an abundance of 7×10^{-6} .

These values are in good agreement with previous estimates of the C₂H₂ abundance (e.g., Keady & Hinkle 1988, Fuente et al. 1998). Our results indicate that the abundance does not change drastically in the regions below and above the dust forming region, i.e., no significant depletion of C₂H₂ onto grains seems to take place in the dust forming region.

Acknowledgments. The authors are grateful to Prof. K. Kawaguchi for providing the laboratory spectra of C₂H₂. I. Y. and L.B.F.M. W. acknowledge financial support from a NWO PIONIER grant.

References

- Cami J., Justtanont K., de Jong T., et al., 1997, First ISO Workshop on Analytical Spectroscopy, A.M. Heras et al. (eds.), ESA SP-419, p. 159
- Cernicharo J., Barlow M.J., González-Alfonso E., et al., 1996, A&A 315, L201
- de Graauw Th., Haser L.N., Beintema D.A., et al., 1996, A&A 315, L49
- Fuente A., Cernicharo J., Omont A., 1998, A&A 330, 232
- Groenewegen M.A.T., de Jong T., van der Blik N.S., Slijkhuis S., Willems F.J., 1992, A&A 253, 150
- Höfner S., Dorfi E.A., 1997, A&A 319, 648
- Justtanont K., Feuchtgruber H., de Jong T., et al., 1998, A&A 330, L17
- Kabbdaj Y., Herman M., Di Lonardo, et al., 1991, J. Mol. Spec. 150, 535
- Keady J.J., Hinkle K.H., 1988, ApJ 331, 539
- Keady J.J., Ridgway S.T., 1993, ApJ 406, 199
- Kessler M., Steinz J.A., Anderegg M.E., et al., 1996, A&A 315, L27
- Loup C., Forveille T., Omont A., Paul J.F., 1993, A&AS 99, 291
- Olofsson H., Carlström U., Eriksson K., et al., 1990, A&A 230, L13
- Palmer K.F., Mickelson M.E., Rao K.N., 1972, J. Mol. Spec. 44, 131
- Rinsland C.P., Baldacci A., Rao K.N., 1982, ApJS 49, 487
- Tsuji T., 1964, Ann. Tokyo Astr. Obs. 2nd Ser. 9, 1
- Weber M., Blass W.E., Halsey G.W., et al., 1994, J. Mol. Spec. 165, 107
- Yamamura I., de Jong T., Justtanont K., et al., 1998, Ap&SS 255, 351