

Polymerase chain reaction in the detection of an ‘outbreak’ of asymptomatic viral infections in a community birth cohort in south India

B. P. GLADSTONE^{1,2}, M. ITURRIZA-GOMARA³, S. RAMANI¹, B. MONICA¹,
I. BANERJEE¹, D. W. BROWN³, J. J. GRAY³, J. MULIYIL² AND G. KANG^{1*}

*Departments of*¹*Gastrointestinal Sciences and*²*Community Health, Christian Medical College, Vellore, India*
³*Enteric Virus Unit, Virus Reference Department, Health Protection Agency, London, UK*

(Accepted 13 April 2007; first published online 24 May 2007)

SUMMARY

Asymptomatic enteric infections are important where sequelae or protection from subsequent illness is an outcome measure. The use of reverse transcription–polymerase chain reaction (RT–PCR) to identify asymptomatic enteric infections in a birth cohort followed for rotaviral infections in a south Indian urban slum is reported. Of 1191 non-diarrhoeal samples from 371 children collected in May–June 2003, 22 (1·9%) were positive by ELISA. A total of 147 (40·6%) of 362 samples tested by VP6 RT–PCR were positive. In those samples that could be typed, a high diversity of G types including G1, G2, G4, G8, G9 and G10, and a high proportion (34·4%) of mixed infections were detected. Noroviruses were identified in 6/28 (21·4%) samples tested. The identification of infections undetectable by conventional techniques indicates the importance of the use of sensitive diagnostic techniques in research studies. Asymptomatically infected children may also act as a source of infection for other susceptible hosts.

INTRODUCTION

Enteric infectious diseases are complex, multifactorial conditions resulting from interactions between numerous factors including pathogen exposure, infectious dose, strain variation, environmental conditions, nutritional state and immune status [1]. These factors are not specific to a single infectious agent and a diagnosis of enteric infection is usually made either when patients present as sporadic cases of gastroenteritis or as part of an outbreak. Asymptomatic enteric infections are only identified as part of research studies, in both case-control and cohort designs, and have been reported from both developed and developing countries [2–4]. Given the range of

pathogens capable of enteric infection and replication within the gastrointestinal tract, the aetiology of both symptomatic and asymptomatic infection can be established only when appropriate laboratory facilities are available [5]. In most developing countries, diagnostic capacity is limited to a few bacterial and parasitic pathogens, contributing to a large diagnostic gap in the identification of pathogens producing both symptomatic and asymptomatic infections [6].

With repeated exposure to enteric pathogens at all ages, most enteric infections are asymptomatic, and the proportion that is asymptomatic increases beyond 2 years of age, possibly owing to the development of active immunity [7]. However, even in children aged <2 years, asymptomatic shedding can be detected for several days or weeks, with stools containing infectious viruses, bacteria, or protozoal cysts [8–11]. Persons with asymptomatic infections play an important role in the spread of many enteric pathogens,

* Author for correspondence: Dr G. Kang, Department of Gastrointestinal Sciences, Christian Medical College, Vellore – 632004, India.
(Email: gkang@cmcvellore.ac.in)

especially as they are unaware of their infection, take no special hygienic precautions and move normally from place to place [12]. Asymptomatic infections in children may also have a major role to play in determination of nutritional status and susceptibility to other infections [13].

In the present study, we highlight the use of molecular diagnostic tools to identify asymptomatic enteric infections, possibly resulting from a contamination or seeding event, in a birth cohort of children in a south Indian urban slum being followed for rotaviral infections. The identification of infections, which would not have been diagnosed by conventional techniques, in a large proportion of the paediatric population under study indicates the importance of the use of sensitive diagnostic techniques in studies where sequelae or protection from subsequent illness is an outcome measure.

METHODS

Study population

A birth cohort of 452 children, resident in an urban slum area in Vellore, was recruited between April 2002 and July 2003 and monitored for a total of 3 years to assess the protective effect of natural rotavirus infection on subsequent homotypic and heterotypic exposure. The study was based at three adjacent urban slums in Vellore: Ramnaickanpalayam, Chinnallapuram and Kaspaa, covering 2.2 km². Vellore municipality had an estimated population density of 1660 per km² in 2001, but density in the slums is higher. Mean annual rainfall in this area is about 1050 mm, peaking between August and November. Water for drinking is supplied at intervals of 2–28 days by the municipality from overhead tanks. During summer, temperatures are above 40 °C. The slum has temporary and permanent housing, poor water supply, large garbage dumps, open drains and poor roads. All households in the area were initially surveyed and information collected on the composition of the household in terms of residents, family structure, occupation and socioeconomic status. Following this baseline survey, all pregnant women were identified and visited at least once a week by the field worker. The family of each child born in the study area was approached for recruitment into the study cohort. Ethical approval was obtained for the study from the Christian Medical College Research Committee and recruitment

depended on parents having given informed written consent.

During the follow-up period each child was visited at home, twice a week, to record any morbidity in the child, or any diarrhoea in the family. Faecal samples were collected every 2 weeks from each child whether or not diarrhoea was reported. Samples were also collected when a child had diarrhoea. Any child who developed diarrhoea was assessed clinically and treated appropriately.

Sample testing

All samples were screened for rotavirus using either of two ELISA kits, based on VP6 detection by a monoclonal or polyclonal antibody (Rotavirus screen EIA, cat. no. Rota 003, Microimmune Ltd, UK; Rota IDEIA, cat. no. K6020, Dako Ltd, Denmark). All diarrhoeal samples were screened by bacterial culture and serogrouping for enteropathogenic *Escherichia coli*, *Salmonella* spp., *Shigella* spp., *Aeromonas* and vibrios and by microscopy for protozoan and helminthic parasites, including coccidian parasites such as *Cryptosporidium* as previously described [5].

Molecular testing for enteric viruses

The study protocol required testing of all ELISA positives by reverse transcription–polymerase chain reaction (RT–PCR) for confirmation of rotavirus by VP6 amplification and for genotyping of VP7 and VP4. In order to evaluate the relative sensitivities of the screening ELISA and RT–PCR for VP6, all 236 surveillance samples collected from asymptomatic children during a 2-week period in May–June 2003 were tested by both methods. When a high proportion of these samples were found to be positive, further studies were initiated.

A subset of 50 VP6 PCR-positive samples was genotyped by determination of VP7 and VP4 types. A subset of 28 surveillance samples and 10 diarrhoeal samples were also tested for noroviruses by screening with Ni and E3 primers as described below.

A 10% faecal suspension in balanced salt solution was used for nucleic acid extraction using guanidinium isothiocyanate/silica. Reverse transcription with random primers, VP6 detection and G- and P-typing semi-nested PCRs were performed as described previously [14–16]. The random priming method has previously been shown to detect 10 000 particles/ml of 10% faecal suspension [16]. The cDNA generated




Fig. 1. Testing of non-diarrhoeal samples for rotavirus by reverse transcription–polymerase chain reaction (RT–PCR) for the VP6 gene in a birth cohort under surveillance (—). The numbers of samples negative (□) and positive (■) for rotavirus VP6 by RT–PCR in each month are shown as proportionate bars.

by reverse transcription was used directly for detecting genogroup II noroviruses using published methods [17].

Additional data collection

When an unexpectedly high proportion of asymptomatic rotavirus infection was noted in the birth cohort, additional data was collected on rainfall and water supply, in addition to hospital laboratory-based data on other potentially waterborne infections, *Cryptosporidium*, cholera, hepatitis A and hepatitis E, presenting to the hospital during 2003, and compared with data for previous years during the period April–July.

RESULTS

A planned comparison of the sensitivity and specificity of ELISA and RT–PCR for VP6 resulted in the identification of a high rate of asymptomatic infections in the birth cohort under follow-up, during the period May–June 2003. The mean age of the cohort at this time was 6 months (s.d. = 3.8 months). Of the 371 children who contributed to the period May–June

2003, 118 (31.8%) had been previously infected with rotavirus. Of a total of 1191 non-diarrhoeal samples from 371 children collected over the 2-month period of May–June 2003, 22 (1.9%) were positive by ELISA and of these, 362 samples were tested by RT–PCR and 147 (40.6%) were positive (Fig. 1). This peak in PCR detection in asymptomatic samples was confined to a 2-week period at the end of May and in early June, since subsequent samples from the same children were negative. During this 2-week period, there was a significantly higher detection of positive samples (127 PCR positives/236 PCR tested, 53.8%) ($P < 0.001$) than during the period of May–June 2003, excluding the 2-week period (20 PCR positive/126 PCR tested, 15.9%). The detection rate by PCR was also higher ($P < 0.001$) than in the previous follow-up period from the beginning of enrolment of the birth cohort for samples tested by PCR (119 PCR positive/585 PCR tested, 20.3%, from 1 April 2002 to 26 May 2003).

Genotyping of the VP7 gene was attempted for 50 samples in order to determine whether this was an outbreak due to a single strain, and VP7 types could be determined for 29 samples. In 19 surveillance samples, a single G type was identified and in 10, there were

Table. Drinking water supply to study areas

Supply area	Average interval Jan.–Mar. 2003	Average interval Apr.–June 2003	Average interval July–Sep. 2003	Average interval Oct.–Dec. 2003
Kaspa	7 days	14 days	9 days	5 days
Ramnaickanpalayam	7 days	14 days	8 days	5 days

multiple G types indicating mixed infection. The G types identified were G1 ($n=25$), G2 ($n=7$), G4 ($n=1$), G9 ($n=4$) and G10 ($n=7$). The VP4 types were P[8] and P[4] and P[11], the latter was found only in association with G10 strains. The proportion of multiple infections was significantly higher ($P < 0.001$) in this 2-week period than in the period up to 26 May 2003, during when 66 of the 119 samples positive by VP6 PCR had G or P type determined and only 1.5% had multiple G or P types.

A total of 6/28 (21.4%) of surveillance samples tested for noroviruses during the period 27 May to 10 June 2003 were positive. Of these, two samples were also positive for rotaviruses. Other enteric pathogens were not tested in the surveillance samples.

During May–June 2003, there were 166 diarrhoeal samples of which three (1.8%) were positive for rotavirus antigen by EIA, and 34 (20.5%) were positive by VP6 PCR, this represented no difference in the 20.3% positivity by PCR that had been seen in diarrhoeal samples from the community tested from the start of follow-up for the birth cohort, where one of 10 diarrhoeal samples examined for noroviruses was positive. Bacteria identified in diarrhoeal samples included *Salmonella* C1 ($n=1$), *Vibrio cholerae* O1 ($n=1$), non-O1 vibrios ($n=3$), and *Aeromonas* ($n=6$).

Of the 70 diarrhoeal samples from the period 27 May to 10 June 2003, mixed enteric infections were found in 6.3% and of the 670 diarrhoeal samples from the period of follow-up prior to 27 May 2003, 3.5% had mixed infections ($P=0.5$).

There was a marked increase in the number of *Cryptosporidium*-associated cases of diarrhoea in the cohort in the period June–August 2003, with 14 infections during this period, whereas only six infections were documented in the period March 2002–May 2003 and seven infections during September 2003–March 2004. Earlier hospital-based data do not indicate any seasonality of cryptosporidial (data not shown) or rotaviral [14] infections in Vellore.

When the age of infected children in the cohort was studied, it was found that during the period of study, the median age of children with diarrhoea was 3.6




Fig. 2. The distribution of samples from asymptomatic (---) and symptomatic (—) children of different ages in a birth cohort tested for rotavirus by reverse transcription–polymerase chain reaction for the VP6 gene during the period April–July 2003. The median age of asymptomatic and symptomatic rotavirus-positive children are indicated by arrows.

months, while the median age of children providing surveillance or asymptomatic samples was 6.3 months (Mann–Whitney U test $P=0.0002$). A similar pattern was seen in children infected with rotavirus where rotavirus diarrhoea was seen at a median age of 3.5 months but asymptomatic rotavirus infections were seen at a median age of 5.5 months (Mann–Whitney U test $P=0.0004$) as shown in Figure 2.

In order to identify a possible cause for the increased incidence of enteric pathogen shedding by the children, the water supply to the area was investigated. Most people in the area utilized the municipal water for drinking, with other water sources reserved for washing. The municipal water supply to the study families was from two main overhead tanks, Ramnaickanpalayam and Kaspa, which supply domestic and public stand pipes. Water supply is intermittent (Table) and during the 92-day period from May to July 2003, water was supplied a total of five times. Since the increased enteric pathogen positivity began after the third week of May 2003, it is interesting to note that water was pumped to the two areas on 21 and 24 May 2003.

Additional data on disease due to other potential waterborne pathogens was collected at the referral hospital and showed no increase in numbers or proportion positive among samples tested for cholera during this time compared to the same months in the previous 3 years, but there were increases in cases of *Cryptosporidium* diarrhoea during June–August 2003 and hepatitis E cases during the following month from Vellore town (individual data not shown).

DISCUSSION

In the present study, we report the use of molecular techniques resulting in the identification of a possible contamination event resulting in widespread infection with a range of enteric pathogens which would have been undetected by conventional laboratory techniques. The only increased detection of pathogens associated with diarrhoea was with *Cryptosporidium* spp., but since this was associated with <10% of cases of diarrhoea, it was not recognized as an outbreak.

However, the use of RT–PCR permitted detection of rotaviruses in 40.6% of surveillance samples obtained during the period May–June 2003. This is much higher than the background prevalence of 5–10% positivity prior to and following this period of increased detection, as shown in Figure 1. In addition to multiple genotypes of rotavirus, noroviruses were also associated with asymptomatic infections during this period. The increased detection of enteric pathogens in the community could indicate a contamination event resulting in widespread infections. Since multiple pathogens and a significantly higher proportion of mixed infections were identified, a water source possibly contaminated with sewage would be the most likely source of infection. Resources for testing water samples for viral pathogens were not available, but previous studies in the area have found that all samples of drinking water are heavily contaminated with faecal coliforms, including those obtained from overhead tanks.

Since the detection of infection was by molecular-based tests applied to surveillance or asymptomatic samples, important questions to consider are the relevance of these results to the individual, to research, and to public health. At the level of the individual, a number of studies have reported the identification of enteric pathogens in asymptomatic individuals in both developed and developing countries [9–11, 18–22]. Reports from southern India have identified

enteric pathogens in up to 50% of controls [5], although more recent studies from the region indicate that the level of asymptomatic excretion of enteric pathogens is declining [23]. Children with diarrhoea and children with rotaviral diarrhoea were significantly younger than children who submitted surveillance specimens ($P=0.0002$), and children with asymptomatic rotavirus infection ($P=0.0004$) respectively, supporting the contention that protection from maternal antibodies may decline rapidly in developing countries. Studies from other developing countries have shown even asymptomatic enteric infections can have a significant negative impact on subsequent growth and development [13, 24]. It can be argued that the impact of reducing enteric infections should be measured taking into account the effects on both short-term consequences such as reduced diarrhoeal morbidity as well as the long-term effects on nutrition and development [25].

The reduction of the entry of enteric pathogens into the environment is key to lowering the levels of subsequent infection. Enteric infectious agents survive well under most environmental conditions, remaining in the environment where they can be transmitted directly through food or water or indirectly by contact with contaminated faeces, fomites such as equipment or mechanical vectors such as flies [26–29]. In the causation of disease, key factors include the number of organisms in the infecting inoculum, their viability and the susceptibility of the host. When most infections in children resulting from a contamination event are asymptomatic, it can be argued that the infectious dose must have been low, because of the lack of disease or that prior exposure to the pathogens or maternal antibody protection resulted in a protective immune response. In this study, diverse genotypes of rotaviruses and noroviruses were detected in asymptomatic infections, in addition to an increased incidence of cryptosporidial infections. Polymicrobial enteric infections have been previously reported mainly in outbreaks of disease [22, 30], but may be a common unrecognized event when water or other sources of infection are contaminated. It is also important to note that children asymptotically excreting virus may also act as a source of infection for other susceptible hosts [31], because their mobility is not restricted by illness.

In considering the implications of detecting the occurrence of asymptomatic infections in this study, ELISA was found to be an inappropriate technique for the detection of asymptomatic infection. Although

ELISA is sensitive and specific for the detection of group A rotavirus gastroenteritis [15] and previous studies on rotavirus epidemiology have used this technique [7, 32] when assessing asymptomatic infection or protection from disease, a detailed and careful investigation of infection that requires detection of asymptomatic infection should have molecular methods for rotavirus screening.

In community-based studies examining infection, disease and protection, it is important to use appropriate tools to ensure that infections are not missed due to a lack of sensitivity of the assay. This paper highlights the fact that the increased sensitivity of molecular tools enhanced the ability to detect infections that would have been missed by conventional ELISA-based techniques or by examination of only symptomatic cases. Understanding the burden and transmission of both infection and disease are important to formulate intervention strategies to prevent enteric infection and their long-term consequences.

ACKNOWLEDGEMENTS

This work was supported by Wellcome Trust Trilateral Cooperative Initiative on Infectious Diseases grant 063144. We thank Dr G. Sridharan and Dr Mary V. Jesudason for providing hospital data on potentially waterborne viral and bacterial infections.

DECLARATION OF INTEREST

None.

REFERENCES

1. Colwell RR. A voyage of discovery: cholera, climate and complexity. *Environmental Microbiology* 2002; **4**: 67–69.
2. Guerrant RL, *et al.* Mechanisms and impact of enteric infections. *Advances in Experimental Medicine and Biology* 1999; **473**: 103–112.
3. Scott TM, *et al.* Stool viruses in babies in Glasgow. 3. Community studies. *Journal of Hygiene (London)* 1979; **83**: 469–485.
4. Kamalratnam CN, *et al.* A prospective study of nosocomial enteric pathogen acquisition in hospitalised children in South India. *Journal of Tropical Paediatrics* 1979; **47**: 46–49.
5. Simpson R, *et al.* Infantile viral gastroenteritis: on the way to closing the diagnostic gap. *Journal of Medical Virology* 2003; **70**: 258–262.
6. Mathew M, *et al.* The relationship of microbial pathogens to acute infectious diarrhoea of childhood. *Journal of Tropical Medicine and Hygiene* 1991; **94**: 253–260.
7. Velazquez FR, *et al.* Rotavirus infections in infants as protection against subsequent infections. *New England Journal of Medicine* 1996; **335**: 1022–1028.
8. Kang G, *et al.* Prevalence of enteroaggregative and other HEp-2 cell adherent *Escherichia coli* in asymptomatic rural south Indians by longitudinal sampling. *Microbios* 1999; **100**: 57–66.
9. Abiodun PO, *et al.* Asymptomatic rotavirus infection in Nigerian day-care centres. *Annals of Tropical Paediatrics* 1985; **5**: 163–165.
10. Huang DB, *et al.* Cryptosporidiosis in children. *Seminars in Pediatric Infectious Diseases* 2004; **15**: 253–259.
11. Kang G, *et al.* Prevalence of intestinal parasites in rural Southern Indians. *Tropical Medicine and International Health* 1998; **3**: 70–75.
12. de Wit MA, *et al.* Risk factors for norovirus, Sapporo-like virus, and group A rotavirus gastroenteritis. *Emerging Infectious Diseases* 2003; **9**: 1563–1570.
13. Steiner TS, *et al.* Enteroaggregative *Escherichia coli* expresses a novel flagellin that causes IL-8 release from intestinal epithelial cells. *Journal of Clinical Investigation* 2000; **105**: 1769–1777.
14. Banerjee I, *et al.* Comparative study of the epidemiology of rotavirus in children from a community-based birth cohort and a hospital in South India. *Journal of Clinical Microbiology* 2006; **44**: 2468–2474.
15. Kang G, *et al.* Quantitation of group A rotavirus by real-time reverse-transcription-polymerase chain reaction: correlation with clinical severity in children in South India. *Journal of Medical Virology* 2004; **73**: 118–122.
16. Iturriza-Gomara M, *et al.* Comparison of specific and random priming in the reverse transcriptase polymerase chain reaction for genotyping group A rotaviruses. *Journal of Virological Methods* 1999; **78**: 93–103.
17. Green J, *et al.* Broadly reactive reverse transcriptase polymerase chain reaction for the diagnosis of SRSV-associated gastroenteritis. *Journal of Medical Virology* 1995; **47**: 392–398.
18. Araya M, *et al.* Acute diarrhoea and asymptomatic infection in Chilean preschoolers of low and high socioeconomic strata. *Acta Paediatrica Scandinavia* 1986; **75**: 645–651.
19. Barron-Romero BL, *et al.* Asymptomatic rotavirus infections in day care centers. *Journal of Clinical Microbiology* 1985; **22**: 116–118.
20. Caprioli A, *et al.* Enteropathogens associated with childhood diarrhoea in Italy. The Italian Study Group on Gastrointestinal Infections. *Pediatric Infectious Diseases Journal* 1996; **15**: 876–883.
21. Gallimore CI, *et al.* Asymptomatic and symptomatic excretion of noroviruses during a hospital outbreak of gastroenteritis. *Journal of Clinical Microbiology* 2004; **42**: 2271–2274.
22. Hellard ME, *et al.* Prevalence of enteric pathogens among community based asymptomatic individuals.

- Journal of Gastroenterology and Hepatology* 2000; **15**: 290–293.
23. **Kang G, et al.** Epidemiological and laboratory investigations of outbreaks of diarrhoea in rural South India: implications for control of disease. *Epidemiology and Infection* 2001; **127**: 107–112.
 24. **Steiner TS, et al.** Infectious diarrhoea: new pathogens and new challenges in developed and developing areas. *Clinical Infectious Diseases* 2006; **43**: 408–410.
 25. **Lima AA, Guerrant RL.** Strategies to reduce the devastating costs of early childhood diarrhoea and its potential long-term impact: imperatives that we can no longer afford to ignore. *Clinical Infectious Diseases* 2004; **38**: 1552–1554.
 26. **Binka FN, et al.** Incidence and risk factors of paediatric rotavirus diarrhoea in northern Ghana. *Tropical Medicine and International Health* 2003; **8**: 840–846.
 27. **Mitra AK, Rabbani F.** The importance of breastfeeding in minimizing mortality and morbidity from diarrhoeal diseases: the Bangladesh perspective. *Journal of Diarrhoeal Diseases Research* 1995; **13**: 1–7.
 28. **Pazzaglia G, et al.** Campylobacter-associated diarrhoea in Egyptian infants: epidemiology and clinical manifestations of disease and high frequency of concomitant infections. *Journal of Diarrhoeal Diseases Research* 1993; **11**: 6–13.
 29. **Prado MS, et al.** Risk factors for infection with *Giardia duodenalis* in pre-school children in the city of Salvador, Brazil. *Epidemiology and Infection* 2003; **131**: 899–906.
 30. **Nimri LF, et al.** Polymicrobial infections in children with diarrhoea in a rural area of Jordan. *FEMS Immunology and Medical Microbiology* 2004; **42**: 255–259.
 31. **Nwachuku N, Gerba CP.** Health risks of enteric viral infections in children. *Reviews in Environmental Contamination and Toxicology* 2006; **186**: 1–56.
 32. **Haffejee IE, Moosa A.** Rotavirus studies in Indian (Asian) South African infants with acute gastroenteritis: I. Microbiological and epidemiological aspects. *Annals of Tropical Paediatrics* 1990; **10**: 165–172.