



	Baseline Periods			Study Period							Total Tnal Period
	11/2019 - 02/2020	11/2020 - 02/2021	05/2021 - 10/2021	21-Dec	Jan - 22	22-Feb	22-Mar	22-Apr	22-May	22-Jun	
Patient Days	1,368	1,676	2,891	459	567	464	382	406	392	353	3,023
MRSA Infections	1	8	8	0	0	0	0	0	0	0	0
MRSA Infection Rate	0.08	0.67	0.41	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
C. diff Infections	1	5	4	0	0	0	0	0	0	0	0
C. diff Infection Rates	0.09	0.46	0.23	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
CLABSI Infections	0	0	2	0	0	0	0	0	0	0	0
CLABSI Infection Rate	0.00	0.00	0.69	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Presentation Type:

Poster Presentation - Poster Presentation

Subject Category: Implementation Science

Comparison of a standard environmental surface sampling method and a composite approach for select healthcare pathogens

Monica Chan; Judith Noble-Wang and Laura Rose

Background: Hospital surfaces are known to contribute to the spread of healthcare-associated infections (HAIs). Environmental sampling is often performed to locate a reservoir or to evaluate intervention strategies in healthcare facilities. Composite sampling is commonly practiced in other fields of environmental sampling and involves collection of multiple samples combined entirely or partially to form a new sample. We compared a standard CDC surface whole-tool sampling method with a composite sampling approach. **Methods:** *Acinetobacter baumannii* (AB), *Klebsiella pneumoniae* that produce *K. pneumoniae* carbapenemase (KPC), vancomycin-resistant *Enterococcus faecalis* (VRE), methicillin-resistant *Staphylococcus aureus* (MRSA), and *Clostridioides difficile* spores were suspended in an artificial soil and deposited as 40 µL droplets (~104 CFU total) onto steel coupons of surface areas 323 cm², 645 cm², or 1,290 cm² and dried for 2 hours. The surfaces were sampled with a single pass of a cellulose sponge—either the larger side of the sponge (face) or the smaller side of the sponge (edge)—and the optimal surface area was determined. Recovery from the optimal surface area with a single pass sampling was compared to the recovery using a standard CDC method in which all sides were used (ie, whole-tool method) to sample a standard area (645 cm²). Recovery was determined by culture and total CFU were determined for each optimal surface area. Theoretical composites were constructed using the mean total CFU of optimal surface area; 2×((face) + (edge)). Significance was set at $P \leq .05$. **Results:** Total CFU recovery using the whole-tool method was significantly greater than the single pass sample recovery for MRSA (18,300 vs 16,600 CFU) and VRE (27,600 vs 26,400 CFU) ($P < .05$). When comparing the theoretical composite method to the standard whole-tool area (625 cm²), the theoretical composite total CFU was significantly greater than the whole-tool method for all organisms. For example, VRE recovery with the standard CDC whole-tool method was 27,600 CFU from 625 cm², yet a

theoretical composite approach recovered 79,800 CFU from an area of 1,290 cm². **Conclusions:** Many factors influence recovery when sampling the environment, and composite sampling is a promising approach when sampling large surface areas. Using a theoretical composite of single-pass samples, the potential for improved detection with composite sampling was demonstrated. A composite sampling approach will reduce time and resources for sampling and sample processing, allowing larger surface areas to be investigated which will improve infection control strategies.

Disclosures: None

Antimicrobial Stewardship & Healthcare Epidemiology 2023;3(Suppl. S2):s74
doi:10.1017/ash.2023.325

Presentation Type:

Poster Presentation - Poster Presentation

Subject Category: Infection Control in Low and Middle-Income Countries
Infection prevention and control perspective and practices among healthcare workers in Bangladesh: A multicenter cross section

Md. Golam Dostogir Harun; Shariful Amin Sumon; Aninda Rahman; Md Mahabub Ul Anwar and Md. Saiful Islam

Background: Infection prevention and control (IPC) is a critical feature of preventing the spread of healthcare-associated infections (HAIs) in hospitals. IPC practices are particularly important in resource-constrained and crowded hospital settings. The successful implementation of infection prevention measures depends on healthcare worker (HCW) knowledge of, attitude toward, and practice (KAP) of IPC. In this project, we assessed the KAP of HCWs and identified factors associated with IPC compliance at tertiary-care hospitals in Bangladesh. **Methods:** From September 2020 to January 2021, we conducted this hospital-based cross-sectional assessment at 11 tertiary-care hospitals. A semistructured questionnaire was used to conduct face-to-face interviews with physicians, nurses, and cleaning staff who were directly involved in patient care. Based on >75% of the total score, each KAP component was divided into adequate knowledge, favorable attitude, and safe practice. We performed descriptive analysis and multivariate logistic regression to determine the KAP score and associated