

Corrigendum

Cite this article: Lillitos PJ, Nassar MS, Tibby SM, Simmonds J (2019). Is the medical treatment for arterial hypertension after primary aortic coarctation repair related to age at surgery? A retrospective cohort study – CORRIGENDUM. *Cardiology in the Young* **29**: 1005. doi: [10.1017/S1047951118000276](https://doi.org/10.1017/S1047951118000276)

First published online: 5 November 2018

Is the medical treatment for arterial hypertension after primary aortic coarctation repair related to age at surgery? A retrospective cohort study – CORRIGENDUM

Peter J Lillitos, Mohamed S. Nassar, Shane M Tibby and Joy Simmonds

doi: [10.1017/S1047951117001019](https://doi.org/10.1017/S1047951117001019), Published by Cambridge University Press, 9 June 2017.

The authors apologise for an error in the caption for Figure 3. The caption currently reads: “Kaplan-Meier plot of risk of requiring future anti-hypertensive medication following primary surgical coarctation repair in neonatal, infant and child groups”.

The caption should read:

“Kaplan-Meier plot of estimated time from surgery free from requiring anti-hypertensive medication following primary surgical coarctation repair in neonatal, infant and child groups”.

Reference

Lillitos PJ, Nassar MS, Tibby SM, Simmonds J. Is the medical treatment for arterial hypertension after primary aortic coarctation repair related to age at surgery? A retrospective cohort study. *Cardiol Young*. Published by Cambridge University Press, 9 June 2017. doi: [10.1017/S1047951117001019](https://doi.org/10.1017/S1047951117001019).