

REFERENCES

- Banerjee SN, Emori TG, Culver DA, Gaynes RP, Jarvis WR, Horan T, et al. Secular trends in nosocomial primary bloodstream infections in the United States. National Nosocomial Infectious Surveillance System. *Am J Med* 1991;91(suppl 3B):865-895.
- Komshian SV, Uwaydah AK, Sobel JD, Crane L. Fungemia caused by *Candida* sp and *Torulopsis glabrata* in hospitalized patients frequency, characteristics and evaluation of factors influencing outcome. *Rev Infect Dis* 1989;3:379-390.
- Wey SB, Motomi M, Pfaller MA, Woollen RF, Wenzel RP. Hospital acquired candidemia: the attributable mortality and excess length of stay. *Arch Intern Med* 1988;148:2642-2647.
- Vazquez JA, Sanchez V, Dmuchowski C, Dembry LM, Sobel JD, Zervos MJ. Nosocomial acquisition of *Candida albicans*: an epidemiologic study. *J Infect Dis* 1993;168:195-201.
- Sanchez V, Vazquez J, Jones DB, Dembry L, Sobel JD, Zervos MJ. Nosocomial acquisition of *Candida parapsilosis*: an epidemiologic study. *Am J Med* 1993;94:577-582.
- Sanchez V, Vazquez JA, Barth-Jones D, Dembry L, Sobel JD, Zervos MJ. Epidemiology of nosocomial acquisition of *Candida lusitanae*. *J Clin Microbiol* 1993;30:3005-3008.
- Vazquez JA, Dembry LM, Sanchez V, Vazquez MA, Sobel JD, Dmuchowski C, Zervos MJ. Nosocomial *Candida glabrata* colonization: an epidemiologic study. *J Clin Microbiol* 1998;36:421-426.
- Isenberg HD, Tucci V, Clintron F, Sincer C, Weinstein GS, Tyras D. Single source outbreak of *Candida tropicalis* complicating coronary bypass surgery. *J Clin Microbiol* 1989;27:2426-2428.
- Jarvis WR. Epidemiology of nosocomial fungal infections, with emphasis on *Candida* species. *Clin Infect Dis* 1995;20:1526-1530.
- Vaudry WL, Tierney AJ, Weinman. Investigating of cluster of systemic *Candida albicans* infections in a neonatal intensive care unit. *J Infect Dis* 1988;158:1375-1379.
- Weems JJ, Chamberland ME, Ward J, Willy M, Payne AA, Solomon SL. *Candida parapsilosis* fungemia associated with parenteral nutrition and contaminated blood pressure transducers. *J Clin Microbiol* 1987;1925:1029-1032.
- Burnie JP, Odds FC, Lee W, Webster C, Williams JD. Outbreak of systemic *Candida albicans* in an intensive care unit caused by cross infection. *BMJ* 1985;290:746-748.
- Berkelman RL, Holland BW, Anderson R. Increased bactericidal activity of dilute preparations of Clinidine solutions. *J Clin Microbiol* 1982;15:635-639.
- Haley CE, Mailing-Casin M, Smith J, Luby J, Mackowiak P. Bactericidal activity of antiseptics against methicillin resistant *Staphylococcus aureus*. *J Clin Microbiol* 1985;21:991-992.
- Russell AD, Ahonkhai I, Rogers DT. Microbiological applications of the inactivation of antibiotics and other antimicrobial agents. *J Appl Bacteriol* 1979;46:207-245.
- Phelps M, Ayliffe GAJ, Babb JR. An outbreak of candidiasis in a special care unit: the use of resitogram typing method. *J Hosp Infect* 1986;7:13-20.
- Maki DG, Alvarado CJ, Hassemr CA, Zilz MA. Relation of the inanimate hospital environment to endemic nosocomial infection. *N Engl J Med* 1982;307:1562-1566.
- Nicoletti G, Boghossian V, Gurevitch F, Borland R, Morganroth P. The antimicrobial activity in vitro of chlorhexidine, a mixture of isothiazolones ('Kathon' CG) and cetyl trimethyl ammonium bromide (CTAB). *J Hosp Infect* 1993;23:87-111.

Parenteral Antibiotic Administration on Persistence of VRE

Gina Pugliese, RN, MS
Martin S. Favero, PhD

Donskey and colleagues from the Veterans Affairs' Medical Center and Case Western Reserve University School of Medicine, Cleveland, Ohio, used a mouse model of vancomycin-resistant *Enterococcus faecium* (VRE) intestinal colonization to study the effect of different subcutaneous antibiotics on persistence and density of VRE colonization. Gastric inoculation of a clinical *vanB* VRE isolate, in conjunction with oral vancomycin in drink-

ing water (250 µg/mL), resulted in high-level VRE colonization (mean, 9.5 log₁₀ colony-forming units [CFU]/g) in all 169 experimental mice. After discontinuation of oral vancomycin, the level of VRE in the stool specimens of mice receiving subcutaneous saline steadily decreased (mean, 3.59 log₁₀ CFU/g at day 19). Subcutaneous vancomycin, clindamycin, piperacillin-tazobactam, ticarcillin-clavulanic acid, metronidazole, cefotetan, ampicillin, and ampicillin-sulbactam all promoted persistent high levels of stool VRE. Subcutaneous ceftriaxone, cefepime, ciprofloxacin, and aztreonam promot-

ed increased VRE density to a lesser degree or not at all.

Thus, in a mouse model, vancomycin and antibiotics with potent anti-anaerobic activity promoted persistent high-density intestinal VRE colonization, whereas antibiotics lacking potent anti-anaerobic activity did not.

FROM: Donskey CJ, Hanrahan JA, Hutton RA, Rice LB. Effect of parenteral antibiotic administration on persistence of vancomycin-resistant *Enterococcus faecium* in the mouse gastrointestinal tract. *J Infect Dis* 1999;180:384-390.