

Climate change

O0082

Climate change and mental health. Polish perspective.

M. Gawrych

Psychology Institute, The Maria Grzegorzewska Univeristy, Warsaw, Poland

doi: 10.1192/j.eurpsy.2024.204

Introduction: World Health Organization estimates that climate changes are expected to cause an additional 250 000 deaths worldwide per year between 2030 and 2050 (1). We do not know, in what extent, population mental health will deteriorate due to climate change. Unfortunately, not all European countries, including Poland collect the evidence-based information about current and possible future risks for mental health.

(1)World Health Organization. Climate change and health; 2018. Available from who.int/news-room/fact-sheets/detail/climate-change-and-health [cited 20 October 2020]

Objectives: The aims of present study are: (1) summarize the available literature through a current review and (2) make recommendations for future actions/ prevention strategy for Poland.

Methods: Medline database (through PubMed) and Polish authorities documents was searched for records published in 2010–2024. Mental health-related descriptors (i.e. „mental health” OR “mental disorders”) and „climate change” and “Poland/ Polish” and “Europe” term were used in particular searches. The results of the screening were included in the final selection list. References of screened full-text articles and reports were manually searched for further literature. Additionally, European and worldwide publications and reports prepared by mental health and/ or climate change organizations were taken into account.

Results: Mental health risks of climate change can stem from climate-related natural disasters (e.g., extreme weather events), slower moving events (e.g., drought), or concern about the phenomenon of climate change itself. Primary mental health impact is related mostly to disasters itself and its consequences: environment of disruption, trauma and grief. Direct consequences include increased rates of high-risk behaviours. Secondary effects of climate change are due to various processes of environmental changes and ecological disruptions. They consist of damages to physical and social infrastructure, physical health effects, food and water shortages, conflict, and displacement. Long-term droughts affect food and water supplies and can subsequently affect the economic and mental wellbeing not only the land-based workers.

Conclusions: A focus on climate change impact on mental health can help enhance the understanding of factors that strengthening psychosocial resilience and adaptation. The future mental health challenges of climate change in Poland cover:

- developing scientific knowledge regarding adaptation process and resilience,
- focusing on high-risk groups (i.e. children, rural workers),
- strengthened community engagement,
- developing available locally strategies for mental health improving.

Disclosure of Interest: None Declared

Intellectual Disability

O0081

Effectiveness of AI-driven Individualized Learning Approach for Children with Autism Spectrum Disorder (ASD)

H. Atturu^{1*} and S. Naraganti²¹Dept of Psychiatry, CARE Hospitals and ²Dept of Clinical Psychology, Rainbow Children's Hospital, Banjara Hills, Hyderabad, India

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.205

Introduction: Autism Spectrum Disorder (ASD) is a condition with varying degrees of social, emotional and behavioural disability. These children require focused and individualised learning plan to facilitate social integration. Robots have been used for this purpose but are not routinely available in several parts of the world. Effective, point of care (POC) digital therapies that can be used anywhere by anyone is the need of the day.

Objectives: To evaluate the effectiveness of Artificial Intelligence (AI) driven individualised learning plans delivered through POC digital platform (CognitiveBotics) for children with ASD.

Methods: After Ethical approval and parental consent, children diagnosed with ASD (Childhood Autism Rating Scale CARS 2) aged 2 years and above were screened for study inclusion and exclusion criteria and enrolled. AI driven individualised learning plan was administered through CognitiveBotics software that could be used on either computer or a tablet. Initially, interactive questions were administered to parents by the AI tool to understand child's functioning. Based on these, an individualised learning plan was assigned. Each task is delivered using either interactive videos, chatbot and/or animated/AI games. The child's progress is captured for attention (attempted questions, retries and timeouts) and retention (first time corrects and corrects) continuously. The initial interactive questions administered to parents were repeated to assess child's progress in real life. Paired 't' test using SPSS version 26 was used to compare initial and final data.

Results: Out of 85 registered children, 41 regularly used the AI tool. Mean age was 43.93 months (range 26 to 72 months). 37 (90.24%) were boys. The baseline mean scores (ranges), were CARS 33.48 (30-39.5); Social quotient 53.4 (27.25-80.32), Developmental quotient 71.35 (45.90-93.33) and IQ 62.34 (36.58-86.83). The base line mean score of parents assessed child function was 115.24 (range 58 to 215). A mean of 15.54 tasks were given (range 5 to 48). At the time of analysis with a mean follow up of 3 months (range 2 to 5 months) the children completed a mean of 10.10 tasks (range 0 to 42). There was significant improvement in child's learning captured by the AI software based on attention and retention parameters ($p < 0.00001$). This improvement was also reflected in parent assessed child function (mean 147.15, (59 to 231)) ($p < 0.00001$). The percentage of improvement in both software captured and parent assessed child function was directly related to the amount of time spent by the child on the software.

Conclusions: AI driven individualised learning approach is effective in teaching skills and promote social integration for children with ASD. Such technology can capture the child's progress on a day-to-day basis and deliver personalised training.

Disclosure of Interest: None Declared

Bipolar Disorders

O0082

The relationship between sleep problems and suicidality in Bipolar Disorder: a Systematic Review and a Meta-analysis

C. Possidente^{1,2,3}, M. Bort^{1,2,4*}, M. De Prisco^{1,2,5}, V. Oliva^{1,2,6,7}, G. Fico^{4,8,9}, L. Bracco^{1,2,10}, C. Sommerhoff^{1,2}, L. Montejo^{1,2,5,7}, A. Murru^{1,2,4} and E. Vieta^{1,2,4,5}

¹Departament de Medicina, Facultat de Medicina i Ciències de la Salut, Universitat de Barcelona; ²Bipolar and Depressive Disorders Unit, Hospital Clinic de Barcelona, Barcelona, Spain; ³Bipolar and Depressive Disorders Unit, University of Bologna, Bologna, Italy; ⁴Institut d'Investigacions Biomèdiques August Pi i Sunyer, Barcelona; ⁵Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Instituto de Salud Carlos III, Madrid, Spain; ⁶Department of Biomedical and Neuromotor Sciences, University of Bologna, Bologna; ⁷Institut d'Investigacions Biomèdiques August Pi i Sunyer; ⁸Departament de Medicina, Facultat de Medicina i Ciències de la Salut, Universitat de Barcelona; ⁹Bipolar and Depressive Disorders Unit, Hospital Clinic de Barcelona, Barcelona and ¹⁰Department of Pathophysiology and Transplantation, University of Milan, Milan, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.206

Introduction: Bipolar disorder (BD) is a multifaceted illness encompassing mood, energy, cognitive and biorhythms alterations. Sleep disturbances are common in prodromic, acute and inter-episodic phases of BD. Suicidality presents a known association with sleep disturbances. However, their interplay in BD remains intricate and not fully elucidated.

Objectives: The aim of the present systematic review (SR) and meta-analysis (MA) is to summarise the available evidence and to provide an estimate of the association between sleep disturbances and suicidality, defined as presence of suicide ideation, behaviour and suicide attempts, in patients with BD.

Methods: We conducted a comprehensive literature search following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines across PubMed, PsycINFO, and SCOPUS databases. We included all studies reporting an association between sleep problems and suicidal behaviour in BD patients. No language restriction was imposed. Effect sizes were calculated as odds ratio (OR) for dichotomic variables, standard mean difference (SMD) for continuous outcomes, and Spearman's coefficient (r) for correlations. Heterogeneity was assessed using the I² statistic. Global inconsistency was evaluated using the Q statistics with the corresponding p-value.

Results: The initial search yielded 911 unique abstracts, of which 62 underwent full-text screening. Fourteen publications were included, comprising twelve cross-sectional and two longitudinal studies. The total sample consisted of 19,601 subjects diagnosed with BD, of which 51.76% were females and 69.52% had a diagnosis of BD type 1. We found that people with BD and sleep disturbances tend to have higher suicidality, both current (SMD=0.79, 95% CI=0.53, 1.05) and lifetime (OR=1.8; 95%CI=1.41, 2.55), when compared with people with BD and no sleep disturbances. Additionally, patients with BD and a history of suicide attempts tend to have more sleep problems (OR=1.37, 95%CI=1.21, 1.55).

Moreover, a positive correlation exists between suicidality and poor sleep quality measured by the Pittsburgh Sleep Quality Index (PSQI) (r= 0.24, 95%CI=0.10, 0.36). No heterogeneity was found, except in the subanalysis of correlation (I²=66.67%, Q p-value=0.01).

Conclusions: Our SRMA outlines a positive relation between sleep disturbances and suicidality in patients with BD. The small number of included studies and the scarcity of longitudinal studies, preventing the inference of a causal relationship, represent the major limitations of this study. Also, studies with objective measures of sleep alterations are currently lacking. The prompt recognition, objective measurement, and treatment of sleep alterations could be crucial in averting or reducing suicidal attempts in BD.

Disclosure of Interest: C. Possidente: None Declared, M. Bort: None Declared, M. De Prisco: None Declared, V. Oliva: None Declared, G. Fico Grant / Research support from: Fellowship from "La Caixa" Foundation (ID 100010434 - fellowship code LCF/BQ/DR21/11880019), L. Bracco: None Declared, C. Sommerhoff: None Declared, L. Montejo: None Declared, A. Murru Grant / Research support from: Spanish Ministry of Science and Innovation (PI19/00672, PI22/00840) integrated into the Plan Nacional de I +D+I and co-financed by the ISCIII-Subdirección General de Evaluación and the Fondo Europeo de Desarrollo Regional (FEDER), E. Vieta Grant / Research support from: Spanish Ministry of Science and Innovation (PI18/00805, PI21/00787) integrated into the Plan Nacional de I+D+I and cofinanced by the ISCIII-Subdirección General de Evaluación and the Fondo Europeo de Desarrollo Regional (FEDER); the Instituto de Salud Carlos III; the CIBER of Mental Health (CIBERSAM); the Secretaria d'Universitats i Recerca del Departament d'Economia i Coneixement (2017 SGR 1365), the CERCA Programme, and the Departament de Salut de la Generalitat de Catalunya for the PERIS grant SLT006/17/00357; the European Union Horizon 2020 research and innovation program (EU.3.1.1. Understanding health, wellbeing and disease: Grant No 754907 and EU.3.1.3. Treating and managing disease: Grant No 945151).

O0083

Evaluation of Neurocognitive Functions of Children and Adolescents Diagnosed with Pediatric Bipolar Disorder and Disruptive Mood Dysregulation and at High Risk for Bipolar Disorder

I. Yenen Sivri^{1*}, E. Çöp², K. C. Can³ and A. S. Yenen Menderes⁴

¹Child and Adolescent Psychiatry, Osmaniye State Hospital; ²Child and Adolescent Psychiatry; ³Adult Psychiatry, Ankara City Hospital and ⁴Child and Adolescent Psychiatry, Keçiören Educational Research Hospital, Ankara, Türkiye

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.207

Introduction: Pediatric Bipolar Disorder (BPD) is a chronic psychiatric disorder that alters normal and psychological development processes among patients. Although cognitive deficits in BPD have identified in recent studies, little is known about the developmental trajectory of these deficits. DMDD is a newly defined diagnosis included in the DSM-V. Since it added a new dimension to the clinical spectrum but few studies conducted on DMDD, there are