

SURVEY

Counting problems from the viewpoint of ergodic theory: from primitive integer points to simple closed curves

FRANCISCO ARANA-HERRERA 

*Department of Mathematics, University of Maryland,
William E. Kirwan Hall, 4176 Campus Dr, College Park, MD 20742, USA
(e-mail: farana@umd.edu)*

(Received 14 March 2022 and accepted in revised form 19 August 2024)

Abstract. In her thesis, Mirzakhani showed that the number of simple closed geodesics of length $\leq L$ on a closed, connected, oriented hyperbolic surface X of genus g is asymptotic to L^{6g-6} times a constant depending on the geometry of X . In this survey, we give a detailed account of Mirzakhani's proof of this result aimed at non-experts. We draw inspiration from classic primitive lattice point counting results in homogeneous dynamics. The focus is on understanding how the general principles that drive the proof in the case of lattices also apply in the setting of hyperbolic surfaces.

Key words: counting, lattice, geodesic, surface, Teichmüller

2020 Mathematics Subject Classification: 57K20 (Primary); 30F60 (Secondary)

Contents

1	Introduction	1
2	Counting primitive lattice points in the Euclidean plane	3
3	Hyperbolic surfaces, Teichmüller spaces, and simple closed curves	12
4	Weil–Peterson volumes and Mirzakhani's integration formulas	23
5	Counting simple closed geodesics on hyperbolic surfaces	32
6	Beyond simple closed geodesics	39
	Acknowledgements	46
	References	46

1. Introduction

Let us begin our discussion by recalling a classic result in number theory: the prime number theorem. According to this theorem, the number $\pi(N)$ of positive prime integers p such that $p \leq N$ satisfies the following asymptotic estimate as $N \rightarrow \infty$,

$$\pi(N) \sim \frac{N}{\log(N)},$$

where the symbol \sim represents the fact that the following identity holds:

$$\lim_{N \rightarrow \infty} \frac{\pi(N)}{N / \log(N)} = 1.$$

Analogous results also hold in the setting of hyperbolic surfaces. A closed geodesic on a hyperbolic surface is said to be primitive if it cannot be represented as a concatenation of multiple copies of a shorter closed geodesic. Given a closed, oriented hyperbolic surface X and a parameter $L > 0$, denote by $c(X, L)$ the number of non-oriented, primitive closed geodesics on X of length $\leq L$. The prime geodesic theorem states that, for any closed, connected, oriented hyperbolic surface X , the following asymptotic estimate holds as $L \rightarrow \infty$,

$$c(X, L) \sim \frac{e^L}{2L}.$$

This result shows in particular that the asymptotics of $c(X, L)$ as $L \rightarrow \infty$ do not depend on the geometry of X nor on its topology. The first proof of this theorem was given by Huber and Selberg using analytic methods. See [Bus92, Ch. 9] for a detailed discussion of this proof. A dynamical proof of this result for more general negatively curved manifolds was given by Margulis in [Mar70]; this proof relates the exponential growth rate of the counting function of periodic orbits to the topological entropy of the geodesic flow on the manifold.

In the setting of hyperbolic surfaces, we can push our curiosity even further. A closed geodesic on a hyperbolic surface is said to be simple if it does not intersect itself. Given a closed, connected, oriented hyperbolic surface X and a parameter $L > 0$, denote by $s(X, L)$ the number of simple closed geodesics on X of length $\leq L$. What are the asymptotics of this quantity as $L \rightarrow \infty$? Do they depend on the geometry of X ? Do they depend on the topology of X ?

These questions, which *a priori* seem very similar to those answered by the prime geodesic theorem, remained out of reach for quite a long time. The analytic perspective of Huber and Selberg was not of much use in this setting as it could not distinguish simple closed geodesics among primitive ones. Some progress toward answering these questions was made by Birman and Series [BS85], McShane and Rivin [MR95a, MR95b], and Rivin [Riv01]. The first major breakthrough would come through the work of Mirzakhani, who, in her thesis [Mir04], proved the following result.

THEOREM 1.1. [Mir08b, Theorem 1.1] *Let X be a closed, connected, oriented hyperbolic surface of genus $g \geq 2$. Then, there exists $s(X) > 0$ such that the following asymptotic estimate holds as $L \rightarrow \infty$:*

$$s(X, L) \sim s(X) \cdot L^{6g-6}.$$

Mirzakhani's proof of Theorem 1.1 uses ergodic theory in a crucial way. Her proof also makes important use of a couple of other breakthroughs in her work: her formulas for the total Weil–Petersson volumes of moduli spaces and her integration formulas over moduli space.

The tools developed by Mirzakhani in her thesis have broad-ranging consequences in many fields of mathematics. Perhaps most well known is her novel proof of Witten's conjecture on intersection numbers of psi classes on moduli space of Riemann surfaces [Wit91]. On a simpler note, let us highlight the following concrete consequence of her work.

THEOREM 1.2. [Mir08b, Corollary 1.4] *On any closed, oriented hyperbolic surface of genus 2, it is 48 times more likely for a random long simple closed geodesic to be non-separating rather than separating.*

The main goal of this survey is to give a detailed account of Mirzakhani's proof of Theorem 1.1 aimed at non-experts. We will draw inspiration from classic primitive lattice point counting results in homogeneous dynamics. Although we will cover the necessary background, the focus will be on understanding how the general principles that drive the proof in the case of lattices also apply in the setting of hyperbolic surfaces. In particular, we will take for granted several fundamental results about Teichmüller spaces and mapping class groups and focus on understanding their applications.

1.1. Organization of this survey. In §2, we study counting problems for primitive lattice points in the Euclidean plane. This discussion will later guide the proof of Theorem 1.1. In §3, we cover the background material on hyperbolic surfaces, Teichmüller spaces, and simple closed curves needed to understand the proof of Theorem 1.1. In §4, we discuss Mirzakhani's famous formulas for the total Weil–Petersson volumes of moduli spaces and her famous integration formulas over moduli space. In §5, we give a complete proof of Theorem 1.1. In §6, we give a brief overview of several counting results for closed curves on surfaces and other related objects that have been proved since the debut of Mirzakhani's thesis.

1.2. Other surveys. For a review of similar topics from a more algebro-geometric perspective, see [Wol13]. For a survey covering the full range of Mirzakhani's outstanding research beyond the few results discussed here, see [Wri20]. For discussions of geodesic counting theorems from the point of view geodesic currents, see [ES22, EU20].

2. Counting primitive lattice points in the Euclidean plane

2.1. Outline of this section. In this section, we study counting problems for primitive lattice points in the Euclidean plane. The techniques introduced in this section will serve as a rough guide for the proof of Theorem 1.1 that will be discussed in §5.

2.2. Counting primitive integer points. Consider the integer lattice $\mathbb{Z}^2 \subseteq \mathbb{R}^2$. In analogy with the definition of prime numbers, a vector $v \in \mathbb{Z}^2$ is said to be primitive if it cannot be written as a non-negative integer multiple of another vector in \mathbb{Z}^2 . Equivalently, a vector $v = (a, b) \in \mathbb{Z}^2$ is said to be primitive if the greatest common divisor of a and b is 1. Denote by $\mathbb{Z}_{\text{prim}}^2 \subseteq \mathbb{Z}^2$ the subset of all primitive vectors of \mathbb{Z}^2 . See Figure 1. Notice




FIGURE 1. Primitive vectors of the integer lattice \mathbb{Z}^2 (colour online).

$\mathbb{Z}^2_{\text{prim}} \subseteq \mathbb{Z}^2$ is not a sublattice, not even a subgroup. Denote by $\|\cdot\|$ the Euclidean norm on \mathbb{R}^2 . For every $L > 0$, consider the counting function

$$p(\mathbb{Z}^2, L) := \#\{v \in \mathbb{Z}^2_{\text{prim}} : \|v\| \leq L\}. \tag{2.1}$$

Just as in the case of the counting function $\pi(N)$ introduced in §1, we are interested in the asymptotics of $p(\mathbb{Z}^2, L)$ as $L \rightarrow \infty$. The main goal of this section is to discuss a rather non-standard proof of the following classic result.

THEOREM 2.1. *The following asymptotic estimate holds as $L \rightarrow \infty$:*

$$p(\mathbb{Z}^2, L) \sim \frac{6}{\pi} \cdot L^2.$$

To prove Theorem 2.1, we begin by rewriting the counting function $p(\mathbb{Z}^2, L)$ in a more convenient way. More concretely, notice that for every $L > 0$,

$$\begin{aligned} p(\mathbb{Z}^2, L) &= \#\{v \in \mathbb{Z}^2_{\text{prim}} : \|v\| \leq L\} \\ &= \#\left\{v \in \mathbb{Z}^2_{\text{prim}} : \left\| \frac{1}{L} \cdot v \right\| \leq 1\right\} \\ &= \#\left\{v \in \frac{1}{L} \cdot \mathbb{Z}^2_{\text{prim}} : \|v\| \leq 1\right\}. \end{aligned}$$

It is natural then to study the rescaled lattice $1/L \cdot \mathbb{Z}^2_{\text{prim}} \subseteq \mathbb{R}^2$ as $L \rightarrow \infty$. We do this by means of a measure theoretic approach. For every $L > 0$, consider the counting measure on \mathbb{R}^2 given by

$$\nu_L^{\text{prim}} := \frac{1}{L^2} \cdot \sum_{v \in \mathbb{Z}^2_{\text{prim}}} \delta_{1/L \cdot v}.$$

Notice that if $B \subseteq \mathbb{R}^2$ denotes the unit ball centered at the origin, then

$$\nu_L^{\text{prim}}(B) = \frac{p(\mathbb{Z}^2, L)}{L^2}.$$

This setup reduces the original problem of proving an asymptotic estimate for the counting function $p(\mathbb{Z}^2, L)$ to the problem of understanding the asymptotic behavior of the measures ν_L^{prim} as $L \rightarrow \infty$.

2.3. *The Lebesgue measure.* To develop some intuition, let us first consider a much simpler family of counting measures on \mathbb{R}^2 . For every $L > 0$, consider the counting measure ν_L on \mathbb{R}^2 given by

$$\nu_L := \frac{1}{L^2} \cdot \sum_{v \in \mathbb{Z}^2} \delta_{1/L \cdot v}. \tag{2.2}$$

Denote by ν the standard Lebesgue measure on \mathbb{R}^2 . Based solely on geometric intuition, one expects that the following weak- \star convergence of measures holds:

$$\lim_{L \rightarrow \infty} \nu_L = \nu. \tag{2.3}$$

Indeed, the area of a subset of \mathbb{R}^2 can be computed by counting the number of points of finer and finer lattices that are contained in the set. Recall that the precise definition of weak- \star convergence requires that, for every continuous, compactly supported function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$, the following identity holds:

$$\lim_{L \rightarrow \infty} \int_{\mathbb{R}^2} f \, d\nu_L = \int_{\mathbb{R}^2} f \, d\nu. \tag{2.4}$$

The proof of equation (2.3) will be left as an exercise toward the end of this section. See Exercise 2.15. Let us point out for the moment that the main idea behind the proof of equation (2.3) is the fact that every weak- \star limit point of the sequence of counting measures $(\nu_L)_{L>0}$ is translation invariant.

Exercise 2.2. What would happen with equation (2.3) if we considered a finite index subgroup of \mathbb{Z}^2 instead of all \mathbb{Z}^2 in the definition of the counting measures $(\nu_L)_{L>0}$ in equation (2.2)?

2.4. *Invariance of counting measures.* The discussion above suggests that to prove an analog of equation (2.3) for the counting measures $(\nu_L^{\text{prim}})_{L>0}$, one should try to understand the behavior of the weak- \star limit points of this sequence. *A priori*, one has no reason to expect the limit points of this sequence will be translation invariant, as we will conclude *a fortiori*. Nevertheless, there is still a relevant notion of invariance present in this setting. Indeed, consider the discrete matrix group

$$\text{SL}(2, \mathbb{Z}) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{Mat}_{2 \times 2}(\mathbb{R}) \mid a, b, c, d \in \mathbb{Z}, ab - cd = 1 \right\}$$

acting on \mathbb{R}^2 by linear transformations. The following crucial exercise will provide the notion of invariance we will use to study the counting measures $(\nu_L^{\text{prim}})_{L>0}$.

Exercise 2.3. Show that the $\text{SL}(2, \mathbb{Z})$ orbit of the vector $(1, 0) \in \mathbb{R}^2$ is precisely $\mathbb{Z}_{\text{prim}}^2 \subseteq \mathbb{Z}^2$. Conclude that any weak- \star limit point of the sequence $(\nu_L^{\text{prim}})_{L>0}$ is $\text{SL}(2, \mathbb{Z})$ -invariant. *Hint:* Use Bézout’s identity for greatest common divisors.

Exercise 2.3 ensures that any weak- \star limit point of the sequence $(\nu_L^{\text{prim}})_{L>0}$ is $\text{SL}(2, \mathbb{Z})$ -invariant. Using ergodic theory, we will show this property greatly constrains the possible weak- \star limit points.

2.5. *Ergodic theory.* Let (X, \mathcal{A}) be a measurable space and G be a countable group acting on (X, \mathcal{A}) by measure-preserving transformations. We say a σ -finite measure μ on (X, \mathcal{A}) is G -invariant if $\mu(g.A) = \mu(A)$ for every measurable subset $A \in \mathcal{A}$ and every $g \in G$. Furthermore, we say a G -invariant measure μ on (X, \mathcal{A}) is G -ergodic if it admits no non-trivial G -invariant subsets, that is, if $\mu(A) = 0$ or $\mu(X \setminus A) = 0$ for every $A \in \mathcal{A}$ such that $g.A = A$ for every $g \in G$. Equivalently, a G -invariant measure μ on (X, \mathcal{A}) is ergodic if every G -invariant measurable function is constant almost everywhere with respect to μ . The following exercise characterizes G -invariant measures on (X, \mathcal{A}) that are absolutely continuous with respect to a G -ergodic measure.

Exercise 2.4. Let (X, \mathcal{A}) be a measurable space and G be a countable group acting on (X, \mathcal{A}) by measure-preserving transformations. Suppose that ν is a G -ergodic measure on (X, \mathcal{A}) and that μ is a G -invariant measure on (X, \mathcal{A}) that is absolutely continuous with respect to ν . Show that μ is a non-negative constant multiple of ν . *Hint: Show that the Radon–Nikodym derivative of μ with respect to ν is G -invariant and use the G -ergodicity of ν to show this derivative is constant.*

In the setting of counting problems for primitive lattice points, the following ergodicity result will be relevant for us. Although we will not prove it in this survey, let us at least mention that this result can be proved using the ergodicity of the horocycle flow on the unit tangent of the modular curve $T^1\mathcal{M}_1 := \mathrm{SL}(2, \mathbb{Z}) \backslash \mathrm{SL}(2, \mathbb{R})$; for a proof of this fact, we refer the reader to [BM00].

THEOREM 2.5. *The Lebesgue measure ν on \mathbb{R}^2 is ergodic with respect to the linear action of $\mathrm{SL}(2, \mathbb{Z})$.*

2.6. *Portmanteau’s theorem.* We now state a classic theorem of Portmanteau that gives useful alternative characterizations to the definition of weak- \star convergence in equation (2.4); for a proof of this result, we refer the reader to [Bog18].

THEOREM 2.6. *Let X be a metric space and $(\mu_L)_{L>0}$ be a sequence of locally finite Borel measures on X converging in the weak- \star topology to a Borel measure μ on X . Then, for every open subset $U \subseteq X$,*

$$\mu(U) \leq \liminf_{L \rightarrow \infty} \mu_L(U).$$

Additionally, for every compact subset $K \subseteq X$ such that $\mu(\partial K) = 0$,

$$\lim_{L \rightarrow \infty} \mu_L(K) = \mu(K). \quad (2.5)$$

Exercise 2.7. Assuming the weak- \star convergence in equation (2.3) holds, find counterexamples to the identity in equation (2.5) in Theorem 2.6 when either $K \subseteq \mathbb{R}^2$ is not compact or does not satisfy $\nu(\partial K) = 0$.

2.7. *Limit points of counting measures.* Using Exercises 2.3 and 2.4, and Theorems 2.5 and 2.6, one can greatly restrict the possible weak- \star limit points of the sequence of counting measures $(\nu_L^{\mathrm{prim}})_{L>0}$.

PROPOSITION 2.8. Every weak- \star limit point ν^{prim} of the sequence of counting measures $(\nu_L^{\text{prim}})_{L>0}$ is of the form $\nu^{\text{prim}} = c \cdot \nu$ for some constant $c \geq 0$.

Proof. Let ν^{prim} be a weak- \star limit point of the sequence $(\nu_L^{\text{prim}})_{L>0}$. Exercise 2.3 guarantees that ν^{prim} is $\text{SL}(2, \mathbb{Z})$ -invariant. Theorem 2.5 ensures that the Lebesgue measure ν on \mathbb{R}^2 is ergodic with respect to the linear action of $\text{SL}(2, \mathbb{Z})$. Thus, by Exercise 2.4, to prove $\nu^{\text{prim}} = c \cdot \nu$ for some constant $c \geq 0$, it is enough to check that ν^{prim} is absolutely continuous with respect to ν .

To prove this, let us consider an arbitrary Borel measurable subset $A \subseteq \mathbb{R}^2$ such that $\nu(A) = 0$. Our goal is to show that $\nu^{\text{prim}}(A) = 0$. Let $\delta > 0$ be arbitrary. The outer regularity of the Lebesgue measure ν guarantees that one can find a countable collection of open squares $\{B_i\}_{i \in \mathbb{N}}$ such that

$$A \subseteq \bigcup_{i \in \mathbb{N}} B_i, \quad \sum_{i \in \mathbb{N}} \nu(B_i) \leq \delta. \tag{2.6}$$

To bound $\nu^{\text{prim}}(A)$, let us first give a rough bound of the measure $\nu^{\text{prim}}(B)$ of an arbitrary open square $B \subseteq \mathbb{R}^2$. Denote by $\epsilon > 0$ the side length of B . Notice that, for every $L > 0$,

$$\nu_L^{\text{prim}}(B) \leq \nu_L(B) = \frac{\#(\mathbb{Z}^2 \cap (L \cdot B))}{L^2}. \tag{2.7}$$

The set $L \cdot B \subseteq \mathbb{R}^2$ is an open square of side length $L \cdot \epsilon$. Suppose $L > 0$ is large enough so that this side length satisfies $L \cdot \epsilon \geq 1$. Consider the set $S \subseteq \mathbb{R}^2$ obtained by taking the union of disjoint open squares of side length $1/2$ centered at every point $v \in \mathbb{Z}^2 \cap (L \cdot B)$. This set is contained in an open square $B' \subseteq \mathbb{R}^2$ of side length $L \cdot \epsilon + 1$. It follows that

$$\#(\mathbb{Z}^2 \cap (L \cdot B)) \cdot (1/4) = \nu(S) \leq \nu(B') = (L \cdot \epsilon + 1)^2 \leq 4 \cdot L^2 \cdot \epsilon^2 = 4 \cdot L^2 \cdot \nu(B). \tag{2.8}$$

Putting together equations (2.7) and (2.8), we deduce

$$\limsup_{L \rightarrow \infty} \nu_L^{\text{prim}}(B) \leq 16 \cdot \nu(B). \tag{2.9}$$

We now bound $\nu^{\text{prim}}(A)$. Using the cover in equation (2.6) and the subadditivity of ν^{prim} , we deduce

$$\nu^{\text{prim}}(A) \leq \sum_{i \in \mathbb{N}} \nu^{\text{prim}}(B_i). \tag{2.10}$$

Theorem 2.6 ensures that, for every $i \in \mathbb{N}$,

$$\nu^{\text{prim}}(B_i) \leq \liminf_{L \rightarrow \infty} \nu_L^{\text{prim}}(B_i) \leq \limsup_{L \rightarrow \infty} \nu_L^{\text{prim}}(B_i). \tag{2.11}$$

The identity in equation (2.9) guarantees that, for every $i \in \mathbb{N}$,

$$\limsup_{L \rightarrow \infty} \nu_L^{\text{prim}}(B_i) \leq 16 \cdot \nu(B_i). \tag{2.12}$$

Putting together equations (2.10), (2.11), and (2.12), and using the inequality in equation (2.6), we deduce

$$\nu^{\text{prim}}(A) \leq \sum_{i \in \mathbb{N}} 16 \cdot \nu(B_i) \leq 16 \cdot \delta.$$

As $\delta > 0$ is arbitrary, we conclude $\nu^{\text{prim}}(A) = 0$, thus finishing the proof. □

2.8. The space of unimodular lattices. All that remains is to show that the constant $c \geq 0$ in the conclusion of Proposition 2.8 is positive, independent of the limit point ν^{prim} , and actually equal to $6/\pi^2$. We do this by considering an averaging argument over the space of unimodular lattices of \mathbb{R}^2 up to rotation.

By definition, a lattice $\Lambda \subseteq \mathbb{R}^2$ is the \mathbb{Z} -span of an \mathbb{R} -basis of \mathbb{R}^2 . A marking of a lattice $\Lambda \subseteq \mathbb{R}^2$ is a choice of positively oriented \mathbb{R} -basis (v_1, v_2) of \mathbb{R}^2 such that $\Lambda = \text{span}_{\mathbb{Z}}(v_1, v_2)$. The covolume of a lattice $\Lambda := \text{span}_{\mathbb{Z}}(v_1, v_2) \subseteq \mathbb{R}^2$ is defined as $\text{covol}(\Lambda) := |\det(v_1, v_2)|$. This definition is independent of the choice of marking (v_1, v_2) . A lattice is said to be unimodular if it has unit covolume.

Notice that the special linear group

$$\text{SL}(2, \mathbb{R}) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{Mat}_{2 \times 2}(\mathbb{R}) \mid a, b, c, d \in \mathbb{R}, ab - cd = 1 \right\}$$

acts transitively on the set of unimodular lattices through its linear action on \mathbb{R}^2 . The stabilizer of the integer lattice $\mathbb{Z}^2 \subseteq \mathbb{R}^2$ is the group $\text{SL}(2, \mathbb{Z}) \subseteq \text{SL}(2, \mathbb{R})$. We can thus identify the space of unimodular lattices of \mathbb{R}^2 with the quotient $T^1\mathcal{M}_1 := \text{SL}(2, \mathbb{R})/\text{SL}(2, \mathbb{Z})$. The corresponding identification maps the equivalence class of matrices $[A] \in T^1\mathcal{M}_1$ to the lattice $A \cdot \mathbb{Z}^2 \subseteq \mathbb{R}^2$.

Notice that the special orthogonal group

$$\text{SO}(2) := \left\{ \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \in \text{Mat}_{2 \times 2}(\mathbb{R}) \mid \theta \in [0, 2\pi] \right\}$$

acts on the space of unimodular lattices through its linear action on \mathbb{R}^2 , or, equivalently, by left multiplication on $T^1\mathcal{M}_1$. The corresponding quotient $\mathcal{M}_1 := \text{SO}(2) \backslash \text{SL}(2, \mathbb{R}) / \text{SL}(2, \mathbb{Z})$ can thus be identified with the space of unimodular lattices of \mathbb{R}^2 up to rotation.

It will be convenient to consider the following alternative description of the space of unimodular lattices of \mathbb{R}^2 up to rotation. Denote the upper half-space by

$$\mathbf{H}^2 := \{z \in \mathbb{C} \mid \Im(z) > 0\}.$$

The space \mathbf{H}^2 can be identified with the space $\text{SO}(2) \backslash \text{SL}(2, \mathbb{R})$ of marked unimodular lattices of \mathbb{R}^2 up to rotation via the map which sends $z \in \mathbf{H}^2$ to the marked unimodular lattice $\Lambda(z) := \text{span}_{\mathbb{Z}}(c_z e_1, c_z z) \subseteq \mathbb{R}^2$, where $c_z := \Im(z)^{-1/2} > 0$; essentially, because rotations are allowed, prescribing any vector $z \in \mathbf{H}^2$ generates an arbitrary marked unimodular lattice together with the vector $(1, 0) \in \mathbb{Z}^2$ after appropriately rescaling. The discrete group $\text{SL}(2, \mathbb{Z})$ acts properly discontinuously on \mathbf{H}^2 by Möbius transformations




FIGURE 2. A fundamental domain of the action of $SL(2, \mathbb{Z})$ on \mathbf{H}^2 and its quotient, the modular curve \mathcal{M}_1 (colour online).

in the following way:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot z = \frac{az + b}{cz + d}, \quad z \in \mathbf{H}^2.$$

The identification above intertwines this action with the action of $SL(2, \mathbb{Z})$ on $SO(2) \backslash SL(2, \mathbb{R})$ by right multiplication. Thus, the quotient $\mathcal{M}_1 := \mathbf{H}^2 / SL(2, \mathbb{Z})$, commonly known as the modular curve, can be identified with the space of unimodular lattices of \mathbb{R}^2 up to rotation. See Figure 2 for a representation of this space using a fundamental domain of the action of $SL(2, \mathbb{Z})$ on \mathbf{H}^2 .

Consider the Lebesgue class measure μ on \mathbf{H}^2 given in coordinates $z = x + iy$ by

$$\mu = \frac{dx \, dy}{y^2}. \tag{2.13}$$

Alternatively, after identifying $\mathbf{H}^2 = SO(2) \backslash SL(2, \mathbb{R})$, the measure μ is obtained by integrating the $SO(2)$ -direction of the Haar measure on $SL(2, \mathbb{R})$. A direct computation shows that μ is preserved by the action of $SL(2, \mathbb{Z})$ on \mathbf{H}^2 ; see Exercise 2.9. Denote by $\widehat{\mu}$ the local pushforward of μ to the quotient $\mathcal{M}_1 := \mathbf{H}^2 / SL(2, \mathbb{Z})$. More explicitly, this is the measure on \mathcal{M}_1 obtained by restricting μ to the fundamental domain of the action of $SL(2, \mathbb{Z})$ on \mathbf{H}^2 in Figure 2. Local pushforwards of measures will be discussed in more detail in §3 following an approach independent of fundamental domains. Nevertheless, the explicit fundamental domain in Figure 2 is useful for computing the total mass $\widehat{\mu}(\mathcal{M}_1)$, as suggested in the following exercise.

Exercise 2.9. Using change of variables, show that the measure μ is preserved by the action of $SL(2, \mathbb{Z})$ on \mathbf{H}^2 . Using Figure 2, show that $\widehat{\mu}(\mathcal{M}_1) = \pi/3$.

2.9. *Averaging over the space of unimodular lattices.* Let $\Lambda \in \mathcal{M}_1$ be a unimodular lattice of \mathbb{R}^2 . A vector $v \in \Lambda$ is said to be primitive if it cannot be written as a non-negative integer multiple of another vector of Λ . Denote by $\Lambda_{\text{prim}} \subseteq \Lambda$ the subset of all primitive vectors of Λ . Recall that $\|\cdot\|$ denotes the Euclidean norm of \mathbb{R}^2 . For every $L > 0$, we

consider the counting function

$$p(\Lambda, L) := \{v \in \Lambda_{\text{prim}} : \|v\| \leq L\}. \tag{2.14}$$

As one might already expect after considering the case of the integer lattice $\mathbb{Z}^2 \subseteq \mathbb{R}^2$, explicitly describing $p(\Lambda, L)$ for a given unimodular lattice $\Lambda \in \mathcal{M}_1$ as a function of L is a particularly hard problem. Nevertheless, and perhaps surprisingly, the average

$$\int_{\mathcal{M}_1} p(\Lambda, L) d\widehat{\mu}(\Lambda)$$

can be computed explicitly, as the next result shows.

PROPOSITION 2.10. *The following integration formula holds:*

$$\int_{\mathcal{M}_1} p(\Lambda, L) d\widehat{\mu}(\Lambda) = 2 \cdot L^2.$$

In §4, we will prove this result using a general local change of variables formula; see Exercise 4.3. From a more classical point of view, Proposition 2.10 is a consequence of Siegel’s famous integration formulas over \mathcal{M}_1 [Sie45].

To prove Theorem 2.1, we will need a more uniform control on the integrability of the counting functions $p(\Lambda, L)$ as $L \rightarrow \infty$. To this end, we consider the function $u : \mathcal{M}_1 \rightarrow \mathbb{R}$ given by

$$u(\Lambda) := \sup_{v \in \Lambda} \frac{1}{\|v\|}.$$

Exercise 2.11. Show there exists a constant $C > 0$ such that for every $\Lambda \in \mathcal{M}_1$ and every $L > 0$,

$$p(\Lambda, L) \leq C \cdot L^2 \cdot u(\Lambda).$$

Additionally, show that the function $u : \mathcal{M}_1 \rightarrow \mathbb{R}$ is integrable with respect to the measure $\widehat{\mu}$, that is,

$$\int_{\mathcal{M}_1} u(\Lambda) d\widehat{\mu}(\Lambda) < \infty.$$

2.10. *Convergence and compactness.* Before proceeding with the proof of Theorem 2.1, let us discuss a couple of basic convergence and compactness criteria. To prove the sequence of counting measure $(\nu_L^{\text{prim}})_{L>0}$ on \mathbb{R}^2 converges in the weak- \star topology, we will use the following convergence criterion.

Exercise 2.12. Let X be a metric space. Show that a sequence $(x_L)_{L>0}$ in X converges to $x \in X$ if and only if every subsequence of $(x_L)_{L>0}$ has a subsubsequence converging to x .

The following compactness criterion for the weak- \star topology serves as a convenient tool for extracting convergent subsequences of locally finite measures on metric spaces. This result is a consequence of the well-known Banach–Alaoglu theorem in functional analysis.

THEOREM 2.13. *Let X be a metric space and $(\mu_n)_{n \in \mathbb{N}}$ be a sequence of locally finite measures on X . Suppose that the sequence $(\mu_n(K))_{n \in \mathbb{N}}$ is bounded for every $K \subseteq X$ compact. Then, the sequence $(\mu_n)_{n \in \mathbb{N}}$ has a subsequence converging in the weak- \star topology to a locally finite measure on X .*

2.11. Equidistribution of primitive integer points. We are finally ready to prove that the sequence of counting measure $(\nu_L^{\text{prim}})_{L>0}$ on \mathbb{R}^2 converges in the weak- \star topology to a constant multiple of the Lebesgue measure ν . We will later deduce Theorem 2.1 directly from this result.

THEOREM 2.14. *With respect to the weak- \star topology for measures on \mathbb{R}^2 ,*

$$\lim_{L \rightarrow \infty} \nu_L^{\text{prim}} = \frac{6}{\pi^2} \cdot \nu.$$

Proof. By Exercise 2.12 applied to the space of regular measures on \mathbb{R}^2 with the weak- \star topology, it is enough to show every subsequence of $(\nu_L^{\text{prim}})_{L>0}$ has a subsubsequence converging to $(6/\pi^2) \cdot \nu$ in the weak- \star topology. Theorem 2.13 and the bound in equation (2.9) ensure that every subsequence of $(\nu_L^{\text{prim}})_{L>0}$ has a subsubsequence $(\nu_{L_k}^{\text{prim}})_{k \in \mathbb{N}}$ converging in the weak- \star topology to a locally finite measure ν^{prim} on \mathbb{R}^2 . By Proposition 2.8, it must be the case that $\nu^{\text{prim}} = c \cdot \nu$ for some constant $c \geq 0$. It is enough then to show that $c = 6/\pi^2$, independent of the subsubsequence considered. To prove this identity holds, we compute the following limit in two different ways:

$$\lim_{k \rightarrow \infty} \int_{\mathcal{M}_1} \frac{p(\Lambda, L_k)}{L_k^2} d\widehat{\mu}(\Lambda).$$

The first computation is immediate. Directly from Proposition 2.10, we deduce

$$\lim_{k \rightarrow \infty} \int_{\mathcal{M}_1} \frac{p(\Lambda, L_k)}{L_k^2} d\widehat{\mu}(\Lambda) = 2. \tag{2.15}$$

The second computation requires more work. For every matrix $A \in \text{SL}(2, \mathbb{R})$, consider the subset

$$B_A := \{v \in \mathbb{R}^2 : \|A \cdot v\| \leq 1\}.$$

Notice that $\nu(B_A) = 1$ for every $A \in \text{SL}(2, \mathbb{R})$. Notice also that for every unimodular lattice $\Lambda := A \cdot \mathbb{Z}^2 \in \mathcal{M}_1$ with $A \in \text{SL}(2, \mathbb{R})$, and every $k \in \mathbb{N}$,

$$\frac{p(\Lambda, L_k)}{L_k^2} = \nu_{L_k}^{\text{prim}}(B_A).$$

In this setting, as $\lim_{k \rightarrow \infty} \nu_{L_k}^{\text{prim}} = c \cdot \nu$ in the weak- \star topology, Theorem 2.6 ensures that

$$\lim_{k \rightarrow \infty} \frac{p(\Lambda, L_k)}{L_k^2} = \lim_{k \rightarrow \infty} \nu_{L_k}^{\text{prim}}(B_A) = c \cdot \nu(B_A) = c \cdot \pi.$$

Exercise 2.11 now allows us to use the dominate convergence theorem to deduce

$$\lim_{k \rightarrow \infty} \int_{\mathcal{M}_1} \frac{p(\Lambda, L_k)}{L_k^2} d\widehat{\mu}(\Lambda) = c \cdot \frac{\pi^2}{3}. \tag{2.16}$$

Putting together equations (2.15) and (2.16), we obtain

$$c \cdot \frac{\pi^2}{3} = 2.$$

Solving for c , we conclude $c = 6/\pi^2$, and thus finish the proof. □

Exercise 2.15. Recall the definition of the sequence of counting measures $(\nu_L)_{L>0}$ on \mathbb{R}^2 in equation (2.2). Using the methods introduced in the proof of Theorem 2.14, show that equation (2.3) holds, that is, show that with respect to the weak- \star topology for measures on \mathbb{R}^2 ,

$$\lim_{L \rightarrow \infty} \nu_L = \nu.$$

2.12. *Counting primitive integer points.* We are now ready to prove Theorem 2.1, the main result of this section. Let us restate this theorem in the following equivalent way.

THEOREM 2.16. *The following asymptotic estimate holds:*

$$\lim_{L \rightarrow \infty} \frac{p(\mathbb{Z}^2, L)}{L^2} = \frac{6}{\pi}.$$

Proof. Recall that if $B \subseteq \mathbb{R}^2$ is the Euclidean unit ball centered at the origin, then, for every $L > 0$,

$$\frac{p(\mathbb{Z}^2, L)}{L^2} = \nu_L^{\text{prim}}(B).$$

Using Theorems 2.14 and 2.6, we conclude

$$\lim_{L \rightarrow \infty} \frac{p(\mathbb{Z}^2, L)}{L^2} = \lim_{L \rightarrow \infty} \nu_L^{\text{prim}}(B) = \frac{6}{\pi^2} \cdot \nu(B) = \frac{6}{\pi}. \quad \square$$

Exercise 2.17. Using Theorem 2.14, show that for every unimodular lattice $\Lambda \in \mathcal{M}_1$,

$$\lim_{L \rightarrow \infty} \frac{p(\Lambda, L)}{L^2} = \frac{6}{\pi}.$$

Remark 2.18. Comparing Exercise 2.17 to Theorem 1.1, one can see that, very different to the case of unimodular lattices, the leading constant in Mirzakhani’s counting formula does depend on the underlying hyperbolic surface. Later, in Exercises 5.13 and 5.14, a more explicit description of this constant will be provided.

3. Hyperbolic surfaces, Teichmüller spaces, and simple closed curves

3.1. *Outline of this section.* In this section, we cover the background material needed to understand the proof of Theorem 1.1. The focus will be in developing geometric intuition rather than on giving complete proofs. Unless otherwise stated, all surfaces considered will be connected and orientable. Two excellent references for the topics that will be covered in this section are [FM12, Mar23].




FIGURE 3. The geodesics of the hyperbolic plane.




FIGURE 4. Right-angled hyperbolic hexagons are rigid.

3.2. *The hyperbolic plane.* By definition, the hyperbolic plane \mathbf{H}^2 is the unique, up to isometry, two-dimensional simply connected Riemannian manifold of constant sectional curvature -1 . Recall that the hyperbolic plane can be modeled on the upper half space $\{z \in \mathbf{C} \mid \Im(z) > 0\}$ by endowing it with the Riemannian metric

$$g := \frac{dx^2 + dy^2}{y^2}.$$

The geodesics of this metric are the lines and half circles of the upper half space perpendicular to the the real axis $\mathbb{R} \subseteq \mathbf{C}$. See Figure 3. The orientation-preserving isometries of this metric can be identified with the group $\text{PSL}(2, \mathbb{R}) = \text{SL}(2, \mathbb{R})/\{\pm \text{Id}\}$ acting on \mathbf{H}^2 by Möbius transformations:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} . z := \frac{az + b}{cz + d} \quad \text{for all } z \in \mathbf{H}.$$

This group acts simply transitively on the unit tangent bundle of \mathbf{H}^2 ; through this action, one can realize \mathbf{H}^2 as the symmetric space $\text{SO}(2)\backslash\text{SL}(2, \mathbb{R})$. Such a large isometry group should hint at a rigid geometry: it becomes hard to distinguish objects up to isometry. The next exercise is an important manifestation of this idea.

Exercise 3.1. Show that for every $(a, b, c) \in (\mathbb{R}_+)^3$, there exists a unique, up to isometry, hyperbolic right-angled hexagon with alternating edge lengths (a, b, c) . *Hint: Consider a configuration as in Figure 4 and study how the length $c(y)$ varies as the parameter y varies.*



(a) Every geodesic has a collar. (b) Shorter geodesics have longer collars.

FIGURE 5. The collar lemma (colour online).

Despite Exercise 3.1, hyperbolic geometry in dimension 2 remains quite flexible, as we will see below. In higher dimensions, the picture becomes incredibly rigid; curious readers are invited to investigate Mostow's rigidity theorem [Mos68].

3.3. Hyperbolic surfaces. A hyperbolic surface is a surface whose geometry is locally modeled on \mathbf{H}^2 . More concretely, a hyperbolic surface X is a surface endowed with an atlas of charts to \mathbf{H}^2 whose transition functions are restrictions of isometries of \mathbf{H}^2 . Pulling back the metric of \mathbf{H}^2 via these charts yields a metric of constant curvature equal to -1 on X . A geodesic of X is a geodesic of this metric. Equivalently, a geodesic of X is a curve which is mapped to geodesics of \mathbf{H}^2 via local charts. Every closed curve on X can be tightened, that is, homotoped, to a unique geodesic representative.

We will not spend much time discussing the many interesting features of hyperbolic surfaces, but let us highlight one fact that is very useful to keep in mind for the sake of geometric intuition. The following fact is known as the collar lemma: every simple closed geodesic on a hyperbolic surface has a collar, that is, an embedded annular neighborhood, whose width goes to infinity as the length of the geodesic goes to zero. See Figure 5. For a more precise version of this fact, see Lemma 5.7.

3.4. Teichmüller space. All hyperbolic surfaces of a given genus fit together nicely into a moduli space. To keep track of how any pair of hyperbolic surfaces looks with respect to each other, we will record more information than just their position in moduli space. This leads us to introduce Teichmüller spaces. We will later see how to recover moduli spaces as quotients of Teichmüller spaces.

Throughout the rest of this survey, we use the words 'homotopy' and 'isotopy' interchangeably. This does not pose an issue because we are working in dimension 2; see [FM12, Ch. 1] for a detailed discussion on this highly non-trivial matter.

For the rest of this lecture, we fix an integer $g \geq 2$ and a connected, oriented, closed surface S_g of genus g . The Teichmüller space \mathcal{T}_g of marked hyperbolic structures on S_g is the space of all pairs (X, φ) , where X is an oriented hyperbolic surface and $\varphi: S_g \rightarrow X$ is an orientation-preserving homeomorphism, modulo the equivalence relation $(X_1, \varphi_1) \sim (X_2, \varphi_2)$ if and only if there exists an orientation-preserving isometry $I: X_1 \rightarrow X_2$ isotopic to $\varphi_2 \circ \varphi_1^{-1}$. In most situations, we will omit the marking of a point $(X, \varphi) \in \mathcal{T}_g$ and denote it simply by $X \in \mathcal{T}_g$.




FIGURE 6. Dehn twist in an annular neighborhood of a simple closed curve γ (in green) (colour online).

Roughly speaking, a point in Teichmüller space does not only keep track of a hyperbolic structure on S_g but also of how S_g is ‘wearing’ that hyperbolic structure. This allows us to communicate marked hyperbolic structures on S_g in a very explicit way. For instance, given a marked hyperbolic structure $(X, \varphi) \in \mathcal{T}_g$, any parameterized closed curve γ on S_g can be canonically identified with the closed geodesic obtained by tightening the closed curve $\varphi(\gamma)$ on X to its unique geodesics representative. We denote the length of this representative by $\ell_\gamma(X)$.

3.5. *The mapping class group.* The mapping class group of S_g , denoted Mod_g , is the group of orientation-preserving homeomorphisms of S_g up to homotopy. More explicitly,

$$\text{Mod}_g := \text{Homeo}^+(S_g)/\text{Homeo}_0(S_g).$$

This group acts naturally on \mathcal{T}_g by changing the markings: for every $(X, \varphi) \in \mathcal{T}_g$ and every $\phi \in \text{Mod}_g$,

$$\phi.(X, \varphi) = (X, \varphi \circ \phi^{-1}).$$

We think of Mod_g acting on \mathcal{T}_g as an analog of $\text{SL}(2, \mathbb{Z})$ acting on \mathbf{H}^2 .

Exercise 3.2. Let $(X, \varphi) \in \mathcal{T}_g$ be a marked hyperbolic structure on S_g . Show there exists a natural one-to-one correspondence between the Mod_g stabilizer of (X, φ) and the set of isometries of X .

A particularly important family of mapping classes are Dehn twists. Given a simple closed curve γ on S_g , the Dehn twist T_γ of S_g along γ is the mapping class which leaves the surface S_g untouched except for an embedded annular neighborhood of γ which gets twisted to the right, with respect to the orientation of S_g , by a full rotation. See Figure 6.

3.6. *Moduli space of hyperbolic surfaces.* As Mod_g acts on \mathcal{T}_g by changing the markings, one should expect that, if one quotients \mathcal{T}_g by this action, one should get a space grouping together all (unmarked) hyperbolic surfaces of genus g . It turns out that Mod_g acts on \mathcal{T}_g properly discontinuously and thus the corresponding quotient behaves nicely. The quotient $\mathcal{M}_g := \mathcal{T}_g/\text{Mod}_g$ is the moduli space of genus g hyperbolic surfaces. By uniformization, this space can be canonically identified with the moduli space of genus g Riemann surfaces you might have seen in algebraic geometry.

3.7. *Fenchel–Nielsen coordinates.* Notice that any orientable topological surface of genus $g \geq 2$ can be constructed by gluing $2g - 2$ pairs of pants, that is, spheres with three




FIGURE 7. Pair of pants decomposition of a genus 2 surface (colour online).




FIGURE 8. Cutting a hyperbolic pair of pants into isometric right-angled hexagons (colour online).

boundary components, along their boundaries. See Figure 7 for an example. A similar construction can also be considered for hyperbolic surfaces. Indeed, cutting an orientable hyperbolic surface X of genus $g \geq 2$ along any collection of $3g - 3$ disjoint simple closed geodesics yields $2g - 2$ hyperbolic pairs of pants with geodesic boundary components. This class of pairs of pants is very rigid, as the following exercise shows.

Exercise 3.3. Show that, for every $a, b, c \in \mathbb{R}_+$, there exists a unique, up to isometry, hyperbolic pair of pants with geodesic boundary components of lengths a, b, c . *Hint:* Cutting a hyperbolic pair of pants with geodesic boundary components along the orthogeodesics joining its boundary components yields a pair of isometric hyperbolic right-angled hexagons. See Figure 8.

By Exercise 3.3, the hyperbolic pairs of pants obtained by cutting an orientable hyperbolic surface X of genus $g \geq 2$ along a maximal collection of disjoint simple closed geodesics are determined, up to isometry, by the lengths of the geodesics one cuts along. Gluing back these pairs of pants following the same original pattern allows us to recover X . One needs to be careful at this point as there are several ways in which one can glue back these pants along their cuffs. Indeed, for each geodesic one cutted X along, there is a full circle worth of different twists with which one can glue back the adjacent pairs of pants. If one is interested in marked hyperbolic surfaces rather than just hyperbolic surfaces, there is actually a full line worth of different twists one can glue with respect to.

More concretely, one can deform (marked) oriented hyperbolic surfaces using the following operation: given a (marked) oriented hyperbolic surface X , a simple closed geodesic γ on X , and $t \in \mathbb{R}$, cut X along γ and glue the resulting surface back along γ with a twist of t units of hyperbolic length to the right with respect to the orientation of X . See Figure 9. This operation is known as a Fenchel–Nielsen twist. Given a simple closed curve γ on S_g and a marked hyperbolic structure on $X \in \mathcal{T}_g$, the point $T_\gamma.X \in \mathcal{T}_g$ is equal




FIGURE 9. Fenchel–Nielsen twist along a simple closed curve (in red) (colour online).

to the marked hyperbolic structure obtained by doing a Fenchel–Nielsen twist of parameter $t = \ell_\gamma(X)$ along the unique geodesic representative of γ on X .

The discussion above leads us to introduce Fenchel–Nielsen coordinates. A pair of pants decomposition of S_g is a maximal collection of disjoint simple closed curves on S_g . Fix a pair of pants decomposition $\mathcal{P} := (\gamma_i)_{i=1}^{3g-3}$ of S_g . Given a marked hyperbolic structure $X \in \mathcal{T}_g$, for every $i \in \{1, \dots, 3g - 3\}$, denote $\ell_i(X) := \ell_{\gamma_i}(X)$ and let $\tau_i(X)$ be the twist of X at the geodesic representative of γ_i . Defining what it means to have zero twist is a rather technical issue which we avoid in this survey; the interested reader is invited to consult [FM12, Ch. 10]. The parameters $(\ell_i, \tau_i)_{i=1}^{3g-3} \in (\mathbb{R}_+ \times \mathbb{R})^{3g-3}$ provide global coordinates for \mathcal{T}_g known as Fenchel–Nielsen coordinates. In particular, \mathcal{T}_g is homeomorphic to an open ball of dimension $6g - 6$. For every $i \in \{1, \dots, 3g - 3\}$, the action of the Dehn twist T_{γ_i} in Fenchel–Nielsen coordinates corresponds to leaving all coordinates constant except for τ_i , which gets changed by the transformation $\tau_i \mapsto \tau_i + \ell_i$.

Exercise 3.4. Using Fenchel–Nielsen coordinates, the collar lemma, Dehn twists, and your geometric intuition, come up with an explanation of the following fact: a marked hyperbolic structure on Teichmüller space escapes to infinity, that is, leaves every compact set, if and only if some fixed geodesic becomes arbitrarily long.

Let us highlight an interesting feature of a geodesic pair of pants decompositions of hyperbolic surfaces. By the work of Bers [Ber85], every closed, connected, oriented hyperbolic surface admits a geodesic pair of pants decompositions whose cuffs have lengths bounded by a linear function of g . Buser conjectured this result can be improved to ensure the cuffs have lengths bounded by a linear function of \sqrt{g} [Bus92]. This conjecture remains open until today.

Exercise 3.5. Show that there are finitely many pairs of pants decompositions of S_g up to homeomorphism. Can you give an asymptotic estimate for the number of such equivalence classes? *Hint:* If I_N denotes the number of isomorphism classes of cubic multigraphs on N vertices, then, as $N \rightarrow \infty$,

$$I_N \sim \frac{e^2}{\sqrt{\pi N}} \cdot \left(\frac{3N}{4e}\right)^{N/2}.$$

Exercise 3.6. Use Bers’s theorem and Exercise 3.5 to show that for every $\epsilon > 0$, the subset $K_\epsilon \subseteq \mathcal{M}_g$ of genus g hyperbolic surfaces, all of whose closed geodesics have length $\geq \epsilon$, is compact. This result is commonly known as Mumford’s compactness criterion. *Hint:* Using

Fenchel–Nielsen coordinates, write $K_\epsilon \subseteq \mathcal{M}_g$ as a union of finitely many projections of compact subsets of \mathcal{T}_g .

3.8. *The Weil–Petersson measure on Teichmüller space.* The discussion above suggests considering the following volume form on Teichmüller space: given a set of Fenchel–Nielsen coordinates $(\ell_i, \tau_i)_{i=1}^{3g-3} \in (\mathbb{R}_+ \times \mathbb{R})^{3g-3}$, denote by v_{wp} the Weil–Petersson volume form on \mathcal{T}_g given by

$$v_{\text{wp}} := \bigwedge_{i=1}^{3g-3} d\ell_i \wedge d\tau_i. \quad (3.1)$$

This volume form is independent of the choice of Fenchel–Nielsen coordinates and, in particular, is mapping class group invariant. This fact is a consequence of a deep result of Wolpert known as Wolpert’s magic formula [Wol83]. We refer to the corresponding measure μ_{wp} as the Weil–Petersson measure of \mathcal{T}_g .

3.9. *Local pushforwards of measures.* In the upcoming discussion, we make use of the following abstract measure theoretic construction. Let X be a locally compact, Hausdorff, second countable topological space endowed with a properly discontinuous action of a discrete group G . Notice that X/G is also locally compact, Hausdorff, and second countable. Let $\pi : X \rightarrow X/G$ be the corresponding quotient map. As the action of G on X is properly discontinuous, one can cover X by open subsets $U \subseteq X$ invariant under the action of finite subgroups $\Gamma_U < G$ and such that $gU \cap U = \emptyset$ for all $g \in G \setminus \Gamma_U$. Open sets $U/\Gamma_U \subseteq X/G$ of this form will be referred to as well covered.

Given a locally finite G -invariant Borel measure μ on X , there exists a unique locally finite Borel measure $\pi_{\#}\mu$ on X/G satisfying the following property: if $U/\Gamma_U \subseteq X/G$ is a well-covered open set,

$$(\pi_{\#}\mu)|_{U/\Gamma_U} = \frac{1}{|\Gamma_U|} \cdot (\pi|_U)_*(\mu|_U), \quad (3.2)$$

where $(\pi|_U)_*(\mu|_U)$ denotes the usual pushforward of the measure $\mu|_U$ through the map $\pi|_U$. We refer to the measure $\pi_{\#}\mu$ as the local pushforward of μ to X/G .

Exercise 3.7. Check that the definition of $\pi_{\#}\mu$ gives rise to a unique well-defined measure on X/G .

3.10. *The Weil–Petersson measure on moduli space.* Recall that the quotient $\mathcal{M}_g := \mathcal{T}_g/\text{Mod}_g$ is the moduli space of genus g hyperbolic surfaces. As the action of Mod_g on \mathcal{T}_g is properly discontinuous and preserves the Weil–Petersson measure μ_{wp} , one can consider the local pushforward of μ_{wp} to \mathcal{M}_g . We denote this measure by $\widehat{\mu}_{\text{wp}}$ and refer to it as the Weil–Petersson measure of \mathcal{M}_g .

Exercise 3.8. Using Bers’s theorem and Exercise 3.5, show that the Weil–Petersson measure $\widehat{\mu}_{\text{wp}}$ on \mathcal{M}_g is finite. Can you give a bound on the total Weil–Petersson measure $\widehat{\mu}_{\text{wp}}(\mathcal{M}_g)$? *Hint: Follow a similar approach as in Exercise 3.6.*

Mirzakhani and Zograf [MZ15] proved asymptotic estimates for the total Weil–Petersson volumes $\widehat{\mu}_{\text{wp}}(\mathcal{M}_g)$ as $g \rightarrow \infty$ that wildly differ from the rough upper bound one can obtain by following the hint in Exercise 3.8. The finiteness of the Weil–Petersson measure on moduli space suggests studying random hyperbolic surfaces sampled according to this measure. This has become a very active field of study in recent years. As a starting point for curious readers, we recommend [LW24, Mir13, Mon22, MP19, MT20, MZ15].

3.11. *Simple closed multi-curves.* We refer to a parameterized simple closed curve on S_g up to homotopy and orientation reversal as a simple closed curve. We will always assume simple closed curves on S_g are homotopically non-trivial. The group Mod_g acts naturally on these equivalence classes. Given a marked hyperbolic structure $X \in \mathcal{T}_g$, every simple closed curve on S_g corresponds to a unique simple closed geodesic on X . Through this perspective, counting problems for simple closed geodesics on X can be easily reformulated as counting problems for simple closed curves on S_g .

Exercise 3.9. The $9g - 9$ theorem, see for instance [FM12, Theorem 10.7], guarantees that a marked hyperbolic structure $X \in \mathcal{T}_g$ is completely determined by its simple marked length spectrum, that is, by the function which to every simple closed curve γ on S_g assigns the length $\ell_\gamma(X)$ of its unique geodesic representative with respect to X . Using this theorem and Dehn–Thurston coordinates, show that the kernel of the action of Mod_g on \mathcal{T}_g is equal to the kernel of the action of Mod_g on the set of simple closed curves on S_g .

More generally, a simple closed multi-curve on S_g is a formal sum $\gamma = \sum_{i=1}^k a_i \gamma_i$ of distinct simple closed curves on S_g that can be realized disjointly, together with real positive weights $a_i \in \mathbb{R}_+$. Given $X \in \mathcal{T}_g$, we define $\ell_\gamma(X) := \sum_{i=1}^k a_i \ell_{\gamma_i}(X)$. An integral simple closed multi-curve on S_g is a simple closed multi-curve all of whose weights are integral. The group Mod_g acts naturally on simple closed multi-curves by acting on each of the components of the formal sum. We think of Mod_g acting on integral simple closed multi-curves as an analog of $\text{SL}(2, \mathbb{Z})$ acting on \mathbb{Z}^2 .

We say two simple closed multi-curves on S_g have the same topological type if they belong to the same mapping class group orbit. More generally, we say two simple closed multi-curves on homeomorphic surfaces have the same topological type if there exists a homeomorphism between the surfaces mapping one simple closed multi-curve to the other. See Figure 10 for examples of simple closed curves of different topological types. We think of integral simple closed multi-curves on S_g of a fixed topological type as an analog of $\mathbb{Z}_{\text{prim}}^2$.

3.12. *Measured geodesic laminations.* Just as the integer lattice \mathbb{Z}^2 sits inside the continuum \mathbb{R}^2 , we would like to have a continuum interpolating between integral simple closed multi-curves on S_g . Although we will not need it explicitly, let us give a geometric description of how objects in this continuum look like. Fix a marked hyperbolic structure $X \in \mathcal{T}_g$. A geodesic lamination on X is a closed subset of X that can be written as a disjoint union of simple geodesics. The most basic example of a geodesic lamination is a union of disjoint simple closed geodesics. See Figure 11 for a more complicated example. Very




FIGURE 10. Simple closed curves of different topological types on a genus 2 surface (colour online).




FIGURE 11. A non-trivial example of a geodesic lamination on a genus 2 surface (colour online).

commonly, the intersection of a geodesic lamination with a transverse arc is a fractal set, making it hard to draw non-trivial examples.

A measured geodesic lamination is a geodesic lamination endowed with a fully supported invariant transverse measure. The transverse measure assigns a finite Borel measure to every arc transverse to the lamination. This assignment is invariant under splitting of arcs and homotopies preserving the leaves of the lamination. Not every lamination admits a fully supported invariant transverse measure.

Exercise 3.10. Show that the geodesic lamination in Figure 11 does not admit a fully supported invariant transverse measure. *Hint:* Consider an arc going across the middle closed geodesic.

The different spaces of measured geodesic laminations obtained as X varies over \mathcal{T}_g can be canonically identified to each other. We will denote by \mathcal{ML}_g any such space and refer to it as the space of measured geodesic laminations on S_g . Simple closed multi-curves embed naturally into \mathcal{ML}_g by considering their geodesic representatives and endowing them with appropriately weighted transverse Dirac measures. We denote by $\mathcal{ML}_g(\mathbb{Z}) \subseteq \mathcal{ML}_g$ the set of integral simple closed multi-curves on S_g .

The space \mathcal{ML}_g can be topologized in such a way that rationally weighted simple closed multi-curves on S_g are dense in it. The action of Mod_g on simple closed multi-curves extends continuously to an action on \mathcal{ML}_g . We think of Mod_g acting on \mathcal{ML}_g as an analog of $\text{SL}(2, \mathbb{Z})$ acting on \mathbb{R}^2 . By the work of Thurston [Thu80], the length $\ell_\lambda(X) > 0$ of a measured geodesic lamination $\lambda \in \mathcal{ML}_g$ with respect to a marked hyperbolic structure $X \in \mathcal{T}_g$ can be defined in a unique continuous way extending the definition on simple closed multi-curves introduced above.

3.13. *Dehn–Thurston coordinates.* In a similar spirit to how hyperbolic surfaces can be constructed by gluing hyperbolic pairs of pants with geodesic boundary components, integral simple closed multi-curves on surfaces can be constructed by gluing simple arc




FIGURE 12. The six isotopy classes of simple arcs in a pair of pants (colour online).

systems on pairs of pants. By an arc on a pair of pants, we will always mean an arc joining two of its boundary components. Every simple arc on a pair of pants is isotopic to exactly one of the six arcs represented in Figure 12; here, we allow the isotopy to move points on the boundary of the pair of pants. Every simple arc system on a pair of pants is isotopic to an arc system constructed by taking disjoint parallel copies of these arcs. Given an assignment of non-negative integers $(a, b, c) \in \mathbb{N}$ to the boundary components of a pair of pants satisfying $a + b + c \in 2\mathbb{N}$, there exists a unique, up to isotopy, arc system realizing (a, b, c) as the number of intersections of the arc system with the boundary components.

Given simple arc systems on pairs of pants glued along two of their boundary components, these arc systems can be glued together if and only if the number of times they intersect the glued boundary components is the same. There exist \mathbb{Z} many different ways of gluing these arc systems depending on how much we twist one with respect to the other before gluing them together.

Let $\mathcal{P} := (\gamma_i)_{i=1}^{3g-3}$ be a pair of pants decomposition of S_g . Following the discussion above, we can parameterize integral simple closed multi-curves on S_g in terms of intersection numbers $m_i \in \mathbb{N}$ and twisting numbers $t_i \in \mathbb{Z}$ with respect to the components of \mathcal{P} . Consider the parameter space

$$\Sigma := \prod_{i=1}^{3g-3} (\mathbb{R}_{\geq 0} \times \mathbb{R}) / \sim,$$

where \sim denotes the equivalence relation on $\mathbb{R}_{\geq 0} \times \mathbb{R}$ identifying $(0, t) \sim (0, -t)$ for every $t \in \mathbb{R}$. The parameters $(m_i, t_i)_{i=1}^{3g-3}$ give rise to a bijection between the set of integral simple closed multi-curves on S_g and the set of integral points in Σ such that for every complementary region R of $S_{g,n} \setminus \mathcal{P}$, the parameters m_i corresponding to components γ_i of \mathcal{P} bounding R add up to an even number; this parity restriction comes from the fact that every strand of a multi-curve that enters a pair of pants must also exit it. If $m_i = 0$ and $t_i = t \in \mathbb{Z}$, the corresponding simple closed multi-curve has $|t|$ disjoint parallel copies of γ_i . We refer to the parameters $(m_i, t_i)_{i=1}^{3g-3}$ as a set of Dehn–Thurston coordinates.

Although we have not defined twisting numbers precisely, we can easily describe the effect of changing them by a given amount. See Figure 13 for an example. In particular, for every $i \in \{1, \dots, 3g - 3\}$, the action of the Dehn twist T_{γ_i} in Dehn–Thurston coordinates corresponds to leaving all coordinates constant except for t_i which gets changed by the transformation $t_i \mapsto t_i + m_i$.



FIGURE 13. The effect of changing a twist coordinate by $t_i \mapsto t_i + 1$ (colour online).

By the work of Thurston [Thu80], the parameterization of $\mathcal{ML}_g(\mathbb{Z})$ provided by any set of Dehn–Thurston coordinates extends to a homeomorphism between \mathcal{ML}_g and the parameter space Σ . This shows in particular that \mathcal{ML}_g is homeomorphic to \mathbb{R}^{6g-6} . For more details on the definition and properties of Dehn–Thurston coordinates, we refer the reader to [PH92, §1.2].

3.14. *The Thurston measure.* The Thurston measure μ_{Thu} on \mathcal{ML}_g is the pullback of the Lebesgue measure on the parameter space Σ of any set of Dehn–Thurston coordinates. This measure is well defined, that is, independent of the choice of Dehn–Thurston coordinates. Indeed, in analogy with the case of the Lebesgue measure, one can characterize μ_{Thu} as the weak- \star limit of a family of rescaled integral simple closed multi-curve counting measures. Consider the natural \mathbb{R}_+ action on \mathcal{ML}_g scaling transverse measures. This action corresponds to the natural scaling action of \mathbb{R}_+ on Σ via the identification by Dehn–Thurston coordinates. For every $L > 0$, consider the rescaled counting measure

$$\mu_L := \frac{1}{L^{6g-6}} \sum_{\gamma \in \mathcal{ML}_g(\mathbb{Z})} \delta_{1/L \cdot \gamma}.$$

Exercise 3.11. Show that the sequence of rescaled counting measures $(\mu_L)_{L>0}$ on \mathcal{ML}_g converges in the weak- \star topology as $L \rightarrow \infty$ to the pullback of the Lebesgue measure on Σ under the identification with \mathcal{ML}_g induced by any set Dehn–Thurston coordinates.

Directly from Exercise 3.11, we deduce μ_{Thu} is invariant with respect to the Mod_g action on \mathcal{ML}_g . The definition of μ_{Thu} in terms of Dehn–Thurston coordinates guarantees it satisfies the following scaling property: for every Borel measurable subset $A \subseteq \mathcal{ML}_g$ and every $t > 0$,

$$\mu_{\text{Thu}}(t \cdot A) = t^{6g-6} \cdot \mu_{\text{Thu}}(A). \tag{3.3}$$

This follows from Exercise 3.11 and the same property for the Lebesgue measure.

Exercise 3.12. Let $f: \mathcal{ML}_g \rightarrow \mathbb{R}$ be a non-negative, continuous function. Suppose that f is homogeneous with respect to the \mathbb{R}_+ scaling action on \mathcal{ML}_g . Using equation (3.3), show that, for every $c > 0$,

$$\mu(f^{-1}(\{c\})) = 0.$$

TABLE 1. Analogies between the lattice and surface settings.

Lattice setting	Surface setting
$SL(2, \mathbb{Z})$	$Mod_g =$ mapping class group of S_g
\mathbb{R}^2	$\mathcal{ML}_g =$ measured geodesic laminations on S_g
$\mathbb{Z}^2_{\text{prim}}$	$Mod_g \cdot \gamma =$ mapping class group orbit
$Leb_{\mathbb{R}^2}$	$\mu_{\text{Thu}} =$ Thurston measure
B_A	$B_X = \{\lambda \in \mathcal{ML}_g \mid \ell_\lambda(X) \leq 1\}$

3.15. *Analogies with the lattice setting.* Let us finish this section by highlighting some analogies that will be useful to have in mind when proving Theorem 1.1. The proof of Theorem 1.1 will follow the same approach as the proof of Theorem 2.1, but will use the analogies in Table 1 together with important results of Mirzakhani that will be discussed in §4.

4. *Weil–Petersson volumes and Mirzakhani’s integration formulas*

4.1. *Outline of this section.* In this section, we discuss two remarkable results of Mirzakhani that will play a crucial role in the proof of Theorem 1.1. The first of these results corresponds to the fact that the total Weil–Petersson volume of any moduli space of hyperbolic surfaces with totally geodesic boundary components is a polynomial on the lengths of the boundary components; see Theorem 4.1. The second of these results corresponds to Mirzakhani’s integration formulas over moduli space; see Exercise 4.7. We focus most of our attention on the second of these results. For the first result, we sketch a proof in the case of tori with one boundary component. An important part of the intuition behind this section comes from analogous facts for the Haar measure; see Proposition 2.10.

4.2. *Surfaces with boundary.* The first result we discuss concerns moduli spaces of hyperbolic surfaces with totally geodesic boundary components and their total Weil–Petersson volumes. To state this result precisely, we first introduce some notation. As many of the definitions that follow are analogous to those discussed in the previous section, we do not spend much time covering them.

Let $g, b \geq 0$ be a pair of non-negative integers such that $2 - 2g - b < 0$. Fix a connected, oriented, compact surface $S_{g,b}$ of genus g with b labeled boundary components β_1, \dots, β_b . Denote by $Mod_{g,b}$ the mapping class group of $S_{g,b}$, that is, the group of orientation-preserving homeomorphisms of $S_{g,b}$ that setwise fix each boundary component, up to homotopy, setwise fixing each boundary component. We warn the reader that different conventions are adopted elsewhere in the literature with regard to mapping classes fixing boundary components either pointwise or setwise; this is not an issue as long as the corresponding definitions are appropriately adapted.

Let $\mathbf{L} := (L_i)_{i=1}^b \in (\mathbb{R}_{>0})^b$ be a vector of positive real numbers. Denote by $\mathcal{T}_{g,b}(\mathbf{L})$ the Teichmüller space of marked, oriented hyperbolic structures on $S_{g,b}$ with labeled geodesic boundary components whose lengths are given by \mathbf{L} , up to isotopy fixing each boundary component setwise. The mapping class group $Mod_{g,b}$ acts properly discontinuously on $\mathcal{T}_{g,b}(\mathbf{L})$ by changing the markings. The quotient $\mathcal{M}_{g,b}(\mathbf{L}) := \mathcal{T}_{g,b}(\mathbf{L})/Mod_{g,b}$ is the moduli space of oriented hyperbolic structures on $S_{g,b}$ with labeled geodesic boundary components whose lengths are given by \mathbf{L} .

4.3. *Weil–Petersson volumes.* The Teichmüller space $\mathcal{T}_{g,b}(\mathbf{L})$ can be parameterized using Fenchel–Nielsen coordinates in a similar way as for closed surfaces. More concretely, for any pair of pants decomposition $\mathcal{P} := (\gamma_i)_{i=1}^{3g-3+b}$ of $S_{g,b}$, the length and twist parameters $(\ell_i, \tau_i)_{i=1}^{3g-3+b}$ of marked hyperbolic structures $X \in \mathcal{T}_{g,b}(\mathbf{L})$ with respect to the components of \mathcal{P} provide a global coordinate system for $\mathcal{T}_{g,b}(\mathbf{L})$; pairs of pants decompositions of surfaces with boundary are not considered to include the boundary components of the surface. The Weil–Petersson volume form v_{wp} of $\mathcal{T}_{g,b}(\mathbf{L})$ can be defined as follows:

$$v_{\text{wp}} := \bigwedge_{i=1}^{3g-3+b} d\ell_i \wedge d\tau_i. \tag{4.1}$$

We refer to the measure μ_{wp} induced by v_{wp} as the Weil–Petersson measure of $\mathcal{T}_{g,b}(\mathbf{L})$.

On the moduli space $\mathcal{M}_{g,b}(\mathbf{L}) := \mathcal{T}_{g,b}(\mathbf{L})/\text{Mod}_{g,b}$, consider the local pushforward $\widehat{\mu}_{\text{wp}}$ of the Weil–Petersson measure μ_{wp} . We refer to this measure as the Weil–Petersson measure of $\mathcal{M}_{g,b}(\mathbf{L})$. The same arguments suggested in Exercise 3.8 can be used to show that the Weil–Petersson measure $\widehat{\mu}_{\text{wp}}$ on $\mathcal{M}_{g,b}(\mathbf{L})$ is finite. The total Weil–Petersson volume of $\mathcal{M}_{g,b}(\mathbf{L})$ will be denoted by

$$V_{g,b}(\mathbf{L}) := \widehat{\mu}_{\text{wp}}(\mathcal{M}_{g,b}(\mathbf{L})).$$

By Exercise 3.3, if the surface $S_{g,b}$ is a pair of pants, that is, if $g = 0$ and $b = 3$, the moduli space $\mathcal{M}_{g,n}^b(\mathbf{L})$ consists of exactly one point. In this case, we adopt the convention

$$V_{0,3}(\mathbf{L}) := 1.$$

The following remarkable theorem of Mirzakhani shows that the total Weil–Petersson volumes $V_{g,b}(\mathbf{L})$ are polynomials on the \mathbf{L} variables. We will later sketch a proof of this result for the case of tori with one boundary component, that is, for the case $g = 1$ and $b = 1$.

THEOREM 4.1. [Mir07a, Theorem 1.1] [Mir07b, Theorem 1.1] *Let $g, b \geq 0$ be non-negative integers such that $2 - 2g - b < 0$. The total Weil–Petersson volume*

$$V_{g,b}(L_1, \dots, L_b)$$

is a polynomial of degree $3g - 3 + b$ on the variables L_1^2, \dots, L_b^2 . Moreover, if

$$V_{g,b}(L_1, \dots, L_b) = \sum_{\substack{\alpha \in (\mathbb{Z}_{\geq 0})^b \\ |\alpha| \leq 3g-3+b}} c_\alpha \cdot L_1^{2\alpha_1} \cdots L_b^{2\alpha_b},$$

where $|\alpha| := \alpha_1 + \dots + \alpha_b$ for every $\alpha \in (\mathbb{Z}_{\geq 0})^b$, then $c_\alpha \in \mathbf{Q}_{>0} \cdot \pi^{6g-6+2b-2|\alpha|}$. In particular, the leading coefficients of $V_{g,b}(L_1, \dots, L_b)$ belong to $\mathbf{Q}_{>0}$.

4.4. *The local change of variables formula.* We now introduce a local change of variables formula that will be used in forthcoming discussions. Let X be a locally compact, Hausdorff, second countable topological space endowed with a properly discontinuous

action of a discrete group G and $f : X \rightarrow \mathbb{R}$ be a measurable, non-negative function. Suppose that f is invariant with respect to a subgroup $H < G$, that is, $f(h.x) = f(x)$ for every $x \in X$ and every $h \in H$. Denote by $\tilde{f} : X/H \rightarrow \mathbb{R}$ the measurable, non-negative function induced by f on X/H . Consider the measurable non-negative function $\hat{f} : X/G \rightarrow \mathbb{R}$ which to every $x \in X/G$ assigns the value

$$\hat{f}(x) := \sum_{g \in G/H} f(g.x).$$

Let μ be a locally finite G -invariant Borel measure on X . Denote by $\tilde{\mu}$ the local pushforward of μ to X/H and by $\hat{\mu}$ the local pushforward of μ to X/G . The following exercise can be interpreted as a local change of variables formula.

Exercise 4.2. Show that, in the setting described above, the following integration formula holds:

$$\int_{X/G} \hat{f}(x) d\hat{\mu}(x) = \int_{X/H} \tilde{f}(y) d\tilde{\mu}(y).$$

Hint: Use a partition of unity of X/G to reduce to the case where f is supported on the preimage of a well-covered neighborhood.

The following exercise, which corresponds to Proposition 2.10, is a classical application of the local change of variables formula in the setting of lattices.

Exercise 4.3. The main goal of this exercise is to prove Proposition 2.10; we refer the reader back to §2 for the definitions and notation that will be used throughout. Let $X := \text{SO}(2) \backslash \text{SL}(2, \mathbb{R})$ and $G := \text{SL}(2, \mathbb{Z})$ acting on X via right multiplication. Denote by $\|\cdot\|$ the Euclidean norm on \mathbb{R}^2 and by $e_1 := (1, 0) \in \mathbb{R}^2$ the first canonical vector of \mathbb{R}^2 . For every $L > 0$, consider the function $f : X \rightarrow \mathbb{R}$ which to every $A \in X$ assigns the value $f(A) := \mathbb{1}_{\|A \cdot e_1\| \leq L}$. Notice that f is invariant with respect to the subgroup $H < G$ generated by the unipotent matrix

$$u_1 := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

(1) Show that for every $\Lambda \in \mathcal{M}_1 := X/G$ and every $L > 0$,

$$\hat{f}(\Lambda) = p(\Lambda, L).$$

(2) Show that if $A \in X$ gets identified with $z \in \mathbf{H}^2$, then

$$\|A \cdot e_1\| = \Im(z)^{-1/2},$$

and, in particular, for every $L > 0$,

$$f(A) = \mathbf{1}_{\Im(z) \geq 1/L^2}.$$

(3) Denote by $\hat{\mu}$ the local pushforward of the measure μ in equation (2.13) to \mathcal{M}_1 as defined in equation (3.2). Notice this definition differs from that used in Proposition 2.10 by a factor of 2. Using Exercise 4.2, the fundamental domain in

Figure 2 and the previous parts show that

$$\int_{\mathcal{M}_1} p(\Lambda, L) d\widehat{\mu}(\Lambda) = L^2.$$

4.5. *The cut and glue fibration.* We now discuss a remarkable observation due to Mirzakhani regarding how the Weil–Petersson measure on certain quotients of Teichmüller space disintegrates along the fibers of natural projections over products of moduli spaces of lower complexity. This observation is nothing more than a simple consequence of Wolpert’s magic formula but its importance cannot be overstated.

For the rest of this discussion, we fix an integer $g \geq 2$ and a connected, oriented, closed surface S_g of genus g . Recall that Mod_g denotes the mapping class group of S_g . Given a simple closed curve α on S_g , denote by $\text{Stab}_0(\alpha) \subseteq \text{Mod}_g$ the subgroup of mapping classes of S_g that fix α up to isotopy together with its orientations. Although α is unoriented, it admits two possible orientations. A mapping class belongs to $\text{Stab}_0(\alpha)$ if it sends each orientation of α back to itself. An ordered simple closed multi-curve on S_g is a tuple $\vec{\gamma} := (\gamma_i)_{i=1}^k$ of pairwise disjoint, pairwise non-isotopic simple closed curves on S_g . Given such a simple closed multi-curve on S_g , denote

$$\text{Stab}_0(\vec{\gamma}) := \bigcap_{i=1}^k \text{Stab}_0(\gamma_i) \subseteq \text{Mod}_g;$$

considering this particular definition of stabilizer allows us to avoid possible inconsistencies in the formulas that could arise from the existence of extra symmetries for specific multi-curves.

For the rest of this discussion, we fix an ordered simple closed multi-curve $\vec{\gamma} := (\gamma_i)_{i=1}^k$ on S_g . Recall that \mathcal{T}_g denotes the Teichmüller space of marked hyperbolic structures on S_g . The quotient $\mathcal{T}_g/\text{Stab}_0(\vec{\gamma})$ fibers naturally over a product of moduli spaces of surfaces with boundary of less complexity than S_g . This fibration, which we refer to as the cut and glue fibration of $\mathcal{T}_g/\text{Stab}_0(\vec{\gamma})$, will play a crucial role in our discussion of Mirzakhani’s integration formula. To describe this fibration in detail, we first introduce some notation.

Let $S_g(\vec{\gamma})$ be the potentially disconnected oriented surface with boundary obtained by cutting S_g along the components of $\vec{\gamma}$. See Figure 14 for an example. Let $c > 0$ be the number of components of $S_g(\vec{\gamma})$. Fixing an orientation on each component of $\vec{\gamma}$, we can keep track of which components of $S_g(\vec{\gamma})$ lie to the left and to the right of each component of $\vec{\gamma}$, so we can label the components of $S_g(\vec{\gamma})$ in a consistent way, say Σ_j with $j \in \{1, \dots, c\}$. As the components of $\vec{\gamma}$ are labeled and oriented, this induces a labeling of the boundary components of each Σ_j . Let $g_j, b_j \geq 0$ with $2 - 2g_j - b_j < 0$ be a pair of non-negative integers such that Σ_j is homeomorphic to S_{g_j, b_j} . Fix a homeomorphism between this pair of surfaces respecting the labeling of their boundary components.

The base of the cut and glue fibration is the space $\Omega_g(\vec{\gamma})$ of pairs (\mathbf{L}, \mathbf{X}) , where

$$\begin{aligned} \mathbf{L} &:= (\ell_i)_{i=1}^k \in (\mathbb{R}_{>0})^k, \\ \mathbf{X} &:= (X_j)_{j=1}^c \in \prod_{j=1}^c \mathcal{M}_{g_j, b_j}(\mathbf{L}_j), \end{aligned}$$




FIGURE 14. Cutting a genus 2 surface along a non-separating simple closed curve (colour online).

such that $\mathbf{L}_j \in (\mathbb{R}_{>0})^{b_j}$ is defined using the vector $\mathbf{L} := (\ell_i)_{i=1}^k \in (\mathbb{R}_{>0})^k$ and the correspondence between the labeling of the boundary components of Σ_j and the labeling of the components of $\vec{\gamma}$.

An ordered simple closed multi-geodesic $\alpha := (\alpha_i)_{i=1}^k$ on a hyperbolic surface X is a tuple of pairwise disjoint simple closed geodesics on X . Every ordered simple closed multi-geodesic is in particular an ordered simple closed multi-curve. Two ordered simple closed multi-curves on homeomorphic surfaces are said to have the same topological type if there exists a homeomorphism between the surfaces mapping one simple closed multi-curve to the other respecting their labelings. Recall that \mathcal{M}_g denotes the moduli space of genus g hyperbolic surfaces. The following exercise will play a crucial role in our description of the cut and glue fibration.

Exercise 4.4. Show there exists a one-to-one correspondence between points in the quotient space $\mathcal{T}_g/\text{Stab}_0(\vec{\gamma})$ and pairs $(X, \vec{\alpha})$, where $X \in \mathcal{M}_g$ is a genus g hyperbolic surface and $\vec{\alpha} := (\alpha_i)_{i=1}^k$ is an ordered simple closed multi-geodesic on X of the same topological type as $\vec{\gamma}$ with a choice of orientation on each of its components, modulo the equivalence relation $(X', \vec{\alpha}') \sim (X'', \vec{\alpha}'')$ if and only if there exists an orientation-preserving isometry $I: X' \rightarrow X''$ sending the components of $\vec{\alpha}'$ to the components of $\vec{\alpha}''$ respecting their labelings and orientations.

Consider the identification of the quotient space $\mathcal{T}_{g,n}/\text{Stab}_0(\vec{\gamma})$ provided by Exercise 4.4. The cut and glue fibration of $\mathcal{T}_{g,n}/\text{Stab}_0(\vec{\gamma})$ is the map $\Psi: \mathcal{T}_{g,n}/\text{Stab}_0(\vec{\gamma}) \rightarrow \Omega_{g,n}(\vec{\gamma})$ which to every equivalence class $(X, \vec{\alpha}) \in \mathcal{T}_{g,n}/\text{Stab}_0(\vec{\gamma})$ assigns the pair $(\mathbf{L}, \mathbf{X}) \in \Omega_g(\vec{\gamma})$ given by

$$\mathbf{L} := (\ell_{\alpha_i}(X))_{i=1}^k, \quad \mathbf{X} := (X(\vec{\alpha}_j)_j)_{j=1}^c,$$

where $X(\vec{\alpha}_j)_j \in \mathcal{M}_{g_j, b_j}(\mathbf{L}_j)$ is the j th component, according to the labeling and orientation of the components of $\vec{\alpha}$, of the hyperbolic surface with geodesic boundary components obtained by cutting X along the components of $\vec{\alpha}$. The fiber $\Psi^{-1}(\mathbf{X}, \mathbf{L})$ above any pair $(\mathbf{L}, \mathbf{X}) \in \Omega_g(\vec{\gamma})$ is given by all possible ways of gluing the components of $\mathbf{X} := (X_j)_{j=1}^c$ along their boundaries respecting the labelings. Given any point $(X, \vec{\alpha}) \in \Psi^{-1}(\mathbf{X}, \mathbf{L})$, the whole fiber $\Psi^{-1}(\mathbf{X}, \mathbf{L})$ can be recovered by considering all possible Fenchel–Nielsen twists of X along the components of $\vec{\alpha}$.

Remark 4.5. Notice that, because of Exercise 3.3, Fenchel–Nielsen coordinates modulo Dehn twists along the components of the corresponding pair of pants decomposition are




FIGURE 15. The isometric involution of a torus with one boundary component (colour online).

an instance of the cut and glue fibration; the base of the fibration is the positive orthant of possible cuff lengths.

Consider a point $(\mathbf{L}, \mathbf{X}) \in \Omega_g(\vec{\gamma})$ on the base of the cut and glue fibration. Let $(X, \vec{\alpha}) \in \Psi^{-1}(\mathbf{X}, \mathbf{L})$ with $\vec{\alpha} := (\alpha_i)_{i=1}^k$. The fiber $\Psi^{-1}(\mathbf{X}, \mathbf{L})$ supports well-defined 1-forms $d\tau_{\alpha_i}$ that measure the infinitesimal Fenchel–Nielsen twists along the components of $\vec{\alpha}$. Wedging these 1-forms yields a natural volume form on $\Psi^{-1}(\mathbf{X}, \mathbf{L})$. For every set of lengths $\mathbf{L} := (\ell_i)_{i=1}^k$ and for Weil–Petersson almost every \mathbf{X} , the total mass of the fiber $\Psi^{-1}(\mathbf{X}, \mathbf{L})$ with respect to this volume form is given by

$$|\Psi^{-1}(\mathbf{X}, \mathbf{L})| = 2^{-\rho_g(\vec{\gamma})} \cdot \ell_1 \cdots \ell_k, \tag{4.2}$$

where $\rho_g(\vec{\gamma})$ is the number of components of $\vec{\gamma}$ that bound (on any of its sides) a component of $S_g(\vec{\gamma})$ that is homeomorphic to a torus with one boundary component. We refer to pairs $(\mathbf{L}, \mathbf{X}) \in \Omega_g(\vec{\gamma})$ such that equation (4.2) holds as generic pairs. The factor $2^{-\rho_g(\vec{\gamma})}$ reflects the observation that every hyperbolic torus with one geodesic boundary component has a non-trivial isometric involution preserving its boundary setwise; see Figure 15. A similar issue arises for higher genera hyperbolic surfaces with extra symmetries, and hence the need to impose the Weil–Petersson almost every condition on \mathbf{X} in equation (4.2).

Denote by μ_{wp} the Weil–Petersson measure on \mathcal{T}_g and by $\tilde{\mu}_{\text{wp}}$ the local pushforward of μ_{wp} to $\mathcal{T}_g/\text{Stab}_0(\vec{\gamma})$. It follows from Wolpert’s magic formula, that is, from equations (3.1) and (4.1), that $\tilde{\mu}_{\text{wp}}$ can be disintegrated along the fibers of the cut and glue fibration $\Psi: \mathcal{T}_{g,n}/\text{Stab}_0(\vec{\gamma}) \rightarrow \Omega_{g,n}(\vec{\gamma})$ as

$$\tilde{\mu}_{\text{wp}} = \sigma_g(\vec{\gamma}) \cdot \prod_{i=1}^k d\tau_{\alpha_i} \otimes \prod_{i=1}^c \widehat{\mu}_{\text{wp}}^j \otimes \prod_{i=1}^k d\ell_i,$$

where $\widehat{\mu}_{\text{wp}}^j$ denotes the Weil–Petersson measure on $\mathcal{M}_{g_j, b_j}(\mathbf{L}_j)$ and $\sigma_g(\vec{\gamma}): \mathcal{T}_g/\text{Stab}_0(\vec{\gamma}) \rightarrow \mathbf{Q}_{>0}$ is the function with non-negative rational values that records the discrepancy between the stabilizer factors, that is, the factors $|\Gamma|$ in equation (3.2), of the measure $\tilde{\mu}_{\text{wp}}$ and the product of measures $\prod_{i=1}^c \widehat{\mu}_{\text{wp}}^j$. One can show that the function $\sigma_g(\vec{\gamma})$ is almost everywhere constant with respect to $\tilde{\mu}_{\text{wp}}$ and is given by

$$\sigma_g(\vec{\gamma}) := \frac{\prod_{j=1}^c |K_{g_j, b_j}|}{|\text{Stab}_0(\vec{\gamma}) \cap K_g|},$$

where $K_{g_j, b_j} \triangleleft \text{Mod}_{g_j, b_j}$ is the kernel of the mapping class group action on \mathcal{T}_{g_j, b_j} and $K_g \triangleleft \text{Mod}_g$ is the kernel of the mapping class group action on \mathcal{T}_g . Indeed, $\widehat{\mu}_{\text{wp}}^j$ almost every hyperbolic surface in $\mathcal{M}_{g_j, b_j}(\mathbf{L})$ has $|K_{g_j, b_j}|$ automorphisms and analogously for elements of $\mathcal{T}_g/\text{Stab}_0(\vec{\gamma})$. For example, if $g = 2$ and $\vec{\gamma}$ is a separating simple closed curve on S_2 , then $\sigma_2(\vec{\gamma}) = 4/2 = 2$. In general,

$$|K_{g,b}| = \begin{cases} 4 & \text{if } (g, b) = (0, 4), \\ 2 & \text{if } (g, b) \in \{(1, 1), (1, 2), (2, 0)\}, \\ 1 & \text{if } (g, b) \notin \{(0, 4), (1, 1), (1, 2), (2, 0)\}. \end{cases}$$

Let us record the most important conclusions of the discussion above in the following theorem.

THEOREM 4.6. *The Weil–Petersson measure $\widetilde{\mu}_{\text{wp}}$ on $\mathcal{T}_g/\text{Stab}_0(\vec{\gamma})$ can be disintegrated along the fibers of the cut and glue fibration $\Psi: \mathcal{T}_{g,n}/\text{Stab}_0(\vec{\gamma}) \rightarrow \Omega_{g,n}(\vec{\gamma})$ in the following way:*

$$\widetilde{\mu}_{\text{wp}} = \sigma_g(\vec{\gamma}) \cdot \prod_{i=1}^k d\tau_{\alpha_i} \otimes \prod_{i=1}^c \widehat{\mu}_{\text{wp}}^j \otimes \prod_{i=1}^k d\ell_i.$$

Generically, the volume of the fiber $\Psi^{-1}(\mathbf{X}, \mathbf{L})$ above a $(\mathbf{L}, \mathbf{X}) \in \Omega_{g,n}(\vec{\gamma})$ with $\mathbf{L} := (\ell_i)_{i=1}^k$ is equal to

$$|\Psi^{-1}(\mathbf{X}, \mathbf{L})| = 2^{-\rho_g(\vec{\gamma})} \cdot \ell_1 \cdots \ell_k.$$

4.6. Mirzakhani’s integration formula. We are now ready to discuss Mirzakhani’s integration formula. Let $\vec{\gamma} := (\gamma_i)_{i=1}^k$ be an ordered simple closed multi-curve on S_g . Consider the stabilizer

$$\text{Stab}(\vec{\gamma}) := \bigcap_{i=1}^k \text{Stab}(\gamma_i) \subseteq \text{Mod}_g.$$

Notice that $\text{Stab}_0(\vec{\gamma})$ is a finite index subgroup of $\text{Stab}(\vec{\gamma})$. Let $c > 0$ and $g_j, b_j \geq 0$ with $j \in \{1, \dots, c\}$ be as above. Consider the total Weil–Petersson volumes

$$V_{g_j, b_j}(\mathbf{x}_j) := \widehat{\mu}_{\text{wp}}^j(\mathcal{M}_{g_j, b_j}(\mathbf{x}_j))$$

of the moduli spaces $\mathcal{M}_{g_j, b_j}(\mathbf{x}_j)$ as functions of the boundary lengths $\mathbf{x}_j \in \mathbb{R}_+^{b_j}$. Recall the definitions of the constants $\rho_g(\vec{\gamma}) \in \mathbb{N}$ and $\sigma_g(\vec{\gamma}) \in \mathbf{Q}_{>0}$ introduced above. Consider the measurable function $V_g(\vec{\gamma}, \cdot): \mathbb{R}_+^k \rightarrow \mathbb{R}_+$ which, to every vector $\mathbf{x} := (x_i)_{i=1}^k$ with positive entries, assigns the value

$$V_g(\vec{\gamma}, \mathbf{x}) := \frac{\sigma_g(\vec{\gamma}) \cdot 2^{-\rho_g(\vec{\gamma})}}{[\text{Stab}(\vec{\gamma}) : \text{Stab}_0(\vec{\gamma})]} \cdot \prod_{j=1}^c V_{g_j, b_j}(\mathbf{x}_j), \tag{4.3}$$

where the vector $\mathbf{x}_j \in \mathbb{R}_+^{b_j}$ is defined using the correspondence between the components of $\vec{\gamma}$ and the boundary components of S_{g_j, b_j} . By Theorem 4.1, $V_g(\vec{\gamma}, \mathbf{x})$ is a polynomial on the \mathbf{x} variables.

Let $f: \mathbb{R}_+^k \rightarrow \mathbb{R}$ be a non-negative, measurable function. Given $X \in \mathcal{T}_g$ and an ordered simple closed multi-curve $\vec{\alpha} := (\alpha_i)_{i=1}^k$ on S_g , denote $\ell_{\vec{\alpha}}(X) := (\ell_{\alpha_i}(X))_{i=1}^k \in \mathbb{R}_+^k$. Consider the non-negative, measurable function $f_{\vec{\gamma}}: \mathcal{T}_g \rightarrow \mathbb{R}$ which, to every $X \in \mathcal{T}_g$, assigns the value

$$f_{\vec{\gamma}}(X) := \sum_{\vec{\alpha} \in \text{Mod}_g \cdot \vec{\gamma}} f(\ell_{\vec{\alpha}}(X)).$$

This function is clearly invariant with respect to the action of Mod_g on \mathcal{T}_g . Denote by $\widehat{f}_{\vec{\gamma}}: \mathcal{M}_g \rightarrow \mathbb{R}$ the corresponding non-negative, measurable function induced on moduli space. Denote by $\widehat{\mu}_{\text{wp}}$ the Weil–Petersson measure on \mathcal{M}_g . On \mathbb{R}_+^k , consider the standard coordinate system $\mathbf{x} := (x_i)_{i=1}^k$ and the Lebesgue class measure $\mathbf{x} \cdot d\mathbf{x} := x_1 \cdots x_k \cdot dx_1 \cdots dx_k$. The following exercise corresponds to Mirzakhani’s integration formula [Mir08b, Theorem 4.1].

Exercise 4.7. Let $\vec{\gamma} := (\gamma_i)_{i=1}^k$ be an ordered simple closed multi-curve on S_g . Using Exercise 4.2 and Theorem 4.6, show that the following integration formula holds:

$$\int_{\mathcal{M}_g} \widehat{f}_{\vec{\gamma}}(X) d\widehat{\mu}_{\text{wp}}(X) = \int_{\mathbb{R}_+^k} f(\mathbf{x}) \cdot V_g(\vec{\gamma}, \mathbf{x}) \cdot \mathbf{x} \cdot d\mathbf{x}.$$

4.7. McShane’s identity. We now sketch a proof of Theorem 4.1 for the case of tori with one boundary component, that is, for the case $g = 1$ and $b = 1$. For the rest of this discussion, we fix a torus with one boundary component $S_{1,1}$ and a parameter $L > 0$. To compute the total Weil–Petersson volume $V_{1,1}(L) := \widehat{\mu}_{\text{wp}}(\mathcal{M}_{1,1}(L))$, one could recall $\mathcal{M}_{1,1}(L) := \mathcal{T}_{1,1}(L)/\text{Mod}_{1,1}$ and try to find a fundamental domain for the action of $\text{Mod}_{1,1}$ on $\mathcal{T}_{1,1}(L)$ that can be described explicitly in Fenchel–Nielsen coordinates. This happens to be quite a formidable task. In the absence of such a fundamental domain, we consider a different approach that relies on Mirzakhani’s integration formula.

For the rest of this discussion, we fix a simple closed curve γ on $S_{1,1}$. Let $f: \mathbb{R}_+ \rightarrow \mathbb{R}$ be a non-negative measurable function. Just as in the case of closed surfaces, consider the non-negative, measurable transform $\widehat{f}_{\gamma}: \mathcal{M}_{1,1}(L) \rightarrow \mathbb{R}$ which, to every $X \in \mathcal{M}_{1,1}(L)$, assigns the value

$$\widehat{f}_{\gamma}(X) = \sum_{\alpha \in \text{Mod}_{1,1} \cdot \gamma} f(\ell_{\alpha}(X)).$$

An analog of Mirzakhani’s integration formula in Exercise 4.7 also holds in this setting. Our immediate goal is to find a non-negative, measurable function $f: \mathbb{R}_+ \rightarrow \mathbb{R}$ for which the transform $\widehat{f}_{\gamma}: \mathcal{M}_{1,1}(L) \rightarrow \mathbb{R}$ is equal to a constant $c(L) > 0$. Indeed, for such a function, we would have

$$c(L) \cdot \widehat{\mu}_{\text{wp}}(\mathcal{M}_{1,1}(L)) = \int_{\mathcal{M}_{1,1}(L)} \widehat{f}_{\gamma}(X) d\widehat{\mu}_{\text{wp}}(X) = \frac{1}{2} \cdot \int_{\mathbb{R}_+} f(x) \cdot x \cdot dx. \tag{4.4}$$

Rearranging the terms in this equation would yield

$$V_{1,1}(L) := \widehat{\mu}_{\text{wp}}(\mathcal{M}_{1,1}(L)) = \frac{1}{2 \cdot c(L)} \cdot \int_{\mathbb{R}_+} f(x) \cdot x \cdot dx. \tag{4.5}$$

Finding a non-negative, measurable function $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ for which $\widehat{f}_\gamma : \mathcal{M}_{1,1}(L) \rightarrow \mathbb{R}$ is constant is the content of McShane’s identity. Consider the function $D : \mathbb{R}^3 \rightarrow \mathbb{R}$ given by

$$D(x, y, z) := 2 \log \left(\frac{e^{x/2} + e^{(y+z)/2}}{e^{-x/2} + e^{(y+z)/2}} \right).$$

THEOREM 4.8. [McS91] *Let γ be a simple closed curve on $S_{1,1}$ and $L > 0$. Then, for every $X \in \mathcal{M}_{1,1}(L)$,*

$$\sum_{\alpha \in \text{Mod}_{1,1} \cdot \gamma} D(L, \ell_\alpha(X), \ell_\alpha(X)) = L. \tag{4.6}$$

Let us give a quick rundown of the main ideas of the proof of Theorem 4.8. For this, we interpret the right-hand side of equation (4.6) as the length of the boundary component of $X \in \mathcal{M}_{1,1}(L)$. For every point on this boundary component, we shoot a geodesic into X in the direction orthogonal to the boundary. Three things can happen at this stage. Either the geodesic remains in X at all times without intersecting itself, the geodesic exits X without intersecting itself, or the geodesic intersects itself. By the work of Birman and Series [BS85], the first case only happens for a measure zero subset of points on the boundary. In each of the two other cases, we consider the simple closed curves α described in Figure 16. Cutting X along the corresponding geodesic representatives yields a hyperbolic pair of pants with boundary lengths $(L, \ell_\alpha(X), \ell_\alpha(X))$. Using the rigidity of such pairs of pants, one can show that for points in exactly two arcs of the original boundary component whose lengths add up to $D(L, \ell_\alpha(X), \ell_\alpha(X))$, the corresponding orthogonal geodesics either intersect themselves or the original boundary component before exiting the pants. Putting these ideas together finishes the proof.

Exercise 4.9. Using equation (4.5) and Theorem 4.8 show that

$$V_{1,1}(L) = \frac{1}{48} \cdot L^2 + \frac{\pi^2}{12}. \tag{4.7}$$

Hint: To simplify the computations, differentiate equation (4.4) with respect to L instead of using equation (4.5) directly.

In her thesis [Mir04], Mirzakhani generalized Theorem 4.8 to arbitrary closed, orientable surfaces [Mir07a, Theorem 1.3]. Using this generalization, Mirzakhani proved a recursive formula for the total Weil–Petersson volumes $V_{g,b}(\mathbf{L})$ [Mir07a, §5]. Using this formula, she ultimately deduced Theorem 4.1. An alternative proof of Theorem 4.1 using symplectic reduction can also be found in Mirzakhani’s thesis [Mir07b, Theorem 1.1]. An excellent reference for all these topics is Do’s survey [Do13].



(a) Geodesic intersects boundary first. (b) Geodesic intersects itself first.

FIGURE 16. The geodesic shot orthogonal to the boundary (in blue) determines a simple closed curve (in green) which tightens to a simple closed geodesic (in red) (colour online).

In subsequent discussions, we will use the following explicit volume polynomial:

$$V_{1,2}(L_1, L_2) = \frac{1}{192} \cdot L_1^4 + \frac{1}{96} \cdot L_1^2 L_2^2 + \frac{1}{192} \cdot L_2^4 + \frac{\pi^2}{12} \cdot L_1^2 + \frac{\pi^2}{12} \cdot L_2^2 + \frac{\pi^2}{4}. \quad (4.8)$$

Exercise 4.10. Recall the definition of the polynomials $V_g(\vec{\gamma}, \mathbf{x})$ in equation (4.3). Let γ_1 and γ_2 be non-separating and separating simple closed curves on S_2 , respectively. Using equations (4.7) and (4.8), show that

$$V_2(\gamma_1, x) = \frac{1}{96} \cdot x^4, \quad V_2(\gamma_2, x) = \frac{1}{4608} \cdot x^4.$$

5. Counting simple closed geodesics on hyperbolic surfaces

5.1. Outline of this section. In this section, we give a complete proof of Theorem 1.1, the main result of this survey. We follow the same approach as in the proof of Theorem 2.1, but use the vocabulary and tools introduced in §§3–4. More concretely, we reduce to the study of an appropriate family of counting measures whose limit points are investigated using ergodic theory and integration formulas. We encourage the reader to keep in mind the analogies described in Table 1 for the rest of this section.

5.2. Counting simple closed geodesics. Recall that Theorem 1.1, the main result of this survey, asserts that the number $s(X, L)$ of unoriented simple closed geodesics of length $\leq L$ on an arbitrary closed, orientable hyperbolic surface X of genus $g \geq 2$ is asymptotic as $L \rightarrow \infty$ to a polynomial of degree $6g - 6$ on L . We aim to rewrite the counting function $s(X, L)$ using the terminology introduced in previous sections; we refer the reader to §§3 and 4 for the notation that will be used throughout.

For the rest of this section, we fix an integer $g \geq 2$ and a connected, oriented, closed surface S_g of genus g . We begin by considering the following combinatorial exercise; compare with Exercise 3.5.

Exercise 5.1. Show there are only finitely many topological types of simple closed curves on S_g . Can you give an exact formula for the number of such equivalence classes?

Recall that if γ is a simple closed curve on S_g and $X \in \mathcal{T}_g$ is a marked hyperbolic structure on S_g , then $\ell_\gamma(X) > 0$ denotes the length of the unique geodesic representative of γ with respect to X . Let γ be a simple closed curve on S_g and $X \in \mathcal{T}_g$. For every $L > 0$,

consider the counting function

$$s(X, \gamma, L) := \#\{\alpha \in \text{Mod}_g \cdot \gamma \mid \ell_\alpha(X) \leq L\}.$$

As every free homotopy class of simple closed curves on S_g has a unique geodesic representative with respect to X , this is exactly the number of unoriented simple closed geodesics on X of the same topological type as γ and length $\leq L$. Hence, we can decompose the counting function $s(X, L)$ as

$$s(X, L) = \sum_{\gamma} s(X, \gamma, L), \tag{5.1}$$

where γ runs over all the finitely many different topological types of simple closed curves on S_g ; see Exercise 5.1. Thus, it is enough for our purposes to study the asymptotics of $s(X, \gamma, L)$ as $L \rightarrow \infty$.

Motivated by this observation, we generalize the definition of the counting function $s(X, \gamma, L)$ to general integral simple closed multi-curves on S_g . Recall that if $\alpha := \sum_{i=1}^k a_i \alpha_i$ is a simple closed multi-curve on S_g and $X \in \mathcal{T}_g$, then $\ell_\alpha(X) := \sum_{i=1}^k a_i \ell_{\alpha_i}(X) > 0$. Let $\gamma := \sum_{i=1}^k a_i \gamma_i$ be an integral simple closed multi-curve on S_g and $X \in \mathcal{T}_g$. For every $L > 0$, consider the counting function

$$s(X, \gamma, L) := \#\{\alpha \in \text{Mod}_g \cdot \gamma \mid \ell_\alpha(X) \leq L\}.$$

Notice the function $s(X, \gamma, L)$ does not depend on the marking of $X \in \mathcal{T}_g$, but only on its underlying hyperbolic structure $X \in \mathcal{M}_g$. We aim to prove the following asymptotic estimate for the counting function $s(X, \gamma, L)$. Theorem 1.1 will later be deduced as a direct consequence of this estimate.

THEOREM 5.2. *Let $\gamma := \sum_{i=1}^k a_i \gamma_i$ be an integral simple closed multi-curve on S_g and $X \in \mathcal{M}_g$. Then, there exists a constant $n(X, \gamma) > 0$ depending only on the topological type of γ and the geometry of X such that the following asymptotic estimate holds as $L \rightarrow \infty$,*

$$s(X, \gamma, L) \sim n(X, \gamma) \cdot L^{6g-6}.$$

5.3. *Counting measures on the space of measured geodesic laminations.* Inspired by the case of lattices, to prove Theorem 5.2, we introduce appropriate families of counting measures. Recall that \mathcal{ML}_g denotes the space of measured geodesic laminations on S_g , that is, the natural $6g - 6$ dimensional completion of the set of simple closed multi-curves on S_g . Let $\gamma := \sum_{i=1}^k a_i \gamma_i$ be an integral simple closed multi-curve on S_g . For every $L > 0$, consider the counting measure on \mathcal{ML}_g given by

$$\mu_L^\gamma := \frac{1}{L^{6g-6}} \cdot \sum_{\alpha \in \text{Mod}_g \cdot \gamma} \delta_{1/L \cdot \alpha}. \tag{5.2}$$

Recall that $\ell_\lambda(X) > 0$ denotes the length of a measured geodesic lamination $\lambda \in \mathcal{ML}_g$ with respect to a marked hyperbolic structure $X \in \mathcal{T}_g$. Given $X \in \mathcal{T}_g$, consider the set

$$B_X := \{\lambda \in \mathcal{ML}_g \mid \ell_\lambda(X) \leq 1\}.$$

This set depends on the marking of $X \in \mathcal{T}_g$. Directly from the definitions, we see that, for every $L > 0$,

$$\mu_L^\gamma(B_X) = \frac{s(X, \gamma, L)}{L^{6g-6}}. \tag{5.3}$$

This reduces the original problem of proving an asymptotic estimate for the counting function $s(X, \gamma, L)$ to the problem of understanding the behavior as $L \rightarrow \infty$ of the sequence of counting measures $(\mu_L^\gamma)_{L>0}$.

5.4. Ergodicity of the mapping class group action. To study the asymptotic behavior of the sequence of counting measures $(\mu_L^\gamma)_{L>0}$, we use the dynamics of the action of the mapping class group on \mathcal{ML}_g . Recall the definition of the Thurston measure μ_{Thu} on \mathcal{ML}_g and its relation to Dehn–Thurston coordinates as described in Exercise 3.11. The following result of Masur is an analog of the fact that the Lebesgue measure is ergodic with respect to the action of $\text{SL}(2, \mathbb{Z})$ on \mathbb{R}^2 . See Theorem 2.5.

THEOREM 5.3. [Mas85] *The measure μ_{Thu} is ergodic with respect to the action of Mod_g on \mathcal{ML}_g .*

In analogy with the case of lattices, the ergodicity of the mapping class group action on \mathcal{ML}_g can be used to study the weak- \star limit points of the sequence of counting measures $(\mu_L^\gamma)_{L>0}$. The following crucial exercise is an analog of Proposition 2.8.

Exercise 5.4. Let $\gamma := \sum_{i=1}^k a_i \gamma_i$ be an integral simple closed multi-curve on S_g . Show that every weak- \star limit point μ^γ of the sequence of counting measures $(\mu_L^\gamma)_{L>0}$ introduced in equation (5.2) is of the form $\mu^\gamma = c \cdot \mu_{\text{Thu}}$ for some constant $c \geq 0$. *Hint:* Follow the same approach as in the proof of Proposition 2.8 but working in Dehn–Thurston coordinates and recalling Exercise 3.11. More concretely, use a rough lattice points counting bound to show every limit point is absolutely continuous with respect to μ_{Thu} and then use Theorem 5.3 and Exercise 2.4 to conclude.

5.5. Integration and integrability. Our next goal is to show that the constant $c \geq 0$ in the conclusion of Exercise 5.4 is positive and independent of the limit point μ^γ . Similar to the case of lattices, we achieve this goal by averaging over moduli space and using Mirzakhani’s integration formula.

Recall that $\widehat{\mu}_{\text{wp}}$ denotes the Weil–Petersson measure on \mathcal{M}_g . Let $\vec{\gamma} := (\gamma_i)_{i=1}^k$ be an ordered simple closed multi-curve on S_g and $\mathbf{a} := (a_i)_{i=1}^k \in \mathbb{R}_+^k$ be a vector of positive weights. On S_g , consider the simple closed multi-curve given by $\mathbf{a} \cdot \vec{\gamma} := \sum_{i=1}^k a_i \gamma_i$. Recall the definition of the polynomial $V_g(\vec{\gamma}, \mathbf{x})$ introduced in equation (4.3). Notice that $\text{Stab}(\vec{\gamma}) \subseteq \text{Mod}_g$ is a finite index subgroup of $\text{Stab}(\mathbf{a} \cdot \vec{\gamma})$. Define

$$V_g(\mathbf{a} \cdot \vec{\gamma}, \mathbf{x}) := [\text{Stab}(\mathbf{a} \cdot \vec{\gamma}) : \text{Stab}(\vec{\gamma})]^{-1} \cdot V_g(\vec{\gamma}, \mathbf{x}).$$

On \mathbb{R}_+^k , consider the standard coordinate system $\mathbf{x} := (x_i)_{i=1}^k$ and the Lebesgue class measure $\mathbf{x} \cdot d\mathbf{x} := x_1 \cdots x_k \cdot dx_1 \cdots dx_k$. The following exercise is an analog of Proposition 2.10 and Exercise 4.3.

Exercise 5.5. Let $\vec{\gamma} := (\gamma_i)_{i=1}^k$ be an ordered simple closed multi-curve on S_g and $\mathbf{a} := (a_i)_{i=1}^k \in \mathbb{R}_+^k$ be a vector of positive weights on the components of γ . Using Exercise 4.7, show that, for every $L > 0$,

$$\int_{\mathcal{M}_g} s(X, \mathbf{a} \cdot \vec{\gamma}, L) d\widehat{\mu}_{\text{wp}}(X) = \int_{\mathbf{a} \cdot \mathbf{x} \leq L} V_g(\mathbf{a} \cdot \vec{\gamma}, \mathbf{x}) \cdot \mathbf{x} \cdot d\mathbf{x}.$$

Using this formula and Theorem 4.1, deduce that the function

$$P(\mathbf{a} \cdot \vec{\gamma}, L) := \int_{\mathcal{M}_g} s(X, \mathbf{a} \cdot \vec{\gamma}, L) d\widehat{\mu}_{\text{wp}}(X) \tag{5.4}$$

is a polynomial of degree $6g - 6$ on L with rational leading coefficient.

Let $\gamma := \mathbf{a} \cdot \vec{\gamma} = \sum_{i=1}^k a_i \gamma_i$ be a simple closed multi-curve on S_g and $P(\gamma, L) = P(\mathbf{a} \cdot \vec{\gamma}, L)$ be as in equation (5.4). Following Exercise 5.5, we define the frequency of γ to be the positive rational number

$$c(\gamma) := \lim_{L \rightarrow \infty} \frac{P(\gamma, L)}{L^{6g-6}}. \tag{5.5}$$

Using Mirzakhani’s recursion for the Weil–Petersson volume polynomials $V_{g,b}(\mathbf{L})$ [Mir07a, §5], the frequency of any simple closed curve can be computed explicitly by means of a recursive algorithm.

Exercise 5.6. Let γ_1 and γ_2 be non-separating and separating simple closed curves on S_2 , respectively. Using Exercise 4.10, show that the frequencies of these simple closed curves are given by

$$c(\gamma_1) = 1/576, \quad c(\gamma_2) = 1/27\,648.$$

We now discuss an analog of the integrability bound in Exercise 2.11 for the function $s(X, \gamma, L)$. Let us first introduce a more precise statement of the collar lemma discussed in §3. A proof of this result can be found in [FM12, Lemma 13.6]. Consider the width function $w: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ given by

$$w(x) := \operatorname{arcsinh}\left(\frac{1}{\sinh(x/2)}\right).$$

LEMMA 5.7. *Let α be a simple closed geodesic on a closed, orientable hyperbolic surface X . Denote by d the metric on X . Then, the subset $N_\alpha \subseteq X$ defined as follows is an embedded annulus in X :*

$$N_\alpha := \{x \in X : d(x, \alpha) < w(\ell_\alpha(X))\}.$$

Let $\mathcal{P} := (\gamma_i)_{i=1}^{3g-3}$ be a pair of pants decomposition of S_g and $(m_i, t_i)_{i=1}^{3g-3}$ be a set of Dehn–Thurston coordinates of \mathcal{ML}_g induced by \mathcal{P} . Given a measured geodesic lamination $\lambda \in \mathcal{ML}_g$ and $X \in \mathcal{T}_g$, define the combinatorial length of λ with respect to X and \mathcal{P} as

$$L_\lambda(X, \mathcal{P}) := \sum_{i=1}^N (m_i(\gamma) \cdot w(\ell_{\gamma_i}(X)) + |t_i(\gamma)| \cdot \ell_{\gamma_i}(X)).$$

For a simple closed curve α on S_g and $X \in \mathcal{T}_g$, this definition has a concrete interpretation: add the width $w(\ell_{\gamma_i}(X))$ of the collar given by Lemma 5.7 every time α intersects γ_i and add the length $\ell_{\gamma_i}(X)$ every time α twists around γ_i . A pair of pants decomposition $\mathcal{P} := (\gamma_i)_{i=1}^{3g-3}$ of S_g is said to be L -bounded with respect to $X \in \mathcal{T}_g$ for some $L > 0$ if $\ell_{\gamma_i}(X) \leq L$ for every $i \in \{1, \dots, 3g - 3\}$. The following result of Mirzakhani shows that combinatorial lengths approximate hyperbolic lengths.

PROPOSITION 5.8. [Mir08b, Proposition 3.5] *For every $L > 0$, there exists $C = C(L) > 0$ such that for every $X \in \mathcal{T}_g$ and every pair of pants decomposition \mathcal{P} of S_g that is L -bounded with respect to X , there exist Dehn–Thurston coordinates $(m_i, t_i)_{i=1}^{3g-3}$ of \mathcal{ML}_g induced by \mathcal{P} such that for every $\lambda \in \mathcal{ML}_g$,*

$$C^{-1} \cdot L_{\mathcal{P}}(X, \lambda) \leq \ell_{\lambda}(X) \leq C \cdot L_{\mathcal{P}}(X, \lambda).$$

To prove the aforementioned integrability bound for the counting function $s(X, \gamma, L)$, we will also use the following stronger version of Bers’s theorem. For a proof, see [FM12, Theorem 12.8].

THEOREM 5.9. *For every $\epsilon > 0$, there exists $L = L(\epsilon) > 0$ with the following property. Let $X \in \mathcal{T}_g$ be a marked hyperbolic structure and $\gamma := (\gamma_i)_{i=1}^k$ be a simple closed multi-curve on S_g such that*

$$\ell_{\gamma_i}(X) < \epsilon \quad \text{for all } i = 1, \dots, k.$$

Then, there exists a completion of γ to a pair of pants decomposition $\mathcal{P} := (\gamma_i)_{i=1}^{3g-3}$ of S_g such that

$$\ell_{\gamma_i}(X) < L \quad \text{for all } i = 1, \dots, 3g - 3.$$

By Lemma 5.7, there exists a constant $\epsilon > 0$ such that on any closed, orientable hyperbolic surface, no two closed geodesics of length $< \epsilon$ intersect. For the rest of this section, we fix such a constant and denote it by $\epsilon > 0$. Consider the measurable function $u: \mathcal{M}_g \rightarrow \mathbb{R}_+$ given for every $X \in \mathcal{M}_g$ by

$$u(X) := \prod_{\alpha: \ell_{\alpha}(X) < \epsilon} \frac{1}{\ell_{\alpha}(X)},$$

where the product runs over all simple closed geodesics α on X of length $\ell_{\alpha}(X) < \epsilon$. We interpret empty products as taking the value 1. The following exercise is an analog of Exercise 2.11.

Exercise 5.10. Let $\gamma := \sum_{i=1}^k a_i \gamma_i$ be an integral simple closed multi-curve on S_g . Show there exists a constant $C > 0$ such that for every $X \in \mathcal{M}_g$ and every $L > 0$,

$$s(X, \gamma, L) \leq C \cdot L^{6g-6} \cdot u(X). \tag{5.6}$$

Additionally, show that the function $u : \mathcal{M}_g \rightarrow \mathbb{R}_+$ is integrable with respect to $\widehat{\mu}_{\text{wp}}$, that is,

$$\int_{\mathcal{M}_g} u(X) d\widehat{\mu}_{\text{wp}}(X) < \infty. \tag{5.7}$$

Hint: To prove equation (5.6), use Proposition 5.8 and Theorem 5.9 to reduce to a lattice point counting problem in Dehn–Thurston coordinates. To prove equation (5.7), follow the same approach as in Exercise 3.8.

For every marked hyperbolic structure $X \in \mathcal{T}_g$, denote

$$B(X) := \mu_{\text{Thu}}(B_X) = \mu_{\text{Thu}}(\{\lambda \in \mathcal{ML}_g \mid \ell_\lambda(X) \leq 1\}). \tag{5.8}$$

As the Thurston measure μ_{Thu} is invariant with respect to the Mod_g action on \mathcal{ML}_g , the value $B(X)$ is independent of the marking of $X \in \mathcal{T}_g$ and depends only on the underlying hyperbolic structure $X \in \mathcal{M}_g$. Thus, equation (5.8) gives rise to a function $B : \mathcal{M}_g \rightarrow \mathbb{R}_+$ known as the Mirzakahani function.

Exercise 5.11. Show there exists a constant $C > 0$ such that for every $X \in \mathcal{M}_g$,

$$B(X) \leq C \cdot u(X). \tag{5.9}$$

Conclude that the function $B : \mathcal{M}_g \rightarrow \mathbb{R}_+$ is integrable with respect to $\widehat{\mu}_{\text{wp}}$, that is,

$$\int_{\mathcal{M}_g} B(X) d\widehat{\mu}_{\text{wp}}(X) < \infty. \tag{5.10}$$

Hint: To prove equation (5.9), use Proposition 5.8, Theorem 5.9, and interpret μ_{Thu} as Lebesgue measure in Dehn–Thurston coordinates using Exercise 3.11. To prove equation (5.10), use Exercise 5.10.

Following Exercise 5.11, we consider the constant $b_g > 0$ defined as

$$b_g := \int_{\mathcal{M}_g} B(X) d\widehat{\mu}_{\text{wp}}(X). \tag{5.11}$$

5.6. Equidistribution of counting measures. We are finally ready to prove that the sequence of counting measure $(\mu_L^\gamma)_{L>0}$ on \mathcal{ML}_g converges in the weak- \star topology to a constant multiple of the Thurston measure μ_{Thu} . The following exercise is an analog of Theorem 2.14.

Exercise 5.12. Let $\gamma := \sum_{i=1}^k a_i \gamma_i$ be an integral simple closed multi-curve on S_g . The goal of this exercise is to show that, with respect to the weak- \star topology for measures on \mathcal{ML}_g ,

$$\lim_{L \rightarrow \infty} \mu_L^\gamma = \frac{c(\gamma)}{b_g} \cdot \mu_{\text{Thu}}.$$

Suppose $(L_k)_{k \in \mathbb{N}}$ is a sequence in \mathbb{R}_+ with $L_k \nearrow +\infty$ as $k \rightarrow \infty$ and such that, for some $c \geq 0$,

$$\lim_{k \rightarrow \infty} \mu_{L_k}^\gamma = c \cdot \mu_{\text{Thu}}.$$

We compute the following limit in two different ways,

$$\lim_{k \rightarrow \infty} \int_{\mathcal{M}_g} \frac{s(X, \gamma, L_k)}{L_k^{6g-6}} d\widehat{\mu}_{\text{wp}}(X).$$

(1) Use Exercise 5.5 to show that

$$\lim_{k \rightarrow \infty} \int_{\mathcal{M}_g} \frac{s(X, \gamma, L_k)}{L_k^{6g-6}} d\widehat{\mu}_{\text{wp}}(X) = c(\gamma).$$

(2) Use Exercise 5.10 and the dominate convergence theorem to show that

$$\lim_{k \rightarrow \infty} \int_{\mathcal{M}_g} \frac{s(X, \gamma, L_k)}{L_k^{6g-6}} d\widehat{\mu}_{\text{wp}}(X) = c \cdot b_g.$$

(3) Use the previous parts and Exercise 5.4 to conclude that

$$\lim_{L \rightarrow \infty} \mu_L^\gamma = \frac{c(\gamma)}{b_g} \cdot \mu_{\text{Thu}}.$$

5.7. *Counting simple closed multi-curves.* We have developed all the tools needed to prove the following more precise version of Theorem 5.2.

Exercise 5.13. Let $\gamma := \sum_{i=1}^k a_i \gamma_i$ be an integral simple closed multi-curve on S_g and $X \in \mathcal{T}_g$ be a marked hyperbolic structure on S_g . Show that the following identity holds:

$$\lim_{L \rightarrow \infty} \frac{s(X, \gamma, L)}{L^{6g-6}} = \frac{c(\gamma) \cdot B(X)}{b_g}.$$

Hint: Follow the same approach as in the proof of Theorem 2.16. Aside from Exercise 5.12, it will be useful to recall the identity in equation (5.3) as well as Exercise 3.12.

Now, we direct our attention to Theorem 1.1, the main result of this survey.

Exercise 5.14. Using equation (5.1) and Exercise 5.13, prove Theorem 1.1, that is, show that for every closed, orientable hyperbolic surface X of genus $g \geq 2$, there exists a constant $n(X) > 0$ such that

$$\lim_{L \rightarrow \infty} \frac{s(X, L)}{L^{6g-6}} = n(X).$$

Let us end this section by following a more precise version of Theorem 1.2.

Exercise 5.15. Let γ_1 and γ_2 be non-separating and separating simple closed curves on S_2 , respectively. Using Exercises 5.6 and 5.13, show that on any genus 2 hyperbolic surface $X \in \mathcal{M}_2$, it is 48 times more likely for a long random simple closed geodesic to be

non-separating rather than separating, that is, show that

$$\lim_{L \rightarrow \infty} \frac{s(X, \gamma_1, L)}{s(X, \gamma_2, L)} = 48.$$

6. *Beyond simple closed geodesics*

6.1. *Outline of this section.* In this section, we give a brief overview of several counting results for closed curves on surfaces and other related objects that have been proved since the debut of Mirzakhani’s thesis. Rather than exhaustively covering the great amount of material available, we aim at giving a landscape picture of the relevance of these results and of the variety of techniques behind their proofs. In particular, although many of the results that follow hold for surfaces with punctures, we focus on the case of closed surfaces. The reader is encouraged to look into the cited references for more details.

6.2. *General length functions.* For the rest of this section, we fix an integer $g \geq 2$ and a connected, oriented, closed surface S_g of genus g . Let $\gamma := \sum_{i=1}^k a_i \gamma_i$ be a simple closed multi-curve on S_g . Consider a continuous function $\ell: \mathcal{ML}_g \rightarrow \mathbb{R}_+$ on the space of measured geodesic laminations that is homogeneous, that is, such that $\ell(t \cdot \lambda) = t \cdot \ell(\lambda)$ for every $t > 0$ and every $\lambda \in \mathcal{ML}_g$. Interesting examples of such functions include the square root of extremal length with respect to a given conformal structure [Ker80] and the length of geodesic representatives with respect to an arbitrary negatively curved metric [Ota90]. For every $L > 0$, consider the counting function

$$s(\ell, \gamma, L) := \#\{\alpha \in \text{Mod}_g \cdot \gamma \mid \ell(\alpha) \leq L\}.$$

Furthermore, consider the finite, positive constant

$$B(\ell) := \mu_{\text{Thu}}(\{\lambda \in \mathcal{ML}_g \mid \ell(\lambda) \leq 1\}).$$

Recall the definition of the constants $c(\gamma) > 0$ and $b_g > 0$ introduced in equations (5.5) and (5.11). A careful consideration of the techniques introduced in the proof of Theorem 5.2 shows that similar asymptotic estimates can be proved in this more general setting. Indeed, the following holds.

Exercise 6.1. Let $\gamma := \sum_{i=1}^k a_i \gamma_i$ be an integral simple closed multi-curve on S_g and $\ell: \mathcal{ML}_g \rightarrow \mathbb{R}_+$ be a continuous, homogeneous function. Show that the following identity holds:

$$\lim_{L \rightarrow \infty} \frac{s(\ell, \gamma, L)}{L^{6g-6}} = \frac{c(\gamma) \cdot B(\ell)}{b_g}.$$

Hint: Follow the same approach as in Exercise 5.13.

6.3. *Non-simple closed geodesics.* One can also consider counting problems for closed geodesics that are not simple. Extending the definition for simple closed curves, we say two closed curves on homeomorphic surfaces have the same topological type if there exists a homeomorphism between the surfaces that identifies the free homotopy classes of the curves. A closed curve is said to be filling if every homotopically non-trivial closed curve on the surface intersects the curve.

Let $X \in \mathcal{T}_g$ be a marked hyperbolic structure and γ be a filling closed curve on S_g . For every $L > 0$, consider the counting function

$$f(X, \gamma, L) := \#\{\alpha \in \text{Mod}_g \cdot \gamma \mid \ell_\alpha(X) \leq L\}.$$

This quantity does not depend on the marking of $X \in \mathcal{T}_g$ and corresponds to the number of closed geodesics on X of the same topological type as γ and length $\leq L$. Consider the finite constant

$$c(\gamma) := \mu_{\text{Thu}}(\{\lambda \in \mathcal{ML}_g \mid i(\gamma, \lambda) \leq 1\}) / \#\text{Stab}(\gamma),$$

where $i(\gamma, \lambda)$ denotes the geometric intersection number between γ and λ , for a precise definition, see for instance [Bon88], and $\text{Stab}(\gamma) \subseteq \text{Mod}_g$ denotes the stabilizer of γ with respect to the natural mapping class group action. Recall the definition of the Mirzakhani function $B: \mathcal{M}_g \rightarrow \mathbb{R}_+$ in equation (5.8). In [Mir16], Mirzakhani introduced novel techniques to prove the following result.

THEOREM 6.2. [Mir16, Theorem 1.1] *Let $X \in \mathcal{M}_g$ be a hyperbolic structure and γ be a filling closed curve on S_g . Then, the following asymptotic formula holds:*

$$\lim_{L \rightarrow \infty} \frac{f(X, \gamma, L)}{L^{6g-6}} = \frac{c(\gamma) \cdot B(X)}{b_g}.$$

Mirzakhani's proof of Theorem 6.2 is inspired by work of Margulis [Mar70] and relies on previous work by herself on the ergodic theory of the earthquake flow [Mir08a]. More concretely, instead of counting the points in a mapping class group orbit of a filling closed curve, one can count the points in a mapping class group orbit of Teichmüller space. Then, the length of a filling closed curve on Teichmüller space is asymptotically a piecewise linear function with respect to Fenchel–Nielsen coordinates. Partitioning Teichmüller space into finer and finer cones with respect to these coordinates, one can study the equidistribution of these cones as the length of the filling closed curve goes to infinity. This can be done by using the earthquake flow, a generalization of Fenchel–Nielsen twists along simple closed geodesics but for measured geodesic laminations, and by following the general approach introduced by Margulis in his thesis.

A result analogous to Theorem 6.2 for closed geodesics of any topological type was later proved by Erlandsson and Souto [ES19] using original arguments introduced in their earlier work [ES16]. The main idea of their approach is to consider counting measures as in equation (5.2), but defined for general closed curves that are not necessarily simple. Technically speaking, such measures are defined on the space of geodesic currents, an analog of the space of measured geodesic laminations, but for closed geodesics with self-intersections; this space will be introduced below. Similar to Mirzakhani's original approach in [Mir08b], the proof then boils down to showing that such counting measures converge in the weak- \star topology to the Thurston measure. The main idea to achieve this is to study the procedure of smoothing a non-simple closed geodesic at its self-intersections.

6.4. *Geodesic currents.* In [Bon88], Bonahon gave a unified treatment of several seemingly unrelated notions of length for closed curves on closed, orientable surfaces using the concept of geodesic currents.

To define geodesic currents, let us endow the surface S_g with an auxiliary hyperbolic metric. The projective tangent bundle $PT S_g$ admits a one-dimensional foliation by lifts of geodesics on S_g . A geodesic current on S_g is a Radon transverse measure of the geodesic foliation of $PT S_g$. Equivalently, a geodesic current on S_g is a $\pi_1(S_g)$ -invariant Radon measure on the space of unoriented geodesics of the universal cover of S_g . Endow the space of geodesic currents on S_g with the weak- \star topology. Different choices of auxiliary hyperbolic metrics on S_g yield canonically identified spaces of geodesic currents [Bon88, Fact 1]. Denote the space of geodesic currents on S_g by \mathcal{C}_g . This space supports a natural \mathbb{R}_+ scaling action and a natural Mod_g action [RS19, §2].

Free homotopy classes of weighted, unoriented closed curves on S_g embed into \mathcal{C}_g by considering their geodesic representatives with respect to any auxiliary hyperbolic metric. By the work of Bonahon [Bon88, Proposition 2], this embedding is dense. Moreover, the geometric intersection number pairing for closed curves on S_g extends in a unique way to a continuous, symmetric, bilinear pairing $i(\cdot, \cdot)$ on \mathcal{C}_g [Bon88, Proposition 3]. In this sense, geodesic currents are to closed curves what measured geodesic laminations are to simple closed curves.

Many different spaces of metrics on S_g embed into \mathcal{C}_g in such a way that the geometric intersection number of a metric with any closed curve is equal to the length of the geodesic representatives of the closed curve with respect to the metric. The geodesic current corresponding to any such metric is usually referred to as its Liouville current. Examples of metrics admitting Liouville currents include hyperbolic metrics [Bon88] and negatively curved Riemannian metrics [Ota90].

A geodesic current $\alpha \in \mathcal{C}_g$ is said to be filling if $i(\alpha, \beta) > 0$ for every non-zero $\beta \in \mathcal{C}_g$. Relevant examples of filling geodesic currents include free homotopy classes of unoriented filling closed curves and the Liouville currents introduced above. Filling geodesic currents $\alpha \in \mathcal{C}_g$ have finite stabilizers $\text{Stab}(\alpha) \subseteq \text{Mod}_g$ with respect to the natural mapping class group action [EM22].

Consider a continuous function $\ell: \mathcal{C}_g \rightarrow \mathbb{R}_+$ that is homogeneous, that is, such that $\ell(t \cdot \alpha) = t \cdot \ell(\alpha)$ for every $t > 0$ and every $\alpha \in \mathcal{C}_g$. Let $\alpha \in \mathcal{C}_g$ be a filling geodesic current. For every $L > 0$, define


$$\text{cur}(\ell, \alpha, L) := \#\{\beta \in \text{Mod}_g \cdot \alpha \mid \ell(\beta) \leq L\}.$$

Furthermore, consider the finite, positive constant

$$c(\alpha) := \mu_{\text{Thu}}(\{\lambda \in \mathcal{ML}_g \mid i(\alpha, \lambda) \leq 1\})/\#\text{Stab}(\alpha).$$

In [RS19], Rafi and Souto proved the following generalization of Theorem 6.2.

THEOREM 6.3. [RS19, Main Theorem] *Let $\ell: \mathcal{C}_g \rightarrow \mathbb{R}_+$ be a continuous, homogeneous function and $\alpha \in \mathcal{C}_g$ be a filling geodesic current. Then, the following asymptotic estimate*



(a) Square-tiled surface. (b) Horizontal core multi-curve. (c) Vertical core multi-curve.

FIGURE 17. Example of a square-tiled surface of genus 2. The horizontal core multi-curve is $\alpha_1 + 2\alpha_2$. The vertical core multi-curve is $\beta_1 + \beta_2 + \beta_3$ (colour online).

holds:

$$\lim_{L \rightarrow \infty} \frac{\text{cur}(\ell, \alpha, L)}{L^{6g-6}} = \frac{B(\ell) \cdot c(\alpha)}{b_g}.$$

Theorem 6.3 was later generalized by Erlandsson and Souto for non-filling geodesic currents [ES22]. The proof of Theorem 6.3 follows a similar approach to the arguments of Erlandsson and Souto described above, but involves other technical hurdles. The real power of this result comes from its applications. As a remarkable application of Theorem 6.3, Rafi and Souto proved an asymptotic formula for the number of points in a mapping class group orbit of Teichmüller space that lie within a ball of given center and large radius with respect to Thurston’s asymmetric metric; Thurston’s asymmetric metric quantifies the minimal Lipschitz constant among Lipschitz maps between marked hyperbolic surfaces [Thu98]. Counting problems of this nature have been extensively studied in homogeneous dynamics and, in the setting of Teichmüller space, were previously studied by Athreya *et al* [ABEM12]; later, we describe this work in more detail.

6.5. *Square-tiled surfaces.* A square-tiled surface is a closed, connected, oriented surface constructed from finitely many disjoint unit area squares on the complex plane, with sides parallel to the real and imaginary axes, by identifying pairs of sides by translation and/or 180° rotation. Any such surface inherits a singular flat metric from the complex plane; the singularities of the metric are cone points with total angle an integer multiple of π . For simplicity, we only consider square-tiled surfaces with no cone points of total angle π . The horizontal core multi-curve of a square tiled-surface is the integrally weighted simple closed multi-curve obtained by concatenating the horizontal segments running through the middle of each square. The vertical core multi-curve of a square-tiled surface is defined in an analogous way. See Figure 17 for an example.

Recall that two integrally weighted simple closed multi-curves on homeomorphic surfaces have the same topological type if there exists a homeomorphism between the surfaces mapping one multi-curve to the other preserving the weights. Let $\alpha := \sum_{i=1}^k a_i \alpha_i$ be an integral simple closed multi-curve on S_g . For every $L > 0$, consider the counting function

$$sq(\alpha, L) := \# \left\{ \begin{array}{l} \text{square-tiled surfaces with horizontal core multi-curve} \\ \text{of the same topological type as } \alpha \text{ and } \leq L \text{ squares} \end{array} \right\} / \sim,$$

where \sim denotes the equivalence relation induced by cut and paste operations. As in the counting problems above, we are interested in the asymptotics of $sq(\alpha, L)$ as $L \rightarrow \infty$. Denote by

$$\epsilon_g := \begin{cases} 2 & \text{if } g = 2, \\ 1 & \text{if } g \neq 2, \end{cases} \tag{6.1}$$

the number of automorphisms of a generic square-tiled surface of genus g . Recall the definition of the constants $c(\gamma) > 0$ introduced in equation (5.5).

THEOREM 6.4. *Let $\alpha := \sum_{i=1}^k a_i \alpha_i$ be an integral simple closed multi-curve on S_g . Then,*

$$\lim_{L \rightarrow \infty} \frac{sq(\alpha, L)}{L^{6g-6}} = \frac{\epsilon_g \cdot c(\alpha)}{2^{2g-3}}.$$

One can also consider more refined counting functions of square-tiled surfaces. Let $\alpha := \sum_{i=1}^k \alpha_i \alpha_i$ and $\beta := \sum_{j=1}^l b_j \beta_j$ be integral simple closed multi-curves on S_g . For every $L > 0$, denote

$$sq(\alpha, \beta, L) := \# \left\{ \begin{array}{l} \text{square-tiled surfaces with horizontal core multi-curve of} \\ \text{the same topological type as } \alpha \text{ and vertical core multi-curve} \\ \text{of the same topological type as } \beta \text{ and } \leq L \text{ squares} \end{array} \right\} / \sim,$$

where \sim denotes the equivalence relation induced by cut and paste operations.

THEOREM 6.5. *Let $\alpha := \sum_{i=1}^k a_i \alpha_i$ and $\beta := \sum_{j=1}^l b_j \beta_j$ be integral simple closed multi-curves on S_g . Then, the following asymptotic formula holds:*

$$\lim_{L \rightarrow \infty} \frac{sq(\alpha, \beta, L)}{L^{6g-6}} = \frac{\epsilon_g \cdot c(\alpha) \cdot c(\beta)}{2^{2g-3}}.$$

Theorems 6.4 and 6.5 were originally proved by Delecroix *et al* using algebro-geometric methods [DGZZ21]. Different proofs of these theorems were later provided in [Ara20]. These proofs made crucial use of the results of Mirzakhani discussed in this survey. Indeed, square-tiled surfaces are parameterized by filling pairs of integral simple closed multi-curves [HM79]. Through this parameterization, it is possible to recast Theorems 6.4 and 6.5 as counting problems for simple closed multi-curves, which one can then tackle using Mirzakhani’s original approach.

6.6. Individual components of multi-curves. Inspired by Mirzakhani’s counting theorems, Wolpert conjectured that analogous results should hold for countings of simple closed multi-geodesics that keep track of the hyperbolic length of individual components, rather than just the total hyperbolic length.

For instance, let $X \in \mathcal{M}_g$ be a closed, connected, oriented hyperbolic surface of genus $g \geq 2$ and $\vec{\gamma} := (\gamma_1, \dots, \gamma_k)$ be an ordered simple closed multi-curve on S_g with $1 \leq k \leq 3g - 3$ components. For every $L > 0$, consider the counting function

$$m(X, \vec{\gamma}, L) := \# \left\{ \begin{array}{l} \text{ordered simple closed multi-geodesics } \vec{\alpha} := (\alpha_1, \dots, \alpha_k) \text{ on } X \\ \text{of the same topological type as } \vec{\gamma} \text{ with } \max_{i=1, \dots, k} \ell_{\alpha_i}(X) \leq L \end{array} \right\}.$$

Exercise 6.6. Let $\vec{\gamma} := (\gamma_1, \dots, \gamma_k)$ be an ordered simple closed multi-curve on S_g . Using Mirzakhani’s integration formulas gives an explicit expression for

$$M(\vec{\gamma}, L) := \int_{\mathcal{M}_g} m(X, \vec{\gamma}, L) d\widehat{\mu}_{\text{wp}}(X)$$

in terms of Weil–Petersson volume polynomials and conclude that $M(\vec{\gamma}, L)$ is a polynomial in L of degree $6g - 6$. *Hint: Follow the same approach as in Exercise 5.5.*

Following Exercise 6.6, we define

$$m(\vec{\gamma}) := \lim_{L \rightarrow \infty} \frac{M(\vec{\gamma}, L)}{L^{6g-6}}.$$

THEOREM 6.7. *Let $X \in \mathcal{M}_g$ be a hyperbolic structure on S_g and $\vec{\gamma} := (\gamma_1, \dots, \gamma_k)$ with $1 \leq k \leq 3g - 3$ be an ordered simple closed multi-curve on S_g . Then, the following asymptotic formula holds:*

$$\lim_{L \rightarrow \infty} \frac{m(X, \vec{\gamma}, L)}{L^{6g-6}} = \frac{m(\vec{\gamma}) \cdot B(X)}{b_g}.$$

In [Ara22], a generalization of Theorem 6.7 for ordered filling closed multi-curves was proved using techniques introduced by Mirzakhani in [Mir16]. A result analogous to Theorem 6.7 for all topological types of closed curves has since been proved by Erlandsson and Souto using original methods [ES22]. Theorem 6.7 was proved independently by the author [Ara22] and Liu [Liu22] using general averaging and unfolding arguments introduced by Margulis in his thesis [Mar70]. The main idea of the proof is to average over the moduli space of hyperbolic surfaces to reduce the original counting problem to an equidistribution problem for ‘horoballs’ in moduli space. This problem, in turn, can be studied using the ergodicity of the earthquake flow established by Mirzakhani in [Mir08a].

6.7. Effective countings. The counting results discussed above only provide asymptotic estimates without explicit error terms. The search for effective estimates for counting problems of simple closed geodesics has only seen progress in recent years. In [EMM22], Eskin, Mirzakhani, and Mohammadi introduced new methods fundamentally based on Teichmüller dynamics to prove the following effective version of Mirzakhani’s simple closed geodesic counting theorem.

THEOREM 6.8. [EMM22] *Let $X \in \mathcal{M}_g$ be a hyperbolic structure on S_g and $\gamma := \sum_{i=1}^k a_i \gamma_i$ be an integral simple closed multi-curve on S_g . Then, for every $L > 0$,*

$$s(X, \gamma, L) = \frac{c(\gamma) \cdot B(X)}{b_g} \cdot L^{6g-6} + O_{X,\gamma}(L^{6g-6-\kappa}),$$

where $\kappa = \kappa(g) > 0$ is a positive constant depending only on the genus $g \geq 2$.

Even more recently, in [Ara24], novel methods were introduced by the author to prove analogous effective estimates for countings of filling closed geodesics of a given topological type.

THEOREM 6.9. [Ara24] *Let $X \in \mathcal{M}_g$ be a hyperbolic structure on S_g and γ be a filling closed curve on S_g . Then, for every $L > 0$,*

$$f(X, \gamma, L) = \frac{c(\gamma) \cdot B(X)}{b_g} \cdot L^{6g-6} + O_{X,\gamma}(L^{6g-6-\kappa}),$$

where $\kappa = \kappa(g) > 0$ is a positive constant depending only on the genus $g \geq 2$.

Forthcoming work of Honaryar [Hon24] extends the techniques introduced in [Ara24] to prove a version of Theorem 6.9 for non-simple, non-filling closed curves. The approach in [Ara24] is based on a so-called ‘tracking method’. The backbone of this method is a ‘tracking principle’ established in [Ara21], which, roughly speaking, shows that the action of the mapping class group on the space of closed curves of a closed surface effectively ‘tracks’ the corresponding action on Teichmüller space in the following sense: for all but quantitatively few mapping classes, the information of how a mapping class moves a given point of Teichmüller space determines, up to a power saving error term, how it changes the geometric intersection numbers of a given closed curve with respect to arbitrary geodesic currents. This principle allows one to reduce Theorem 6.9 to an effective orbit counting problem in Teichmüller space. This problem was studied in the prequel [Ara23] building on previous work of Athreya *et al* [ABEM12], as we now explain.

6.8. Orbits in Teichmüller space. The minimal quasiconformal distortion among quasiconformal maps between marked hyperbolic surfaces is measured by the Teichmüller distance; this is a metric on Teichmüller space. Denote by $B_R(X) \subseteq \mathcal{T}_g$ the ball of radius $R > 0$ centered at $X \in \mathcal{T}_g$ with respect to the Teichmüller metric. Inspired by analogous results in homogeneous dynamics, see for instance [EM93], Athreya *et al* proved the following result in [ABEM12].

THEOREM 6.10. *There exists a positive constant $C_g > 0$ such that for every $X, Y \in \mathcal{T}_g$,*

$$\lim_{R \rightarrow \infty} \frac{\#(\text{Mod}_g \cdot Y \cap B_R(X))}{e^{(6g-6)R}} = C_g.$$

Building on the work of Athreya *et al*, the author proved the following effective version of [Ara23, Theorem 6.10].

THEOREM 6.11. *There exists a positive constant $C_g > 0$ such that for every $X, Y \in \mathcal{T}_g$,*

$$\#(\text{Mod}_g \cdot Y \cap B_R(X)) = C_g \cdot e^{(6g-6)R} + O_{X,Y}(e^{(6g-6-\kappa)R}),$$

where $\kappa = \kappa(g) > 0$ is a positive constant depending only on the genus $g \geq 2$.

The main idea behind the proof of Theorem 6.10 can already be seen in some of the results discussed above. Following Margulis’s thesis [Mar70], it is possible to reduce Theorem 6.10 to a problem about equidistribution of Teichmüller metric balls on moduli space. This problem can, in turn, be studied using the exponential mixing property of the Teichmüller geodesic flow [AG13, AGY06, AR12]. The effectivization of Theorem 6.10 in Theorem 6.11 requires a very fine analysis of the geometry and dynamics of Teichmüller space. Let us also highlight that the ‘tracking method’ discussed above can also be used to

prove an effective version of Rafi and Souto's asymptotic estimate for the number of points in a mapping class group orbit of Teichmüller space that lie within a ball of given center and large radius with respect to Thurston's asymmetric metric [Ara24].

Acknowledgments. The author is very grateful to Alex Wright and Steve Kerckhoff for their invaluable advice, patience, and encouragement. The author would also like to thank Alex Wright, Anton Zorich, and Howard Masur for their helpful comments on an earlier version of this survey. This survey got started as a set of notes for a minicourse taught by the author at the CMI-HIMR Dynamics and Geometry Online Summer School. The author is very grateful to Viveka Erlandsson, John Mackay, and Jens Marklof for giving him the chance to teach this minicourse. This survey was finished while the author was a member of the Institute for Advanced Study (IAS). The author is very grateful to the IAS for its hospitality. This material is based upon work supported by the National Science Foundation under Grant No. DMS-1926686.

REFERENCES

- [ABEM12] J. Athreya, A. Bufetov, A. Eskin and M. Mirzakhani. Lattice point asymptotics and volume growth on Teichmüller space. *Duke Math. J.* **161**(6) (2012), 1055–1111.
- [AG13] A. Avila and S. Gouëzel. Small eigenvalues of the Laplacian for algebraic measures in moduli space, and mixing properties of the Teichmüller flow. *Ann. of Math. (2)* **178**(2) (2013), 385–442.
- [AGY06] A. Avila, S. Gouëzel and J.-C. Yoccoz. Exponential mixing for the Teichmüller flow. *Publ. Math. Inst. Hautes Études Sci.* **104** (2006), 143–211.
- [AR12] A. Avila and M. J. Resende. Exponential mixing for the Teichmüller flow in the space of quadratic differentials. *Comment. Math. Helv.* **87**(3) (2012), 589–638.
- [Ara20] F. Arana-Herrera. Counting square-tiled surfaces with prescribed real and imaginary foliations and connections to Mirzakhani's asymptotics for simple closed hyperbolic geodesics. *J. Mod. Dynam.* **16** (2020), 81–107.
- [Ara21] F. Arana-Herrera. Effective mapping class group dynamics II: geometric intersection numbers. *Preprint*, 2021, [arXiv:2104.01694](https://arxiv.org/abs/2104.01694).
- [Ara22] F. Arana-Herrera. Counting multi-geodesics on hyperbolic surfaces with respect to the lengths of individual components. *Geom. Topol.* **26**(3) (2022), 1291–1347.
- [Ara23] F. Arana-Herrera. Effective mapping class group dynamics I: counting lattice points in Teichmüller space. *Duke Math. J.* **172**(8) (2023), 1437–1529.
- [Ara24] F. Arana-Herrera. Effective mapping class group dynamics III: counting filling closed curves on surfaces. *Can. J. Math.* **12**(3) (2024), 563–622.
- [Ber85] L. Bers. An inequality for Riemann surfaces. *Differential Geometry and Complex Analysis*. Ed. I. Chavel and H. M. Farkas. Springer, Berlin, 1985, pp. 87–93.
- [BM00] M. B. Bekka and M. Mayer. *Ergodic Theory and Topological Dynamics of Group Actions on Homogeneous Spaces (London Mathematical Society Lecture Note Series, 269)*. Cambridge University Press, Cambridge, 2000.
- [Bog18] V. I. Bogachev. *Weak Convergence of Measures (Mathematical Surveys and Monographs, 234)*. American Mathematical Society, Providence, RI, 2018.
- [Bon88] F. Bonahon. The geometry of Teichmüller space via geodesic currents. *Invent. Math.* **92**(1) (1988), 139–162.
- [BS85] J. S. Birman and C. Series. Geodesics with bounded intersection number on surfaces are sparsely distributed. *Topology* **24**(2) (1985), 217–225.
- [Bus92] P. Buser. *Geometry and Spectra of Compact Riemann Surfaces (Progress in Mathematics, 106)*. Birkhäuser Boston, Inc., Boston, MA, 1992.
- [DGZZ21] V. Delecroix, É. Goujard, P. Zograf and A. Zorich. Masur–Veech volumes, frequencies of simple closed geodesics, and intersection numbers of moduli spaces of curves. *Duke Math. J.* **170**(12) (2021), 2633–2718.

- [Do13] N. Do. Moduli spaces of hyperbolic surfaces and their Weil–Petersson volumes. *Handbook of Moduli. Volume 1 (Advanced Lectures in Mathematics (ALM), 24)*, International Press, Somerville, MA, 2013, pp. 217–258.
- [EM93] A. Eskin and C. McMullen. Mixing, counting, and equidistribution in Lie groups. *Duke Math. J.* **71**(1) (1993), 181–209.
- [EM22] V. Erlandsson and G. Mondello. Ergodic invariant measures on the space of geodesic currents. *Ann. Inst. Fourier (Grenoble)* **72**(6) (2022), 2449–2513.
- [EMM22] A. Eskin, M. Mirzakhani and A. Mohammadi. Effective counting of simple closed geodesics on hyperbolic surfaces. *J. Eur. Math. Soc.* **24**(9) (2022), 3059–3108.
- [ES16] V. Erlandsson and J. Souto. Counting curves in hyperbolic surfaces. *Geom. Funct. Anal.* **26**(3) (2016), 729–777.
- [ES19] V. Erlandsson and J. Souto. Mirzakhani’s curve counting. *Preprint*, 2019, [arXiv:1904.05091](https://arxiv.org/abs/1904.05091).
- [ES22] V. Erlandsson and J. Souto. *Mirzakhani’s Curve Counting and Geodesic Currents (Progress in Mathematics, 345)*. Birkhäuser, Cham, 2022.
- [EU20] V. Erlandsson and C. Uyanik. Length functions on currents and applications to dynamics and counting. *In the Tradition of Thurston*. Ed. K. Ohshika and A. Papadopoulos. Springer, Cham, 2020.
- [FM12] B. Farb and D. Margalit. *A Primer on Mapping Class Groups (Princeton Mathematical Series, 49)*. Princeton University Press, Princeton, NJ, 2012.
- [HM79] J. Hubbard and H. Masur. Quadratic differentials and foliations. *Acta Math.* **142**(3–4) (1979), 221–274.
- [Hon24] P. Honaryar. Effective counts for closed curves in hyperbolic surfaces. Manuscript in preparation, 2024.
- [Ker80] S. P. Kerckhoff. The asymptotic geometry of Teichmüller space. *Topology* **19**(1) (1980), 23–41.
- [Liu22] M. Liu. Length statistics of random multicurves on closed hyperbolic surfaces. *Groups Geom. Dyn.* **16**(2) (2022), 437–459.
- [LW24] M. Lipnowski and A. Wright. Towards optimal spectral gaps in large genus. *Ann. Probab.* **52**(2) (2024), 545–575.
- [Mar70] G. A. Margulis. On some aspects of the theory of Anosov systems. *PhD Thesis*, 1970; published in *On Some Aspects of the Theory of Anosov Systems (Springer Monographs in Mathematics)*. Springer-Verlag, Berlin, 2003.
- [Mar23] B. Martelli. *An Introduction to Geometric Topology* (B. Martelli, 2023).
- [Mas85] H. Masur. Ergodic actions of the mapping class group. *Proc. Amer. Math. Soc.* **94**(3) (1985), 455–459.
- [McS91] G. McShane. A remarkable identity for lengths of curves. *PhD Thesis*, University of Warwick, UK, ProQuest LLC, Ann Arbor, MI, 1991.
- [Mir04] M. Mirzakhani. Simple geodesics on hyperbolic surfaces and the volume of the moduli space of curves. *PhD Thesis*, Harvard University, ProQuest LLC, Ann Arbor, MI, 2004.
- [Mir07a] M. Mirzakhani. Simple geodesics and Weil–Petersson volumes of moduli spaces of bordered Riemann surfaces. *Invent. Math.* **167**(1) (2007), 179–222.
- [Mir07b] M. Mirzakhani. Weil–Petersson volumes and intersection theory on the moduli space of curves. *J. Amer. Math. Soc.* **20**(1) (2007), 1–23.
- [Mir08a] M. Mirzakhani. Ergodic theory of the earthquake flow. *Int. Math. Res. Not. IMRN* **2008**(3) (2008), Article ID rnm116.
- [Mir08b] M. Mirzakhani. Growth of the number of simple closed geodesics on hyperbolic surfaces. *Ann. of Math. (2)* **168**(1) (2008), 97–125.
- [Mir13] M. Mirzakhani. Growth of Weil–Petersson volumes and random hyperbolic surfaces of large genus. *J. Differential Geom.* **94**(2) (2013), 267–300.
- [Mir16] M. Mirzakhani. Counting Mapping class group orbits on hyperbolic surfaces. *Preprint*, 2016, [arXiv:1601.03342](https://arxiv.org/abs/1601.03342).
- [Mon22] L. Monk. Benjamini–Schramm convergence and spectrum of random hyperbolic surfaces of high genus. *Anal. PDE* **15**(3) (2022), 727–752.
- [Mos68] G. D. Mostow. Quasi-conformal mappings in n -space and the rigidity of hyperbolic space forms. *Publ. Math. Inst. Hautes Études Sci.* **34** (1968), 53–104.
- [MP19] M. Mirzakhani and B. Petri. Lengths of closed geodesics on random surfaces of large genus. *Comment. Math. Helv.* **94**(4) (2019), 869–889.
- [MR95a] G. McShane and I. Rivin. A norm on homology of surfaces and counting simple geodesics. *Int. Math. Res. Not. IMRN* **1995**(2) (1995), 61–69.
- [MR95b] G. McShane and I. Rivin. Simple curves on hyperbolic tori. *C. R. Math. Acad. Sci. Paris* **320**(12) (1995), 1523–1528.

- [MT20] L. Monk and J. Thomas. The tangle-free hypothesis on random hyperbolic surfaces. *Preprint*, 2021, arXiv:2008.09363.
- [MZ15] M. Mirzakhani and P. Zograf. Towards large genus asymptotics of intersection numbers on moduli spaces of curves. *Geom. Funct. Anal.* **25**(4) (2015), 1258–1289.
- [Ota90] J.-P. Otal. Le spectre marqué des longueurs des surfaces à courbure négative. *Ann. of Math. (2)* **131**(1) (1990), 151–162.
- [PH92] R. C. Penner and J. L. Harer. *Combinatorics of Train Tracks (Annals of Mathematics Studies, 125)*. Princeton University Press, Princeton, NJ, 1992.
- [Riv01] I. Rivin. Simple curves on surfaces. *Geom. Dedicata* **87**(1–3) (2001), 345–360.
- [RS19] K. Rafi and J. Souto. Geodesic currents and counting problems. *Geom. Funct. Anal.* **29**(3) (2019), 871–889.
- [Sie45] C. L. Siegel. A mean value theorem in geometry of numbers. *Ann. of Math. (2)* **46** (1945), 340–347.
- [Thu80] W. P. Thurston. *Geometry and Topology of Three-Manifolds (Lecture Notes)*. Princeton University Press, Princeton, NJ, 1980.
- [Thu98] W. P. Thurston. Minimal stretch maps between hyperbolic surfaces. *Preprint*, 1998, arXiv:math/9801039.
- [Wit91] E. Witten. Two-dimensional gravity and intersection theory on moduli space. *Surveys in Differential Geometry (Cambridge, MA, 1990)*. Lehigh University, Bethlehem, PA, 1991, pp. 243–310.
- [Wol83] S. Wolpert. On the symplectic geometry of deformations of a hyperbolic surface. *Ann. of Math. (2)* **117**(2) (1983), 207–234.
- [Wol13] S. A. Wolpert. Mirzakhani’s volume recursion and approach for the Witten–Kontsevich theorem on moduli tautological intersection numbers. *Moduli Spaces of Riemann Surfaces (IAS/Park City Mathematics Series, 20)*. American Mathematical Society, Providence, RI, 2013, pp. 221–266.
- [Wri20] A. Wright. A tour through Mirzakhani’s work on moduli spaces of Riemann surfaces. *Bull. Amer. Math. Soc. (N.S.)* **57**(3) (2020), 359–408.