

The potential role of small fish species in improving micronutrient deficiencies in developing countries: building evidence

Nozomi Kawarazuka*† and Christophe Béné‡

Policy, Economics and Social Sciences, The WorldFish Center, Consultative Group on International Agricultural Research (CGIAR), Jalan Batu Maung, Batu Maung, 11960 Bayan Lepas, Penang, Malaysia

Submitted 26 July 2010; Accepted 9 March 2011; First published online 20 May 2011

Abstract

Objective: To build a comprehensive overview of the potential role of fish in improving nutrition with respect to certain micronutrient deficiencies in developing countries.

Design: A comprehensive literature review was completed. For this the electronic library databases ASFA, CABD and Scopus were systematically searched and relevant references cited in these sources were carefully analysed. The search terms used were 'fish', 'small fish species', 'micronutrients', 'food-based strategies', 'fish consumption' and 'developing countries'. The quality of data on nutritional analyses was carefully reviewed and data that lacked proper information on methods, units and samples were excluded.

Results: The evidence collected confirmed the high levels of vitamin A, Fe and Zn in some of the small fish species in developing countries. These small fish are reported to be more affordable and accessible than the larger fish and other usual animal-source foods and vegetables. Evidence suggests that these locally available small fish have considerable potential as cost-effective food-based strategies to enhance micronutrient intakes or as a complementary food for undernourished children. However, the present review shows that only a few studies have been able to rigorously assess the impact of fish consumption on improved nutritional status in developing countries.

Conclusions: Further research is required in areas such as determination of fish consumption patterns of poor households, the nutritional value of local fish and other aquatic animals and the impact of fish intake on improved nutritional status in developing countries where undernutrition is a major public health problem.

Keywords
 Micronutrient deficiency
 Small fish species
 Small-scale fisheries
 Aquaculture
 Developing countries

At present, more than two billion people worldwide, in particular in developing countries, are estimated to be deficient in essential vitamins and minerals, especially in vitamin A, Fe and Zn^(1,2). Micronutrient deficiencies occurring at particular stages of human life (pregnancy, breast-feeding, childhood) can severely affect health and development, leading in some cases to irreversible effects.

Fish can potentially contribute to reducing these micronutrient deficiencies. A few studies investigating this issue have been published in recent years. However, most of these studies are isolated stand-alone analyses, focusing on specific aspects of the problem. The overall contribution of fish to nutritional security is yet to be fully assessed. The purpose of the present study was to address this gap. Our main objective was to review and collate this scattered and relatively scarce literature in order to produce the first

global overview of the role played by fish in improving nutrition in developing countries. In doing so, the quality of the data was carefully reviewed and data that lacked proper information on methods, units or sample size were excluded. Particular effort was made to highlight not only the information recently generated but also the gap in knowledge where more research is needed. Our focus was on developing countries where the largest proportion of people exposed to risk of undernutrition is found and where 95% of the population depends on small-scale fisheries or small-scale aquaculture for their livelihood⁽³⁾.

Understanding the nutritional importance of fish in developing countries

Fish as a major animal source of food in food-deficient countries

At the global level, fish consumption has increased from an average of 10.1 kg/capita per year in 1965 to 16.4 kg in 2005⁽⁴⁾. However, the amount of fish consumption varies

† Correspondence address: School of International Development, University of East Anglia, Norwich, NR4 7TJ, UK.

‡ Present address: Vulnerability and Poverty Reduction team, Institute of Development Studies, University of Sussex, East Sussex, UK.

among regions. Figure 1 shows changes in fish consumption between 1965 and 2005 for different regions of the world. In all regions, fish consumption per capita has increased, except in sub-Saharan Africa and Latin America/the Caribbean, where it has stagnated from the early 1970s⁽⁴⁾.

FAO food balance sheets were used to estimate the contribution of fish to total protein and total animal protein at the country level⁽⁴⁾. Among the thirty countries in the world where fish contribute more than one-third of the total animal protein supply (Fig. 2a), twenty-two are officially referred to as low-income food-deficient (LIFD) countries⁽⁵⁾. In other words, a large majority (73%) of the countries where fish is an important source of food are poor and food deficient. However, when other sources of protein (i.e. plant) are considered, the contribution of fish

to total protein consumption is substantially low (Fig. 2b), indicating that in LIFD countries the majority of protein comes from plant-source foods. These statistics based on protein quantity at the national level, however, overlook the contribution of fish in terms of quality of protein and other nutrients.

Fish consumption pattern of the poor

Overall, data on fish consumption are scarce and poorly reflected in national statistics. However, the few field data that are available at household/community levels reveal the nutritional importance of fish among the poor. In Kapasia, Bangladesh, for example, the mean fish intake was as high as 83–96 g/person per d for whole fish in 1998–1999⁽⁶⁾, which was more than three times the national level (estimated by fish supply) in the same year⁽⁴⁾. In Cambodia, surveys conducted in Svag Rineng province in 1997–1998 showed an average fish intake of 70 g/person per d for raw, cleaned parts (adjusted for cleaning loss of 30% weight of raw, whole fish) and an intake of 9 g/person per d for raw, cleaned parts of other aquatic animals (*n* 66)⁽⁷⁾, whereas the latest national statistics in Cambodia show only an average of 6.84 g/person per d⁽⁴⁾. More recent surveys focusing on women in Kompong Chhnang, Prey veng and Kampong speu Provinces (Cambodia) suggest that the mean intake of fish could be as high as 103 g/d for raw, cleaned parts (*n* 163)⁽⁸⁾.

Regarding species consumed by the poor in rural areas in Asia, a variety of small indigenous species account for 50–80% of the total amount of fish consumed^(9–11). This higher dependence on smaller fish is explained by the fact that poor people can afford only comparatively




Fig. 1 Changes in fish consumption per capita (estimated by per capita availability of fish as food) for different developing regions (■, world; —, Latin America and the Caribbean; —, Oceania developing countries; —, South and South-East Asia; —, sub-Saharan Africa; data were calculated from the FAO food balance sheet⁽⁴⁾)




Fig. 2 (a) Fish as a percentage of animal protein consumption; (b) total protein consumption in g/capita per d (□, plant-source protein; ■, other animal protein; ■, fish protein; data were calculated from the FAO food balance sheet⁽⁴⁾)

cheaper fish species, whereas the better-off households purchase larger, medium-sized fish species, which they prefer because of the fact that they have fewer bones, more flesh and taste better⁽⁸⁾. In addition to wild small fish, data from Laos and Cambodia indicate that other aquatic animals such as frogs, freshwater molluscs and snails are frequently included in the everyday diet among the poor^(8,12).

Similar trends seem to exist in sub-Saharan Africa. In Nigeria, household data indicate values of fish intake of approximately 217 g/person per d for whole fish in households in the inland state of Niger and 124 g/person per d for whole fish in the coastal state of Lagos⁽¹³⁾, whereas the national statistics indicate a consumption level of 24.6 g/person per d⁽⁴⁾. In East Africa, the small indigenous species *dagaa/omena/mukene* (*Rastrineobola argentea*) from Lake Victoria is one relatively cheap fish that is consumed dried among the poor households^(14–16). Similarly, in Lubumbashi, Democratic Republic of the Congo (Congo DR), the small dried fish *kapenta* (*Limnothrissa miodon*) from Lake Kariba was reported to be the most consumed fish product by the lowest-income households⁽¹⁷⁾.

In general, one would expect the population that benefits from fish nutritional intake to be found along the coasts. However, data show that such benefits are also found among the inland population, as fish is a highly commercialised commodity that 'migrates' extensively within and between countries, even after it is caught. In sub-Saharan Africa, for instance, dried and smoked fish from Lake Chad travel more than 1000 km through a very efficient truck transportation network to be traded in urban markets in the south of Nigeria⁽¹⁸⁾. Similarly, in Ghana, smoked fish from coastal villages are traded as far as 600 km to Northern Ghana and sometimes even further north to Burkina Faso⁽¹⁹⁾. In Lubumbashi, Congo DR, a large part of the smoked, dried and salted fish that are sold in markets come from Zambia, Tanzania, Zimbabwe, Malawi or even Mozambique⁽¹⁷⁾.

In summary, the data available from household surveys suggest that small fish, other aquatic animals and processed fish of low market value play a very important role in the diet of the poor not only through subsistence fishing but also through extensive market networks. The scattered evidence from the literature indicates an average fish intake of approximately 70–150 g/person per d for cleaned parts, confirming the nutritional significance of fish for the poor in Asia and sub-Saharan Africa.

The contribution of fish intake to improving nutrition

The contribution of fish to human nutrition and its impact on health have been examined from different perspectives in both developed and developing countries.

In developed countries, the major focus has been on PUFA from fish and fish oil, which lowers blood pressure, reduces the risk of heart disease⁽²⁰⁾ and boosts infant growth and cognitive development⁽²¹⁾. In contrast, in developing countries, focus has been on the role of fish in tackling undernutrition and micronutrient deficiencies.

Nutrients in fish

The nutrient content of fish varies with species. There are relatively limited data on the specific nutritional composition of the fish species consumed in developing countries, with some exceptions. In Bangladesh, the micronutrient content of approximately twenty small indigenous species has been examined and several nutrient-dense fish have been identified^(22,23). Partial information is also available on twenty-nine small indigenous species in Cambodia^(24,25). In north-east Thailand, fatty acid composition in five fish species and in a type of prawn that inhabits rice fields was established⁽²⁶⁾. Table 1 summarizes the data found in the literature, grouped into three categories of differential nutritional significance: large freshwater fish, small freshwater fish and marine fish. Data for the latter group were extracted from the US Department of Agriculture⁽²⁷⁾. For the purpose of comparison, the nutrient content of some other food items is also displayed. Bold indicates high content values.

Macronutrients in fish

Protein

Protein from fish contributes to the overall protein intake significantly as the digestibility of protein from fish is approximately 5–15% higher than that from plants⁽²⁸⁾. Furthermore, protein from fish helps in the absorption of protein from plants. Staple foods such as rice or maize contain only a small amount of lysine, an essential amino acid, limiting the total absorption of protein. In contrast, animal sources of food such as fish have more balanced concentrations of all essential amino acids, and the concentration of lysine is particularly high⁽²⁸⁾. When fish is added to a plant-based diet, the total protein intake increases as lysine in fish compensates for the shortage of lysine in the rest of the diet. Therefore, fish play an important role in plant-based diets in LIFD countries.

Lipids

The lipid composition of fish is unique, having PUFA in the form of arachidonic acid (20:4n-6), EPA (20:5n-3) and DHA (22:6n-3), with many potential beneficial effects for adult health⁽²⁰⁾ and child development⁽²¹⁾. The amount of PUFA in large freshwater fish such as carp and tilapia is relatively low, whereas the amount in smaller indigenous species is yet to be determined. Among fish species that are cheaper and traded in developing countries, small pelagic forage fish such as anchovies and sardines⁽²⁹⁾ are perhaps some of the richest sources of PUFA⁽²⁷⁾.

Table 1 The nutrient content of fish and other foods (per 100 g)*

Group	Scientific name/common name (local name/common name)	Protein (g)	Fat					Ca (mg)	Fe (mg)	Zn (mg)	Vitamin A (RAE)†	Notes (per 100 g)	Source	
			Total lipid (fat; g)	Total saturated fat (g)	Total PUFA (g)	EPA (g)	DHA (g)							
Large freshwater fish and prawn	Carp	17·83	5·60	1·08	1·431	0·238	0·114	41	1·24	1·48	9	Raw, edible	(27)	
	Catfish	15·60	7·59	1·77	1·568	0·067	0·207	9	0·50	0·74	15	Farmed, raw, edible	(27)	
	<i>Channa striatus</i> (Snakehead)		0·99	0·34	0·475	<0·001	0·133					Raw, whole, Thailand	(26)	
	Tilapia	20·80	1·70	0·77	0·476	0·007	0·113	10	0·56	0·33	0	Raw, edible	(27)	
	<i>Macrobrachium nipponense</i> (Prawn)		1·13	0·37	0·020	0·008	0·061					Raw, whole, Thailand	(26)	
Small freshwater fish	<i>Amblypharyngodon mola</i> (Mola)							776	5·70	3·20	> 2680	Raw, edible, Bangladesh	(23)	
	<i>Esomus danricus</i> (Darkina)							775	12·00	4·00	500–1500	Raw, edible, Bangladesh	(23)	
	<i>Esomus longimanus</i> (Chanwa phlieng)							350	45·10	20·30	100–500	Raw, edible, Cambodia	(24,25)	
	<i>Helostoma temmincki</i> (Kanthrawb)							432‡	5·3‡	6·5‡	100–500	Raw, edible, Cambodia	(24,25)	
	<i>Puntius ticto</i> (Puti)							992	3·00	3·10	500–1500	Raw, edible, Bangladesh	(23)	
	<i>Rasbora tornieri</i> (Changwa mool)							700‡	0·70‡	2·7‡	> 1500	Raw, edible, Cambodia	(24,25)	
	<i>Anabas testudineus</i> (Climbing perch)		0·99	0·34	0·384	<0·001	0·088					Raw, whole, Thailand	(26)	
	<i>Puntius brevis</i> (Swamp barb)		0·90	0·31	0·314	0·000	0·047					Raw, whole, Thailand	(26)	
	<i>Rasbora borapensis</i> (Blackline rasbora)		0·86	0·33	0·319	0·002	0·083					Raw, whole, Thailand	(26)	
	Marine fish	Anchovy	20·35	4·84	1·28	1·637	0·538	0·911	147	3·25	1·72	15	Raw, edible, European	(27)
Herring		16·39	9·04	2·04	2·423	0·969	0·689	83	1·12	0·99	32	Raw, edible, Pacific	(27)	
Mackerel		18·60	13·89	3·26	3·350	0·898	1·401	12	1·63	0·63	50	Raw, edible	(27)	
Milkfish		20·53	6·73	1·67	1·840			51	0·32	0·82	30	Raw, edible, Philippines	(27)	
Sardine		24·60	11·45	1·53	5·148	0·470	0·509	382	2·92	1·31	33	Canned in oil, drained solids with bone	(27)	
Other animal-source foods		Beef ground	14·30	30·00	11·29	0·696			24	1·64	3·57	0	Raw, ground, 70% lean meat 30% fat	(27)
	Chicken breast	14·70	15·75	3·26	3·340			19	1·11	0·78	0	Breast tender, uncooked	(27)	
	Chicken egg	35·60	9·94	3·10	7·555	0·004	0·037	171	3·23	1·11	140	Raw, whole	(27)	
	Chicken liver	16·90	4·83	1·56	1·306			8	8·99	2·67	3292	All classes, raw	(27)	
	Cow's milk	3·28	3·66	2·28	0·136			119	0·05	0·37	33	3·7% milk fat	(27)	
Plant-source foods	Cassava	1·40	0·28	0·28	0·048			16	0·27	0·34	1	Raw	(27)	
	Rice	2·69	0·28	0·28	0·323			10	1·20	0·49	0	White, long-grained, regular, cooked	(27)	
	Kidney beans	8·67	0·09	0·09	0·278			35	2·22	0·86	0	Mature, cooked	(27)	
	Carrot	0·93	0·17	0·04	0·117			33	0·30	0·24	835	Raw	(27)	
	Kale	3·30	0·70	0·70	0·338			135	1·70	0·44	769	Raw	(27)	
	Spinach	2·86	0·39	0·39	0·165			99	2·71	0·53	469	Raw	(27)	
	High content	> 15·00			>2·000	>0·400	>0·400	>100	>3·00	>3·50	>500			

RAE, retinol activity equivalents.

Bold indicates high content values.

*Authors' own compilation. Nutritional composition data from references listed.

†See Food and Nutrition Board, Institute of Medicine⁽⁶¹⁾.

‡Raw, cleaned parts.

Fish intake influences the PUFA levels in the breast milk of lactating women. In China the level of DHA in the breast milk of women living in coastal regions has been shown to be higher than the level in other regions⁽³⁰⁾. Similarly, in Tanzania, women with high intakes of freshwater fish had levels of arachidonic acid and DHA in their breast milk that were above the present recommendations for infant formulae⁽³¹⁾. However, it is still not clear how the PUFA in breast milk contributes to fetal and infant development, and further investigations are required into the quantities and nutritional significance of the fatty acids in fish species commonly consumed by the poor^(32,33).

Small fish as a source of micronutrients

Little attention has been given so far to the role of fish as a source of micronutrients. However, recent research suggests that small fish species that are consumed whole with bones, heads and viscera play a critical role in micronutrient intakes, as these parts are where most micronutrients are concentrated. Small fish also offer other nutritional advantages: they can be processed and stored for a long period; they are more affordable for the poor as they can be purchased in small quantities; and they can also be more evenly divided among household members⁽²²⁾. The contribution of fish to micronutrient intakes is therefore determined not only by the nutrient content of the species but also by the local processing methods and eating patterns. As a consequence, several studies have indicated the actual nutrient content of the edible part by reflecting the local methods used to clean and prepare the fish for the meal (e.g. leaving or cutting off the head, removing a part of the viscera) and correcting the calculation for plate waste after meals^(8,23–25,32) (see Table 2 for details).

Although small fish are rich in Ca and some forms of marine fish are rich in iodine, we focus specifically on vitamin A, Fe and Zn, deficiencies of which are widely spread and for which sustainable solutions have not been found yet.

Vitamin A

Dark-green, orange and yellow vegetables, which contain provitamin A carotenoids, have long been considered a major source of vitamin A and are often used in food-based interventions in order to increase vitamin A intake. However, as indicated in Table 1, some small fish are also very rich in vitamin A. In Bangladesh, two species, *mola* (*Amblypharyngodon mola*) and *chanda* (*Parambassis baculis*), were identified as having a vitamin A content as high as 2500 and 1500 µg retinol activity equivalents (RAE)/100 g raw edible parts, respectively⁽²³⁾. In Cambodia, the small indigenous species *chanteas pbluk* (*Parachela sianensis*) and *changwa mool* (*Rasbora tornieri*) were reported to contain >1500 µg RAE/100 g raw edible parts⁽²⁴⁾. Fish often exceed vegetables with regard to both

the amount and frequency of consumption in some rural areas in Asia. For example, in Cambodia, children consume more fish than vegetables during all three seasons (mean intake 65.2 g/d per raw, cleaned parts for fish and 19 g/d for vegetables, *n* 163). At the household level, there is little difference in the amount of consumption between fish and vegetables (mean intake 654.6 g/d for raw, cleaned parts for fish and 765.1 g/d for vegetables) while fish are consumed more frequently than vegetables – 54.6% of the surveyed households consumed fish 7 d/week, *v.* 47.9% consumed vegetables⁽⁸⁾. In Kishoreganj, Bangladesh, field data show that daily consumption of small fish contributes 40% of the total daily requirement of vitamin A at the household level⁽³⁴⁾. Finally, vitamin A being a fat-soluble vitamin, a small quantity of fat (e.g. 5 g fat/person per meal) is sufficient to ensure adequate bioefficacy⁽³⁵⁾. Fish that are rich in vitamin A, cooked with some vegetables and some vegetable oils, are therefore an ideal combination to enhance vitamin A intake and bioefficacy.

There are two problems regarding vitamin A in fish. First, vitamin A content is species specific. The content may therefore be very different among fish species which belong to very close taxonomic groups. Similarly there is no direct relationship with habitat, and species living in the same habitat may have totally different vitamin A contents⁽³⁶⁾. A former study that examined vitamin A in the flesh of 157 species found that nearly 85% of the species contained little vitamin A (<60 µg RAE/100 g flesh), whereas a few others were extremely rich (e.g. >18 000 µg RAE/100 g flesh for the highest)⁽³⁶⁾. Identifying particular species and promoting the consumption and conservation of those species continue to be a big challenge in many developing countries. Second, vitamin A in freshwater fish exists mainly in the form of 3,4-dehydroretinol. In Bangladesh, a study examined the efficacy of the intake of the local small fish *mola* in the daily diet (9 weeks, 6 d/week) in improving the vitamin A status of children measured through marginal serum retinol concentration in blood⁽³⁷⁾. However, no significant effect was found in the group of children fed fish curry, suggesting that vitamin A from *mola*, of which 80% is 3,4-dehydroretinol, was not converted to retinol or was converted at an insufficient rate. Further studies on the effect of 3,4-dehydroretinol in man are clearly needed to confirm or refute this result, as well as to develop and test other indicators and approaches to assessing vitamin A status.

Fe

Some fish (especially small species) are rich in Fe (Tables 1 and 2); however, this nutrient is usually concentrated in the fish head and viscera. In a study conducted in Cambodia, the species *chanwa pblheng* (*Esomus longimanus*) was found to have a high content of Fe in its edible parts, even after the viscera had been removed through traditional cleaning methods. A serving of the sour soup made with this type of fish, eaten with boiled

Table 2 The micronutrient contents of small indigenous species in Bangladesh and Cambodia* (per 100 g, raw, edible†)

	Scientific name	Common name/ local name	Fe			Zn (mg)	Vitamin A (RAE)§	Source	
			Ca (mg)‡	T-Fe (mg)	Hm-Fe (mg)				Hm-Fe/T-Fe (%)
Bangladesh									
Small fish	<i>Amblypharyngodon mola</i>	Mola	776	5·7		3·2	> 2680	(23)	
	<i>Chanda beculis</i>	Chanda	349	0·8		1·8	> 1500	(23)	
	<i>Chanda nama</i>	Chanda	863	2·1		2·0	100–500	(23)	
	<i>Chanda ranga</i>	Chanda	1061	2·1		2·6	100–500	(23)	
	<i>Chanda ssp.</i>	Chanda	879	1·8		2·3		(23)	
	<i>Channa punctuatus</i>	Taki	199	1·8		1·5		(23)	
	<i>Corica soborna</i>	Kaski	443	2·8		3·1	< 100	(23)	
	<i>Esomus danricus</i>	Darkina	775	12·0		4·0	500–1500	(23)	
	<i>Gudusia chapra</i>	Chapila	786	7·6		2·1	< 100	(23)	
	<i>Mascrognathus ssp.</i>	Chikra	203	2·4		1·2	< 100	(23)	
	<i>Mastocembelus aculeatus</i>	Chikra	201	2·5		1·2	< 100	(23)	
	<i>Mastocembelus armatus</i>	Chikra	198	1·9		1·1		(23)	
	<i>Mastocembelus pancholus</i>	Chikra	216	2·7		1·3	< 100	(23)	
	<i>Mystus vittatus</i>	Tengra	481	4·0		3·1		(23)	
	<i>Puntius chola</i>	Puti	750	4·1		3·1	< 100	(23)	
	<i>Puntius ticto</i>	Puti	992	3·4		3·8	500–1500	(23)	
	<i>Putius ssp.</i>	Puti	785	3·0		3·1	< 100	(23)	
	<i>Putius sophore</i>	Puti	698	2·2		2·9	< 100	(23)	
Large fish	<i>Chirrhinus mrigala</i>	Mrigal	0	2·5		1·5	< 100	(23)	
	<i>Hilsa ilisha</i>	Hilsa	0				69	(62)	
	<i>Hypophthalmichthys molitrix</i>	Silver carp	4	4·4		1·4	< 100	(23)	
	<i>Labeo rohita</i>	Rui	317				27	(62)	
Juvenile	<i>Colisha lalius</i>	Tilapia					19	(62)	
	<i>H. molitrix</i>	Silver carp					13	(62)	
Cambodia									
Indigenous species common in commercial catches	<i>Anguilla bicolor</i>	Chlok					< 100	(24)	
	<i>Channa marulius</i>	Ros/Ptuok/Raws	604	6·2 	1·3	77	6·1 	100–500¶	(24,25)
	<i>Channa micropeltes</i>	Diep/Chhaur	453	5·2 	4·0	76	6·0 	< 100	(24,25)
	<i>Cyclocheilichthys apogon</i>	Srawka kdam	483	2·9	2·2	71	8·7 	< 100	(24,25)
	<i>Cyclocheilichthys armatus</i>	Pka kor					< 100	(24)	
	<i>Dangia lineata</i>	Khnaung veng					< 100	(24)	
	<i>Dangia spilopleura</i>	Arch kok	325	7·6 	5·4	70	7·1	< 100	(24,25)
	<i>Henicorhynchus siamensis</i>	Kantrawb						100–500¶	(24)
	<i>Notopterus notopterus</i>	Slat						< 100	(24)
	<i>Osteochilus hasselti</i>	Kros	414	4·2	2·2	54	6·8 	< 100	(24,25)
	<i>Parambassis wolffi</i>	Kantrang preng	466	5·7 	4·6	78	6·7 	100–500	(24,25)
	<i>Puntioplites proctozystron</i>	Chra keng	267	3·4	2·3	66	5·2 	500–1500	(24,25)
	<i>Thynnichthys thynnoides</i>	Linh						500–1500	(24)
Small fish species with low market value	<i>Dermogenys pusilla</i>	Phtong	416	3·6	2·1	56	11·0	< 100	(24,25)
	<i>Helostoma temmincki</i>	Kanthrawb	432	5·3 	3·7	71	6·5 	100–500¶	(24,25)
	<i>Parachela siamensis</i>	Chunteas phluk	243	5·0 	3·4	67	9·1	100–500	(24,25)
	<i>P. siamensis</i> (juvenile)	Chunteas phluk						< 100	(24)
	<i>Trichogaster microlepis</i>	Kamphleanh phluk	373	5·0	3·3	67	6·5	100–500¶	(24,25)
	<i>Trichogaster trichopterus</i>	Kawmphleanh samrei						100–500¶	(24)
Other small non-commercial species common in rice fields	<i>Clupeoides borneensis</i>	Bawndol ampeou						100–500	(24)
	<i>Corica laciniata</i>	Bawndol ampeou						100–500¶	(24)
	<i>Esomus longimanus</i>	Chanwa phlieng	350	45·1 	36·0	78	20·3 	100–500	(24,25)
	<i>Euryglossa panoides</i>	Andat chhke veng	439	5·2	3·9	72	7·1	100–500¶	(24,25)

Table 2 Continued

Scientific name	Common name/ local name	Fe					Source
		Ca (mg)†	T-Fe (mg)	Hm-Fe (mg)	Hm-Fe/T-Fe (%)	Zn (mg)	
<i>Luciosoma setigerum</i>	Changwa ronaung	304	2-7	2-0	72	11-4	100-500* (24)
<i>Rasbora tornieri</i>	Changwe mool						100-500 (24)
<i>Trichopsis pumila</i>	Kroem tun sai						100-500* (24)
<i>Barbodes altus</i>	Kahe	216	3-4	2-7	76	4-1	<100 (24,25)
<i>Barbodes gonionotus</i>	Chhpin	204	3-4	2-6	76	4-4	100-500 (24,25)
<i>Osteochilus melanopleus</i>	Krum						100-500* (24)
	High content	>700	>5.0			>4.0	>500

T-Fe, total Fe; Hm-Fe, haem Fe; RAE, retinol activity equivalents.
 Bold indicates high content values.
 *Authors' own compilation. Nutritional composition data from references listed.
 †Edible parts were estimated by employing local women to clean the fish according to traditional practices^(23,32).
 ‡The Ca content in edible parts was calculated from plate waste in Bangladesh⁽²³⁾ (p. 53). The Ca content in Cambodian fish is calculated from raw, cleaned parts, except for *E. longimanus*, in which all fish bones were found to be consumed and therefore the Ca intake from fish was proportional to the measured Ca content⁽²⁵⁾ (p. 1231).
 §1 RAE = 1 µg all-trans retinol = 1 RE (retinol equivalent)⁽⁶¹⁾. All-trans 3,4-dehydroretinol and all-cis 3,4-dehydroretinol found in examined fish were calculated as having 40% activity in relation to all-trans retinol, and 16% activity in relation to β-carotene^(23,32).
 ||Raw, cleaned parts.
 ¶Raw, whole fish.

rice – the most common dish in the study area – was shown to supply on average 45% of the daily requirement of Fe in women of childbearing age and 42% of that in children⁽²⁴⁾. Another important point with regard to Fe is the fact that the composition of Fe in fish is different from that found in plant-source foods: fish contain large amounts of haem Fe (Table 2), which is characterized by high bioavailability as opposed to non-haem Fe.

Zn

Cereals and legumes contain inhibitors of Zn absorption, such as phytate⁽³⁸⁾. The habitual diets of the poor, which are dominated by staple foods, therefore reduce Zn (as well as other minerals) bioavailability, and little Zn intake is expected from such diets. Yet, the daily Zn requirement in women in the third semester of pregnancy and among those lactating is as high as 20 mg/d⁽³⁹⁾. Likewise, children require 8.3–11.2 mg Zn/d depending on their body weight. These daily requirements are difficult to meet unless a significant amount of additional Zn is taken every day (unlike vitamin A, Zn cannot be stored in the human body⁽³⁹⁾ and is therefore needed in everyday diet).

Small fish are very rich in Zn compared with other animal-source foods and large fish species (Tables 1 and 2). The small low-market-value species *chanwa philieng*, for instance, which is commonly consumed by poor people in Cambodia, was found to contain 20.3 mg Zn/100 g raw edible part⁽²⁵⁾. A serving of the local sour soup dish contains on average 49 g of cleaned *chanwa philieng*, thus covering approximately 50% of the daily Zn requirement for pregnant and lactating women⁽²⁵⁾. Another survey in the same country shows that fish contribute 33% and 39% of the total daily requirement of Zn in children and women, respectively⁽⁸⁾. Small fish in a plant-based diet is therefore expected to increase Zn intake considerably and to compensate for the low bioavailability induced by the phytate of the staple foods.

Although its deficiency is not prevalent, small fish consumed with bones are a very efficient source of Ca. Their bioavailability is as high as that of milk⁽⁴⁰⁾, whereas the concentration is approximately eight times higher than that of milk⁽⁴¹⁾ (Table 2). According to a study conducted in Kishoreganj district in Bangladesh, an average daily small fish consumption of 65 g/person for edible part (cleaning and plate waste) can meet 31% of the average daily requirement of Ca⁽³⁴⁾, whereas in Cambodia fish contribute 53% of the daily requirement in children⁽⁸⁾. Small fish with bones are therefore introduced as a complementary food for children where milk is not available or affordable⁽⁴²⁾.

Fish processing and its effects on the nutritional value of products

Nutritional loss through processing is a specific issue with fish, whereas that through cooking is common to

many other foods including vegetables. Although in developing countries fish is often sun-dried, salted, fermented or smoked for preservation purposes, protein, fat and minerals remain stable even after processing. In contrast, vitamin A is sensitive to sunlight and heat. A study in Thailand found that boiled and sun-drying processing methods destroy 90% of the vitamin A content in small fish. In contrast, steaming and oven-drying were shown to result in a 50% loss only⁽⁴³⁾. In Bangladesh it was found that nearly all vitamin A in small fish is destroyed after sun-drying⁽²³⁾.

These results, however, need to be considered with a series of caveats. First, unlike other minerals and water-soluble vitamins, vitamin A can be stored in the human liver for 3–4 months⁽⁴⁴⁾. In Bangladesh and Cambodia, people mainly consume sun-dried fish only during the low-productive season^(8,23); therefore, vitamin A absorbed from small fish that are consumed fresh or with minimum processing will still be present in the human body weeks after the fish have been consumed, thus contributing to meeting long-term nutritional needs. Second, although they destroy certain vitamins, processing techniques still contribute greatly to extending the period during which fish can be consumed (by up to 4 or 5 months), thereby prolonging the time during which people who consume these processed fish can benefit from the remaining protein and micronutrients. Third, as mentioned above, loss of nutritional content through cleaning and cooking is not exclusive to fish. Vegetables that are boiled or fried also lose part of their nutritional value for the same reason.

The effects of fish in improving nutritional status

Fish as a complementary food for undernourished children

As part of the food-based approaches discussed in the literature, dietary diversification strategies suggest improving micronutrient intakes by promoting production and consumption of locally available nutritious foods⁽⁴⁵⁾. In that context, utilising locally available small fish has considerable potential as a cost-effective food-based strategy to enhance micronutrient intakes. Earlier studies conducted in the 1990s were reviewed by Caulfield *et al.*⁽⁴⁶⁾. Although fish was not the main focus in these experiments, recent studies in sub-Saharan Africa used small fish to increase intakes of Fe, Zn and Ca. In Malawi, for instance, Gibson *et al.*⁽⁴²⁾ introduced fermented porridge mixed with whole-dried fish with bones and fruit as a complementary food. In addition, information on nutrition was also provided to mothers. After 12 months, the children in the intervention group showed lower incidence of common infectious illnesses and anaemia compared with control children, although no significant changes in hair Zn concentration and growth were found.

In Ghana, a study reported that fish powder from smoked anchovies mixed with fermented maize porridge supports rates of infant growth comparable to those obtained from a cereal–legume blend with vitamin- and mineral-fortified supplements, indicating the potential role of local fish in the improvement of infant growth⁽⁴⁷⁾.

Finally, a study in Uganda used local dried fish *mukene* as an ingredient of low-cost supplement porridge to feed undernourished children. The experiment shows better outcomes in weight growth and mortality compared with the diets of imported skimmed milk that are usually used for undernourished children in hospitals⁽⁴⁸⁾.

Challenges

Several challenges still remain in assessing and improving interventions on the contribution of fish to human nutritional status.

Measuring nutritional outcomes

Fish intake may improve the micronutrient content of a person's diet; yet, this does not necessarily mean that it will improve the nutritional status of that person or that this impact can be appropriately measured. Studies in Ghana, Malawi and Bangladesh, for instance, that tested the effect of fish intake on micronutrient status using biochemical indicators such as ferritin score, hair Zn concentration and retinol score found no statistically significant effects on the tested groups^(37,42,47,49), raising the question of whether appropriate indicators were used. However, the issue related to the choice of responsive nutritional outcomes to measure the impact of increased fish intake is not specific to fish; it is common to all experiments and interventions aimed at measuring the impact of changes in the intake of any food (e.g. meat intake, biofortified foods, etc.). Overall, problems continue to persist with regard to 'demonstrating' the impact on micronutrient status or other functional outcomes (infections, growth, etc.) and these problems apply to all food-based approaches, not only to fish-related ones. The reason for this is that the entire diet of the tested groups cannot be completely controlled, unless trials are conducted in clinical settings. The issue is that if people consume, say, more fish they may 'compensate' and eat less of other foods like chicken or other meat products. Demonstrating impact is therefore complex and does not depend only on having the right indicators; it also depends on having the right evaluation design. The latter, rather than the former, has been the biggest problem in evaluating food-based approaches up to now.

Sustainable supply of nutrient-dense fish

Data from Bangladesh and Cambodia confirm the potential of small indigenous species to improve nutritional status. For instance, for the vitamin A-rich small fish *mola* which is already present in 1.3 million of the small and seasonal fish

Table 3 Summary of nutritional role by fish groups and gaps in knowledge

Group	Main nutrients	Species (examples)	Identified contribution to improving nutritional status	The nutritional loss by processing method	Need for further research
Freshwater small fish	Protein, vitamin A, Ca, Fe and Zn (PUFA)	Very high in vitamin A <i>Parachela siamensis</i> (Cambodia) <i>Rasbora tornieri</i> (Cambodia) <i>Amblypharyngodon mola</i> (Bangladesh)	Consumed frequently in poor households, contributes significantly to daily nutritional needs for micronutrients	Sun-drying destroys nearly all vitamin A No effect on protein and minerals	Document food habits, culture and other factors affecting consumption and dietary intake at individual and household levels Develop more appropriate indicators to measure the effects of fish intake on improved nutritional status Develop and test new indicators to assess vitamin A status Quantify contribution of fish-based diet in increasing response of people living with HIV/AIDS to anti-retroviral therapy
		Very high in Ca <i>Puntius ticto</i> (Bangladesh) <i>Chanda ranga</i> (Bangladesh)	Enhances the content and bioavailability of Fe, Zn and Ca of plant-based diets		
		Very high in Fe <i>Esomus longimanus</i> (Cambodia) <i>Esomus danricus</i> (Bangladesh)	Increases bioavailability of protein from staple food Increases bioefficacy of vitamin A		
		Very high in Zn <i>E. longimanus</i> (Cambodia) <i>R. tornieri</i> (Cambodia)	Complementary food for malnourished children/people living with HIV/AIDS		
Freshwater large fish	Protein, PUFA	A source of PUFA Carps Catfish Tilapia	Contributes to daily nutritional requirements for PUFA	Mostly traded as fresh fish	Systematic evaluation of nutrient content of most commonly consumed fish species in developing countries
Marine fish	Protein, PUFA (Ca; Fe)	Rich in PUFA Anchovy Herring Mackerel Sardine	Complementary food for malnourished children		Document the role of fish PUFA in fetal and infant development

ponds in Bangladesh, it was estimated that production of only 10 kg/pond per year could meet the annual recommended intake for two million children⁽¹⁰⁾. From an ecological perspective, these small indigenous species are self-recruiting; that is, they reproduce and grow in natural water bodies, fish ponds and rice fields without breeding technology or any additional investment, although the productivity is usually low compared with fish aquaculture. Fish supplied by these common pool resources have, however, been reported to be declining in many areas in recent years. This trend may be one of the reasons for the decrease in fish consumption that has been observed, particularly among the rural poor, in some countries such as Bangladesh^(14,24,50), Laos⁽¹²⁾ or Malawi⁽⁵¹⁾. Since it is not clear to what extent increased fish consumption through aquaculture can compensate for these declines (as aquaculture tends to favour the production of larger fish⁽⁵²⁾ with higher value in markets but lower nutritional value), there is a strong need to ensure that the fishing of small fish that takes place in common pool resources (ponds, floodplains, rivers) remains sustainable.

Addressing the underlying factors that determine the fish consumption patterns of the poor

Although fish is relatively cheap and accessible for the poor in many developing countries, recent studies in Asia indicate that low-income households consume less fish compared with rich households^(9,53–55). Studies on socio-economic factors affecting Fe deficiency also show that the low-income group has lower intakes of Fe from fish and meat compared with better-off households, leading to a higher prevalence of Fe deficiency^(56,57). Many poor households do not have enough food stocks all year round, even for their basic needs of staple foods (energy deficits)^(9,58,59). For these populations, fish caught by household members are often used to compensate for their requirement of staple foods to meet their daily energy needs. This responsive behaviour is one of the reasons why many countries in which people rely on fish as their main animal-source food are also low-income food-deficient countries. However, those are also the countries where micronutrient deficiencies are the highest and where interventions promoting fish as a nutrient-dense food are the most needed to maximize the benefits for the poor.

At the intra-household level, women and children, whose micronutrient needs are higher than those of men, unfortunately often consume less fish. For instance, a study in Nigeria found that male heads of households consumed 59% more fish by weight compared with their wives and children and that when a single (large) fish was shared there was a tendency to distribute the body to the man, the tail to his wife and the head to the children on seven to eight out of ten occasions⁽¹³⁾. In Bangladesh, fish are likely to be distributed more evenly among household members compared with other animal-source foods; yet,

adult men and pre-school boys still receive a larger share than their mother and/or sisters⁽⁶⁰⁾. Understanding social, economic and cultural contexts is therefore a necessary primary condition to improve the nutritional benefits accruing from fish for the population most in need.

Synthesis

The different sections above have summarized the information available on the nutrient contents of fish and the effects of processing them, as well as the bioavailability and efficacy of fish in improving the nutritional status of the poor. The main findings related to these points are summarized in Table 3, along with the areas in which requirements for further research on these issues have emerged.

Conclusion

On account of its high nutrient content, fish could be used as a key component in strategies aimed at reducing micronutrient deficiencies in developing countries. A few studies have been published in relation to this issue; however, evidence is still sparse and fragmented and the overall potential contribution of fish to nutritional security is yet to be fully understood. The objective of the present study was to take the first step towards resolving this issue, by conducting a comprehensive overview of the existing literature and documenting more systematically the different nutritional roles that fish could play in preventing and controlling micronutrient deficiencies.

The review confirms that small indigenous species caught and/or traded locally do represent a major source of micronutrients in the everyday diet of the poor in developing countries because of their high nutrient content and bioavailability. Existing data suggest that adding fish to plant-based diets boosts protein absorption from food staples. Small fish, frequently consumed by the poor, are therefore nutritionally important, even in small quantities.

Currently, in most developing countries the production of small fish species is highly dependent on the activities of small-scale fisheries and to a lesser extent on aquaculture. A sustainable supply of these species should be prioritised and the production and consumption of nutrient-dense fish should be promoted in order to make full use of the capacity of these species in reducing micronutrient deficiencies. For this, conservation of wild stocks and dissemination of aquaculture techniques producing these small indigenous species are needed.

Further research is required in areas such as nutrient composition, cleaning/cooking methods, impact of declining fish catches on the fish consumption patterns of the poor and seasonal availability in order to derive appropriate policies and effective food-based approaches.

More technically, research protocols need to be developed to better quantify the outcomes of increased

nutritional intake. In the specific case of fish, the present review revealed that so far very few studies have examined the nutritional outcomes of fish-related interventions. Research methods should also be developed to better monitor and assess aquaculture and small-scale fishery activities and their impact on the nutritional status of households.

Accumulating this type of information would certainly help in increasing our understanding of the mechanisms through which poor households can improve their dietary intake and nutritional status through fisheries and aquaculture activities, although we also recognise that primary causes of undernutrition might be multifaceted and that other factors such as market and income constraints, disease incidence and effects, health service provision and health environment, gender inequity and child care are likely to have equally important effects on nutritional status.

Acknowledgements

The present research was supported by the WorldFish Center. The authors have no conflict of interest to declare. N.K. conducted the initial literature search, whereas both authors contributed to the writing of the paper and therefore have primary responsibility for its final content. Institutional support from Helen Leitch and useful comments by Shakuntala H. Thilsted, Edward H. Allison and Marie Ruel on earlier drafts of the present paper are gratefully acknowledged.

References

- World Health Organization (2001) *The World Health Report 2001: Reducing Risks, Promoting Healthy Life*. Geneva: WHO.
- The Micronutrient Initiative (2009) Investing in the future. A united call to action on vitamin and mineral deficiencies. Global Report 2009. http://www.unitedcalltoaction.org/documents/Investing_in_the_future.pdf (accessed April 2010).
- Food and Agriculture Organization of the United Nations & The WorldFish Center (2009) *Big Number Program. Intermediate Report*. Rome: FAO and WorldFish Center.
- Food and Agriculture Organization of the United Nations (2009) FAOSTAT. Calculated from food balance sheets. <http://faostat.fao.org/site/368/DesktopDefault.aspx?PageID=368#ancor> (accessed April 2010).
- Food and Agriculture Organization of the United Nations (2009) FAO country profiles. Low-income food-deficit countries. <http://www.fao.org/countryprofiles/lifdc.asp> (accessed November 2009).
- Thompson PM, Sultana P, Nuruzzaman M *et al.* (2000) *Fisheries Extension Evaluation Project Final Report*. Dhaka: ICLARM and Department of Fisheries.
- Toft M (2001) The importance of fish and other aquatic animals for food and nutrition security in the Lower Mekong Basin. MSc Thesis, The Royal Veterinary and Agricultural University.
- Chamnan C, Thilsted HS, Roitana B *et al.* (2009) *The Role of Fisheries Resources in Rural Cambodia: Combating Micronutrient Deficiencies in Women and Children*. Phnom Penh: Department of Fisheries Post-harvest Technologies and Quality Control, Fisheries Administration, Ministry of Agriculture, Forestry and Fisheries.
- Islam FU (2007) Self-recruiting species (SRS) in aquaculture: their role in rural livelihoods in two areas of Bangladesh. PhD Thesis, University of Stirling.
- Roos N, Islam MM & Thilsted SH (2003) Small indigenous fish species in Bangladesh: contribution to vitamin A, calcium and iron intakes. *J Nutr* **133**, Suppl. 11, 4021S–4026S.
- Hels O (2002) Strengthening the reliability of data for describing the pathway from food intake to food and nutrition security in rural Bangladesh. PhD Thesis, The Royal Veterinary and Agricultural University.
- Meusch E, Yhoun-Aree J, Friend R *et al.* (2003) *The Role and Nutritional Value of Aquatic Resources in the Livelihoods of Rural People: A Participatory Assessment in Attapeu Province, Lao PDR*. Bangkok: FAO.
- Gomna A & Rana K (2007) Inter-household and intra-household patterns of fish and meat consumption in fishing communities in two states in Nigeria. *Br J Nutr* **97**, 145–152.
- Food and Agriculture Organization of the United Nations (2008) *Present and Future Markets for Fish and Fish Products from Small-Scale Fisheries – Case Studies from Asia, Africa and Latin America*. FAO Fisheries Circular no. 1033. Rome: FAO.
- Geheb K, Kalloch S, Medard M *et al.* (2008) Nile perch and the hungry of Lake Victoria: gender, status and food in an East African fishery. *Food Policy* **33**, 85–98.
- Jansen EG (1997) *Rich Fisheries – Poor Fisherfolk: Some Preliminary Observations about the Effects of Trade and Aid in the Lake Victoria Fisheries: IEA Programme*. Nairobi: IUCN Eastern Africa Regional Office.
- Mujinga W, Lwamba J, Mutala S *et al.* (2009) *An Inventory of Fish Species at the Urban Markets in Lubumbashi, Democratic Republic of Congo. Regional Programme Fisheries and HIV/AIDS in Africa: Investing in Sustainable Solutions. Project Report no. 1983*. Lusaka: The World-Fish Center.
- Ladu BMB, Ovie SI, Erinne EC *et al.* (2000) *Definition and Characterisation of the Southern Nigerian Fish Market. A Second Phase Report of the Project Sustainable Development of African Continental Fisheries: A Regional Study of Policy Options and Policy Formation Mechanisms for the Lake Chad Basin*. New Bussa, Nigeria: NIFFR.
- Overa R (1993) Wives and traders. *Maritime Anthropology Stud* **6**, 110–135.
- Wang C, Harris W, Chung M *et al.* (2006) n-3 Fatty acids from fish or fish-oil supplements, but not α -linolenic acid, benefit cardiovascular disease outcomes in primary- and secondary-prevention studies: a systematic review. *Am J Clin Nutr* **84**, 5–17.
- Koletzko B, Cetin I & Brenna JT (2007) Dietary fat intakes for pregnant and lactating women. *Br J Nutr* **98**, 873–877.
- Thilsted SH, Roos N & Hassan N (1997) The role of small indigenous fish species in food and nutrition security in Bangladesh. *WorldFish Centre Q* **22**, 82–84.
- Roos N (2001) Fish consumption and aquaculture in rural Bangladesh: nutritional contribution and production potential of culturing small indigenous fish species (SIS) in pond polyculture with commonly cultured carps. PhD Thesis, The Royal Veterinary and Agricultural University.
- Roos N, Chamnan C, Loeung D *et al.* (2007) Freshwater fish as a dietary source of vitamin A in Cambodia. *Food Chem* **103**, 1104–1111.
- Roos N, Thorseng H, Chamnan C *et al.* (2007) Iron content in common Cambodian fish species: perspectives for dietary iron intake in poor, rural households. *Food Chem* **104**, 1226–1235.
- Karapangiotidis LT, Yakupitiyage A & Little DC (2010) The nutritional value of lipids in various tropical aquatic animals from rice-fish farming systems in northeast Thailand. *J Food Compos Anal* **23**, 1–8.

27. US Department of Agriculture (2005) National nutrient data base. http://fnic.nal.usda.gov/nal_display/index.php?info_center=4&tax_level=2&tax_subject=279&topic_id=1387 (accessed October 2009).
28. World Health Organization (1985) *Energy and Protein Requirements*. Geneva: WHO.
29. Tacon AGJ & Metian M (2009) Fishing for feed or fishing for food: increasing global competition for small pelagic forage fish. *Ambio* **38**, 294–302.
30. Ruan C, Liu X, Man H *et al.* (1995) Milk composition in women from five different regions of China: the great diversity of milk fatty acids. *J Nutr* **125**, 2993–2998.
31. Muskiet FA, van Goor SA, Kuipers RS *et al.* (2006) Long-chain polyunsaturated fatty acids in maternal and infant nutrition. *Prostaglandins Leukot Essent Fatty Acids* **75**, 135–144.
32. Roos N, Wahab MA, Chamnan C *et al.* (2007) The role of fish in food-based strategies to combat vitamin A and mineral deficiencies in developing countries. *J Nutr* **137**, 1106–1109.
33. Dewailly E, Chateau-Degat L & Suhas E (2008) Fish consumption and health in French Polynesia. *Asia Pac J Clin Nutr* **17**, 86–93.
34. Roos N, Wahab MA, Hossain MAR *et al.* (2007) Linking human nutrition and fisheries: incorporating micronutrient-dense, small indigenous fish species in carp polyculture production in Bangladesh. *Food Nutr Bull* **28**, 281–293.
35. Michaelsen KF, Hoppe C, Roos N *et al.* (2009) Choice of foods and ingredients for moderately malnourished children 6 months to 5 years of age. *Food Nutr Bull* **30**, Suppl. 3, S343–S404.
36. Hirao S, Yamada J & Kikuchi R (1959) Vitamin A in fish flesh. *Bull Tokai Reg Fish Res Lab* **25**, 49–65.
37. Kongsbak K, Thilsted SH & Wahed MA (2008) Effect of consumption of the nutrient-dense, freshwater small fish *Amblypharyngodon mola* on biochemical indicators of vitamin A status in Bangladeshi children: a randomised, controlled study of efficacy. *Br J Nutr* **99**, 581–597.
38. International Zinc Nutrition Consultative Group (2004) Assessment of the risk of zinc deficiency in populations and options for its control. *Food Nutr Bull* **25**, Suppl. 1, S99–S123.
39. World Health Organization & Food and Agriculture Organization of the United Nations (2004) *Vitamin and Mineral Requirements in Human Nutrition*, 2nd ed. Geneva: WHO/FAO.
40. Hansen M, Thilsted SH, Sandstrom B *et al.* (1998) Calcium absorption from small soft-boned fish. *J Trace Elem Med Biol* **12**, 148–154.
41. Larsen T, Thilsted SH, Kongsbak K *et al.* (2000) Whole small fish as a rich calcium source. *Br J Nutr* **83**, 191–196.
42. Gibson R, Yeudall F, Drost N *et al.* (2003) Experiences of a community-based dietary intervention to enhance micronutrient adequacy of diets low in animal source foods and high in phytate: a case study in rural Malawian children. *J Nutr* **133**, Suppl. 11, 3992S–3999S.
43. Chittchang U, Jittinandana S, Sungpuag P *et al.* (1999) Recommending vitamin A-rich foods in southern Thailand. *Food Nutr Bull* **20**, 238–242.
44. Olson JA (1996) Biochemistry of vitamin A and carotenoids. In *Vitamin A Deficiency: Health, Survival, and Vision*, pp. 221–250 [A Sommer and KP West Jr, editors]. New York: Oxford University Press.
45. Ruel MT (2001) *Can Food-Based Strategies Help Reduce Vitamin A and Iron Deficiencies?: A Review of Recent Evidence*. Washington, DC: IFPRI.
46. Caulfield LE, Huffman SL & Piwoz EG (1999) Interventions to improve intake of complementary foods by infants 6 to 12 months of age in developing countries: impact on growth and on the prevalence of malnutrition and potential contribution to child survival. *Food Nutr Bull* **20**, 183–200.
47. Lartey A, Manu A, Brown KH *et al.* (1999) A randomized, community-based trial of the effects of improved, centrally processed complementary foods on growth and micronutrient status of Ghanaian infants from 6 to 12 mo of age. *Am J Clin Nutr* **70**, 391–404.
48. Greco L, Balungi J, Amono K *et al.* (2006) Effect of a low-cost food on the recovery and death rate of malnourished children. *J Pediatr Gastroenterol Nutr* **43**, 512–517.
49. Lin C, Manary M, Maketa K *et al.* (2008) An energy-dense complementary food is associated with a modest increase in weight gain when compared with a fortified porridge in Malawian children aged 6–18 months. *J Nutr* **138**, 593–598.
50. Thompson PM, Roos N, Sultana P *et al.* (2002) Changing significance of inland fisheries for livelihoods and nutrition in Bangladesh. In *Food Systems for Improved Human Nutrition: Linking Agriculture, Nutrition and Productivity*, pp. 249–317 [PK Katakai and SC Babu, editors]. Binghamton, NY: Haworth Press.
51. Dey MM, Kambewa M, Prein D *et al.* (2006) Impact of development and dissemination of integrated aquaculture-agriculture (IAA) technologies in Malawi. *NAGA WorldFish Centre Q* **29**, 28–35.
52. Kawarazuka N & Béné C (2010) Linking small-scale fisheries and aquaculture to household nutritional security: an overview. *Food Security* **2**, 343–357.
53. Dey MM, Rab MA, Paraguas FJ *et al.* (2005) Fish consumption and food security: a disaggregated analysis by types of fish and classes of consumers in selected Asian countries. *Aquac Econ Manag* **9**, 89–111.
54. Hossain M, Naher F & Shahabuddin Q (2005) Food security and nutrition in Bangladesh: progress and determinants. *J Agric Dev Econ* **2**, 103–132.
55. Bose ML & Dey MM (2007) Food and nutritional security in Bangladesh: going beyond carbohydrate counts. *Agric Econ Res Rev* **20**, 203–225.
56. Bhargava A, Bouis HE & Scrimshaw NS (2001) Dietary intakes and socioeconomic factors are associated with the hemoglobin concentration of Bangladeshi women. *J Nutr* **131**, 758–764.
57. Keskin Y, Moschonis G, Dimitriou M *et al.* (2005) Prevalence of iron deficiency among schoolchildren of different socio-economic status in urban Turkey. *Eur J Clin Nutr* **59**, 64–71.
58. Béné C, Neiland A, Jolley T *et al.* (2003) Inland fisheries, poverty and rural livelihoods in the Lake Chad basin. *J Asian African Stud* **38**, 17–51.
59. Karim M (2006) The livelihood impacts of fishponds integrated within farming systems in Mymensingh district, Bangladesh. PhD Thesis, University of Stirling.
60. Bouis H & Novenario-Reese MJG (1997) *The Determinants of Demand for Micronutrients: An Analysis of Rural Households in Bangladesh*. FCND Discussion Paper no. 32. Washington, DC: IFPRI.
61. Food and Nutrition Board, Institute of Medicine (2000) Vitamin A. In *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc*, pp. 162–196 [Medical Research Council, editor]. Washington, DC: FNB, IOM.
62. Thilsted SH & Roos N (1999) Policy issues on fisheries in relation to food and nutrition security. In *Fisheries Policy Research in Developing Countries: Issues, Priorities and Needs*, pp. 61–69 [M Ahmend, C Delgado, S Sverdrup-Jensen *et al.*, editors]. Makati City, Philippines: ICLARM.