
INSTRUCTION TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology,
London School of Hygiene and Tropical Medicine,
Keppel Street, London, WC1E 7HT,
England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief introduction (unheaded); Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration may be printed is 13.5×18 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale should normally be given on the figure. Photocopies of illustrations should also be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 13.5×18 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In *Ecology and Physiology of parasites* (ed A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. and WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49–59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
Evidence for increased susceptibility of <i>Brugia pahangi</i> -infected jirds (<i>Meriones unguiculatus</i>) to subsequent homologous infections. T. R. KLEI, J. W. McCALL and J. B. MALONE	161 - 166
Infectivity in rodents of <i>Schistosoma mansoni</i> cercariae of human and murine origin. C. COMBES and D. IMBERT-ESTABLET	167 - 171
The effect of gamma irradiation on <i>Nematospiroides dubius</i> . Factors affecting the survival of worms in a primary infection in mice. J. M. BEHNKE, H. A. PARISH and P. HAGAN	173 - 182
The distribution of microfilariae in the skin of Guatemalan onchocerciasis patients: an evaluation of diagnostic potentials. M. KAWABATA. <i>et al.</i>	183 - 190
<i>Opisthorchis viverrini</i> : Partial success in adoptively transferring immunity with spleen cells and serum in the hamster. D. J. FLAVELL, K. PATTANAPANYASAT and S. U. FLAVELL	191 - 197
The anthelmintic effects of albendazole on <i>Brugia pahangi</i> . Research Note. D. A. DENHAM, D. A. LIRON and E. BRANDT	199 - 200
Susceptibility of different species of <i>Lymnaea</i> snails to miracidia of <i>Fasciola gigantica</i> and <i>F. hepatica</i> in Iran. Research Note. J. MASSOUD and S. SADJADI	201 - 202
The interaction of some environmental factors influencing <i>Schistosoma mansoni</i> cercarial host-finding. Research Note. N. Ø. CHRISTENSEN, F. FRANSEN and P. NANSEN	203 - 205
<i>Castroia kamariae</i> sp. nov. and <i>Limatulum kuziai</i> sp. nov. (Lecithodendriidae) from the free-tailed bat (<i>Tadarida mops</i> de Blainville) from West Malaysia. J. R. PALMIERI, M. KRISHNASAMY and J. T. SULLIVAN	207 - 213
Presence of cholesterol in the neutral lipids of three sheep cestodes. S. C. NIGAM and G. PREMVATI	215 - 218
Histochemical studies on <i>Raillietina (Raillietina) johri</i> (Cestoda: Davaineidae). III. Esterases. T. K. ROY	219 - 222
The migration of <i>Strongyloides papillosus</i> in rabbits following infection by the oral and subcutaneous routes. O. C. NWAORGU and R. M. CONNAN	223 - 232