

in understanding and addressing the needs of women with OF: clinical care, academia, international health, civil society, and government. Twenty-one individuals were interviewed about their perceptions of IDs for OF self-management and their implementation. Interviews were audio-recorded and transcribed. The Consolidated Framework for Implementation Research (CFIR) guided data collection and analysis. Thematic analyses were carried out within Nvivo v.12. RESULTS/ANTICIPATED RESULTS: Determinants of implementation of an ID for OF self-management (by CFIR domain) include: (1) intervention characteristics—*relative advantage* and *cost*; (2) individual characteristics—*knowledge and beliefs about the innovation*; (3) inner setting—*organizational culture, implementation climate, tension for change, and compatibility*; (4) outer setting—*patient needs and resources* and *external policy and incentives*; (5) process—*opinion leaders* and *collaboration*. Facilitators include: tension for change for low-cost, accessible IDs; relative advantage over existing tools; development of partnerships; and identification of implementation champions. Barriers include: need for educational strategies to encourage clinical provider acceptability; lack of evidence of the optimal beneficiary. DISCUSSION/SIGNIFICANCE OF IMPACT: Tools for therapeutic OF self-management could be integrated into comprehensive OF programming. Employing the CFIR as an overarching typology allows for comparison across contexts and settings where OF care occurs and may be useful for clinicians, researchers, and policy-makers interested in implementing IDs for OF self-management in LMICs. CONFLICT OF INTEREST DESCRIPTION: I am working with colleagues at the non-profit Restore Health on developing an insertable cup for therapeutic self-management of obstetric fistula in LMICs

4432

Transportation Barriers and Preferences Among Drivers with Developmental Disabilities in Southeast

Austin Svancara¹, Rajesh Kana², Benjamin McManus¹, Haley Bednarz¹, Gabriela Sherrod¹, and Despina Stavrinou¹

¹University of Alabama at Birmingham; ²University of Alabama

OBJECTIVES/GOALS: Transportation may be a barrier for individuals with Autism Spectrum Disorder (ASD). More individuals with ASD utilize public transportation compared to typically developing (TD) individuals. This study seeks to elucidate the transportation needs of individuals with ASD in the Southeast. METHODS/STUDY POPULATION: Sixty-one licensed drivers with a diagnosis of ASD (n = 21), Attention-Deficit/Hyperactivity Disorder (ADHD; n = 19), or no diagnosis (TD; n = 21) were recruited and were matched across diagnosis groups by age (16-30 years old), gender, and IQ. Participants completed an adapted version of the Barriers to Care Scale and a survey assessing transportation preferences and quality of life. Means and frequencies were obtained. Chi-square analyses were conducted to estimate associations between diagnosis and transportation preferences. RESULTS/ANTICIPATED RESULTS: Nearly all of the sample had access to a car (98.4%). Yet, only 71.4% of drivers ASD preferred to use their own car compared to 89.5% and 90.5% of the ADHD and TD groups respectively. The use of public transportation (6.6%) and ride-hailing services (18%) for general transportation needs was very low across the

groups. There was a significant association between group type and the reliance on others for transportation ($\chi^2(2,61) = 9.9, p < .01$). Only 21.1% of those with ADHD relying on others for transportation needs, compared to 61.9% of TD and 66.7% of individuals with ASD. 23.8% of ASD drivers, 10.5% of ADHD drivers, and 9.5% of TD drivers believe transportation proved as an obstacle. DISCUSSION/SIGNIFICANCE OF IMPACT: The proportion of ASD drivers who believed transportation to be a barrier appeared slightly higher than other groups. Public transportation use may be low due to lower accessibility to such services in the Southeast. The travel patterns of individuals with ASD and ADHD merits further exploration.

4345

Two-step Algorithm for Clostridioides difficile is Inadequate for Differentiating Infection from Colonization in Children

Maribeth R Nicholson¹, Jacob M Parnell, Irtiqqa Fazili, Sarah C. Bloch, D. Borden Lacy, Eric Skaar, and Kathryn M Edwards

¹Vanderbilt University Medical Center

OBJECTIVES/GOALS: In 2017, new guidelines recommended multi-step algorithms for CDI diagnosis, and clinical centers rapidly implemented changes despite limited pediatric data. We assessed a multi-step algorithm using NAAT followed by EIA for ability to differentiate symptomatic CDI from colonization in children. METHODS/STUDY POPULATION: We prospectively enrolled pediatric patients with cancer, cystic fibrosis, or inflammatory bowel disease who were not being tested or treated for CDI and obtained a stool sample for NAAT. If positive by NAAT (colonized), EIA was performed. Children with symptomatic CDI who tested positive by NAAT via the clinical laboratory were also enrolled and EIA performed on residual stool. A functional cell cytotoxicity neutralization assay (CCNA) was performed in addition. RESULTS/ANTICIPATED RESULTS: Of the 138 asymptomatic children enrolled, 24 (17%) were colonized. An additional 37 children with symptomatic CDI were enrolled. Neither EIA positivity (41% versus 21%, $P = 0.11$) or CCNA positivity (49% versus 46%, $P = 0.84$) were significantly different between symptomatic versus colonized children. When both EIA and CCNA were positive, children were more commonly symptomatic than colonized (33% versus 13%, $P = 0.04$). DISCUSSION/SIGNIFICANCE OF IMPACT: A multi-step testing algorithm with NAAT and EIA failed to differentiate symptomatic CDI from colonization in our pediatric cohort. As multi-step algorithms are moved into clinical care, pediatric providers will need to be aware of the continued limitations in diagnostic testing.

4151

Understanding barriers and solutions towards access to mental health among rural adolescents

Brandy Davis¹, Kimberly B. Garza², Salisa Westrick², Edward Chou², and Cherry Jackson²

¹University of Alabama at Birmingham; ²Auburn University

OBJECTIVES/GOALS: There are two objectives: 1) To identify healthcare providers' (HCP) barriers and potential solutions towards