

Research Paper

Cite this article: Blasco-Costa I, Seppälä K, Feijen F, Zajac N, Klappert K, Jokela J (2019). A new species of *Atriophallophorus* Deblock & Rosé, 1964 (Trematoda: Microphallidae) described from *in vitro*-grown adults and metacercariae from *Potamopyrgus antipodarum* (Gray, 1843) (Mollusca: Tateidae). *Journal of Helminthology* **94**, 1–15. <https://doi.org/10.1017/S0022149X19000993>

Received: 28 August 2019

Revised: 19 October 2019

Accepted: 22 October 2019


Key words:

Microphallus; *in vitro* culture; phylogeny; rRNA genes; mitochondrial genes; life cycle; fresh water; mud snail; waterfowl; New Zealand

Author for correspondence:

I. Blasco-Costa, E-mail: isa.blasco.costa@gmail.com

A new species of *Atriophallophorus* Deblock & Rosé, 1964 (Trematoda: Microphallidae) described from *in vitro*-grown adults and metacercariae from *Potamopyrgus antipodarum* (Gray, 1843) (Mollusca: Tateidae)

I. Blasco-Costa^{1,2} , K. Seppälä^{3,4,5}, F. Feijen^{3,4}, N. Zajac^{3,4}, K. Klappert^{3,4} and J. Jokela^{3,4}

¹Natural History Museum of Geneva, PO Box 6434, CH-1211 Geneva 6, Switzerland; ²Department of Arctic and Marine Biology, UiT The Arctic University of Norway, Langnes, PO Box 6050, 9037 Tromsø, Norway; ³Department of Aquatic Ecology, Swiss Federal Institute of Aquatic Science and Technology (EAWAG), Dübendorf, Switzerland; ⁴Institute of Integrative Biology, ETH-Zürich, Zürich, Switzerland and ⁵Research Department for Limnology, University of Innsbruck, 5310 Mondsee, Austria

Abstract

The adult and metacercaria life stages of a new species of the microphallid genus *Atriophallophorus* Deblock & Rosé, 1964 are described from specimens collected at Lake Alexandrina (South Island, New Zealand). In addition to molecular analyses of ribosomal and mitochondrial genes, metacercariae of *Atriophallophorus winterbourni* n. sp. from the snail host *Potamopyrgus antipodarum* (Gray) were grown *in vitro* to characterize internal and external morphology of adults using light and scanning electron microscopy and histological techniques. *Atriophallophorus winterbourni* n. sp. is readily distinguishable from *Atriophallophorus coxiellae* Smith, 1973 by having a different structure of the prostatic chamber, sub-circular and dorsal to genital atrium, rather than cylindrical, fibrous, elongate and placed between the seminal vesicle and the genital atrium. The new species is most similar to *Atriophallophorus minutus* (Price, 1934) with regards to the prostatic chamber and the morphometric data, but possesses elongate-oval testes and subtriangular ovary rather than oval and transversely oval in *A. minutus*. Phylogenetic analyses including sequence data for *A. winterbourni* n. sp. suggested a congeneric relationship of the new species to a hitherto undescribed metacercariae reported from Australia, both forming a strongly supported clade closely related to *Microphallus* and *Levinseniella*. In addition, we provide an amended diagnosis of *Atriophallophorus* to accommodate the new species and confirm the sinistral interruption of the outer rim of the ventral sucker caused by the protrusion of the dextral parietal atrial scale at the base of the phallus.

Introduction

A microphallid species reported as *Microphallus* sp. from New Zealand has been largely used as a model species in parasitology and evolutionary studies for more than three decades (e.g. Lively, 1987; Dybdahl & Lively, 1998; Lively & Dybdahl, 2000; Jokela *et al.*, 2009; Gibson *et al.*, 2016a). Broad knowledge has accumulated on the role of this parasite in coevolutionary dynamics with the host populations. Particularly, in this case the coevolutionary dynamics are suggested to favour sexual reproduction by the host (i.e. ‘parasite hypothesis’ or ‘Red Queen hypothesis’ for maintenance of sex; Jaenike (1978); Hamilton (1980); Lively (2016)). However, basic knowledge such as the identity of the species, its genetic diversity and phylogenetic affinities are still lacking, with the consequences that it may be difficult to judge the source of heterogeneity in infection experiments and field surveys.

Microphallids are a diverse and cosmopolitan group of very small worms, typically found in the intestine of birds. Second intermediate hosts tend to be crustaceans, but several species present abbreviated life cycles where the metacercariae encyst within the mollusc first intermediate host. Their metacercariae are usually identical to the adult stage in the definitive host. In the definitive host, microphallids tend to mature in a few days and live only for several weeks (Galaktionov & Dobrovolskij, 2003). This feature of microphallids makes them amenable to *in vitro* culture as a method to obtain the mature adults and be able to characterize and describe new species. Such an alternative is, nowadays, desirable, considering the limitations of obtaining permits to collect potential vertebrate definitive hosts, which are in many cases protected by national or international regulations (Blasco-Costa & Poulin, 2017).

The aim of this study is to morphologically characterize and describe the adult and metacercaria life stages of this microphallid, which represents a new species of the genus *Atriophallophorus* Deblock & Rosé, 1964 instead of *Microphallus* Ward, 1901 as had been previously reported. Metacercariae of microphallid specimens from the snail host *Potamopyrgus antipodarum* (Gray) were grown *in vitro* and the adults were analysed with light microscopy, scanning electron microscopy (SEM) and histological techniques to characterize their internal and external morphology. In addition, this study provides an amended diagnosis of *Atriophallophorus*, molecular data for two nuclear and two mitochondrial markers of the new species and an evaluation of its relationship with other microphallids.

Material and methods

Specimens

Potamopyrgus antipodarum snails were collected using a kick net and snorkelling from two sites in Lake Alexandrina (South Island, New Zealand), site 'Swamp' (−43.962102, 170.441728) and site 'JMS' (−43.937199, 170.459495). Live metacercariae were dissected from four infected snails from each locality and allowed to excyst in Tyrode's salt solution (Sigma, Buchs, Switzerland) with pancreatin (3 mg/ml; Sigma) and penicillin–streptomycin–neomycin solution (8% v/v; Sigma) at 39°C. Excysted juvenile worms were washed twice with Tyrode's salt solution supplemented with penicillin–streptomycin–neomycin solution (8% v/v). Then, juveniles from the same locality were pooled and transferred to culture medium of RPMI 1640 (Gibco, ThermoFisher Scientific, Waltham (MA) U.S.) supplemented with horse serum (20% v/v, Gibco), penicillin–streptomycin–neomycin solution (8% v/v), HEPES buffer (25 mM; Gibco) and Amphotericin B solution (0.25 mg/ml; Sigma). Juveniles were incubated at 39°C and cultured for up to three days (72 h). Fresh culture media was changed daily. Metacercariae right after hatching and *in vitro*-grown adults after 24 h, 48 h and 72 h of culturing were fixed in hot saline and preserved in 75% ethanol for later morphological examination. A subsample of specimens was preserved in hot formalin for examination using SEM and histological analysis. Additionally, five metacercariae from each site (from a pool of four snails each) were preserved in 100% ethanol for molecular analyses.

Morphological data

Metacercariae and adult specimens grown *in vitro* were stained using iron acetocarmine, dehydrated through a graded ethanol series, cleared in dimethyl phtalate and examined as permanent mounts in Canada balsam. Figures were made using a drawing tube mounted on a Zeiss light compound microscope at ×1600 magnification. Measurements of the specimens were taken from drawings at ×640 magnification. Five specimens were dehydrated in a graded ethanol series, critical point-dried and sputter-coated with gold for SEM examination using a Zeiss DSM 940A (Zeiss AG, Oberkochen, Germany) at an accelerating voltage of 5 kV. Three specimens were used for histology after dehydration through a graded ethanol series, followed by propylene oxide and immersed in Epon resin. Histological sections of 1 µm were made with an ultramicrotome Reichert-Jung Ultracut E (Leica Microsystems, Heerbrugg, Switzerland), stained with toluidine blue 0.75% and examined using the light microscope mentioned previously. All measurements in the text are in micrometres unless

otherwise stated, and are given as the range followed by the mean ± standard deviation. Permanent mounts of the type material and SEM preparations are deposited in the Platyhelminthes collection of the Natural History Museum of Geneva, the Institute of Parasitology of the Academy of Sciences of the Czech Republic and the Otago Museum, Dunedin, New Zealand.

Molecular data

Four metacercariae isolates from Swamp and five from JMS originating from a pool of four infected *P. antipodarum* from each site were characterized molecularly. Due to the small size of these specimens, it was impossible to keep holonophores of the sequenced specimens. Nonetheless, the specimens used for morphological description herein represent paragenophores and likely genetic clones of the sequenced material. Genomic DNA was extracted from ethanol-fixed metacercariae isolates in 200 µl of a 5% suspension of Chelex® in deionized water and containing 0.1 mg/ml proteinase K followed by incubation at 56°C for 5 h, boiling at 90°C for 8 min and centrifugation at 14,000 g for 10 min. Partial fragments were amplified of the large ribosomal subunit (28S) (1800 bp; primers U178F: 5'-GCA CCC GCT GAA YTT AAG-3' and L1642R: 5'-CCA GCG CCA TCC ATT TTC A-3'; Lockyer *et al.*, 2003) and internal transcribed spacer 2 (ITS2) (500 bp; primers 3S: 5'-GTA CCG GTG GAT CAC GTG GCT AGT G-3' and ITS2:2: 5'-CCT GGT TAG TTT CTT TTC CTC CGC-3' (Morgan & Blair, 1995; Cribb *et al.*, 1998)). Additionally, two fragments of mitochondrial genes (mt) were amplified: cytochrome oxidase subunit I (*cox I*) (1050 bp; primers JB3: 5'-TTTTTTGGGCATCCTGAGGTTTAT -3' and microph_rev: 5'-AAT CAT GAT GCA AAA GG-3' (Bowles *et al.*, 1993; newly designed)) and nicotinamide adenine dinucleotide dehydrogenase subunit 5 (NADH5) (744 bp; primers F2_micND5: 5'-CTTCAACCTTGCTGCTGCC-3' and R2_micND5: 5'-TCCC AACGAAACCTAAAACCTGC-3' (newly designed)).

Polymerase chain reaction (PCR) amplifications were performed in 20 µl reactions containing 2 µl of extraction supernatant (~10–20 ng of template DNA), 2× MyFi™ Mix (Bioline France, France; containing DNA polymerase, dNTPs, MgCl₂ and enhancers at optimal concentrations) and 0.4 µM of each primer combination. Thermocycling conditions used for amplification of the rDNA regions followed Galaktionov *et al.* (2012). The following thermocycling profile was used for amplification of the mt *cox I* and NADH5 fragments: denaturation (95°C for 3 min); 38 cycles of amplification (94°C for 50 s, 52°C for 30 s and 72°C for 1 min); and 4 min extension step at 72°C. PCR amplicons were purified prior to sequencing using exonuclease I and shrimp alkaline phosphatase enzymes (Werle *et al.*, 1994). Amplicons were cycle-sequenced from both strands using PCR primers and an internal primer for the 28S fragment (L1200R: 5'-GCA TAG TTC ACC ATC TTT CGG-3'; Littlewood *et al.*, 2000) at the commercial facility Macrogen (Amsterdam, The Netherlands). Contiguous sequences were assembled and edited using Geneious® (v. 8.1 Biomatters Ltd., Auckland, New Zealand) and submitted to GenBank (see accession numbers in table 1).

Molecular analyses

Newly generated sequences for the 28S rDNA and the ITS2 fragments were aligned in two independent datasets, together with the published sequences of other microphallids from GenBank (see accession numbers in table 1). The sequences were aligned

Table 1. List of taxa included in the phylogenetic analyses, GenBank accession numbers and references.

Taxa	28S rDNA	ITS2	References
Haematoloechidae			
<i>Haematoloechus longiplexus</i>	AF387801		Snyder & Tkach (2001)
Lecithodendriidae			
<i>Paralecithodendrium parvouterus</i>	AY220617		Tkach <i>et al.</i> (2003)
<i>Pycnopus heteroporus</i>	AF151918		Tkach <i>et al.</i> (2000)
Microphallidae			
<i>Atriophallophorus winterbourni</i> n. sp.	MN342153– MN342154	MN342155	This study
<i>Levinsiella</i> sp.	MG783585	MG783580	Galaktionov & Blasco-Costa (2018)
<i>Longiductotrema tethepae</i>	KX712084	KX712086	Kudlai <i>et al.</i> (2016)
<i>Maritrema arenaria</i>	AY220629	HM584171	Galaktionov <i>et al.</i> (2012)
<i>Maritrema brevisacciferum</i>	KT355818	KT355824	Kudlai <i>et al.</i> (2015)
<i>Maritrema corai</i>	KT880222		Hernandez-Orts <i>et al.</i> (2016)
<i>Maritrema deblocki</i>	KJ144173		Presswell <i>et al.</i> (2014)
<i>Maritrema eroliae</i>	JF826247	HQ650132	Al-Kandari <i>et al.</i> (2011)
<i>Maritrema heardi</i>	AY220632		Tkach <i>et al.</i> (2003)
<i>Maritrema madrynense</i>		KF575167	Diaz & Cremona (2010)
<i>Maritrema neomi</i>	AF151927		Tkach <i>et al.</i> (2000)
<i>Maritrema novaezealandense</i>	KJ144178	KJ540203	Presswell <i>et al.</i> (2014)
<i>Maritrema oocysta</i>	AY220630	HM584170	Tkach <i>et al.</i> (2003); Galaktionov <i>et al.</i> (2012)
<i>Maritrema poulini</i>	KJ144175		Presswell <i>et al.</i> (2014)
<i>Maritrema prosthometra</i>	AY220631		Tkach <i>et al.</i> (2003)
<i>Maritrema subdolum</i>	AF151926	HM584172	Tkach <i>et al.</i> (2000); Galaktionov <i>et al.</i> (2012)
<i>Maritrema</i> sp. 1		KC012521	Gilardoni <i>et al.</i> (2011)
<i>Maritrema</i> sp. 2		KC222022	Gilardoni <i>et al.</i> (2011)
Microphallidae gen. sp.	KT355820	KT355826	Kudlai <i>et al.</i> (2015)
Microphallidae gen. sp.	AB974360		Kakui (2014)
<i>Microphallus</i> (?) fusiformis	AY220633		Tkach <i>et al.</i> (2003)
<i>Microphallus abortivus</i>	AY220626	HM584173	Tkach <i>et al.</i> (2003); Galaktionov <i>et al.</i> (2012)
<i>Microphallus basodactylophallus</i>	AY220628		Tkach <i>et al.</i> (2003)
<i>Microphallus calidris</i>	HM584125	HM584183	Galaktionov <i>et al.</i> (2012)
<i>Microphallus kurilensis</i>	HM584140	HM584185	Galaktionov <i>et al.</i> (2012)
<i>Microphallus minutus</i>	KT355822	KT355828	Kudlai <i>et al.</i> (2015)
<i>Microphallus ochotensis</i>	HM584142	HM584175	Galaktionov <i>et al.</i> (2012)
<i>Microphallus piriformes</i>	HM584122	HM584181	Galaktionov <i>et al.</i> (2012)
<i>Microphallus primas</i>	AY220627		Tkach <i>et al.</i> (2003)
<i>Microphallus pseudopygmaeus</i>	HM584126	HM584198	Galaktionov <i>et al.</i> (2012)
<i>Microphallus pygmaeus</i>	HM584133	HM584190	Galaktionov <i>et al.</i> (2012)
<i>Microphallus similis</i>	HM584138	HM584178	Galaktionov <i>et al.</i> (2012)
<i>Microphallus triangulatus</i>	HM584139	HM584195	Galaktionov <i>et al.</i> (2012)
<i>Microphallus</i> sp.	KJ868216		O'Dwyer <i>et al.</i> (2014)
<i>Probolocoryphe uca</i>		GQ377842	Al-Kandari & Al-Bustan (2010)
Plagiorchiidae			
<i>Plagiorchis vespertilionis</i>	AF151931		Tkach <i>et al.</i> (2000)

(Continued)

Table 1. (Continued.)

Taxa	28S rDNA	ITS2	References
Pleurogenidae			
<i>Parabascus duboisi</i>	AY220618		Tkach et al. (2003)
<i>Pleurogenes claviger</i>	AF151925		Tkach et al. (2000)
Prosthogonimidae			
<i>Prosthogonimus ovatus</i>	AF151928		Tkach et al. (2000)
Telorchidae			
<i>Telorchis assula</i>	AF151915		Tkach et al. (2000)

using default parameters of MAFFT implemented in Geneious®, and the extremes of the alignment were trimmed to match the shortest sequences. The 28S dataset (1280 bp long) included 14 representative sequences of *Microphallus* spp., 12 of *Maritrema* spp., one each of *Levinseniella*, *Longiductotrema* and an unidentified microphallid of Kudlai et al. (2015) and a sequence labelled as *Microphallus fusiformis* (which should be disregarded as a species of *Microphallus*; see Kudlai et al., 2015) retrieved from GenBank (table 1). Additionally, five sequences of species belonging to sister families of the Microphallidae – i.e. Lecithodendriidae, Pleurogenidae and Prosthogonimidae in the Microphalloidea – and three sequences of species in the Plagiorchioidea were retrieved from GenBank and included as outgroups. The ITS2 dataset (401 bp long) included ten representative sequences of *Microphallus* spp.; nine sequences of *Maritrema*; one each of *Levinseniella*, *Probolocoryphe* and *Longiductotrema*; and one of an unidentified microphallid of Kudlai et al. (2015). The phylogenetic analyses were run on the two datasets individually under the maximum likelihood (ML) and Bayesian inference (BI) criteria, employing the nucleotide substitution model GTR+ Γ . ML analyses were conducted using the program RAXML v. 8.2 (Stamatakis, 2014). All model parameters, bootstrap nodal support values (1000 repetitions) and an extended majority-rule consensus topology were estimated using RAXML. BI trees were constructed using MrBayes v. 3.2 (Ronquist et al., 2012), running two independent Markov Chain Monte Carlo runs of four chains with standard settings for 10^7 generations and sampling tree topologies every 10^3 generation. Burn-in periods were set automatically to 25% generations, ensuring the remaining trees were obtained after values for standard deviation of split frequencies were <0.01 . A majority-rule consensus topology and nodal support estimated as posterior probability values (Huelsenbeck et al., 2001) were calculated from the remaining trees. All MrBayes and RAXML analyses were performed on the computational resource CIPRES (Miller et al., 2010). Genetic divergences amongst taxa were calculated as uncorrected *p*-distances for each gene region using MEGA v. X (Kumar et al., 2018).

Results

Family: Microphallidae Ward, 1901

Atriophallophorus Deblock & Rosé, 1964

Atriophallophorus winterbourni n. sp.

Taxonomic summary

Synonyms. Metacercaria A of Winterbourn (1974); *Microphallus* sp. of Lively (1987, 1989), Lively & McKenzie (1991), Jokela &

Lively (1995), Dybdahl & Lively (1996, 1998), Levri & Lively (1996), Lively & Jokela (1996), Levri (1999), Krist et al. (2000, 2004), Levri & Fisher (2000), Jokela et al. (2003, 2009), Dybdahl & Krist (2004), Lively et al. (2004, 2008), Osnas & Lively (2004, 2006, 2011), Levri et al. (2005), Fromme & Dybdahl (2006), Koskella & Lively (2007, 2009), Lagrue et al. (2007), Dybdahl et al. (2008), Lagrue & Poulin (2008), Lively et al. (2008), King et al. (2009, 2011a, b), Koskella et al. (2011), Vergara et al. (2013, 2014, 2017), Paczesniak et al. (2014, 2019), Gibson et al. (2016a, b, 2018), McKone et al. (2016), Bankers et al. (2017), Bankers & Neiman (2017); *Microphallus* sp. 'lively' of Hofmann et al. (2016); *Microphallus* sp. 'lively' of Hechinger (2012), Soper et al. (2014), Bankers & Neiman (2017); *Microphallus livelyi* (*nomen nudum*) of Bankers & Neiman (2017).

Type host. *Potamopyrgus antipodarum* (Gray) (Mollusca: Tateidae; first and second intermediate host).

Definitive host. Waterfowl (see Osnas & Lively, 2011).

Type locality. Lake Alexandrina, South Island, New Zealand (site 'Swamp': $-43.962102, 170.441728$).

Other localities. Lake Alexandrina, South Island, New Zealand (site 'JMS': $-43.937199, 170.459495$).

Site in host. Intermediate host: metacercaria encysted in gonads.

Type material. Holotype (MHNG-PLAT-129859), 22 paratypes (MHNG-PLAT-129860–MHNG-PLAT-129865) and seven metacercariae vouchers (MHNG-PLAT-129866–MHNG-PLAT-129867) deposited in the Platyhelminthes collection of the Natural History Museum of Geneva (NHMG), two paratypes (IPCAS D-803) deposited in the helminthological collection of the Institute of Parasitology, Česke Budějovice, Czech Republic (IPCAS) and four paratypes (OMNZ IV110293–OMNZ IV110295) and two voucher metacercariae (OMNZ IV110296) deposited at the Otago Museum (OM), Dunedin, New Zealand.

Representative DNA sequences. 28S rDNA, four sequences of specimens from Swamp and five of specimens from JMS (GenBank MN342153–MN342154); ITS2 rDNA, eight identical sequences (four of specimens from Swamp and four from JMS; GenBank MN342155); mt *cox I*, eight identical sequences (three of specimens from Swamp and five from JMS; GenBank MN342156); and mt NADH5, four sequences of specimens from Swamp and five of specimens from JMS; GenBank MN342157–MN342158).

ZooBank registration. To comply with the regulations set out in article 8.5 of the amended 2012 version of the International Code of

Zoological Nomenclature, details of the new species have been submitted to ZooBank. The Life Science Identifier for *A. winterbourni* n. sp. is urn:lsid:zoobank.org:pub:E0E7C4F3-430E-42BD-9FEC-2D11A159DD12.

Etymology. The species is named after Professor Michael Winterbourn in recognition for his contribution to the field of freshwater ecology – in particular, our knowledge of freshwater invertebrates of New Zealand, including the first mention of this species.

Description of adult (figs 1a–c and 2a–h; table 2; supplementary table S1)

(Based on whole mounts of 29 gravid specimens grown *in vitro* culture for 24–72 h and SEM preparations. Measurements provided as range and mean \pm standard deviation for the type series, variation associated with specimens from each age class (duration of culture) are provided separately in supplementary table S1.)

Body minute, triangular, often curved concave ventrally (body-width-to-length ratio 1:1.3–1.9 (1:1.6 \pm 0.4)) with maximum width at posterior level of testes, 145–200 (167 \pm 12) \times 85–115 (100 \pm 8). Tegument bears spines, glands and sensory papillae. Spines palmate, smooth, 5–9 prongs; present in lateral margins and along-side midline, separated by two narrow ventro-lateral regions devoid of spines, sparser towards posterior extremity on dorsal side; anterior forebody spines width 1.5–1.8, inter-spine space 0.6–0.8; mid-forebody spines width 1.9–2.2, inter-spines space 0.6–1; anterior dorsal spines width 2.2–2.3, inter-spines space 1.0–1.5; mid-dorsal spines width 1.5–1.8, inter-spines space 2.1–4.1; lateral posterior dorsal spines width 1.6–1.8, inter-spines space 1.2–1.5. Forebody 81–125 (97 \pm 9) long, representing 52–63% (58 \pm 2%) of body length. Glands and sensory papillae at anterior extremity, surrounding oral sucker (>14) and lateral margins of body.

Oral sucker subterminal, spherical, 21–29 (25 \pm 2) \times 21–29 (25 \pm 2). Ventral sucker at two-thirds of body length, subspherical, complete, 24–30 (27 \pm 1) \times 22–33 (28 \pm 2); outer rim crescent, interrupted sinistrally by genital pore, bearing spines and nine glands; oral-sucker-to-ventral-sucker-length ratio 1: 0.9–1.2 (1:1.1 \pm 0.2), width ratio 1: 0.9–1.3 (1:1.1 \pm 0.1). Pre-pharynx absent. Pharynx small, oval 11–17 (14 \pm 2) \times 11–15 (13 \pm 1). Pharynx-length-to-oral-sucker-length ratio 1:1.5–2.4 (1:1.7 \pm 0.2). Oesophagus 29–46 (36 \pm 5) long. Intestinal bifurcation pre-equatorial, immediately anterior to seminal vesicle. Caeca as long as oesophagus, widely divergent, extend to anterior margin of testes.

Testes two, postovarian, symmetrical, lateral, somewhat diagonal, smooth, slightly elongate-oval, right testis 18–29 (25 \pm 4) \times 17–28 (20 \pm 3); left testis 18–33 (25 \pm 5) \times 14–24 (19 \pm 3). Seminal vesicle arcuate, transversely oval, intercaecal in mid-body, overlapping anterior margin of ventral sucker dorsally, 12–24 (17 \pm 3) \times 24–39 (32 \pm 3). Seminal-vesicle-length-to-ventral-sucker-length ratio 1:1.1–2.2 (1:1.7 \pm 0.3). Seminal proximal duct long, entering prostatic chamber (i.e. phallophorus). Prostatic chamber subspherical, with loose fibres, sinistral to ventral sucker, dorsal to genital atrium, reaching proximal part of seminal vesicle. Prostatic glands not observed. Ejaculatory duct sinuous, enters prostatic chamber, opens to small papillae dorso-sinistral in genital atrium, together with prostatic ducts running through to periphery of male duct. Phallus of 'microphalloid'-type, glabrous, turgid, evaginable, with ejaculatory duct in axis, with large dextral triangular distal scale at the base. Genital atrium surrounding phallus, 16–24 (20 \pm 3), genital pore large, sinistral to ventral sucker.

Ovary dextral to ventral sucker, pre-testicular, ventral to caeca, adjacent to or slightly overlapping ventral sucker laterally,

subtriangular with large cells, 13–26 (21 \pm 3) \times 17–34 (25 \pm 4). Oötype inter-testicular, slightly dextral posterior to ovary and ventral sucker. Mehlis' gland surrounding oötype posteriorly. Laurer's canal not observed. Uterus confined posterior to mid-level of ventral sucker, overlapping testes ventrally. Metraterm thin-walled, with widened opening into sinistral wall of genital atrium. Vitellarium in two compact clusters of follicles, disaggregated in older specimens (after 24 h cultured), para-, post-testicular or overlapping testes, converging into seminal receptacle next to oötype. Eggs few, large. Excretory vesicle obscured by vitellarium masses and eggs. Flame-cell formula not observed.

Description of metacercaria (fig 1d; table 2; supplementary table S1)

Overall form highly developed and consistent with adult anatomy, except for the absence of eggs.

(Measurements based on 18 encysted specimens; measurements provided as range followed by mean \pm standard deviation.)

Metacercaria folded within small spherical, translucent cyst, 114–130 \times 106–120 (123 \times 113 \pm 4). Cyst wall consisting of three or more hyaline layers, 5–6 (5 \pm 0.5) thick. Metacercaria encysted in the gonads of *P. antipodarum*.

Genetic affinities

Two distinct genotypes for the 28S rRNA gene were obtained from the sequences of the nine specimens analysed. The two genotypes differed in two transitions at the nucleotide positions 281, a cytosine in the sequence of specimens originated from Swamp and a thymine in specimens from JMS and 1358, adenine and guanine, respectively. However, all specimens shared the same sequence for the ITS2 region and the mt *cox I*. Independent phylogenetic analyses of the 28S rDNA and ITS2 regions using BI and ML methods showed congruent results (figs 3 and 4), with the Microphallidae as monophyletic and *A. winterbourni* embedded within. *A. winterbourni* appeared as sister taxon to the sequence of an undescribed microphallid from Australia by Kudlai *et al.* (2015). In the 28S phylogenetic tree (fig. 3), the sequence of *M. fusiformis* appeared closely related to *A. winterbourni* and the undescribed microphallid with strong nodal support, although considerably divergent. In all trees, *Atriophallophorus* appears closely related to *Microphallus* and *Levinseniella*, but the sister relationship among these three genera could not be established due to the low support of an internal node within the clade (figs 3 and 4).

Genetic divergence between *A. winterbourni* and the unidentified microphallid from Australia was 1.2–1.3% for the 28S and 3.2% for the ITS2, which are within the lower limit of the range observed among congeneric species of *Microphallus* (28S: 0.8–9.1%; ITS2: 1.0–9.9%) and *Maritrema* (28S: 0.6–9.1%; ITS2: 0.3–12.3%). The 28S sequences of *A. winterbourni* and the unidentified microphallid diverged 10.5–10.8% from the sequence of *M. fusiformis*, which fell within range of intergeneric distances in the 28S region of microphallids (*Maritrema*–*Microphallus*: 7.5–12.6%; *Microphallus*–*Levinseniella*: 5.5–10.0%; *Maritrema*–*Levinseniella*: 9.1–11.4%; *Longiductotrema*–*Levinseniella*: 7.7%; *Longiductotrema*–*Maritrema*: 7.8–11.1%; *Longiductotrema*–*Microphallus*: 6.5–10.6%).

Sequence data for the COI marker resulted in identical sequences for all specimens (five from JMS and three from Swamp; one sample from swamp did not amplify and another one produced chromatograms with double peaks), whereas sequences for the NADH5 marker showed variability at the




Fig. 1. *Atriophallophorus winterbourni* n. sp. (a) Illustration of the holotype, 24 h *in vitro* grown adult in ventral view. (b) Microphotographs of the terminal genitalia of the holotype using light microscopy. (c) Histological oblique section of a paratype at the level of the ventral sucker. (d) Microphotograph of an encysted metacercaria ex *Potamopyrgus antipodarum* (Gray). Abbreviations: c, caeca; ga, genital atrium; pc, prostatic chamber; ph, phallus; sv, seminal vesicle; vs, ventral sucker.

position 191, with specimens from Swamp having a thymine and specimens from JMS a cytosine.

Discussion

Species differentiation

Following the key to the Microphallidae provided by Deblock (2008), the new species described from *in vitro* grown adults

conforms to the general morphology of the Microphallidae in having a typically very small body, longer than broad, densely covered with squamous spines, suckers well-separated and sub-equal, short digestive tract with divergent caeca not extending posteriorly beyond ventral sucker. Ovary pretesticular, in the opposite side of the body to the genital pore. Two testes, lateral and symmetrical, with male terminal genitalia intercaecal and anterior to ventral sucker. These specimens fit to the diagnosis of the supersubfamily Microphallidi Ward, 1901 by having the




Fig. 2. Scanning electron micrographs of *Atriophallophorus winterbourni* n. sp. (a) Adult. (b) Adult with phallus protruded. (c) Palmate spines on the ventral surface of the body. (d) Detail of the oral sucker, arrows point at a gland opening and a sensory papilla surrounding the oral sucker. (e) Detail of the outer rim of the ventral sucker with spination and the parietal atrial scale at the basis of the phallus. (f) Outer rim of the ventral sucker interrupted sinistrally by the opening for the genital pore with arrows pointing at glands. (g) Tip of the phallus evaginated, which appears as a flower-like structure when invaginated. (h) Detail of the configuration of the protruded phallus and the sinistrally interrupted outer rim of the ventral sucker.

terminal genitalia free in the parenchyma, and the subfamily Microphallinae Ward, 1901 by presenting a genital atrium that closely envelops the phallus. However, the opening of the genital atrium and the presence of the scale at the base of the phallus of the new species were found to alter the outer rim of the ventral sucker. Whereas the latter feature is characteristic of the monotypic Endocotylineae, our specimens differ from the genus

Endocotyle in lacking a connection between the cavity of the sucker and the genital pore, and presenting a scale at the base of a phallus of the 'microphalloid-type'. Our specimens fall within the Microphallini tribe by having a fleshy and muscular phallus, and the opening of the metraterm on the sinistral wall of the genital atrium. The new species exhibits features consistent with the genus *Atriophallophorus*: body of triangular shape; male genital

Table 2. Comparative metrical data for *Atriophallophorus* spp.

Species	<i>Atriophallophorus winterbourni</i> n. sp.				<i>Atriophallophorus minutus</i> (Price, 1934) Deblock & Rosé, 1964			<i>Atriophallophorus coxiellae</i> Smith, 1973	
	Adult		Metacercaria		Adult	Adult	Adult and metacercaria	Metacercaria	
Stage									
Hosts	<i>In vitro</i> cultured		<i>Potamopyrgus antipodarum</i> (Mollusca: Tateidae)		<i>Anas platyrhynchos</i> L. (Aves: Anatidae)	<i>Aythya affinis</i> (Eyton) (Aves: Anatidae)	<i>Ecriobia truncata</i> (Vanatta) and <i>Amnicola limosa</i> (Say) (Mollusca: Hydrobiidae) and mice (experimental)	<i>Coxiella badgerensis</i> (Johnston) (Mollusca: Tomichiidae); <i>Charadrius cucullatus</i> Vieillot and <i>C. ruficapillus</i> Temminck (Aves: Charadriidae)	
Distribution	New Zealand				North America, Caribbean Islands and Europe (France)			Tasmania	
Source	Present study		Present study		Deblock & Rosé, 1964	Price (1934)	Stunkard 1958	Smith, 1973	
	Range (n = 29)	Mean ± SD	Range (n = 9)	Mean ± SD	Range or mean		Range	Range	Mean
Body length	145–200 (28)	167 ± 12	165–193 (9)	175 ± 9	170–200	153–180	120–200	143–229	195
Body width	85–115 (28)	100 ± 8	98–118 (9)	106 ± 7	90–100	105–112	90–120	72–125	90
Forebody	81–125 (28)	97 ± 9	97–115 (9)	104 ± 5					
Oral sucker length	21–29 (29)	25 ± 2	23–28 (9)	25 ± 2	29 ^a		27 ^a		
Oral sucker width	21–29 (28)	25 ± 2	21–28 (9)	24 ± 2	22–25	23–25	23–30	26–34	29
Pharynx length	11–17 (26)	14 ± 2	12–14 (9)	14 ± 1	12 ^a		12 ^a	16–44	20
Pharynx width	11–15 (26)	13 ± 1	11–15 (9)	13 ± 1	12–15	10–18	10–18	15–20	17
Oesophagus	29–46 (22)	36 ± 5	40–50 (8)	44 ± 4		18–30	16–40		
Ventral sucker length	24–30 (28)	27 ± 1	28–35 (9)	31 ± 2	23 ^a		26 ^a		
Ventral sucker width	22–33 (28)	28 ± 2	24–31 (9)	28 ± 2	19–25	22–27	20–28	26–31	29
Right testis length	18–29 (13)	25 ± 4	26–34 (8)	30 ± 3					
Right testis width	17–28 (12)	20 ± 3	24–29 (9)	26 ± 2	20–30	28	12–29	36–40	39
Left testis length	18–33 (10)	25 ± 5	26–35 (7)	30 ± 3					
Left testis width	14–24 (10)	19 ± 3	23–29 (9)	26 ± 2	25–30				
Seminal vesicle length	12–24 (27)	17 ± 3	9–15 (2)	13 ± 3	13–19		14–20		
Seminal vesicle width	24–39 (28)	32 ± 3	35–44 (3)	40 ± 6	13–26		26 ^a	29–36	34
Ejaculatory duct length	21–34 (11)	27 ± 3			20–25				
Ejaculatory duct width	1.5–2.3 (13)	2.0 ± 0.4			2.5				
Genital atrium width	16–24 (10)	20 ± 3	22–23 (2)	22 ± 1	20–25		20–35		

Ovary length	13–26 (20)	21 ± 3	23–31 (9)	25 ± 2	19–25	18	15–20		
Ovary width	17–34 (20)	25 ± 4	17–31 (9)	26 ± 5	13–16	22	21–24	30–48	35
Right vitellarium length	15–26 (4)	20 ± 5	20–28 (9)	25 ± 3	20		5–6 follicles		
Right vitellarium width	23–35 (4)	29 ± 5	28–54 (8)	37 ± 9	30				
Left vitellarium length	15–28 (4)	21 ± 5	20–23 (9)	22 ± 1					
Left vitellarium width	23–37 (4)	28 ± 6	20–35 (9)	31 ± 5					
Egg number	1–12 (29)	5 ± 3			20–25		10–30		
Egg length × width	20–26 × 11–15				21–25 × 10–13	22 × 13	20–22 × 12–13		
Oral-to-ventral-sucker-length ratio	0.9–1.2	1.1 ± 0.2	1.1–1.4	1.3 ± 0.1	0.8 ^a		1.0 ^a	0.9 ^a	
Oral-to-ventral-sucker-width ratio	0.9–1.3	1.1 ± 0.1	0.9–1.3	1.1 ± 0.1	1.1		0.7 ^a	0.9 ^a	
Body-width-to-length ratio	1.3–1.9	1.6 ± 0.4	1.5–1.8	1.7 ± 0.1	1.8 ^a		–	1.6 ^a	
Seminal-vesicle-to-ventral-sucker-length ratio	1.1–2.2	1.7 ± 0.3	2.1–3.8	2.8 ± 0.9	0.7 ^a		1.0 ^a	1.9 ^a	
Pharynx-to-oral-sucker-length ratio	1.5–2.4	1.7 ± 0.2	1.7–2.0	1.8 ± 0.1	2.4 ^a		2.3 ^a	1.3 ^a	
Forebody as percentage of body length	52–63	58 ± 2	57–62	60 ± 1	62 ^a		–	63 ^a	
Metacercaria cyst length			114–130	123 ± 4					
Metacercaria cyst width			106–120	113 ± 4			≤100	88–125	
Metacercaria cyst wall			5–6	5 ± 0.5			8.0–10		

^aEstimated from the published illustration. *n*, number of specimens measured; SD, standard deviation of the mean.




Fig. 3. Phylogenetic relationships for representatives of the family Microphallidae, inferred by maximum likelihood analysis of 28S rDNA sequence data. The newly generated sequences are indicated in bold. Values on the branches correspond to posterior probabilities >0.95 followed by bootstrap support >60. Values below these thresholds were not reported.

pouch absent; proximal ejaculatory ducts very long and entering a large prostatic chamber (described as 'phallophorus' by Deblock & Rosé (1964)), subcircular and similar in size to the ventral sucker and dorsal to the genital atrium, with long prostatic ducts running through the periphery of the male duct; genital atrium containing a phallus of the 'microphalloid-type' and, at its base, a large dextral parietal atrial scale often protruding from the genital pore; a metraterm that widens at its opening into the sinistral wall of the genital atrium; and few but large eggs.

Currently the genus contains only two species, *A. minutus* (Price, 1934) and *A. coxiellae* Smith, 1973. *Atriophallophorus*

coxiellae was described from metacercariae infecting *Coxiella badgerensis* in a freshwater lake in Tasmania (Smith, 1973). However, the metacercaria of *A. winterbourni* n. sp. is readily distinguishable from *A. coxiellae* by having a different structure of the prostatic chamber, sub-circular and dorsal to genital atrium, rather than cylindrical, fibrous, elongate and placed between the seminal vesicle and the genital atrium, as well as smaller pharynx and testes, and somewhat smaller oral sucker and ovary, which barely overlap the lower limit of the range of *A. coxiellae*.

The new species is most similar to *A. minutus* with regards to the prostatic chamber and the morphometric data. But *A.*




Fig. 4. Phylogram for representatives of the family Microphallidae, inferred by maximum likelihood analysis of sequence data for the internal transcribed spacer 2 of the rRNA genes. The newly generated sequences are indicated in bold. Values on the branches correspond to posterior probabilities >0.95 followed by bootstrap support >60. Values below these thresholds were not reported.

minutus possesses oval testes and a transversely oval ovary, versus elongate-oval and subtriangular, respectively, in *A. winterbourni*. In addition, the original description of *A. minutus*, based on adult specimens from *Aythya affinis* from the Caribbean (Price, 1934), reported *A. minutus* had a shorter oesophagus than the new species, although this feature might vary greatly depending on the position of the specimen. Morphometric data provided by Stunkard (1958) for the redescription of *A. minutus* were based on adults experimentally grown in white mice from metacercariae infecting *Hydrobia minuta* and *Amnicola limosa* along the east coast of the USA. However, Stunkard's specimens were measured alive, whereas our measurements were taken from fixed, stained and mounted worms. Therefore, the morphometric data are not directly comparable. Furthermore, Stunkard noted

that fixed specimens were slightly smaller than alive, which suggests that if the measurements of *A. minutus* were taken after mounting they would be slightly smaller than those of *A. winterbourni*. Nonetheless, live specimens of *A. minutus* also show smaller dimensions for several metrical features (body length, ventral sucker width and testes width) extending outside the lower range for *A. winterbourni*, and eggs three times more numerous and with size overlapping the lower range of variation of the new species.

Metrical data of *A. minutus* described by Deblock & Rosé (1964) from France supports that *A. minutus* differs from *A. winterbourni*. Whereas the specimens examined by Deblock & Rosé (1964) were of comparable size to the largest in our sample, their ovary width is smaller than that of *A. winterbourni*, and the ventral sucker width

overlapped and extended below the lower range of variation of *A. winterbourni*. *Atriophallophorus minutus* also shows smaller values for the ratios oral sucker length to ventral sucker length and seminal vesicle length to ventral sucker length (estimated from the original illustrations) than *A. winterbourni*.

In vitro cultivation of the metacercaria and adult allowed us to observe developmental changes through time in the shape of the vitellarium, the seminal vesicle and testes (see also supplementary table S1). In metacercariae, the testes are well visible, the seminal vesicle is small, indistinct and quite empty, and the vitellarium forms two masses of tight follicles. After 24 h of growth, the adults are already gravid, testes become indistinct, the seminal vesicle is full and extended, and the follicles constituting the vitellarium masses start to disaggregate slightly. After 48 h or more, the testes are faintly visible, the seminal vesicle stays fully extended and the vitellarium follicles are drawn further apart in the posterior region of the body. When descriptions are based on specimens collected from the wild birds, this variation in the sample of specimens should be taken into account.

Previous researchers working on *Atriophallophorus* spp. have emphasized the difficulty of observing the features of specimens of such tiny size. Thus, it is highly recommended to describe new species with the support of genetic data and using a holistic biological approach (Blasco-Costa *et al.*, 2016a). Despite *A. winterbourni* showing subtle morphological differences from *A. minutus*, we consider them sufficient to distinguish the new species given the intrinsic difficulties of this group. Molecular results showed, convincingly, that specimens of *A. winterbourni* belong to a different microphallid genus to those already represented by sequence data. The new species appeared sister to an unidentified microphallid recovered from *Posticobia brazieri* (Smith) (Gastropoda, Tateidae) and *Caridina indistincta* Calman (Decapoda, Atyidae) in Australia by Kudlai *et al.* (2015). These authors suggested that their specimens were likely closely related to (or even conspecific) with the material reported as *Microphallus* sp. 'livelyi' by Hechinger (2012), which is considered a junior synonym of *A. winterbourni* herein. Divergence between our sequences and those of the Australian microphallid of Kudlai *et al.* (2015) suggests that they represent two distinct but congeneric species, as the authors anticipated.

The examination of specimens of *A. winterbourni* with SEM has allowed us to confirm a sinistral interruption of the outer rim of the ventral sucker caused by the opening of the genital pore and protrusion of the dextral parietal atrial scale, while light microscopy and histological sections show that, internally, the ventral sucker is complete. Deblock & Rosé (1964) mentioned the possibility of the scale being united to the rim of the ventral sucker (p. 229). However, this feature went unnoticed and was never confirmed by a later redescription or new species description for the genus, nor was it mentioned in the most recent diagnosis provided by Deblock (2008). Since the three species known for the genus so far are characterized by the presence of the large atrial scale, it is likely that in all three cases it has resulted in the same modification of the ventral sucker. Thus, we consider this feature of diagnostic value for the genus and amend herein the generic diagnosis of *Atriophallophorus* as follows.

Atriophallophorus Deblock & Rosé, 1964

Generic diagnosis

Body piriform or triangular, small (150–200 µm). Resembles *Microphallus*. Ventral sucker postequatorial, outer rim

interrupted sinistrally by genital pore. Oesophagus medium or short. Caeca short, divergent, in mid-body. Testes symmetrical in hindbody. Male genital pouch absent. Seminal vesicle intercaecal in mid-body, ovoid; prostatic gland caps distal part of seminal vesicle; proximal ejaculatory duct very long, supported by envelope acting as large prostatic chamber, either: (i) subcircular with diameter of ventral sucker, dorsal to genital atrium (formation described as 'phallophorus' (apparently bearing phallus)) and with long prostatic duct running through to periphery of male duct inside phallus; or (ii) cylindrical, fibrous, elongate between seminal vesicle and genital atrium, adjacent to margin of ventral sucker; bundle of long prostatic ducts enter as far as mid-part of cylindrical prostatic chamber (not inside phallus). Phallus of 'microphalloid'-type, more or less turgid, with ejaculatory duct in axis. Genital atrium present, envelopes phallus, with enormous dextral parietal atrial scale; genital pore sinistral to ventral sucker. Ovary dextral to ventral sucker. Uterus postcaecal, with few coils around testes. Metraterm long, with widened opening into sinistral wall of genital atrium. Eggs not numerous, relatively large. Vitellarium formed of two clusters of follicles, paratesticular and post-testicular in hindbody; vitelline ducts short, arched, post-testicular. Excretory vesicle short, Y-shaped, post-testicular. In intestine of birds (Anseriformes, Charadriiformes); cosmopolitan.

Type species. *Atriophallophorus minutus* (Price, 1934).

Synonym. *Atriophallophorus samarae* Deblock & Rosé, 1964.

Synonym lapsus calami. *Atriophallus samarae* Deblock & Rosé, 1964 (Fig. 1).

Life cycle and putative definitive hosts

Compared to other microphallid genera, the known species diversity of *Atriophallophorus* is quite low. So far, the four members of this genus (including the yet undescribed but molecularly characterized lineage from Australia of Kudlai *et al.* (2015)) show an abbreviated life cycle with the absence of the cercarial stage. Furthermore, *A. winterbourni* also lacks daughter sporocyst parthenitae-bearing germ cells (or has a very reduced life span), so that germ balls and embryonic metacercariae are observed free in the visceral mass of the snail host and appear to develop directly into encysted metacercariae, as reported by Krist & Lively (1998).

Based on the experimental exposure of snails to the faeces of different waterfowl species from Lake Alexandrina, Osnas & Lively (2011) concluded that the likely definitive hosts of *A. winterbourni* are mallard ducks (*Anas platyrhynchos* L.), grey ducks (*Anas superciliosa* Gmelin), their hybrids and the New Zealand scaup (*Aythya novaeseelandiae* Gmelin). Their conclusion agrees with the known distribution of both the putative hosts and *A. winterbourni* throughout the South Island. Furthermore, recent studies of the parasite fauna of mallards have discovered new species to science of microphallids and strigeids in New Zealand (Presswell *et al.*, 2014; Blasco-Costa *et al.*, 2016b). Altogether, these results highlight the still scarce knowledge on the parasites of birds, the most diverse group of vertebrates native to New Zealand, and the need for more biodiversity studies to address this gap.

Supplementary material. To view supplementary material for this article, please visit <https://doi.org/10.1017/S0022149X19000993>.

Acknowledgements. We thank Pilar Ruga Fahy from the University Medical Centre of the Pôle Facultaire de Microscopie Ultrastructurale at the University of Geneva for carrying out the histological sections and procedures, Janik Pralong (Natural History Museum of Geneva) for preparation of the permanent mounts and Dr André Piuze (Natural History Museum of Geneva) for his help with the SEM. We are also grateful to Dr Rodney Bray and Tomáš Scholz for discussions on the interpretation of the morphology, and one anonymous reviewer and the editor for very useful comments that helped us improve the manuscript.

Financial support. This work was supported by the Natural History Museum of Geneva, and indirectly by the Swiss National Science Foundation (IBC, grant number 31003A_169211/1).

Conflicts of interest. None.

Ethical standards. The authors assert that all procedures contributing to this work comply with the ethical standards of Switzerland and New Zealand and our institutional guides on the care and use of wild invertebrate animals.

References

- Al-Kandari WY and Al-Bustan SA (2010) Molecular identification of *Probolocoryphe uca* (Sarkisian, 1957; Digenea: Microphallidae) from Kuwait Bay using ITS1 and ITS2 sequences. *Parasitology Research* **106**, 1189–1195.
- Al-Kandari WY, Al-Bustan SA and Alnaqeeb M (2011) Ribosomal DNA sequence characterization of *Maritrema* cf. *eroliae* Yamaguti, 1939 (Digenea: Microphallidae) and its life cycle. *Journal of Parasitology* **97**, 1067–1074.
- Bankers L and Neiman M (2017) De novo transcriptome characterization of a sterilizing trematode parasite (*Microphallus* sp.) from two species of New Zealand snails. *G3—Genes Genomes Genetics* **7**, 871–880.
- Bankers L, Fields P, McElroy KE, Boore JL, Logsdon JM and Neiman M (2017) Genomic evidence for population-specific responses to co-evolving parasites in a New Zealand freshwater snail. *Molecular Ecology* **26**, 3663–3675.
- Blasco-Costa I and Poulin R (2017) Parasite life-cycle studies: a plea to resurrect an old parasitological tradition. *Journal of Helminthology* **91**, 647–656.
- Blasco-Costa I, Cutmore SC, Miller TL and Nolan MJ (2016a) Molecular approaches to trematode systematics: ‘best practice’ and implications for future study. *Systematic Parasitology* **93**, 295–306.
- Blasco-Costa I, Poulin R and Presswell B (2016b) Species of *Apatemon* Szidat, 1928 and *Australapatemon* Sudarikov, 1959 (Trematoda: Strigeidae) from New Zealand: Linking and characterising life cycle stages with morphology and molecules. *Parasitology Research* **115**, 271–289.
- Bowles J, Hope M, Tiu WU, Liu X and McManus DP (1993) Nuclear and mitochondrial genetic markers highly conserved between Chinese and Philippine *Schistosoma japonicum*. *Acta Tropica* **55**, 217–229.
- Cribb TH, Anderson GR, Adlard RD and Bray RA (1998) A DNA-based demonstration of a three-host life-cycle for the Bivesiculidae (Platyhelminthes: Digenea). *International Journal for Parasitology* **28**, 1791–1795.
- Deblock S (2008) The Microphallidae Ward, 1901. pp. 451–492 in Bray RA, Gibson DI and Jones A (Eds) *Keys to the Trematoda*. Vol. 3. London, CAB International and The Natural History Museum.
- Deblock S and Rosé F (1964) Contribution à l'étude des Microphallidae Travassos, 1920 (Trematoda) des oiseaux de France. VIII. Création du genre *Atriophallophorus*, parasite de canard sauvages. *Bulletin de la Société Zoologique de France* **89**, 225–230.
- Diaz JI and Cremonte F (2010) Development from metacercaria to adult of a new species of *Maritrema* (Digenea: Microphallidae) parasitic in the kelp gull, *Larus dominicanus*, from the Patagonian coast, Argentina. *Journal of Parasitology* **96**, 740–745.
- Dybdahl MF and Krist AC (2004) Genotypic vs. condition effects on parasite-driven rare advantage. *Journal of Evolutionary Biology* **17**, 967–973.
- Dybdahl MF and Lively CM (1996) The geography of coevolution: comparative population structures for a snail and its trematode parasite. *Evolution* **50**, 2264–2275.
- Dybdahl MF and Lively CM (1998) Host-parasite coevolution: evidence for rare advantage and time-lagged selection in a natural population. *Evolution* **52**, 1057–1066.
- Dybdahl MF, Jokela J, Delph LF, Koskella B and Lively CM (2008) Hybrid fitness in a locally adapted parasite. *American Naturalist* **172**, 772–782.
- Fromme AE and Dybdahl MF (2006) Resistance in introduced populations of a freshwater snail to native range parasites. *Journal of Evolutionary Biology* **19**, 1948–1955.
- Galaktionov KV and Blasco-Costa I (2018) *Microphallus ochotensis* sp. nov. (Digenea, Microphallidae) and relative merits of two-host microphallid life cycles. *Parasitology Research* **117**, 1051–1068.
- Galaktionov KV and Dobrovolskij AA (2003) *The biology and evolution of trematodes*. p. 592. Dordrecht, The Netherlands, Kluwer Academic Publishers.
- Galaktionov KV, Blasco-Costa I and Olson PD (2012) Life cycles, molecular phylogeny and historical biogeography of the ‘pygmaeus’ microphallids (Digenea: Microphallidae): widespread parasites of marine and coastal birds in the Holarctic. *Parasitology* **139**, 1346–1360.
- Gibson AK, Jokela J and Lively CM (2016a) Fine-scale spatial covariation between infection prevalence and susceptibility in a natural population. *The American Naturalist* **188**, 1–14.
- Gibson AK, Xu JY and Lively CM (2016b) Within-population covariation between sexual reproduction and susceptibility to local parasites. *Evolution* **70**, 2049–2060.
- Gibson AK, Delph LF, Vergara D and Lively CM (2018) Periodic, parasite-mediated selection for and against sex. *American Naturalist* **192**, 537–551.
- Gilardoni C, Etchegoin J, Diaz JI, Ituarte C and Cremonte F (2011) A survey of larval digeneans in the commonest intertidal snails from Northern Patagonian coast, Argentina. *Acta Parasitologica* **56**, 163.
- Hamilton WD (1980) Sex versus non-sex versus parasite. *Oikos* **35**, 282–290.
- Hechinger RF (2012) Faunal survey and identification key for the trematodes (Platyhelminthes: Digenea) infecting *Potamopyrgus antipodarum* (Gastropoda: Hydrobiidae) as first intermediate host. *Zootaxa* **3418**, 1–27.
- Hernandez-Orts JS, Pinacho-Pinacho CD, Garcia-Varela M and Kostadinova A (2016) *Maritrema corai* n. sp. (Digenea: Microphallidae) from the white ibis *Eudocimus albus* (Linnaeus) (Aves: Threskiornithidae) in Mexico. *Parasitology Research* **115**, 547–559.
- Hofmann H, Blasco-Costa I, Knudsen R, Matthaei CD, Valois A and Lange K (2016) Parasite prevalence in an intermediate snail host is subject to multiple anthropogenic stressors in a New Zealand river system. *Ecological Indicators* **60**, 845–852.
- Huelsenbeck JP, Ronquist F, Nielsen R and Bollback JP (2001) Bayesian inference of phylogeny and its impact on evolutionary biology. *Science* **294**, 2310–2314.
- Jaenike J (1978) A hypothesis to account for the maintenance of sex in populations. *Evolutionary Theory* **3**, 191–194.
- Jokela J and Lively CM (1995) Spatial variation in infection by digenetic trematodes in a population of freshwater snails (*Potamopyrgus antipodarum*). *Oecologia* **103**, 509–517.
- Jokela J, Lively CM, Dybdahl MF and Fox JA (2003) Genetic variation in sexual and clonal lineages of a freshwater snail. *Biological Journal of the Linnean Society* **79**, 165–181.
- Jokela J, Dybdahl MF and Lively CM (2009) The maintenance of sex, clonal dynamics, and host-parasite coevolution in a mixed population of sexual and asexual snails. *The American Naturalist* **174**, S43–S53.
- Kakui K (2014) A novel transmission pathway: first report of a larval trematode in a tanaidacean crustacean. *Fauna Ryukyuan* **17**, 13–22.
- King KC, Delph LF, Jokela J and Lively CM (2009) The geographic mosaic of sex and the red queen. *Current Biology* **19**, 1438–1441.
- King KC, Delph LF, Jokela J and Lively CM (2011a) Coevolutionary hotspots and coldspots for host sex and parasite local adaptation in a snail-trematode interaction. *Oikos* **120**, 1335–1340.
- King KC, Jokela J and Lively CM (2011b) Parasites, sex, and clonal diversity in natural snail populations. *Evolution* **65**, 1474–1481.

- Koskella B and Lively CM (2007) Advice of the rose: experimental coevolution of a trematode parasite and its snail host. *Evolution* **61**, 152–159.
- Koskella B and Lively CM (2009) Evidence for negative frequency-dependent selection during experimental coevolution of a freshwater snail and a sterilizing trematode. *Evolution* **63**, 2213–2221.
- Koskella B, Vergara D and Lively CM (2011) Experimental evolution of sexual host populations in response to sterilizing parasites. *Evolutionary Ecology Research* **13**, 315–322.
- Krist AC and Lively CM (1998) Experimental exposure of juvenile snails (*Potamopyrgus antipodarum*) to infection by trematode larvae (*Microphallus* sp.): infectivity, fecundity compensation and growth. *Oecologia* **116**, 575–582.
- Krist AC, Lively CM, Levri EP and Jokela J (2000) Spatial variation in susceptibility to infection in a snail-trematode interaction. *Parasitology* **121**, 395–401.
- Krist AC, Jokela J, Wiehn J and Lively CM (2004) Effects of host condition on susceptibility to infection, parasite developmental rate, and parasite transmission in a snail-trematode interaction. *Journal of Evolutionary Biology* **17**, 33–40.
- Kudlai O, Cutmore SC and Cribb TH (2015) Morphological and molecular data for three species of the Microphallidae (Trematoda: Digenea) in Australia, including the first descriptions of the cercariae of *Maritrema brevissaciferum* Shimazu et Pearson, 1991 and *Microphallus minutus* Johnston, 1948. *Folia Parasitologica (Praha)* **62**, 053.
- Kudlai O, Cribb TH and Cutmore SC (2016) A new species of microphallid (Trematoda: Digenea) infecting a novel host family, the Muraenidae, on the northern Great Barrier Reef, Australia. *Systematic Parasitology* **93**, 863–876.
- Kumar S, Stecher G, Li M, Knyaz C and Tamura K (2018) MEGA X: molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution* **35**, 1547–1549.
- Lagroe C and Poulin R (2008) Lack of seasonal variation in the life-history strategies of the trematode *Coitocaecum parvum*: no apparent environmental effect. *Parasitology* **135**, 1243–1251.
- Lagroe C, McEwan J, Poulin R and Keeney DB (2007) Co-occurrences of parasite clones and altered host phenotype in a snail-trematode system. *International Journal for Parasitology* **37**, 1459–1467.
- Levri EP (1999) Parasite-induced change in host behavior of a freshwater snail: parasitic manipulation or byproduct of infection? *Behavioral Ecology* **10**, 234–241.
- Levri EP and Fisher LM (2000) The effect of a trematode parasite (*Microphallus* sp.) on the response of the freshwater snail *Potamopyrgus antipodarum* to light and gravity. *Behaviour* **137**, 1141–1151.
- Levri EP and Lively CM (1996) The effects of size, reproductive condition, and parasitism on foraging behaviour in a freshwater snail, *Potamopyrgus antipodarum*. *Animal Behaviour* **51**, 891–901.
- Levri EP, Dillard J and Martin T (2005) Trematode infection correlates with shell shape and defence morphology in a freshwater snail. *Parasitology* **130**, 699–708.
- Littlewood DTJ, Curini-Galletti M and Herniou EA (2000) The interrelationships of Proseriata (Platyhelminthes: Seriata) tested with molecules and morphology. *Molecular Phylogenetics and Evolution* **16**, 449–466.
- Lively CM (1987) Evidence from a New Zealand snail for the maintenance of sex by parasitism. *Nature* **328**, 519.
- Lively CM (1989) Adaptation by a parasitic trematode to local populations of its snail host. *Evolution* **43**, 1663–1671.
- Lively CM (2016) Coevolutionary epidemiology: disease spread, local adaptation, and sex. *The American Naturalist* **187**, E77–E82.
- Lively CM and Dybdahl MF (2000) Parasite adaptation to locally common host genotypes. *Nature* **405**, 679.
- Lively CM and Jokela J (1996) Clinal variation for local adaptation in a host-parasite interaction. *Proceedings of the Royal Society B-Biological Sciences* **263**, 891–897.
- Lively CM and McKenzie JC (1991) Experimental infection of a freshwater snail, *Potamopyrgus antipodarum*, with a digenetic trematode, *Microphallus* sp. *New Zealand Natural Sciences* **18**, 59–62.
- Lively CM, Dybdahl MF, Jokela J, Osnas EE and Delph LF (2004) Host sex and local adaptation by parasites in a snail-trematode interaction. *American Naturalist* **164**, S6–S18.
- Lively CM, Delph LF, Dybdahl MF and Jokela J (2008) Experimental test for a co-evolutionary hotspot in a host-parasite interaction. *Evolutionary Ecology Research* **10**, 95–103.
- Lockyer AE, Olson PD and Littlewood DTJ (2003) Utility of complete large and small subunit rRNA genes in resolving the phylogeny of the Neodermata (Platyhelminthes): implications and a review of the Cercomer theory. *Biological Journal of the Linnean Society* **78**, 155–171.
- McKone MJ, Gibson AK, Cook D, Freymiller LA, Mishkind D, Quinlan A, York JM, Lively CM and Neiman M (2016) Fine-scale association between parasites and sex in *Potamopyrgus antipodarum* within a New Zealand lake. *New Zealand Journal of Ecology* **40**, 330–333.
- Miller MA, Pfeiffer W and Schwartz T (2010), Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In *Proceedings of the Gateway Computing Environments Workshop (GCE)*, New Orleans, LA18.
- Morgan JAT and Blair D (1995) Nuclear rDNA ITS sequence variation in the trematode genus *Echinostoma*: an aid to establishing relationships within the 37-collar-spine group. *Parasitology* **111**, 609–615.
- O'Dwyer K, Blasco-Costa I, Poulin R and Faltnykova A (2014) Four marine digenetic parasites of *Austrolittorina* spp. (Gastropoda: Littorinidae) in New Zealand: morphological and molecular data. *Systematic Parasitology* **89**, 133–152.
- Osnas EE and Lively CM (2004) Parasite dose, prevalence of infection and local adaptation in a host-parasite system. *Parasitology* **128**, 223–228.
- Osnas EE and Lively CM (2006) Host ploidy, parasitism and immune defence in a coevolutionary snail-trematode system. *Journal of Evolutionary Biology* **19**, 42–48.
- Osnas EE and Lively CM (2011) Using definitive host faeces to infect experimental intermediate host populations: waterfowl hosts for New Zealand trematodes. *New Zealand Journal of Zoology* **38**, 83–90.
- Paczynski D, Adolfsen S, Liljeroos K, Klappert K, Lively CM and Jokela J (2014) Faster clonal turnover in high-infection habitats provides evidence for parasite-mediated selection. *Journal of Evolutionary Biology* **27**, 417–428.
- Paczynski D, Klappert K, Kopp K, Neiman M, Seppälä K, Lively CM and Jokela J (2019) Parasite resistance predicts fitness better than fecundity in a natural population of the freshwater snail *Potamopyrgus antipodarum*. *Evolution* **73**, 1634–1646.
- Presswell B, Blasco-Costa I and Kostadinova A (2014) Two new species of *Maritrema* Nicoll, 1907 (Digenea: Microphallidae) from New Zealand: morphological and molecular characterisation. *Parasitology Research* **113**, 1641–1656.
- Price EW (1934) New trematode parasites of birds. *Smithsonian Miscellaneous Collections* **91**, 1–6.
- Ronquist F, Teslenko M, van der Mark P, et al. (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* **61**, 539–542.
- Smith SJ (1973) Three new microphallid trematodes from Tasmanian birds. *Papers and proceedings of the Royal Society of Tasmania* **107**, 197–205.
- Snyder SD and Tkach VV (2001) Phylogenetic and biogeographical relationships among some Holarctic frog lung flukes (Digenea: Haematoloecidae). *Journal of Parasitology* **87**, 1433–1440.
- Soper DM, King KC, Vergara D and Lively CM (2014) Exposure to parasites increases promiscuity in a freshwater snail. *Biology Letters* **10**, 20131091.
- Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313.
- Stunkard HW (1958) The morphology and life-history of *Levinseniella minuta* (Trematoda: Microphallidae). *The Journal of Parasitology* **44**, 225–229.
- Tkach V, Pawlowski J and Mariaux J (2000) Phylogenetic analysis of the sub-order Plagiorchiata (Platyhelminthes, Digenea) based on partial lsrDNA sequences. *International Journal for Parasitology* **30**, 83–93.
- Tkach VV, Littlewood DTJ, Olson PD, Kinsella JM and Swiderski Z (2003) Molecular phylogenetic analysis of the Microphalloidea Ward, 1901 (Trematoda: Digenea). *Systematic Parasitology* **56**, 1–15.
- Vergara D, Lively CM, King KC and Jokela J (2013) The geographic mosaic of sex and infection in lake populations of a New Zealand snail at multiple spatial scales. *American Naturalist* **182**, 484–493.

- Vergara D, Jokela J and Lively CM** (2014) Infection dynamics in coexisting sexual and asexual host populations: support for the Red Queen Hypothesis. *American Naturalist* **184**, S22–S30.
- Vergara D, Fuentes JA, Stoy KS and Lively CM** (2017) Evaluating shell variation across different populations of a freshwater snail. *Molluscan Research* **37**, 120–132.
- Werle E, Schneider C, Renner M, Volker M and Fiehn W** (1994) Convenient single-step, one tube purification of PCR products for direct sequencing. *Nucleic Acids Research* **22**, 4354–4355.
- Winterbourn MJ** (1974) Larval trematoda parasitising the New Zealand species of *Potamopyrgus* (Gastropoda : Hydrobiidae). *Mauri Ora* **2**, 17–30.