

SP0008

Predictors of Adherence and Response to Exercise Interventions in Schizophrenia

L. Röll

Psychiatry and Psychotherapy, LMU Hospital, Munich, Germany
doi: 10.1192/j.eurpsy.2024.53

Abstract: Exercise can be considered a feasible and efficient add-on treatment in schizophrenia. However, there remain two major challenges with regard to feasibility and efficiency: First, drop-out rates during exercise programs appear to be relatively high, ranging between 30 to 80 percent. Second, only between 30 and 50 percent of patients clinically respond to exercise interventions. Hence, we aimed to identify factors that predict adherence and response to exercise programs in people with schizophrenia. Based on data from 180 patients with schizophrenia enrolled in the Enhancing Schizophrenia Prevention and Recovery through Innovative Treatments (ESPRIT) C3 study, we examined clinical baseline characteristics that may predict study completion and number of attended trainings (adherence), as well as clinically relevant improvements in symptomatology and functioning (response). We found that only levels of functioning at baseline, but not symptom severity, cognitive functioning, or physical health, predicted adherence. Further, we provide preliminary evidence suggesting that patients with higher cognitive abilities and higher education who performed regular exercise already prior to the study participation were more likely to respond. To conclude, our findings indicate that exercise is particularly helpful for a subgroup of patients characterized by higher levels of functioning, higher cognitive abilities and education, and more pronounced affinity to exercise. Future studies should additionally include environmental, genetic, and neural data to predict adherence and response to exercise.

Disclosure of Interest: None Declared

SP0007

Telepsychiatry as catalysts for equitable Mental Health Care in Intellectual Disabilities

K. Krysta

Department of Rehabilitation Psychiatry, Medical University of Silesia, Katowice, Poland
doi: 10.1192/j.eurpsy.2024.54

Abstract: Telepsychiatry has emerged as a transformative force in the field of mental health care, addressing disparities in service delivery and increasing access to care. This exploration focuses on the role of telepsychiatry in achieving equitable mental health care for individuals with intellectual disabilities (ID). Intellectual disabilities affect millions globally, posing significant public health challenges. This vulnerable population encounters numerous barriers in accessing quality mental health care, including geographical isolation, limited transportation options, and a shortage of specialized providers. Telepsychiatry offers a promising solution,

leveraging technology to overcome these challenges. The presentation reviews the current landscape of mental health care for individuals with intellectual disabilities and the specific barriers they encounter. It highlights the potential benefits of telepsychiatry, including increased availability of specialized care, reduced geographical barriers, and enhanced caregiver support. Ethical considerations and best practices associated with implementing telepsychiatry in the context of intellectual disabilities are discussed. Case studies and success stories illustrate how telepsychiatry positively impacts individuals with intellectual disabilities and their families. In conclusion, telepsychiatry plays a promising role in promoting equitable mental health care for individuals with intellectual disabilities. Embracing technology and adopting best practices pave the way for a more inclusive and accessible mental health care system, leaving no one behind.

Disclosure of Interest: None Declared

SP0008

Teleassistance Telerehabilitation Services for urgent mental health needs of people with Intellectual and Developmental disabilities

M. O. Bertelli

CREA (Research and Clinical Centre), San Sebastiano Foundation, Misericordia di Firenze, Florence, Italy
doi: 10.1192/j.eurpsy.2024.55

Abstract: Persons with intellectual disability (PwID) and/or autism spectrum disorder with high support needs (ASD-HSN) have resulted to be among the most vulnerable populations to COVID-19 and distress factors associated to the measures for containing its spread. Many health, rehabilitation, and assistance needs were managed through the use of telemedicine, specifically teleassistance (TA) and telerehabilitation (TR), with regard to the prevention and treatment of the epidemic illness as well as the continuity of care required for the condition of developmental disability and co-occurring physical or mental disorders. TA and TR can function either directly or indirectly with the PwID/ASD; in the latter case, a family member, a regular caregiver, or a technician provides local mediation. This paper examines the most common TA and TR activities, along with their requirements, applications, and goals. All of these activities should be in line with the overarching goal of each customized therapy and rehabilitation plan, which is to enhance and support the quality of life for people with intellectual and developmental disabilities.

Studies on TA and TR efficacy for PwID/ASD are limited, especially concerning adulthood. The scant research that is currently available demonstrates efficacy in maintaining or marginally enhancing cognitive, adaptive, and vocational skills. In addition to managing both routine and unusual activities as well as critical episodes, family members and other caregivers reported feeling more empowered about their educational and interpersonal abilities with the PwID/ASD. The primary benefits over traditional in-person services have been found to be greater

accessibility and availability as well as a reduction in both physical and psychological distance. The primary limits include the absence of all elements of the therapeutic alliance pertaining to face-to-face communication, possible poor ability to use technology, accessibility of the technology itself, concerns regarding privacy, and variables that divert attention connected to the household setting.

Even while telemedicine has proven to be feasible and beneficial thus far, it is doubtful that traditional techniques will be able to be replaced, at least not anytime soon. Telemedicine could, nonetheless, serve as a useful addition, integration, or short-term substitute. Future studies should provide light on the indications, contextual deployment, efficacy evaluation, and operational stability over time of certain TA and TR activities in addition to the use of artificial intelligence, machine learning, and interactive avatars.

Disclosure of Interest: None Declared

SP0009

DefiGame, a serious game to discover neurodevelopmental disorders

C. Bonnet*, C. IMMESOETE and Specialists and parents affiliated to the french rare disease network DéfiScience - Professeur Vincent des Portes, Mme Marie-Pierre Reymond, Mme Caroline Immesoete, Mme Fanny Forel, Mme ANTOINE Odile, Mme AUPETIT Laure, Mme BENOIT Céline, Docteur BROSSEA

Filière DéfiScience, LYON, France

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.56

Abstract: Défigame is a training tool designed in collaboration with parents and specialists from the French Rare Diseases Network DéfiScience.

In this serious game, you take on the role of a general practitioner treating four young patients whose developmental trajectories raise questions. Interactively and with the help of concrete tools, you'll learn about the recommendations for coordinating an appropriate course of prescription, care and support for a family, from the search for a diagnosis to early management of a Neurodevelopmental Disorder (NDD).

WHO IS DEFIGAME FOR?

- Any European doctor questioning the etiology of a neurodevelopmental disorder, prescribing genetic tests or wishing to update their knowledge in the field of NDD, particularly in relation to a rare disease.
- Any other healthcare professional concerned with etiological diagnosis or support for people with NDD and their families.

Disclosure of Interest: None Declared

SP0010

Patients' perceptions and preferences regarding telemedicine for addictive disorders

S. Achab

Psychiatry, Clinical and Sociological Research Unit, Faculty of Medicine, Geneva University, Geneva, Switzerland

doi: 10.1192/j.eurpsy.2024.57

Abstract: Telemedicine is an emerging treatment option having been heavily used during covid lockdowns, in order to maintain treatment access including for addictive disorders.

In the present talk, we first present data published on the challenges met at ReConnecte the treatment facility for Addictive behaviors during pandemic.

We second, present results of a survey we conducted on preferences of telemedicine use in patients and doctors in our Geneva University Hospitals.

We finally illustrate findings by clinical cases of patients suffering from addictive behaviors and their specific needs and preferences in terms of telemedicine (phone or Visioconsultation).

Preferences and ehealth tools elicited depended of their psychosocial profiles, their specific needs and expected benefits from online sessions of psychotherapy.

One of the ingredients of successful psychotherapy for addictive behaviors, is the purposeful use of telemedicine as an integrated treatment modality.

Disclosure of Interest: None Declared

SP0011

How digital technology can contribute to timely and effective recognition and response to opioid overdose events

A. M. Baldacchino*

DigitAS St Andrews

Medicine, St Andrews University, St Andrews, United Kingdom

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.58

Abstract

Objectives: To discuss novel approaches in the development early detection, response and interventions of drug overdoses.

Rationale: There is an urgent need to research and develop novel strategies to rapidly and accurately detect, respond, and treat them with the ultimate goal of reducing drug deaths secondary to fatal drug overdose incidents. This should be additional to supporting communities and networks able to intervene utilizing novel public health approaches.

Methods: We will describe technologies and associated systems that are able to accelerate detection and result in a timely response to potential overdose with effective and timely intervention to these