



Research Paper

*These authors contributed equally to this work.

Cite this article: Xu L, Liu H, Li B, Li G, Liu R, Li D (2024). SlCarE054 in Spodoptera litura (Lepidoptera: Noctuidae) showed direct metabolic activity to beta-cypermethrin with stereoselectivity. Bulletin of Entomological Research 114, 482–490. https://doi.org/10.1017/S0007485324000282

Received: 22 December 2023
Revised: 26 March 2024
Accepted: 12 April 2024
First published online: 6 May 2024

Keywords: heterologous expression; metabolic activity; SlCarE054; stereoselectivity; beta-cypermethrin

Corresponding authors: Runqiang Liu; Email: liurqhist@126.com; Dongzhi Li; Email: lidongzhi1989@163.com

SlCarE054 in Spodoptera litura (Lepidoptera: Noctuidae) showed direct metabolic activity to beta-cypermethrin with stereoselectivity

Li Xu1,2,* , Hongyu Liu1,2,* , Bo Li1,2 , Guangling Li1,2 , Runqiang Liu1,2 ID and Dongzhi Li1,2 ID

1Henan Engineering Research Center of Green Pesticide Creation & Intelligent Pesticide Residue Sensor Detection, Henan Institute of Science and Technology, Xinxiang 453003, China and 2College of Resources and Environment, Henan Institute of Science and Technology, Xinxiang 453003, China

Abstract

Carboxylesterases (CarEs) is an important detoxification enzyme system in phase I participating in insecticides resistance. In our previous study, SlCarE054, a CarEs gene from lepidoptera class, was screened out to be upregulated in a pyrethroids and organophosphates resistant population. Its overexpression was verified in two field-collected populations of Spodoptera litura (Lepidoptera: Noctuidae) resistant to pyrethroids and organophosphates by qRT-PCR. Spatiotemporal expression results showed that SlCarE054 was highly expressed in the pupae stage and the digestive tissue midgut. To further explore its role in pyrethroids and organophosphates resistance, its metabolism activity to insecticides was determined by UPLC. Its recombinant protein showed significant metabolism activity to cyhalothrin and fenvalerate, but not to phoxim or chlorpyrifos. The metabolic activity of SlCarE054 to beta-cypermethrin showed stereoselectivity, with higher metabolic activity to theta-cypermethrin than the enantiomer alpha-cypermethrin. The metabolite of beta-cypermethrin was identified as 3-phenoxybenzaldehyde. Further modelling and docking analysis indicated that beta-cypermethrin, cyhalothrin and fenvalerate could bind with the catalytic triad of the 3D structure of SlCarE054. The interaction of beta-cypermethrin with SlCarE054 also showed the lowest binding energy. Our work provides evidence that SlCarE054 play roles in beta-cypermethrin resistance in S. litura.

Introduction

Spodoptera litura (Lepidoptera: Noctuidae) is an important agricultural pest feeding on host plants covering more than 120 species, including many commercial crops, cotton, soybean, tobacco, cruciferous vegetable, and so on (https://www.cabidigitallibrary.org/doi/10.1079/cabicompendium.44520). S. litura is also a gluttonous insect with high fecundity and mobility, and easy to break out. As the frequent chemical control, S. litura has developed resistance to 42 chemical insecticides (https://www.pesticideresistance.org/display.php?page=species&arId=282), especially to pyrethroids, organophosphates and carbamates (Ahmad et al., 2008; Wan et al., 2008; Saleem et al., 2016; Wang et al., 2018; Xu et al., 2020; Zhang et al., 2022). Pyrethroids showed strong insecticidal activity, low toxicity to mammals, broad spectrum and excellent stability. Cyhalothrin, beta-cypermethrin and fenvalerate are conventional pyrethroids widely used in controlling pests in agriculture, public health and graziery. Metabolism resistance mediated by the detoxification enzyme of cytochrome P450 systems (P450s), carboxylesterases (CarEs) and glutathione S-transferases (GSTs) are proved to be an important reason leading to the resistance to pyrethroids, organophosphates, and carbamates (Li et al., 2007).

CarEs genes in insects are divided into 14 clades (A-N) showing functions in three aspects, dietary detoxification, pheromone/hormone degradation and neurodevelopment (Oakeshott et al., 2005). The main CarEs clades included alpha-esterases, juvenile hormone esterases, beta-esterases, integument esterase, gliotactins, acetylcholinesterases, neurotactins, neuroligins, glutactin, and uncharacterised group (Oakeshott et al., 2005). CarEs genes in alpha-esterase, juvenile hormone esterase and beta-esterase families accounted for the majority of catalytically active CarEs (Ranson et al., 2002). CarEs genes in neurotactin, neuroligin, gliotactin, and glutactin are considered to be noncatalytic, and play essential roles in cell-to-cell interactions (Oakeshott et al., 2005), pheromone degradation in moths (Vogt, 2005) and hydrolysis of the neurotransmitter acetylcholine and juvenile hormone (Taylor and Radic, 1994; Riddiford et al., 2003). As a detoxification enzyme in insects, CarEs function as phase I enzyme and could catalytic hydrolysis insecticides contain carboxyl ester by gene amplification, upregulation and coding sequence mutations (Bass et al., 2014; Wang et al., 2015; Hopkins et al., 2017). The CarEs involved hydrolyse reaction is divided into two steps.



Firstly, a nucleophilic attack by the oxygen of serine residue at the active site is taken on the carbonyl carbon of the substrate, releasing an alcohol product and forming an acyl-enzyme. Secondly, another nucleophilic attack is taken by water, releasing an acid product and forming the free enzyme (Oakeshott *et al.*, 2005).

In *S. litura*, multiple studies have implied the involvement of CarEs in insecticide resistance by synergist experiments (Armes *et al.*, 1997; Ahmad *et al.*, 2007; Huang and Han, 2007), increased enzyme activities (Li *et al.*, 2021a), or overexpression of CarEs genes in resistant strains (Xu *et al.*, 2020). Only a few researches provided the evidence proving the involvement of CarEs genes in insecticide resistance (Shi *et al.*, 2022; Li *et al.*, 2023). The function of each specific CarEs genes in insecticides resistance is still lacking.

In our previous study, the addition of CarEs synergist, triphenyl phosphate (TPP), significantly increased the mortality of larvae from a field-collected population after pyrethroids and organophosphates treatment, suggesting a possible role of CarEs genes in pyrethroids and organophosphates resistance (Li *et al.*, 2021b). A series CarEs genes were found to show significantly higher reads in the field-collected population resistant to pyrethroids and organophosphates than the susceptible population by RNA-Seq (data derived from NCBI Sequence Read Archive database with accession number PRJNA843172, Table S1, Xu *et al.*, 2022). *SlCarE054* was selected to heterologous expressed due to its high expression level in the resistant population. Its metabolism activity to pyrethroids and organophosphates was determined by ultra performance liquid chromatography (UPLC), and the metabolite was identified by Gas Chromatography-Mass Spectrometer (GC-MS). The interaction between insecticides and *SlCarE054* was also conducted by molecular docking. Our results enriched the knowledge of the role of CarEs genes in insecticides resistance, and explained the CarEs involved molecular mechanism to pyrethroids and organophosphates resistance in *S. litura*.

Materials and methods

Insects and chemicals

The susceptible population (GX) was firstly collected from Nanning, Guangxi province, China and provided by Guangxi Tianyuan Biochemistry Corp., Ltd. QJ population, showing extremely high-level resistance to fenvalerate (Resistance ratio, $RR > 25,000$), β -cypermethrin ($RR = 3659$ - fold), cyhalothrin ($RR > 50,000$), phoxim ($RR = 295$ - fold) and chlorpyrifos ($RR = 207$ - fold) compared with GX population (Li *et al.*, 2021b), was firstly collected from Qianjiang, Hubei province, China at 2019. KX population was collected from Xinxiang, Henan province, China at 2022. The rear condition was: $26 \pm 1^\circ\text{C}$, 60–70% relative humidity, 14:10 h of light:dark photoperiod.

The technical grade of fenvalerate (93.4%) was provided by Jiangsu Changlong Chemicals Co., Ltd. The technical grade of β -cypermethrin (95.0%), cyhalothrin (98.4%), phoxim (91.0%) and chlorpyrifos (95.0%), were obtained from Beijing Huarong Biochemical Co., Ltd.

Bioassay was determined by topical application described in Xu *et al.* (2020). Briefly, a series diluted insecticides were prepared in acetone and one microliter of insecticides was applied on the thoracic dorsum of third instar larvae of *S. litura* from GX and KX populations. The mortality of larvae was checked 48 h after treatment. Application of acetone was used as control. Each concentration consisted of three replicates.

Cloning the coding sequence of *SlCarE054* and its bioinformatics analysis

The total RNA from third instar larvae was extracted by TaKaRa MiniBEST Universal RNA Extraction Kit (Takara, Dalian, China). cDNA was synthesised using the FastQuant RT Kit (with gDNase) (Tiangen, Beijing, China). The coding sequence of *SlCarE054* was cloned with primer pairs in Table S2. The PCR product was purified by TIANgel Midi Purification Kit (Tiangen) and inserted to pLB vector (Tiangen), then transformed into *E. coli* DH5 α competent cells. Positive clones were selected and sequenced at Sangon Biotech (Shanghai, China).

The protein molecular weight and theoretical isoelectric point (pI) of *SlCarE054* were predicted by Compute pI/Mw tool (https://web.expasy.org/compute_pi/). The signal peptide and protein glycosylation were detected by SignalP-6.0 (<https://services.healthtech.dtu.dk/services/SignalP-6.0/>) and NetNGlyc1.0 server (<http://www.cbs.dtu.dk/services/NetNGlyc/>), respectively. The amino acid sequence of *SlCarE054* was blasted with CarEs genes in other lepidoptera insects using GENEDEC software (<https://www.nrbcs.org/gfx/genedec/>). The conserved motifs of *SlCarE054* were analysed by CD-search in NCBI (<https://www.ncbi.nlm.nih.gov/cdd/?term=>).

Determination the overexpression of CarEs genes and spatiotemporal expression of *SlCarE054*

The expression of CarEs genes in QJ, KX, and GX population was validated by quantitative RT-PCR (qRT-PCR). RNA samples from the third instar larvae of QJ, KX, and GX populations were extracted and the relative expression level of *SlCarE054* was quantified using SuperReal PreMix Plus (SYBR Green, Tiangen) on ABI QuantStudio 6 PCR System (Applied Biosystems by Life Technologies, Foster, CA, U.S.A.).

To determine the spatiotemporal expression of *SlCarE054*, RNA samples from the first, second, third, fourth, fifth, sixth instar larvae and pupae, tissues including head, cuticle, foregut, midgut, hindgut, malpighian tubules, fat body from the fifth instar larvae of QJ population were extracted and also quantified by qRT-PCR. The reaction system consisted of $2 \times$ SuperReal PreMix solution ($10 \mu\text{l}$), $50 \times$ ROX reference dye ($0.4 \mu\text{l}$), each of an F and R primer ($10 \mu\text{mol l}^{-1}$, $0.6 \mu\text{l}$, Table S2), cDNA ($1 \mu\text{l}$), and ddH $_2\text{O}$ ($7.4 \mu\text{l}$). The thermal cycling was conducted according to the protocols. *EF-1 α* and *RPL10* were used as house-keeping genes (Lu *et al.*, 2013). The reaction of each gene was repeated with 3 independent mRNA samples. The relative expression of *SlCarE054* in each sample was quantified by $2^{-\Delta\Delta\text{Ct}}$ method.

The heterologous expression of *SlCarE054* in *E. coli*

The coding sequence of *SlCarE054* (deleting the signal peptide region) was digested by *Bam*H I and *Xho* I (Takara, Dalian, China), and then subcloned into the expression vector pET-30a (+). The recombinant plasmid pET-30a(+)/*SlCarE054* was transformed into *E. coli* BL21 (DE3) and induced by isopropyl β -D-thiogalactoside (IPTG, 1.0 mmol l^{-1}) at 18°C , 160 rpm for 48 h. The mixture was centrifuged at 4°C , 6000 rpm for 10 min and the cells were transferred on ice to lyse by ultrasonic (Sonic and Materials, Inc., U.S.A.). The crude enzyme was obtained from the supernatant after centrifuging at 4°C , 16,000 rpm for 30 min and further purified by ProteinIso[®] Ni-NTA Resin (TransGen Biotech, Beijing, China) with 100 and 200

mmol l⁻¹ imidazole, desalted by Zeba™ Spin Desalting Columns (Thermo, Shanghai, China).

The extracts of empty vector and recombinant protein SiCarE054 were checked by SDS-PAGE. The enzymatic activity of recombinant protein SiCarE054 was detected using α -naphthyl acetate (α -NA) as the model substrate. The stock solution of α -NA was dissolved in acetone. A series concentration of α -NA working solution (12.5, 25, 50, 100, 200 μ M) was diluted by Fast Blue RR salt. The purified recombinant enzyme (33 μ l, 0.5 mg ml⁻¹) was added with PBS (297 μ l), α -NA (330 μ l) and mixed well immediately. The mixture was transferred to 96-well microplates, and the absorption at 450 nm was determined every 1 min at 30°C for 5 min in a time-driver model by microplate reader (Biotek, U.S.A.). K_m and V_{max} values of recombinant protein SiCarE054 were determined according to the Michaelis–Menten equation with SigmaPlot 12.0 (Systat Software, San Jose, CA, U.S.A.).

Insecticides metabolism by recombinant protein SiCarE054 and the identification of metabolites

Purified recombinant protein SiCarE054 (50 μ l, 0.5 mg ml⁻¹) was added with the mixture containing PBS (400 μ l, 0.1 mol l⁻¹, pH 7.0) and insecticides (450 μ l, 50 mg l⁻¹) diluted with PBS. Boiled purified recombinant protein SiCarE054 was used as control. The reaction was incubated at 30°C for 2 h and stopped by adding acetonitrile (900 μ l). The residual insecticides were extracted by adding NaCl to saturation, shaking at 160 rpm for 2 h, vortexing for 5 min, and centrifuging at 12,000 rpm for 8 min. The supernatant was filtered by 0.22 μ m membrane and quantified by UPLC (Waters Acquity UPLC system, U.S.A.) using the detection method in Li et al. (2021b). Briefly, acetonitrile/water (80:20, v/v) was used as mobile phase with flow rate at 0.4 ml min⁻¹. The injection volume was 3 μ l. The residual quantity of fenvalerate, β -cypermethrin, cyhalothrin, phoxim, and chlorpyrifos was detected using a PDA detector at 220, 220, 230, 280, and 290 nm, respectively.

The metabolites were identified using GC/MS (Agilent 7890B/5977A, Santa Clara, CA, U.S.A.) equipped with a capillary column (HP-5MS, 0.25 μ m \times 250 μ m \times 30 m) and mass database of NIST14.L. The injection volume was 1 μ l in splitless mode using carrier gas of helium (\geq 99.999%) with flow rate at 1 ml min⁻¹. The temperature of inlet was 290°C. The initial oven

temperature was set at 60°C for 1 min, with a subsequent temperature gradient of 20°C per minute until a final temperature of 290°C. The transfer line temperature was 290°C. The conditions of mass spectrometry were as follows: the ion source was in the EI mode, 70 eV, 230°C. The temperature of quadrupole was 150°C, and the mass scanning range was 30–500 MHZ.

Homology modelling and molecular docking

The 3D protein structure of SiCarE054 was constructed online using Swiss model server (<https://swissmodel.expasy.org/>, Bienert et al., 2017; Waterhouse et al., 2018). The carboxylic ester hydrolase from *S. frugiperda* (PDB ID: A0A2H1V6U9.1.A) with 84.95% sequence identity was selected as template. The 3D structure of fenvalerate, β -cypermethrin, cyhalothrin, phoxim, and chlorpyrifos were sketched and obtained by ChemDraw 20.0 (PerkinElmer, U.S.A.) and Chem3D (PerkinElmer, U.S.A.). Autodock 4.2 (Autodock, San Diego, CA, U.S.A., Morris et al., 2009) was used to perform the molecular docking with the default parameters. The possible interactions and the binding free energy between ligand and protein were predicted by Autodock 4.2. The interaction between ligand and protein was further predicted using PLIP (Protein-Ligand Interaction Profiler, <https://plip-tool.biotec.tu-dresden.de/plip-web/plip/index>). The results of modelling and docking were visualised by PyMOL (New York, NY, U.S.A.).

Statistical analysis

The expression of CarEs genes in QJ, KX, and GX populations, and the *in vitro* metabolism activity of SiCarE054 were evaluated by Students' *t*-test. The relative expression of SiCarE054 in different development stage and tissues were analysed by One-way analysis of variation (ANOVA) followed by Tukey's HSD multiple comparisons test. $P < 0.05$ was regarded to be statistically significant.

Results

Kx population showed high resistance level to pyrethroids and organophosphates

As shown in table 1, the LD₅₀ values of KX population to fenvalerate, cyhalothrin, β -cypermethrin, chlorpyrifos, and phoxim

Table 1. The resistance ratio of *S. litura* from KX and GX populations to pyrethroids and organophosphates.

Insecticide	Population	Regression equation	LD ₅₀ (95% FL) (μ g/larva)	χ^2	Freedom degree	<i>P</i> values	RR ^a
Fenvalerate	GX	$Y = 1.92 + 0.90X$	0.007 (0.004–0.012)	2.72	4	0.61	1
	KX	$Y = -8.93 + 1.77X$	108.200 (82.433–165.551)	1.37	4	0.71	15,457.14
Cyhalothrin	GX	$Y = 3.07 + 0.98X$	0.001 (0.000–0.001)	1.54	4	0.82	1
	KX	$Y = -8.39 + 1.72X$	73.000(59.519–93.804)	0.71	3	0.95	73,000.00
β -Cypermethrin	GX	$Y = 3.48 + 1.29X$	0.002 (0.001–0.003)	1.54	4	0.82	1
	KX	$Y = -5.95 + 1.69X$	3.295(2.418–4.189)	1.91	3	0.75	1647.50
Chlorpyrifos	GX	$Y = 4.61 + 3.42X$	0.045 (0.039–0.052)	4.33	4	0.36	1
	KX	$Y = -7.52 + 1.87X$	10.34 (7.717–13.585)	10.26	4	0.32	229.78
Phoxim	GX	$Y = 2.22 + 1.46X$	0.031 (0.023–0.042)	4.95	4	0.29	1
	KX	$Y = -5.93 + 1.70X$	3.090(1.939–4.208)	2.18	4	0.54	99.68

^aRR = LD₅₀ value of KX population/LD₅₀ value of GX population.

were 108.200, 73.000, 3.295, 10.340, and 3.090 $\mu\text{g larva}^{-1}$, respectively. Compared with GX population, the resistance ratios of KX population to fenvalerate, cyhalothrin, β -cypermethrin, chlorpyrifos, and phoxim were 15,457.14-, 73,000.00-, 1657.50-, 229.78-, and 99.68- fold, respectively.

The sequence information of *SlCarE054* and its bioinformatics

The cDNA full length of *SlCarE054* was 1677 bp, encoding 558 amino acids (Table S3). The protein molecular weight and theoretical pI of *SlCarE054* were predicted to be 61.04 kDa and 5.45, respectively. According to the SignalP-6.0, the first 18 amino acids at the N-termini of *SlCarE054* were signal peptides (fig. S1). There are three predicted N-glycosylation sites: Asn116-Leu117-Ser118-Val119, Asn256-Phe257-Thr258-Asp259 and Asn459-Leu460-Thr461-His462. The conserved domain analysis identified the catalytic triad (Ser205, Glu334, and His447), the substrate binding pocket (Gly125, Gly126, Gly127, Tyr204, Ser205, Ala206, Ala209, Phe356, Asp360, Leu361, Phe398, Cys448, Ile452), and the pentapeptide motifs (Gly203-X-Ser205-X-Gly206, fig. 1).

Validation of the overexpression of *CarEs* genes and the spatiotemporal expression of *SlCarE054* in QJ population

A total of 11 significantly up-regulated *CarEs* genes in QJ population were selected from the RNA-Seq result. The expression of five *CarEs* genes (*SlCarE054*, *SlCarE056*, *SlCarE004*, *SlCarE020*, *SlCarE077*) were both overexpressed in QJ and KX populations validated by qRT-PCR (fig. 2). *SlCarE054* showed the highest relative expression level in QJ population (7.60 ± 1.25) and was thus selected for further study.

The relative expression of *SlCarE054* in the larvae and pupae stage of QJ population was determined. The expression of *SlCarE054* showed no significant change during the larvae stage, while increased significantly in the pupae, which were 2.8- fold than that in the first instar larvae (fig. 3A). Among the tissues in the fifth instar larvae, *SlCarE054* showed the lowest expression in the head, cuticle, and fat body. Its expression in the midgut and foregut were significantly higher than that in head, which were 46.3- and 21.3-fold, respectively (fig. 3B). The expression of *SlCarE054* in the hindgut and malpighian tubules were significantly higher than that in head or fat body, and showed no significant difference with that in cuticle.



Figure 1. The alignment of *SlCarE054* with Ha001A (AMO44416.1), Ha001 G (AMO44418.1), Ha001H (AMO44417.1) in *Helicoverpa armigera*, PxaE8 (XP_048485366.1), PxCCE016b (AIN76405.1) in *Plutella xylostera*llra, and SlCarE074 (XP_022837633.1) in *S. litura*. The signal peptide sequences of *SlCarE054* were indicated in underline with blue line, the catalytic triad residues were marked with triangles, the highly conserved pentapeptide residues are marked with green horizontal rectangle, the potential N-glycosylation was underlined with green lines, the substrate binding sites are labelled with red underlines.




Figure 2. The relative expression level of CarEs genes in QJ, KX, and GX populations validated by qRT-PCR. “*” indicates significant difference compared with GX population analysed by Students’ *t*-test ($P < 0.05$).

Identification of recombinant protein SICarE054 and its enzymatic activity to α -NA

The extracts of recombinant protein were identified by SDS-PAGE. The bands of extracts induced by IPTG and purified recombinant protein were shown nearly to 66.2 kDa in the lanes (fig. 4A). This result was well coincided with the predicted molecular weight of 61.04 kDa. The quantity of recombinant protein purified by imidazole at 100 mmol l^{-1} is more abundant than that of 200 mmol l^{-1} . Thus, imidazole at 100 mmol l^{-1} was used for recombinant protein purification.

The esterase activity of purified recombinant protein SICarE054 to α -NA was determined. The V_{max} and K_m values of SICarE054 were calculated to be $38.1 \pm 6.1 \text{ OD min}^{-1} \text{ mg}^{-1}$ and $59.5 \pm 8.3 \mu\text{M}$, respectively (fig. 4B).

The metabolic activity of SICarE054 to pyrethroids and organophosphates

The metabolic activity of SICarE054 to pyrethroids and organophosphates was determined using the purified recombinant protein by UPLC following incubation with insecticides for 2 h. Results showed that the insecticides residue area of the determined pyrethroids incubated with purified recombinant protein SICarE054 decreased significantly than that incubated with boiled SICarE054. As a chiral insecticide, β -cypermethrin was resolved into α -cypermethrin and θ -cypermethrin under the detection condition. The depletion rate of SICarE054 to θ -cypermethrin was 33.8%, the highest among all the pyrethroids, while 7.3% to α -cypermethrin, obviously lower than the enantiomer θ -cypermethrin. The depletion rate of SICarE054 to cyhalothrin




Figure 3. The spatiotemporal expression of SICarE054 in QJ population. (A) The expression of SICarE054 in the developmental stages of larvae. (B) The relative expression level of SICarE054 in different tissues of the fifth instar larvae. Different lowercase letters above the bars indicated significant difference analysed by ANOVA followed by Tukey’s HSD test ($P < 0.05$). Developmental stages: the first instar larvae (L1), the second instar larvae (L2), the third instar larvae (L3), the fourth instar larvae (L4), the fifth instar larvae (L5), the sixth instar larvae (L6), pupae (PU). Tissues: head (H), cuticle (C), foregut (FG), midgut (MG), hindgut (HG), malpighian tubules (MT), fat body (FB).




Figure 4. Characterisation of the recombinant protein SlCarE054 by SDS-PAGE (A) and the kinetic parameters of recombinant protein SlCarE054 (B). Marker: protein marker; pET30a(+), extract of vector only; IPTG-, extract of the recombinant protein without IPTG induction; IPTG +, extract of the recombinant protein induced by IPTG; 100 mM, the purified recombinant protein induced by IPTG eluted by 100 mM imidazole; 200 mM, the purified recombinant protein induced by IPTG eluted by 200 mM imidazole. K_m and V_{max} values of SlCarE054 were calculated using α -NA as substrate with Michaelis-Menten plot.

and fenvalerate were 6.4 and 5.2%, respectively (fig. 5). After incubated with SlCarE054, the insecticides residue area of phoxim or chlorpyrifos remain unchanged.

The metabolites of β -cypermethrin after incubation with SlCarE054 were determined by GC/MS. According to the peak, the possible metabolites and degradation pathways were deduced. The ester bond of β -cypermethrin was hydrolysed and the corresponding alcohol was generated. The alcohol was further metabolised to 3-phenoxybenzaldehyde (fig. S7).

Predication on the interaction of SlCarE054 with pyrethroids and organophosphates by molecular docking

The 3D structure of SlCarE054 included 14 α -helixes, 13 β -sheets and connected by several loops. An obvious loop at the N-terminal indicated the signal peptide (fig. 6). β -cypermethrin could bind to the residues of catalytic triad Ser205 and His447 of SlCarE054, and form hydrogen bonds with them. Another hydrogen bond was predicted to bind with residue Gly126 with 3.4 Å.

Hydrophobic interactions were predicted to bind with the residues Leu460, Tyr204, Phe285, Tyr341, Met337, His447 with distance 3.0–4.5 Å. Molecular docking results showed that cyhalothrin and fenvalerate bind to the sites near to the catalytic triad of SlCarE054. The interaction to fenvalerate included a hydrogen bond with residue Leu460, a π -stacking with residue His447, and hydrophobic interactions with residues Thr458, Phe472, Asp449, Met337, His447, Ile452 with distance 3.2–3.7 Å. The interaction with cyhalothrin included a hydrogen bond with residue Asn446, and hydrophobic interactions with residues Thr434, Pro435, Phe472, Asp449, Leu336, Leu339 with distance 3.3–4.0 Å. These residues were mostly near to the catalytic triad, His447 and Glu334.

The residues of SlCarE054 interacted with phoxim or chlorpyrifos were clearly different with those of pyrethroids. The interaction with chlorpyrifos included a hydrogen bond with Met135 and a π -stacking with Trp132. For phoxim, there are two hydrogen bonds with Trp132 and Leu460, a π -stacking with Trp132, hydrophobic interactions with Asp456, Leu460, and Pro463. These residues were far away from the catalytic triad.

The free binding energy of SlCarE054 to β -cypermethrin, fenvalerate, cyhalothrin, phoxim, and chlorpyrifos were predicted to be -7.19 , -6.94 , -6.78 , -6.27 , and -5.18 Kcal mol⁻¹, respectively.




Figure 5. The metabolic activity of recombinant protein SlCarE054 to organophosphates and pyrethroids. “*” indicates significant difference analysed by Students’ t-test ($P < 0.05$). The number above bars indicated the depletion rate of insecticides metabolised by SlCarE054.

Discussion

In the genome of *S. litura*, a total of 110 CarEs genes have been identified. CarEs genes from lepidoptera and α classes showed large expansions. Most CarEs genes from lepidoptera and α classes could be induced by xanthotoxin and imidacloprid. Silencing of two CarEs genes from lepidoptera class by siRNA injection, larvae treated by imidacloprid showed increased mortality compared with control, indicating CarEs genes expansion contributed to the increased detoxification (Cheng *et al.*, 2017). Meanwhile, five (*SlituCOE009*, *SlituCOE090*, *SlituCOE050*, *SlituCOE093*, and *SlituCOE074*) CarEs genes from α -esterases, lepidopteran esterase and integument esterase in the indoxacarb-resistant strains were validated to make contribution to indoxacarb resistance by transcriptional up-regulation (Shi *et al.*, 2022). In this study, five out of eleven selected CarEs genes was found both overexpressed in two field collected *S. litura*




Figure 6. The interactions of SICarE054 with pyrethroids and organophosphates. β -cypermethrin, cyhalothrin, fenvalerate, chlorpyrifos, and phoxim were indicated by yellow, green, fuchsia, blue, and peridot, respectively. The blue lines indicate the hydrogen bond; the gray lines indicated the hydrophobic interaction; the green lines indicate the π -stacking.

populations (QJ and KX). The relative expression of *SICarE054*, *SICarE056*, *SICarE077* in QJ and KX populations were agree with their resistance level to determined pyrethroids and organophosphates. *SICarE054*, derived from lepidoptera class, showed the highest expression in QJ population among the selected CarEs genes. Its amino acid sequence included a signal peptide with 18 amino acids at the N terminal, which indicated that it was a secreted extracellular protein. The predicted N glycosylation site might enhance protein stability and catalytic efficiency.

SICarE054 was expressed in all the determined development stages. The expression of *SICarE054* was significantly higher in the digestive tissue midgut. This result was similar with the CarEs genes of *Pxae14* (Li et al., 2022), *Pxae18*, and *Pxae28* in *Plutella xylosterlla* (Xie et al., 2017), and *CarE001H* in *H. armigera* (Li et al., 2020), which all showed metabolic activity to insecticides, suggesting a possible role of *SICarE054* in insecticide detoxification. While its expression in pupae is significantly higher than that in larvae, which is different from most of the above CarEs genes, similar with that of *Pxae14* (Li et al., 2022). This result suggested that *SICarE054* might also play roles in insect physiology during the pupae stage.

CarEs was reported to metabolise pyrethroids and organophosphates by hydrolysing in insects (Tang et al., 2020). In *Musca domestica*, the recombinant protein MdaE17, MdaE7, MdbE2, and MdIntE showed significant metabolic activity to permethrin, but no metabolic activity to its metabolite, 3-phenoxybenzylalcohol or 3-phenoxybenzaldehyde, indicating that these CarEs genes only participated in the first step of permethrin degradation (Feng et al., 2018; Feng and Liu, 2020). *BoaE1* in *Bradysia odoriphaga* possesses hydrolase activity towards malathion and participates in the detoxification activity of malathion (Tang et al., 2020). CarEs

was also reported to participate in organophosphates resistance by sequestering the insecticide away from its target site (tight binding/low turnover, Wheelock et al., 2005). In *Culex quinquefasciatus*, *Cqestb2* confers metabolic resistance by sequestering the organophosphates rather than proceeding via catalytic hydrolysis (Hopkins et al., 2017). *LcaE-7* in *Lucilia cuprina* was reported to sequester the insecticide molecule and then detoxify it slowly, thus mediating a wide range of insecticides resistance (Yan et al., 2009; Jackson et al., 2013). The metabolic activity of *SICarE054* to pyrethroids and organophosphates was also determined. Only pyrethroids could be metabolised by *SICarE054*, while not the tested organophosphates. *SICarE054* showed the highest metabolic activity to β -cypermethrin, which was similar with the result of *CarE001A* and *CarE001H* in *H. armigera* (Li et al., 2020). The metabolic activity of *SICarE054* to fenvalerate and cyhalothrin were relatively low. This might be explained as that *SICarE054* contribute more to the resistance to β -cypermethrin than that of fenvalerate or cyhalothrin. Also, the metabolic ability of *SICarE054* to β -cypermethrin showed stereoselectivity. The depletion rate of *SICarE054* to θ -cypermethrin is much higher than that of α -cypermethrin. This is similar with the report of stereoselectivity difference of CarEs E4 in *Myzus persicae*. The purified CarEs E4 could only hydrolyse the 1S-trans-permethrin rapidly, but not the other three isomers (Devonshire and Moores, 1982). In *M. domestica*, docking analysis also found that the isomer of permethrin, 1S-trans-permethrin, fit most snugly within the binding cavities of four CarEs, MdaE7, MdaE17, Md β E2, and MdIntE7 with the lowest binding energy (Feng and Liu, 2020). The metabolite 3-phenoxybenzaldehyde of β -cypermethrin was identified by GC/MS after incubated with *SICarE054* or boiled *SICarE054*. The abundant of 3-phenoxybenzaldehyde after

incubated with S1CarE054 was much higher than that of boiled S1CarE054, and the abundant of parent compound β -cypermethrin was lower after incubated with S1CarE054 than that of boiled S1CarE054 (fig. S7). Thus, we deduced that S1CarE054 could metabolite β -cypermethrin directly by hydrolysing the ester bond, which was similar with the metabolic pathway of pyrethroids reported in previous studies (Feng *et al.*, 2018; Feng and Liu, 2020). The residual area of phoxim and chlorpyrifos showed no significant change compared with the boiled S1CarE054, suggesting S1CarE054 could not participate in their resistance by direct metabolism or sequestering. As reported by Li *et al.* (2021a), the overexpression of *PxaE8* in the multi-insecticide resistant *P. xylostella* populations is involved in the resistance to both β -cypermethrin and phoxim. The overexpression of *PxaE14* in *P. xylostella* participated in the multiple resistance to β -cypermethrin, bifenthrin, chlorpyrifos, fenvalerate, malathion, and phoxim. Also, *PxaE14* showed higher metabolic activity to pyrethroids than that of organophosphates, which indicated that *PxaE14* may prefer to metabolise carboxyl esters in pyrethroids than phosphate esters in organophosphates (Li *et al.*, 2022). This preference might be related with the different structure of proteins.

The molecular docking was further conducted to explain the possible reason for the different metabolic activity to pyrethroids and organophosphates. β -cypermethrin could interact with the residues of conserved catalytic triad (Ser205 and His447) of S1CarE054 and the oxyanion hole (Gly126) by forming hydrogen bonds. Fenvalerate could form π -stacking with residue of catalytic triad His447, and hydrogen bond with Leu460. Cyhalothrin could only form hydrogen bond with Asn446. The binding of pyrethroids with S1CarE054 were also surrounded with several hydrophobic residues around the conserved catalytic triad. These results indicated that pyrethroids could bind to or close to the catalytic triad or substrate binding site, which is beneficial to the occurrence of the first step of the catalytic reaction. The binding free energy of S1CarE054 with pyrethroids follows the order of β -cypermethrin < fenvalerate < cyhalothrin, which could explain the difference in metabolic activity of S1CarE054 to these pyrethroids. The binding free energy of S1CarE054 with organophosphates were much higher than that of pyrethroids, and the interaction sites were away from that of catalytic triad or substrate binding sites.

In conclusion, our results suggested that the overexpression of S1CarE054 in pyrethroids and organophosphates resistant populations could contribute to the resistance to β -cypermethrin by direct metabolism. S1CarE054 showed stereoselectivity to the metabolism of β -cypermethrin.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0007485324000282>.

Acknowledgements. This work is supported by the Key Scientific and Technological Research Project of Henan Province (222102110044), the Postdoctoral Research Grant in Henan Province (202103118), Young Talent Lifting Project of Henan Province (2023HYTP003), the Henan Provincial Science and Technology Major Project (221100110100), the special fund project for central guiding Henan province local development (Z20221343034).

Competing interest. None.

References

Ahmad M, Arif MI and Ahmad M (2007) Occurrence of insecticide resistance in field populations of *Spodoptera litura* (Lepidoptera: Noctuidae) in Pakistan. *Crop Protection* 26, 809–817.

- Ahmad M, Sayyed AH, Saleem MA and Ahmad M (2008) Evidence for field evolved resistance to newer insecticides in *Spodoptera litura* (Lepidoptera: Noctuidae) from Pakistan. *Crop Protection* 27, 1367–1372.
- Armes NJ, Wightman JA, Jadhav DR and Rao R (1997) Status of insecticide resistance in *Spodoptera litura* in Andhra Pradesh, India. *Pesticide Science* 50, 240–248.
- Bass C, Puinean AM, Zimmer CT, Denholm I, Field LM, Foster SP, Gutbrod O, Nauen R, Slater R and Williamson MS (2014) The evolution of insecticide resistance in the peach potato aphid, *Myzus persicae*. *Insect Biochemistry and Molecular Biology* 51, 41–51.
- Bienert S, Waterhouse A, de Beer TAP, Tauriello G, Studer G, Bordoli L and Schwede T (2017) The SWISS-MODEL repository - new features and functionality. *Nucleic Acids Research* 45, D313–D319.
- Cheng TC, Wu J, Wu Y, Chilukuri RV, Huang L, Yamamoto K, Feng L, Li WS, Chen ZW, Guo HZ, Liu JQ, Li SL, Wang XX, Peng L, Liu DL, Guo YB, Fu BH, Li ZQ, Liu C, Chen YH, Tomar A, Hilliou F, Montagné N, Jacquin-Joly E, Alençon E, Seth RK, Bhatnagar RK, Jouraku A, Shiotsuki T, Kadono-Okuda K, Promboon A, Smaghe G, Arunkumar KP, Kishino H, Goldsmith MR, Feng QL, Xia QU and Mita K (2017) Genomic adaptation to polyphagy and insecticides in a major East Asian noctuid pest. *Nature Ecology Evolution* 1, 1747–1756.
- Devonshire AL and Moores GDA (1982) Carboxylesterase with broad substrate specificity causes organophosphorus, carbamate and pyrethroid resistance in peach-potato aphids (*Myzus persicae*). *Pesticide Biochemistry and Physiology* 18, 235–246.
- Feng X and Liu NN (2020) Functional analyses of house fly carboxylesterases involved in insecticide resistance. *Frontiers in Physiology* 11, 595009.
- Feng X, Li M and Liu NN (2018) Carboxylesterase genes in pyrethroid resistant house flies, *Musca domestica*. *Insect Biochemistry and Molecular Biology* 92, 30–39.
- Hopkins DH, Fraser NJ, Mabbitt PD, Carr PD, Oakeshott JG and Jackson CJ (2017) Structure of an insecticide sequestering carboxylesterase from the disease vector *Culex quinquefasciatus*: what makes an enzyme a good insecticide sponge? *Biochemistry* 56, 5512–5525.
- Huang SJ and Han ZJ (2007) Mechanisms for multiple resistances in field populations of common cutworm, *Spodoptera litura* (Fabricius) in China. *Pesticide Biochemistry and Physiology* 87, 14–22.
- Jackson CJ, Liu JW, Carr PD, Younus F, Coppin C, Meirelles T, Lethierb M, Pandey G, Ollisa DL, Russellc RJ, Weikb M and Oakeshottc JG (2013) Structure and function of an insect alpha-carboxylesterase (alpha Esterase7) associated with insecticide resistance. *Proceedings of the National Academy of Science of the United States of America* 110, 10177–10182.
- Li XC, Schuler MA and Berenbaum MR (2007) Molecular mechanisms of metabolic resistance to synthetic and natural xenobiotics. *Annual Review of Entomology* 52, 231–253.
- Li YQ, Bai LS, Zhao CX, Xu JJ, Sun ZJ, Dong YL, Li, DX, Liu XL and Ma ZQ (2020) Functional characterization of two carboxylesterase genes involved in pyrethroid detoxification in *Helicoverpa armigera*. *Journal of Agricultural and Food Chemistry* 68, 3390–3402.
- Li R, Zhu B, Shan JQ, Li LH, Liang P and Gao XW (2021a) Functional analysis of a carboxylesterase gene involved in beta-cypermethrin and phoxim resistance in *Plutella xylostella* (L. *Pest Management Science* 77, 2097–2105.
- Li DZ, He CS, Xie LF, Kong FB, Wu YB, Shi MW, Liu RQ and Xu L (2021b) Functional analysis of SIGSTE12 in pyrethroid and organophosphate resistance in *Spodoptera litura*. *Journal of Agricultural and Food Chemistry* 69, 5840–5848.
- Li R, Zhu B, Hu XP, Shi XY, Qi LL, Liang P and Gao XW (2022) Overexpression of *PxaE14* contributing to detoxification of multiple insecticides in *Plutella xylostella* (L.). *Journal of Agricultural and Food Chemistry* 70, 5794–5804.
- Li J, Lv Y, Liu Y, Bi R, Pan Y and Shang Q (2023) Inducible gut-specific carboxylesterase S1COE030 in polyphagous pests of *Spodoptera litura* conferring tolerance between nicotine and cyantraniliprole. *Journal of Agricultural and Food Chemistry* 71, 4281–4291.
- Lu YH, Yuan M, Gao XW, Kang TH, Zhan S, Wan H and Li JH (2013) Identification and validation of reference genes for gene expression analysis

- using quantitative PCR in *Spodoptera litura* (Lepidoptera: Noctuidae). *PLoS ONE* **8**, e68059.
- Morris GM, Huey R, Lindstrom W, Sanner MF, Belew RK, Goodsell DS and Olson AJ** (2009) AutoDock4 and AutoDockTools4: automated docking with selective receptor flexibility. *Journal of Computational Chemistry* **30**, 2785–2791.
- Oakeshott JG, Claudianos C, Campbell PM, Newcomb RD and Russell RJ** (2005) Biochemical genetics and genomics of insect esterases. In Gilbert LI, Iatrou K, Gill SS (eds), *Comprehensive Molecular Insect Science*, vol. 5. Elsevier Ltd: Oxford, pp. 309–361.
- Ranson H, Claudianos C, Orтели F, Abgrall C, Hemingway J, Sharakhova MV, Unger MF, Collins FH and Feyereisen R** (2002) Evolution of multiple gene families associated with insecticide resistance. *Science (New York, N.Y.)* **298**, 179–181.
- Riddiford LM, Hiruma K, Zhou X and Nelson CA** (2003) Insights into the molecular basis of the hormonal control of molting and metamorphosis from *Manduca sexta* and *Drosophila melanogaster*. *Insect Biochemistry and Molecular Biology* **33**, 1327–1338.
- Saleem M, Hussain D, Ghouse G, Abbas M and Fisher SW** (2016) Monitoring of insecticide resistance in *Spodoptera litura* (Lepidoptera: Noctuidae) from four districts of Punjab, Pakistan to conventional and new chemistry insecticides. *Crop Protection* **79**, 177–184.
- Shi Y, Li WL, Zhou YL, Liao XL and Shi L** (2022) Contribution of multiple overexpressed carboxylesterase genes to indoxacarb resistance in *Spodoptera litura*. *Pest Management Science* **78**, 1903–1914.
- Tang BW, Dai W, Qi L J, Du SK and Zhang CN** (2020) Functional characterization of an alpha-esterase gene associated with malathion detoxification in *Bradysia odoriphaga*. *Journal of Agricultural and Food Chemistry* **68**, 6076–6083.
- Taylor P and Radic Z** (1994) The cholinesterases: from genes to proteins. *Annual Review of Pharmacology and Toxicology* **34**, 281–320.
- Vogt RG** (2005) Molecular basis of pheromone detection in insects. In Gilbert LI, Iatrou K and Gill SS (eds), *Comprehensive Insect Physiology, Biochemistry, Pharmacology and Molecular Biology*. Elsevier Ltd: Oxford, vol. 3, pp. 753–804.
- Wan P, Wu K, Huang M, Yu D and Wu J** (2008) Population dynamics of *Spodoptera litura* (Lepidoptera: Noctuidae) on Bt cotton in the Yangtze River Valley of China. *Environmental Entomology* **37**, 1043–1048.
- Wang LL, Huang Y, Lu XP, Jiang XZ, Smagghe G, Feng ZJ, Yuan GR, Wei D and Wang JJ** (2015) Overexpression of two α -esterase genes mediates metabolic resistance to malathion in the oriental fruit fly, *Bactrocera dorsalis* (Hendel). *Insect Molecular Biology* **24**, 467–479.
- Wang XG, Huang Q, Hao Q, Ran S, Wu Y, Cui P, Yang J, Jiang CX and Yang QF** (2018) Insecticide resistance and enhanced cytochrome P450 monooxygenase activity in field populations of *Spodoptera litura* from Sichuan, China. *Crop Protection* **106**, 110–116.
- Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, Heer FT, de Beer TAP, Rempfer C, Bordoli L, Lepore R and Schwede T** (2018) SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic Acids Research* **46**, W296–W303.
- Wheelock CE, Shan G and Ottea J** (2005) Overview of carboxylesterases and their role in the metabolism of insecticides. *Journal of Pesticide Science* **30**, 75–83.
- Xie M, Ren NN, You YC, Chen WJ, Song QS and You MS** (2017) Molecular characterisation of two alpha-esterase genes involving chlorpyrifos detoxification in the diamondback moth, *Plutella xylostella*. *Pest Management Science* **73**, 1204–1212.
- Xu L, Mei Y, Liu RQ, Chen XL, Li DZ and Wang CJ** (2020) Transcriptome analysis of *Spodoptera litura* reveals the molecular mechanism to pyrethroids resistance. *Pesticide Biochemistry and Physiology* **169**, 104649.
- Xu L, Li B, Liu HY, Zhang HW, Liu RQ, Yu H and Li DZ** (2022) CRISPR/Cas9-mediated knockout reveals the involvement of CYP304F1 in β -cypermethrin and chlorpyrifos resistance in *Spodoptera litura*. *Journal of Agricultural and Food Chemistry* **70**, 11192–11200.
- Yan S, Cui F and Qiao C** (2009) Structure, function and applications of carboxylesterases from insects for insecticide resistance. *Protein and Peptide Letters* **16**, 1181–1188.
- Zhang Z, Gao B, Qu C, Gong J, Li W, Luo C and Wang R** (2022) Resistance monitoring for six insecticides in vegetable field-collected populations of *Spodoptera litura* from China. *Horticulturae* **8**, 255.