



p -adic Simpson correspondences for principal bundles in abelian settings

Ben Heuer , Annette Werner, and Mingjia Zhang

Abstract. We explore generalizations of the p -adic Simpson correspondence on smooth proper rigid spaces to principal bundles under rigid group varieties G . For commutative G , we prove that such a correspondence exists if and only if the Lie group logarithm is surjective. Second, we treat the case of general G on ordinary abelian varieties, in which case we prove a generalization of Faltings’ “small” correspondence to general rigid groups. On abeloid varieties, we also prove an analog of the classical Corlette–Simpson correspondence for principal bundles under linear algebraic groups.

1 Introduction

Let K be a complete algebraically closed non-archimedean field over \mathbb{Q}_p . Let X be a connected smooth proper rigid space over K and fix a point $x_0 \in X(K)$. Let $\pi_1^{\text{ét}}(X, x_0)$ be the étale fundamental group of X , i.e., the symmetry group of all connected finite étale covers of X . The starting point of p -adic non-abelian Hodge theory is the question how to relate continuous representations

$$\pi_1^{\text{ét}}(X, x_0) \rightarrow \text{GL}_n(K)$$

to Higgs bundles on X . The guiding question of this article is to what extent one can generalize this theory from GL_n to more general groups. We provide evidence that the correct generality is given by rigid group varieties, the p -adic analogs of complex Lie groups. The question we set out to investigate is the following.

Question 1.1 *For which rigid groups G can we find a reasonable condition “??” on G -Higgs bundles so that there is a “ p -adic Corlette–Simpson correspondence with G -coefficients”*

$$\left\{ \begin{array}{l} \text{continuous homomorphisms} \\ \pi_1^{\text{ét}}(X, x_0) \rightarrow G(K) \end{array} \right\} \overset{\sim}{\rightarrow} \left\{ \begin{array}{l} G\text{-Higgs bundles on } X \\ \text{satisfying ??} \end{array} \right\}?$$

For the definition of G -Higgs bundles in this setting, see Definition 2.18. Based on known results for $G = \text{GL}_n$, one can divide this problem into two steps, using Scholze’s v -site X_v :

Received by the editors February 26, 2024; revised November 20, 2024; accepted December 02, 2024.

This project was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), TRR 326 *Geometry and Arithmetic of Uniformized Structures*, project number 444845124. During the project, the third named author was funded by the Hausdorff Center for Mathematics and Max-Planck Institute for Mathematics.

AMS subject classification: 14G22, 14G45, 14F20.

Keywords: p -adic Simpson correspondence, G -bundles.



Question 1.2 For which rigid groups G can we expect to have an equivalence of categories

$$(1) \quad \{G\text{-torsors on } X_v\} \xrightarrow{\sim} \{G\text{-Higgs bundles on } X\}?$$

In a second step, the relation to the fundamental group is furnished by the profinite-étale universal cover $\tilde{X} \rightarrow X$, a $\pi_1^{\text{ét}}(X, x_0)$ -torsor in X_v . Indeed, forming the pushout of $\tilde{X} \rightarrow X$ along continuous representations of $\pi_1^{\text{ét}}(X, x_0)$ defines a natural functor

$$\left\{ \begin{array}{l} \text{continuous homomorphisms} \\ \pi_1^{\text{ét}}(X, x_0) \rightarrow G(K) \end{array} \right\} \rightarrow \{G\text{-torsors on } X_v\}.$$

Question 1.3 For which rigid groups G is this functor fully faithful?

As many aspects of p -adic non-abelian Hodge theory are still conjectural (even for $G = \text{GL}_n$), understanding specific cases gives important guidance for further investigations. In this article, we achieve progress in several directions with an abelian flavor: First, we answer all three questions for commutative G . Second, when X is an abeloid variety, we answer Question 1.1 for linear algebraic G and Question 1.2 for “small” objects and general rigid G . We also show that Question 1.2 has a positive answer for linear algebraic G on any smooth proper X .

This gives some evidence for what to expect for general G : We can deduce from our results conditions on the logarithm for G which are necessary to understand the Betti side of the desired theory, as well as sufficient conditions that answer Question 1.3 in general. This gives some concrete indications of how to formulate a conjectural p -adic Corlette–Simpson correspondence for principal bundles. We now outline our results in more detail.

1.1 Previous results

Our study is motivated by the following previous results:

1.1.1 The case $G = \mathbb{G}_a$

In the case $G = \mathbb{G}_a$, the answers to these questions are given by the Hodge–Tate short exact sequence of p -adic Hodge theory of Faltings and Scholze [32, Section 3]

$$(2) \quad 0 \rightarrow H_{\text{ét}}^1(X, \mathbb{G}_a) \rightarrow \text{Hom}_{\text{cts}}(\pi_1^{\text{ét}}(X, x_0), \mathbb{G}_a(K)) \rightarrow H^0(X, \Omega_X^1(-1)) \rightarrow 0.$$

It is easy to see that \mathbb{G}_a -Higgs bundles are classified by $H_{\text{ét}}^1(X, \mathbb{G}_a) \times H^0(X, \Omega_X^1(-1))$, so any splitting of this sequence induces an equivalence as in Question 1.1: Here for $G = \mathbb{G}_a$, there is no condition on the Higgs bundle side. Second, we have $\text{Hom}_{\text{cts}}(\pi_1^{\text{ét}}(X, x_0), \mathbb{G}_a(K)) = H_v^1(X, \mathbb{G}_a)$ in this case by the Primitive Comparison Theorem, answering Question 1.2. We therefore think of Question 1.1 as generalizing (2) to more general coefficients G . This explains the name “ p -adic non-abelian Hodge theory”.

1.1.2 The case $G = \text{GL}_n$

For vector bundles, i.e., $G = \text{GL}_n$, an answer to Question 1.2 has recently been given in [18], where it is shown that there is a *p*-adic Simpson correspondence

$$(3) \quad \{\text{vector bundles on } X_v\} \xrightarrow{\sim} \{\text{Higgs bundles on } X\}.$$

Here a Higgs bundle is a pair (E, θ) of an analytic vector bundle E on X and an \mathcal{O}_X -linear homomorphism $\theta : E \rightarrow E \otimes \Omega_X^1(-1)$ such that $\theta \wedge \theta = 0$. Towards Question 1.1, following Faltings’ influential work [10], one can embed continuous homomorphisms of $\pi_1^{\text{ét}}(X, x)$ into the category of “generalized representations”, which in the language of Scholze’s diamonds can be interpreted as the vector bundles on X_v . The equivalence (3) thus leads to a fully faithful functor

$$(4) \quad \left\{ \begin{array}{l} \text{continuous homomorphisms} \\ \pi_1^{\text{ét}}(X, x) \rightarrow \text{GL}_n(K) \end{array} \right\} \hookrightarrow \{\text{Higgs bundles on } X\}.$$

Conjecturally, the essential image should admit a description in terms of a *p*-adic replacement for the condition in the complex theory that the Higgs bundle is semi-stable with vanishing Chern classes, thus yielding a “*p*-adic Corlette–Simpson correspondence”. Known instances include the case $n = 1$ where $G = \mathbb{G}_m$ [19, Theorem 1.1], and the case when X is an abeloid variety [20, Theorem 1.1], where the essential image is the category of pro-finite-étale Higgs bundles: Here, a G -bundle is called pro-finite-étale if it is trivialised by the cover $\tilde{X} \rightarrow X$, and a G -Higgs bundle is pro-finite-étale if its underlying G -bundle is. In general, however, the problem of determining the essential image of (4) currently seems wide open.

One reason why it is fruitful to study (4) by passing to *v*-vector bundles is that both categories in (3) localize, so that it makes sense to analyze their relation for not necessarily proper X . This reinterprets *p*-adic non-abelian Hodge theory as the study of *v*-topological G -torsors on rigid spaces.

1.1.3 The case $G = \text{GL}_n(\mathcal{O}^+)$

An interesting special case is the rigid group $\text{GL}_n(\mathcal{O}^+) \subseteq \text{GL}_n(\mathcal{O})$. In this case, G -torsors on X_v are equivalent to locally finite free \mathcal{O}^+ -modules, which are also called “integral *v*-vector bundles”. Such objects have recently been studied by Min–Wang [28] to investigate integral structures in *p*-adic non-abelian Hodge theory.

1.1.4 General rigid G

In [17, Theorem 1.2], it is shown that a sheafified version of (1) holds for any rigid group G on any smooth rigid space X . Moreover, [17, Theorem 6.5] gives a local version of (1). This produces some evidence that *p*-adic non-abelian Hodge theory for rigid groups is a promising new line of investigation, which was our initial motivation for studying Questions 1.1, 1.2, and 1.3.

1.1.5 Vanishing Higgs field

Beyond $G = \text{GL}_n$, Question 1.1 has previously only been studied in the case of Higgs field $\theta = 0$ for reductive groups G when X is a curve: In this case, Hackstein has constructed a functor from certain G -bundles to continuous representations $\pi_1^{\text{ét}}(X, x) \rightarrow G(K)$ in [13], based on the work of Deninger–Werner [9].

1.2 The case of commutative G

It is clear that an equivalence as in (1) cannot exist for *any* rigid group G : For example, for the open subgroup $\mathbb{G}_a^+ \subseteq \mathbb{G}_a$ given by the open unit disc, it is known that the “integral Hodge–Tate sequence”

$$0 \rightarrow H_{\text{ét}}^1(X, \mathbb{G}_a^+) \rightarrow \text{Hom}_{\text{cts}}(\pi_1^{\text{ét}}(X, x), \mathbb{G}_a^+(K)) \rightarrow H^0(X, \Omega_X^1(-1))$$

is only left-exact, and that describing the image on the right is in general difficult. With this in mind, our first main result is a sufficient and necessary condition for commutative G under which we can answer Question 1.2 and Question 1.3: We call a commutative rigid group G over K *locally p -divisible* if it admits an open subgroup $U \subseteq G$ such that $[p] : U \rightarrow U$ is surjective (see Definition 6.8). For example, G is locally p -divisible whenever $[p] : G \rightarrow G$ is surjective. In particular, any commutative connected algebraic group is locally p -divisible.

Theorem 1.4 (Theorem 7.4) *Let X be a connected smooth proper rigid variety over K . Let G be a commutative rigid group over K that is locally p -divisible.*

- (i) *Choices of an exponential map for K and a flat B_{dR}^+/ξ^2 -lift of X induce an equivalence*

$$\{G\text{-torsors on } X_v\} \xrightarrow{\sim} \{G\text{-Higgs bundles on } X\}.$$

- (ii) *The above functor restricts to an essentially surjective functor*

$$\left\{ \begin{array}{l} \text{continuous homomorphisms} \\ \pi_1^{\text{ét}}(X, x) \rightarrow G(K) \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{pro-finite-étale} \\ G\text{-Higgs bundles on } X \end{array} \right\}.$$

This is an equivalence if moreover G admits an injective morphism into GL_n .

The assumption on G to be locally p -divisible is also necessary. The crucial point is that the morphism $\log_G : \widehat{G} \rightarrow \text{Lie } G$ has to be surjective.

To our surprise, the functor in (ii) fails to be fully faithful in general: There is an unexpected relation to the phenomenon of pro-finite-étale uniformization of abeloids, see Proposition 5.19. This shows that already for commutative G , Question 1.3 can be very subtle.

As an important tool in the proof of Theorem 1.4, we prove a rigidity lemma for the universal cover \widetilde{X} (Lemma 5.7), and we show that any pointed map from a connected smooth proper rigid space to a rigid group G factors through a maximal abeloid subvariety of G (Theorem 5.8).

The choices in Theorem 1.4 (1) are the same as in [18]. The choice of a flat B_{dR}^+/ξ^2 -lift can be thought of as a choice of splitting of (2). There is a canonical such choice if X comes with a model over a local field. The exponential map is a choice of a

continuous group homomorphism $\exp : K \rightarrow 1 + \mathfrak{m}_K$ splitting the *p*-adic logarithm, see Definition 6.1. As we show in Theorem 6.12, this induces compatible splittings of \log_G for all locally *p*-divisible commutative *G*, generalizing Faltings’ result that this holds in the linear algebraic case.

1.3 The Tannakian approach

The obvious problem in answering Question 1.1 is that currently, little is known about the condition “??” even in the case of $G = \mathrm{GL}_n$. However, one case which is understood is when *X* is an abeloid variety: By [20, Theorem 6.1], there is an equivalence of categories between finite-dimensional continuous representations of the Tate module of *X* and pro-finite-étale Higgs bundles on *X*, depending on similar choices as in Theorem 1.4. The first natural strategy to go beyond GL_n in this case is to leverage the Tannakian formalism, following Simpson’s approach in complex non-abelian Hodge theory [39, Section 6]: In fact, Simpson shows that one can deduce from the Corlette–Simpson correspondence for GL_n a generalization for linear algebraic groups *G* by using the Tannakian formalism. We show that a Tannakian argument also works in *p*-adic geometry: For any linear algebraic group *G* over *K*, there is an equivalence of categories

$$\{G\text{-Higgs bundles on } X_{\text{ét}}\} \cong \left\{ \begin{array}{l} \text{exact tensor functors} \\ \mathrm{Rep}_K(G) \rightarrow \mathrm{Higgs}(X) \end{array} \right\}.$$

where $\mathrm{Rep}_K(G)$ is the category of algebraic representations of *G* and $\mathrm{Higgs}(X)$ is the category of Higgs bundles on *X*. We show that in the *p*-adic case, one needs to impose some further conditions on *X* to ensure that the notion of “pro-finite-étale” bundles can be captured by a Tannakian perspective, but these are satisfied for abeloid varieties.

We use this to deduce from [20, Theorem 1.1] the following result.

Theorem 1.5 (Theorem 3.1) *Let *X* be an abeloid variety and let *G* be a linear algebraic group over *K*, considered as a rigid group. Then choices of an exponential map and a flat B_{dR}^+/ξ^2 -lift of *X* induce an equivalence of categories*

$$\left\{ \begin{array}{l} \text{continuous homomorphisms} \\ \pi_1^{\text{ét}}(X, 0) \rightarrow G(K) \end{array} \right\} \cong \left\{ \begin{array}{l} \text{pro-finite-étale} \\ G\text{-Higgs bundles on } X \end{array} \right\}.$$

The same strategy will also work e.g., for curves once the correct condition ?? in Question 1.1 is worked out in this case. Recently, Xu has made progress in this direction [43].

1.4 The *p*-adic Simpson correspondence for small bundles

When translated to the analytic setting, the first step in Faltings’ *p*-adic Simpson correspondence for GL_n on curves [10, Section 2] would be to consider the local case that *X* is an affinoid smooth rigid space with a toric chart. In this situation, he constructs a “local *p*-adic Simpson correspondence”, which, translated to our *v*-topological setup, can be interpreted as an equivalence of categories

$$(5) \quad \{ \text{small } v\text{-vector bundles on } X \} \xrightarrow{\sim} \{ \text{small Higgs bundles on } X \}.$$

Here, on either side, smallness is a technical condition which roughly means that the objects are p -adically close to trivial. This correspondence does not extend beyond “small” objects.

In a “globalization” step, Faltings then chooses a flat lift of a semi-stable model of X to A_{inf}/ξ^2 and reinterprets the local correspondence in terms of this lift. This allows him to glue the correspondences on a small covering under certain conditions.

This “small correspondence” in terms of a lift has since been the subject of extensive studies: At least in the case of semi-stable reduction, it can be regarded as being completely understood following the work of Abbes–Gros and Tsuji [1, 41], and more recently Liu–Zhu [24] and Wang [42] in the rigid analytic setting.

The local correspondence in terms of toric charts admits a generalisation to G -torsors for general rigid groups G [17, Theorem 6.5]. In order to produce evidence that this can be globalized, we also prove a special case of the “small correspondence” for general G .

Theorem 1.6 *Let X be an ordinary abeloid variety over K . Then any choice of anti-canonical p -divisible subgroup $D \subseteq X[p^\infty]$ induces an equivalence of categories*

$$\{\text{small } G\text{-torsors on } X_v\} \xrightarrow{\sim} \{\text{small } G\text{-Higgs bundles on } X\}.$$

For this, we use that the datum of D induces a particularly well-behaved kind of global perfectoid Galois cover $X_\infty \rightarrow X$ which locally plays the same role as Faltings’ toric chart.

This provides evidence that a “small correspondence” exists more generally for any rigid space X and any rigid group variety G . In general, this seems difficult to construct with current methods: For example, in [1, 24, 42], the correspondence is constructed in terms of period sheaves, and we do not know how to adapt these to general G . That being said, when G has good reduction, the approaches of [27, 2] might offer a geometric strategy to construct such a functor, by considering G -bundles on Bhatt–Lurie’s Hodge–Tate stack.

Notations and conventions

- p : a fixed prime number.
- K : a complete algebraically closed non-archimedean field extension of \mathbb{Q}_p . We write \mathcal{O}_K for its ring of integers and \mathfrak{m}_K for the maximal ideal of \mathcal{O}_K .
- $\text{Spa}(K)$: the adic space $\text{Spa}(K, \mathcal{O}_K)$. Similarly, for an adic space $\text{Spa}(R, R^+)$, we suppress the integral subring R^+ from notation if it equals R° , the subring of power bounded elements in R .
- \mathbb{B} : the closed unit disc over K , i.e., $\text{Spa}(K\langle X \rangle, \mathcal{O}_K\langle X \rangle)$. We write \mathbb{D} for the open unit disc $\mathbb{B}(|X| < 1)$ over K .
- $\text{Perf}_{K, \tau}$: the category of perfectoid spaces over $\text{Spd } K$, with the τ -topology, where τ can be ét, proét or v. Let $\text{Perf}_{K, \tau}^{\text{aff}}$ be the subcategory of affinoid perfectoid spaces.
- By a rigid (analytic) space over K , we mean an adic space locally of topologically finite type over $\text{Spa}(K)$. By [22, (1.1.11)], there is an equivalence of categories r from quasi-separated (resp. separated) rigid spaces in the sense of Tate (which we shall call “classical rigid spaces”) to quasi-separated (resp. separated) adic spaces in the

sense of Huber. As separatedness will never be an issue in our considerations, this usage of the terminology of “rigid spaces” will be harmless.

- Let $f : X \rightarrow Y$ be a morphism of classical rigid spaces in the sense of Kiehl. If f is proper in the sense of Kiehl [4, Section 9.6.2], then $r(f) : r(X) \rightarrow r(Y)$ is a proper morphism of adic spaces in the sense of Huber [22, Definition 1.3.2], see [22, Remark 1.3.19(iii)]. Since we wish to use some results from the classical theory, we shall for simplicity define a “proper morphism of rigid spaces” to be one that is locally on the base of the form $r(f)$ for a (Kiehl-)proper morphism of classical rigid spaces. This convention is harmless, because one can show that $r(f)$ is proper if and only if f is proper. This direction is more difficult, and as we do not know a complete reference for it in the literature, we sketch the argument (c.f. [33, footnote 3 on p. 12], [22, Remarks 1.3.18-19]): Without loss of generality, f admits a formal model F . Suppose that $r(f)$ is proper, then the arguments in [21, Bemerkung 3.11.17, Bemerkung 3.11.18, Proposition 3.12.5] do not require the Noetherian assumption and show that F is proper. By [40, Corollaries 4.4 and 4.5], it follows that f is proper. That being said, we will not use the fact that “ $r(f)$ proper implies f proper” in this article, and only mention it to justify our convention.
- Scholze’s diamond functor $(\cdot)^\diamond$ of [36, Section 10.2] attaches to any analytic adic space over K a v -sheaf on $\text{Perf}_{K,v}$, and this assignment is fully faithful on seminormal rigid spaces over $\text{Spa } K$, c.f. [36, Proposition 10.2.3]. We will therefore freely switch back and forth between smooth rigid spaces and their associated diamonds. In particular, we will often drop the diamond symbol from notation for simplicity, unless confusion might arise.
- X_v (X being an adic space over K): the v -site of the diamond X^\diamond , or equivalently the slice category $\text{Perf}_{K,v|X^\diamond}$. We denote by v the natural map $X_v \rightarrow X_{\text{ét}}$.
- All G -torsors are equipped with left G -actions.
- We use Hom_{cts} to denote the set of continuous group homomorphisms and correspondingly we use H_{cts}^* to denote continuous group cohomology sets.
- For a commutative rigid group G over K , we use $G[p^n]$ to denote the sub-rigid group of p^n -torsion elements, and we denote by $G[p^\infty]$ the v -sheaf $\varinjlim_n G[p^n]$.

2 Rigid group theory

We start with structural results on rigid group varieties that we will need throughout. These are of some independent interest. We begin by recalling some technical background.

2.1 Local structure, Lie algebra, and logarithm

Definition 2.1 A rigid group G over K is a group object in the category of adic spaces of locally topologically finite type over $\text{Spa}(K)$.

We refer to [11, Section 1] and [15, Section 3] for some background on rigid group varieties. One relevant fact is that since K has characteristic 0, any rigid group variety is automatically smooth [11, Proposition 1]. For this reason, it is harmless in the following to switch back and forth between the setting of adic spaces over K and that of locally spatial diamonds over K .

Example 2.2 (1) For any algebraic group G_0 over K , the adic analytification G_0^{an} is a rigid group over K . In particular, we have the rigid groups \mathbb{G}_a^{an} , \mathbb{G}_m^{an} , and GL_n^{an} for any n . As we will work exclusively in the analytic category, we will drop the $-^{an}$ from notation throughout when this is clear from context.

(2) For any smooth admissible formal group scheme \mathfrak{G} over \mathcal{O}_K , the adic generic fibre G is a rigid group. We say that G has good reduction if it is of this form. For example, the adic generic fibre of the affine formal additive group is an open subgroup $\mathbb{G}_a^+ \subseteq \mathbb{G}_a$ that can be described as the closed unit disc \mathbb{B} with additive structure.

(3) For any finite dimensional K -vector space V , we have a rigid group $V \otimes \mathbb{G}_a$ defined as the rigid analytification of $\text{Spec}(K[V])$. This defines a fully faithful functor

$$- \otimes_K \mathbb{G}_a : \{\text{finite dimensional } K\text{-vector spaces}\} \rightarrow \{\text{rigid groups over } K\}.$$

We call a rigid group a rigid vector group if it is in the essential image.

(4) A proper connected rigid group is called an abeloid variety. These are the p -adic analogs of complex tori. There is a good structure theory of abeloid varieties due to Lütkebohmert [25], generalising Raynaud’s rigid analytic structure theory of abelian varieties.

Definition 2.3 For any rigid group G over K , we define its Lie algebra to be

$$\text{Lie } G := \ker(G(K[X]/X^2) \rightarrow G(K)).$$

This is a finite dimensional K -vector space which inherits the structure of a Lie algebra of dimension $\dim_K \text{Lie } G = \dim G$. We will denote the associated rigid vector group by

$$\mathfrak{g} := \text{Lie } G \otimes_K \mathbb{G}_a.$$

This represents the v -sheaf on Perf_K defined by $Y \mapsto \text{Lie } G \otimes_K \mathcal{O}(Y)$.

For any $r \in |K|$, let us denote by $\mathbb{B}(r)$ the closed rigid disc of radius r . Then we have the following result about the local structure of rigid groups.

Proposition 2.4 ([15, Corollary 3.8.]) *Let G be any rigid group over K . Then G has a neighbourhood basis of the identity $(G_k)_{k \in \mathbb{N}}$ of open subgroups $G_k \subseteq G$ of good reduction whose underlying inverse system of rigid spaces is isomorphic to $(\mathbb{B}(|p^k|)^d)_{k \in \mathbb{N}}$ for $d = \dim G$.*

We can describe the system G_k more explicitly using the Lie algebra exponential and logarithm. We now summarize the construction, for a more detailed discussion, see [15, Section 3.2]. Let G be a rigid group over K , and let G_0 be any open rigid subgroup of good reduction. Then the Lie algebra of the formal model of G_0 induces an integral subgroup $\mathfrak{g}^+ \subseteq \mathfrak{g}$ of the Lie algebra of G whose underlying rigid space is isomorphic to $\mathbb{B}(1)^d$.

For any $k \in \mathbb{Q}_{>0}$, let $\mathfrak{g}_k := p^k \mathfrak{g}^+$, then by [15, Proposition 3.5] there is $\alpha > 0$ such that for any $k > \alpha$ we have the Lie algebra exponential

$$\exp : \mathfrak{g}_k \rightarrow G$$

of rigid groups. We denote by

$$G_k := \exp(\mathfrak{g}_k)$$

the image of \mathfrak{g}_k , this is an open subgroup of G . Then:

Lemma 2.5 ([15, Lemma 4.20]) $(G_k)_{k \in \mathbb{Q}_{\geq \alpha}^+}$ is a neighbourhood basis of the unit section, consisting of open sub-rigid groups for which \exp has an inverse mapping

$$\log : G_k \xrightarrow{\sim} \mathfrak{g}_k.$$

Recall that a linear algebraic group is an algebraic group that admits a faithful algebraic representation, i.e., a Zariski-closed homomorphism into GL_n for some n . We have the following analog in rigid geometry.

Definition 2.6 We call a rigid group H linear analytic if there is an injective homomorphism of rigid groups $H \rightarrow GL_n$ for some n .

Using Ado’s Theorem, one can show that every rigid group is locally linear analytic.

Proposition 2.7 ([15, Corollary 3.9]) *Let G be any rigid group. Then there is a rigid open subgroup $G_0 \subseteq G$ of good reduction that is linear analytic. More precisely, we can find a homomorphism of rigid groups $G_0 \rightarrow GL_n$ that is a locally closed immersion and that identifies the subgroups $(G_k)_{k > \alpha}$ from Lemma 2.5 with the preimages of $1 + p^k M_n(\mathcal{O}^+)$.*

This sometimes allows us to reduce local statements about G to the case of GL_n .

2.2 The center

As for algebraic groups, there is a good notion of a central subgroup in rigid group theory.

Definition 2.8 For any connected rigid group G over K , the center $Z(G)$ is defined as the kernel of the adjoint morphism (see [15, Section 3.1])

$$\text{ad}: G \rightarrow \text{Aut}(\mathfrak{g})$$

By definition, $Z(G) \subseteq G$ is a Zariski-closed normal subgroup of G .

Lemma 2.9 *Let G be a connected rigid group.*

- (1) $Z(G)$ is commutative and $Z(G)(K)$ is the center of $G(K)$.
- (2) There exists a linear analytic group H for which there is a left-exact sequence

$$1 \rightarrow Z(G) \rightarrow G \rightarrow H.$$

Proof (2) is immediate from the definition, since $\text{Aut}(\mathfrak{g}) \cong GL_n$. For (1), let $g \in G(K)$ and consider the homomorphism

$$c_g : G \rightarrow G, \quad h \mapsto ghg^{-1}.$$

This is the identity if and only if g is in the center of $G(K)$. Since G is connected, c_g is the identity if and only if it is the identity in a neighbourhood of $1 \in G$. By [15, Theorem 3.4], it follows that $c_g = \text{id}_G$ if and only if the induced morphism on tangent spaces $\text{ad}(g) = \text{Lie}(c_g) : \text{Lie } G \rightarrow \text{Lie } G$ is the identity, i.e., if $g \in Z(G)(K)$.

To see that $Z(G)$ is commutative, it suffices to show that the following morphism of rigid spaces is constant:

$$Z(G) \times Z(G) \rightarrow Z(G), \quad g, h \mapsto ghg^{-1}h^{-1}.$$

This we can check on K -points, where it follows from the first part. ■

Remark 2.10 The connectedness assumption is necessary: For example, if G is any étale rigid group, not necessarily commutative, then ad is trivial.

2.3 Rigid groups as v-sheaves of topological groups

Let G be any rigid group. Then its associated v-sheaf inherits the structure of a v-sheaf of groups on Perf_K , and thus more generally on the “big” category of small v-sheaves. In this subsection, we show that one can in fact endow this with the structure of a sheaf of topological groups in a natural way.

Let $G_0 \subseteq G$ be an open subgroup of good reduction with an embedding $G_0 \hookrightarrow \text{GL}_n$ as in Proposition 2.7, inducing a cofinal system of open neighbourhoods $(G_k)_{k>\alpha}$. Without loss of generality, by replacing G_0 with G_k , we can assume that we already have an embedding

$$\rho : G_0 \hookrightarrow \text{GL}_n(\mathcal{O}^+).$$

For any quasi-compact small v-sheaf T , we can define a topology on $G(T)$ as follows: for each $m \in \mathbb{N}$, endow $\text{GL}_n((\mathcal{O}^+/p^m)(T))$ with the discrete topology and

$$\text{GL}_n(\mathcal{O}^+(T)) = \varprojlim_{m \in \mathbb{N}} \text{GL}_n((\mathcal{O}^+/p^m)(T))$$

with the inverse limit topology. The subspace topology on $G_0(T)$ inherited from $\text{GL}_n(\mathcal{O}^+(T))$ extends to a topological group structure on $G(T)$ in a natural way.

Proposition 2.11 *Let T be a quasi-compact small v-sheaf. There is a unique structure of a topological group on $G(T)$ such that $G_0(T)$ is an open subgroup. This topological structure makes G into a v-sheaf of topological groups on $\text{Perf}_{K,v}^{\text{aff}}$.*

We note that this group structure is independent of the choice of ρ . This follows from the fact that ρ is a locally closed immersion. But we will not need this.

Remark 2.12 The quasi-compactness assumption is necessary to get the correct topology: For example, for $G = \mathbb{G}_a$ and $T = \mathbb{G}_a$, the natural topology on $\mathbb{G}_a(\mathbb{G}_a) = \mathcal{O}(\mathbb{G}_a)$ is not the linear one induced by $\mathbb{G}_a^+(\mathbb{G}_a) = \mathbb{G}_a^+(K) = \mathcal{O}_K$, which would be too fine.

The proof of Proposition 2.11 relies on the following Lemma.

Lemma 2.13 *The system of open subgroups $(G_k)_{k \in \mathbb{N}}$ of G satisfies the following: For any $k > \alpha$, any quasi-compact v-sheaf T and any $s \in G(T)$, there exists some $j \in \mathbb{N}$ such that*

$$G_j(T) \subseteq sG_k(T)s^{-1}.$$

Proof The statement is true for $T = \text{Spa}(K)$ since then $c_s : G \rightarrow G, h \mapsto sgs^{-1}$ is a homomorphism of rigid groups. By [15, Lemma 3.10.3], we more precisely know that

c_s sends G_j into G_k if $\text{ad}(s)$ sends \mathfrak{g}_j into \mathfrak{g}_k . This is because $\text{ad}(s)(\mathfrak{g}_j) \subseteq \mathfrak{g}_k$ implies that the following diagram commutes:

$$\begin{array}{ccc} \mathfrak{g}_j & \xrightarrow{\text{ad}(s)} & \mathfrak{g}_k \\ \downarrow \text{exp} & & \downarrow \text{exp} \\ G_j & \xrightarrow{c_s} & G. \end{array}$$

It follows that also for more general affinoid perfectoid T , the inclusion $\text{ad}(s)(\mathfrak{g}_j(T)) \subseteq \mathfrak{g}_k(T)$ implies that $sG_j(T)s^{-1} \subseteq G_k(T)$, since this is a statement that we can check on K -points. The same statement for quasi-compact small v -sheaves follows by covering T by finitely many affinoid perfectoids. The Lemma now follows from the fact that any choice of \mathcal{O}_K -basis of $\mathfrak{g}_0(K) \cong \mathcal{O}_K^d$ induces an isomorphism $\text{Aut}(\mathfrak{g})(T) \cong \text{GL}_d(\mathcal{O}(T))$, and any $\mathcal{O}(T)$ -linear map $f : \mathcal{O}(T)^d \rightarrow \mathcal{O}(T)^d$ is continuous, hence $p^j\mathcal{O}^+(T)^d \subseteq f^{-1}(p^k\mathcal{O}^+(T)^d)$ for some j . ■

Proof of Proposition 2.11 We endow each $G(T)$ with the topology for which a system of open neighbourhoods of the identity is given by $(G_k(T))_{k \in \mathbb{N}}$. To see that this defines the structure of a topological group, we need to see that the preimage W of $g \cdot G_k(T)$ under $m : G(T) \times G(T) \rightarrow G(T)$ is open. For any $(r, s) \in W$, let j be as in Lemma 2.13 for the section s , then

$$rG_j(T) \cdot sG_k(T) \subseteq rsG_k(T)G_k(T) \subseteq gG_k(T).$$

Thus W contains the open neighbourhood $rG_j(T) \times sG_k(T)$ of $(r, s) \in W$.

The continuity of the inverse follows similarly from Lemma 2.13 by

$$(sG_k(T))^{-1} = G_k(T)s^{-1} \supseteq s^{-1}G_j(T).$$

Hence $G(T)$ is a topological group. It is then immediate from the construction that G , regarded as a diamond, is a presheaf of topological groups which is a sheaf on the level of groups. It therefore suffices to see that for any quasi-compact small v -sheaf and any v -cover $\{\tilde{T} \rightarrow T\}$ by quasi-compact small v -sheaves, the short exact sequence of groups

$$0 \rightarrow G(T) \rightarrow G(\tilde{T}) \rightarrow G(\tilde{T} \times_T \tilde{T})$$

is an equalizer in the category of topological groups, i.e., that $G(T)$ carries the subspace topology of $G(\tilde{T})$. This subspace topology is defined by the open subgroup of $G(T)$ obtained by pullback of $G_0(\tilde{T}) \subseteq G(\tilde{T})$. But this is precisely $G_0(T)$ by the sheaf property of G_0 . It thus suffices to verify the statement for $G = G_0$, for which we can reduce to GL_n , where the statement is clear. ■

Lemma 2.14 For any compact Hausdorff space S and any quasi-compact v -sheaf T , we have

$$\text{Map}(\underline{S} \times T, G) = \text{Map}_{\text{cts}}(S, G(T))$$

where $G(T)$ is endowed with the topology from Proposition 2.11.

Proof Evaluation defines a map

$$\text{ev} : \text{Map}(\underline{S} \times T, G) \rightarrow \text{Map}(S, G(T)).$$

For any map $\psi : \underline{S} \times T \rightarrow G$, to see that its image $\phi : S \rightarrow G(T)$ is continuous, it suffices to check this for profinite S since any compact Hausdorff space admits a surjection by a profinite set (automatically a quotient map). Note that the preimage $\phi^{-1}(G_k(T))$ is given by those $s \in S$ such that the restriction of ψ to $\{s\} \times T$ factors through G_k . Assume s is such a point, then the preimage $\psi^{-1}(G_k)$ is an open sub- v -sheaf $U \subseteq \underline{S} \times T$ containing $\{s\} \times T$. We need to see that there is an open neighbourhood $s \in V \subseteq S$ such that $\underline{V} \times T \subseteq U$.

For this we use that $\{s\} \times T = \bigcap_V \underline{V} \times T$, for V running through open and closed neighbourhoods of s in S . Then

$$(\bigcap_V \underline{V} \times T) \cap (\underline{S} \times T \setminus U) = \emptyset.$$

Since $\underline{S} \times T$ is quasi-compact, there is some V as above such that $\underline{V} \times T \subseteq U$. This shows that $\phi^{-1}(G_k(T))$ is open. We deduce the case of $\phi^{-1}(gG_k(T))$ for any $g \in G(T)$ using instead the morphism $\underline{S} \times T \rightarrow G$ defined by $\psi \cdot g^{-1}$. Hence ϕ is continuous.

We now construct an inverse of ev on the set of continuous maps $\phi : S \rightarrow G(T)$. Write $G(T) = \sqcup g_i G_0(T)$ for a set of coset representatives g_i of $G(T)/G_0(T)$, then ϕ is the disjoint union of maps $S_i \rightarrow g_i G_0(T)$ where $S_i := \phi^{-1}(g_i G_0(T))$ is such that $S = \sqcup S_i$. By translation, we can again reduce to the case of maps $S \rightarrow G_0(T)$. Here the result follows from $G_0 \cong \mathbb{B}^d$ by the following Lemma. ■

Lemma 2.15 *The following evaluation map is bijective:*

$$\text{ev} : \text{Map}(\underline{S} \times T, \mathbb{B}) \rightarrow \text{Map}_{\text{cts}}(S, \mathcal{O}^+(T)) = \varprojlim_n \text{Map}_{\text{cts}}(S, \mathcal{O}^+/p^n(T))$$

Proof It is clear from universal properties that the last map is bijective. For the first, it suffices to prove this for affinoid perfectoid T . Using first the universal property of \mathbb{B} in adic spaces, and then the explicit description of fibre products in perfectoid spaces, we have

$$\text{Map}(\underline{S} \times T, \mathbb{B}) = \mathcal{O}^+(\underline{S} \times T) \stackrel{d}{=} \mathcal{O}^+(\underline{S}) \hat{\otimes}_{\mathcal{O}_K} \mathcal{O}^+(T) = \text{Map}_{\text{cts}}(S, \mathcal{O}^+(T)).$$

To see that this almost isomorphism is an honest isomorphism, recall that a function f in $\mathcal{O}(\underline{S} \times T)$ is in $\mathcal{O}^+(\underline{S} \times T)$ if and only if it is ≤ 1 at every point of $\underline{S} \times T$. Since we have $|\underline{S} \times T| = |\underline{S}| \times |T|$, this is equivalent to asking that for every $s \in S$, the specialization $f(s) \in \mathcal{O}(T)$ is in $\mathcal{O}^+(T)$, or in other words that f lies in $\text{Map}_{\text{cts}}(S, \mathcal{O}^+(T))$.

It remains to see that the mappings are inverse to each other. This is because two maps in $\text{Map}(\underline{S} \times T, G)$ agree if and only if they agree at every $s \in S$, or in other words, there is an injection $G(\underline{S} \times T) \rightarrow \prod_{s \in S} G(s \times T)$. Indeed, let S' be the set S endowed with the discrete topology. Then the morphism of adic spaces $h : \underline{S}' \times T \rightarrow \underline{S} \times T$ induces an injection $\mathcal{O}_{\underline{S} \times T} \rightarrow h_* \mathcal{O}_{S' \times T}$. ■

2.4 G-torsors and G-Higgs bundles

Throughout this section, let G be a rigid group over $\text{Spa} K$ as before. Since G is smooth, it is harmless to identify G with its associated locally spatial diamond. We

may therefore also regard G as a (small) sheaf on the v -site of locally spatial diamonds over K .

Definition 2.16

- (1) Let X be any locally spatial diamond, and let $\tau = v$ or $\tau = \acute{e}t$. A G -torsor on X_τ is a sheaf E on X_τ with a left action $m : G \times E \rightarrow E$ of the group G such that locally on X_τ , there is a G -equivariant isomorphism

$$G \rightarrow E$$

where G acts via left-translation on itself. The morphisms of G -bundles are the G -equivariant morphisms of sheaves on X_τ . We also call E a G -bundle on X_τ .

- (2) We denote the category of G -bundles on X_τ by $\text{Bun}_G(X_\tau)$.

There is also a geometric notion of G -bundles, which is equivalent, see Appendix to Lecture 19 in [36]. The set of G -bundles on X_τ up to isomorphism is naturally isomorphic to $H^1_\tau(X, G)$. We refer to [15, Section 3.3, Proposition 3.6] for more background on the notion of G -torsors on diamonds.

In order to go from G -bundles to Higgs bundles, we moreover need a good notion of differentials. We therefore specialize to smooth rigid spaces.

Definition 2.17 For any smooth rigid space X over K , we set $\tilde{\Omega} := \Omega^1_{X|K}(-1)$, where (-1) is a Tate twist, considered as a sheaf on $X_{\acute{e}t}$. The reason to include this Tate twist is that there is then a canonical identification $\tilde{\Omega} = R^1 v_* \mathcal{O}$ for $v : X_v \rightarrow X_{\acute{e}t}$.

Definition 2.18 Let X be a smooth rigid space and G a rigid group over $\text{Spa } K$. A G -Higgs bundle on X is a pair (E, θ) consisting of

- a G -bundle E on $X_{\acute{e}t}$, and
- an element $\theta \in H^0(X, \tilde{\Omega} \otimes_{\mathcal{O}} \text{ad}(E))$ such that $\theta \wedge \theta = 0$,

where $\text{ad}(E) := \mathfrak{g} \times^G E$ is the adjoint bundle of E , with G acting on its Lie algebra \mathfrak{g} via the adjoint representation. The condition $\theta \wedge \theta = 0$ can be understood as saying that in terms of any local choice of basis of $\tilde{\Omega}$, the coefficients of θ commute with each other with respect to the Lie bracket on $\text{ad}(E)$ induced by \mathfrak{g} .

A morphism of G -Higgs bundles $(E_1, \theta_1) \rightarrow (E_2, \theta_2)$ is a morphism of the underlying G -torsors $\varphi : E_1 \rightarrow E_2$ such that the induced map $\tilde{\Omega} \otimes \text{ad}(E_1) \rightarrow \tilde{\Omega} \otimes \text{ad}(E_2)$ sends θ_1 to θ_2 .

Remark 2.19

- (1) Note that since any morphism between G -bundles is an isomorphism, it follows that any morphism of G -Higgs bundles is an isomorphism.
- (2) In the case of $G = \text{GL}_n$, the category of GL_n -Higgs bundles has the same objects as the category of Higgs bundles of rank n , but the morphisms are given only by the isomorphisms of Higgs bundles.
- (3) For the reader who would like more motivation for the definition of G -Higgs bundles, we refer to the proof of Proposition 3.5.

If the rigid group is commutative, the notion of G -Higgs bundles simplifies substantially, because the underlying torsor and the Higgs field can be disentangled.

Lemma 2.20 *Let G be a commutative rigid group. Let X be any smooth rigid space. Then:*

- (1) *A G -Higgs bundle on X is a pair (E, θ) consisting of a G -torsor E on $X_{\acute{e}t}$ and a section $\theta \in H^0(X, \mathfrak{g} \otimes \widetilde{\Omega})$.*
- (2) *The set of morphisms between two G -Higgs bundles (E_1, θ_1) and (E_2, θ_2) is empty unless $\theta_1 = \theta_2$, when it is the set of morphisms between the G -torsors E_1 and E_2 .*
- (3) *For any G -Higgs bundle (E, θ) , we have $\underline{\text{Aut}}(E, \theta) = G$.*

Proof (1) Since G is commutative, the adjoint representation $G \rightarrow \text{GL}(\mathfrak{g})$ is trivial, hence

$$\mathfrak{g} \times^G E = G \backslash (\mathfrak{g} \times E) = \mathfrak{g} \times (G \backslash E) = \mathfrak{g}.$$

- (2) This follows from part (1) since the induced morphism $\mathfrak{g} \times^G E_1 \rightarrow \mathfrak{g} \times^G E_2$ is identified with the identity on \mathfrak{g} .
- (3) This follows from (2) which implies that $\underline{\text{Aut}}(E, \theta) = \underline{\text{Aut}}(E) = G$. Here the last identification uses again that G is commutative. ■

For the rest of this subsection, let us assume that G is commutative. Then for any locally spatial diamond X and $\tau = \acute{e}t$ or v , the set $H^1_\tau(X, G)$ of G -torsors on X_τ up to isomorphism has a natural group structure. This comes from the following functorial construction.

Definition 2.21 Let E_1 and E_2 be G -bundles on X_τ . Following [37, Proposition 7], we consider the product $E_1 \times E_2$. This carries the natural structure of a $G \times G$ -torsor. We let

$$E_1 \otimes^G E_2 := G \times^{G \times G} (E_1 \times E_2)$$

be the pushout of $E_1 \times E_2$ along the multiplication map $m : G \times G \rightarrow G$. This defines a natural symmetric tensor bifunctor,

$$- \otimes^G - : \text{Bun}_G(X_\tau) \times \text{Bun}_G(X_\tau) \rightarrow \text{Bun}_G(X_\tau),$$

natural in G and in X . For any G -torsor E let E^{-1} be the pushout of E along the inverse map $-^{-1} : G \rightarrow G$, then there is a natural isomorphism $E \otimes^G E^{-1} = G$.

2.5 The pro-finite-étale universal cover

The reason why we have set up the formalism of G -torsors not only for smooth rigid spaces but for more general locally spatial diamonds is that we naturally encounter examples of the latter when dealing with Question 1.1. Namely, we will need the p -adic analog of the complex universal cover which we now recall from [16, Definition 4.6].

Definition 2.22 (1) Let X be a connected smooth proper rigid space over K and fix a base point $x_0 \in X(K)$. The pro-finite-étale universal cover of X is the diamond

$$\widetilde{X} := \varprojlim_{X' \rightarrow X} X'$$

where the projective limit is indexed over the category of connected finite étale covers $X' \rightarrow X$ together with a lift $x' \in X'(K)$ of x_0 . Then $\tilde{X} \rightarrow X$ is a pro-étale torsor under $\pi_1^{\text{ét}}(X, x_0)$, the étale fundamental group of X . It comes with a canonical lift of x_0 that we also denote by x_0 .

- (2) We call a G -bundle E on $X_{\text{ét}}$ or X_v pro-finite-étale if the pullback of E along $\tilde{X} \rightarrow X$ is a trivial G -bundle. We call a G -Higgs bundle (E, θ) pro-finite-étale if the underlying G -bundle E is pro-finite-étale.

Example 2.23 If $X = A$ is an abeloid variety, then $\tilde{A} = \varprojlim_{[N], N \in \mathbb{N}} A$ is represented by a perfectoid space [3, Theorem 1]. The statement about torsors means in this case that in the category of diamonds, we have $A = \tilde{A}/TA$, where $TA = \pi_1^{\text{ét}}(A, 0)$ is the adelic Tate module.

Example 2.24 By [19, Theorem 1.1], an analytic line bundle on an abeloid variety A is pro-finite-étale if and only if it is translation invariant and the associated point $x \in A^v(K)$ in the dual abeloid variety A^v is topologically torsion. This means that there is $N \in \mathbb{N}$ such that $x^{Np^n} \rightarrow 1$ for $n \rightarrow \infty$ for the natural non-archimedean topology on $A^v(K)$.

3 The Tannakian approach for linear algebraic groups

In complex geometry, the non-abelian Hodge correspondence of Corlette and Simpson can be extended from GL_n to any reductive group. In fact, for any linear algebraic group G , the equivalence of categories between G -representations of the topological fundamental group and semi-stable G -Higgs bundles with vanishing Chern classes can be deduced from the statement for GL_n by the Tannakian formalism [39, Section 6].

The goal of this section is to show that the Tannakian formalism can also be leveraged to deduce various instances of the p -adic non-abelian Hodge correspondence for linear algebraic G from the (in part still conjectural) case of GL_n . In particular, we obtain a p -adic Corlette–Simpson correspondence for G whenever the correspondence for GL_n is known. We note that this approach only stands a chance to work for linear algebraic groups, and does not generalize to more general algebraic or rigid analytic groups (for example, any homomorphism from an abelian variety to GL_n is clearly trivial).

Our main result in this section is the following partial answer to Question 1.1 for abeloid varieties, which we will deduce using the Tannakian formalism from the case of GL_n treated in [20]

Theorem 3.1 (Corlette–Simpson correspondence for G -bundles) *Let X be an abeloid variety. Let G be the analytification of a linear algebraic group G^{alg} over K . Then choices of a flat B_{dR}^+/ξ^2 -lift \mathbb{X} of X and of an exponential (see Definition 6.1) induce an equivalence of categories*

$$\left\{ \begin{array}{l} \text{continuous homomorphisms} \\ \pi_1^{\text{ét}}(X, 0) \rightarrow G(K) \end{array} \right\} \cong \left\{ \begin{array}{l} \text{pro-finite-étale} \\ G\text{-Higgs bundles on } X \end{array} \right\},$$

where $G(K)$ is equipped with the topology induced from that on K .

For the proof, we begin by setting up the Tannakian formalism in this context.

- Definition 3.2** (1) For any rigid group G which is the analytification of a linear algebraic group G^{alg} , we denote by $\text{Rep}_K(G)$ the tensor category of finite dimensional algebraic representations of G^{alg} .
- (2) For any topological group π , let $\text{Rep}_K(\pi)$ be the tensor category of continuous representations of π on finite dimensional K -vector spaces.

Proposition 3.3 *Let X be any sousperfectoid space in the sense of [36, Section 6.3.1], or any rigid space. Then for any G as in Definition 3.2.(1), there is an equivalence of categories*

$$\{G\text{-bundles on } X_{\text{ét}}\} \xrightarrow{\sim} \{ \text{exact tensor functors } \text{Rep}_K(G) \rightarrow \text{Bun}(X_{\text{ét}}) \}$$

natural in X , where $\text{Bun}(X_{\text{ét}})$ is the tensor category of vector bundles on $X_{\text{ét}}$.

Proof In the sousperfectoid case, this is [36, Theorem 19.5.2]. In fact, the proof in *loc. cit.* only uses the sousperfectoid assumption to guarantee sheafness of the structure sheaf on the étale site, hence it still works in the Noetherian setting of rigid spaces. ■

Proposition 3.4 *Let π be any topological group. Then there is an equivalence of categories*

$$\left\{ \begin{array}{l} \text{continuous homomorphisms} \\ \pi \rightarrow G(K) \end{array} \right\} \cong \left\{ \begin{array}{l} \text{exact tensor functors} \\ \text{Rep}_K(G) \rightarrow \text{Rep}_K(\pi) \end{array} \right\}$$

Here on the left hand side, the set of morphisms $\rho_1 \rightarrow \rho_2$ between any two continuous representations is given by the set of $g \in G(K)$ such that $g\rho_1g^{-1} = \rho_2$.

Proof Given a continuous homomorphism $f : \pi \rightarrow G(K)$, we obtain an exact tensor functor on the right hand side by sending any representation $(V, \rho : G^{\text{alg}} \rightarrow \text{GL}(V))$ to

$$(V, \pi \xrightarrow{f} G(K) \xrightarrow{\rho^{\text{an}}(K)} \text{GL}(V)(K))$$

where $G(K)$ and $\text{GL}(V)(K)$ are endowed with their topologies from Proposition 2.11.

Conversely, denote by $K\text{-Vect}$ the category of finite dimensional topological K -vector spaces. Then for any exact tensor functor η in the set on the right hand side, its composition with the forgetful functor to $K\text{-Vect}$ gives an exact tensor functor

$$F_G : \text{Rep}_K(G) \xrightarrow{\eta} \text{Rep}_K(\pi) \xrightarrow{\text{For}} K\text{-Vect} \cong \text{Bun}(\text{Spa}(K)_{\text{ét}}).$$

By Proposition 3.3, this amounts to a G -bundle E on $\text{Spa}(K)$.

Since E is necessarily trivial, any element of π gives an automorphism of this tensor functor and hence an automorphism of E .

This defines a map of sets

$$f : \pi \rightarrow \text{Aut}_G(E).$$

Since the group structure on π is compatible with compositions of such automorphisms, f is in fact a group homomorphism. Since E is trivial, we have $\text{Aut}_G(E) \cong G$, but there is no canonical such isomorphism. Instead, we can regard $G' := \text{Aut}_G(E)$ as

an inner form of G . But note that we have a natural equivalence of groupoids

$$\left\{ \text{homomorphisms } \pi \rightarrow G(K) \right\} \xrightarrow{\sim} \left\{ \begin{array}{l} \text{homomorphisms } \pi \rightarrow G'(K) \\ \text{into inner forms } G' \text{ of } G \end{array} \right\}$$

as the natural functor is clearly fully faithful and essentially surjective. All in all, we have thus defined functors

$$\begin{aligned} & \left\{ \begin{array}{l} \text{cts. homomorphisms} \\ \pi \rightarrow G(K) \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{exact tensor functors from} \\ \text{Rep}_K(G) \text{ to } \text{Rep}_K(\pi) \end{array} \right\} \\ & \xrightarrow{F_G} \left\{ \text{homomorphisms } \pi \rightarrow G(K) \right\}. \end{aligned}$$

The construction is clearly natural in G . We can therefore now choose any closed embedding $\rho : G^{\text{alg}} \hookrightarrow \text{GL}_n$ to compare F_G with F_{GL_n} . For GL_n , it is clear that the first arrow is an equivalence, with quasi-inverse given by evaluating at the identity $\text{GL}_n \rightarrow \text{GL}_n$. Thus the essential image of F_{GL_n} is given by the continuous homomorphisms. Since ρ is a closed immersion, its K -points $\rho(K) : G(K) \hookrightarrow \text{GL}_n(K)$ exhibit $G(K)$ as a closed subspace of $\text{GL}_n(K)$, and it follows that a homomorphism $\pi \rightarrow G(K)$ is continuous if and only if the composition $\pi \rightarrow G(K) \hookrightarrow \text{GL}_n(K)$ is. By naturality in G , this shows that the essential image of F_G is given by the continuous homomorphisms.

That the other composition of the above functors is also equivalent to the identity easily follows from the observation that η can be reconstructed from the data of F_G and f . ■

We have the following rigid analytic analog of the discussion in [39, Section 6, p. 68]:

Proposition 3.5 *Let X be a smooth rigid space. Then there is an equivalence of categories*

$$\left\{ G\text{-Higgs bundles on } X_{\text{ét}} \right\} \cong \left\{ \begin{array}{l} \text{exact tensor functors} \\ \text{Rep}_K(G) \rightarrow \text{Higgs}(X) \end{array} \right\}.$$

where $\text{Higgs}(X)$ is the tensor category of Higgs bundle on X .

Proof There is clearly a functor from left to right defined by pushout of any given G -Higgs bundle along representations $G \rightarrow \text{GL}_n$. To construct an inverse, since both sides satisfy étale descent, we may localize and assume that $X = \text{Spa}(R)$ is affinoid and Ω_X^1 is finite free, with a chosen basis $\delta_1, \dots, \delta_d$. Let \mathcal{E} be an object on the right hand side and consider the composition

$$\mathcal{F} : \text{Rep}(G) \xrightarrow{\mathcal{E}} \text{Higgs}(X) \rightarrow \text{Bun}(X_{\text{ét}})$$

with the forgetful functor. By Proposition 3.3, this defines a G -bundle F on $X_{\text{ét}}$.

The datum of the Higgs field given by \mathcal{E} amounts to assigning to each δ_i an endomorphism $\theta_{i,V} : \mathcal{F}(V) \rightarrow \mathcal{F}(V)$ for any representation $G \rightarrow \text{GL}(V)$, functorial in V and compatible with tensor products, such that $\theta_{i,V}$ and $\theta_{j,V}$ commute for all i, j and all V . This is the same as the datum of commutative endomorphisms $\theta_i : \mathcal{F} \rightarrow \mathcal{F}$ of the tensor functor \mathcal{F} . We can identify these endomorphisms with automorphisms $1 + \theta_i \varepsilon$ over $R[\varepsilon]/\varepsilon^2$. Using that automorphisms of \mathcal{F} correspond to elements of $G' :=$

$\text{Aut}_G(F)$, we see that such data correspond to elements in the kernel of the reduction map $G'(R[\varepsilon]/\varepsilon^2) \rightarrow G'(R)$, hence to elements of the Lie algebra of $G' = \text{Aut}_G(F)$. One easily checks that this Lie algebra is $\text{Lie}(G') = F \times^G \mathfrak{g} = \text{ad}(F)$. This shows that we may regard $\theta_1, \dots, \theta_d$ as commutative elements of $\tilde{\Omega} \otimes \text{ad}(F)$, which is the same as a G -Higgs field on F . ■

Definition 3.6 We denote by $\text{Higgs}^{\text{profét}}(X)$ the category of pro-finite-étale Higgs bundles on X , and by $\text{Higgs}_G^{\text{profét}}(X)$ the category of pro-finite-étale G -Higgs bundles on $X_{\text{ét}}$.

The following is then the rigid analytic analog of Simpson’s Tannakian description of the category of semi-stable Higgs bundles with vanishing Chern classes.

Lemma 3.7 *Let X be any connected smooth proper rigid space for which \tilde{X} from Definition 2.22 is sousperfectoid, e.g., X could be an abeloid variety. Then there is an equivalence of categories*

$$\text{Higgs}_G^{\text{profét}}(X) \rightarrow \left\{ \begin{array}{l} \text{exact tensor functors} \\ \text{Rep}_K(G) \rightarrow \text{Higgs}^{\text{profét}}(X) \end{array} \right\}.$$

Proof It suffices to prove that the pro-finite-étale assumptions on both sides are identified under the equivalence in Proposition 3.5. It thus suffices to prove the statement for G -Higgs bundles replaced by G -bundles. For this we use Proposition 3.3 for both X and \tilde{X} . By functoriality, this says that the morphism $\tilde{X} \rightarrow X$ induces a commutative diagram:

$$\begin{array}{ccc} \left\{ G\text{-bundles on } X_{\text{ét}} \right\} & \xrightarrow{\sim} & \left\{ \begin{array}{l} \text{exact tensor functors} \\ \text{Rep}_K(G) \rightarrow \text{Bun}(X_{\text{ét}}) \end{array} \right\} \\ \downarrow & & \downarrow \\ \left\{ G\text{-bundles on } \tilde{X}_{\text{ét}} \right\} & \xrightarrow{\sim} & \left\{ \begin{array}{l} \text{exact tensor functors} \\ \text{Rep}_K(G) \rightarrow \text{Bun}(\tilde{X}_{\text{ét}}) \end{array} \right\}. \end{array}$$

Taking essential preimages of the identity objects in the bottom row, we deduce that the top row identifies pro-finite-étale G -bundles with tensor functors into pro-finite-étale objects in $\text{Bun}(X_{\text{ét}})$, as we wanted to see. ■

We now have everything in place to prove Theorem 3.1.

Proof of Theorem 3.1 When $G^{\text{alg}} = \text{GL}_n$ for some $n \in \mathbb{N}$, the theorem holds by [20, Theorem 1.1], which moreover says that we have an exact tensor equivalence

$$\text{Rep}_K(\pi_1^{\text{ét}}(X, 0)) = \text{Higgs}^{\text{profét}}(X).$$

The general case now follows from the combination of Proposition 3.4 and Lemma 3.7. ■

Finally, let us mention that the Tannakian formalism also applies to G -torsors on X_v .

Proposition 3.8 *Let X be any diamond over K , then there is an equivalence of categories*

$$\left\{ G\text{-bundles on } X_v \right\} \cong \left\{ \begin{array}{l} \text{exact tensor functors} \\ \text{Rep}_K(G) \rightarrow \text{Bun}(X_v) \end{array} \right\}.$$

where $\text{Bun}(X_v)$ is the category of v -vector bundles on X .

Proof It is clear that we have a functor from left to right defined by pushout. To see that this is an equivalence, we use that by definition, any diamond admits a *v*-cover $\pi : Y \rightarrow X$ such that $Y \times_X Y$ is again perfectoid.

Let $F : \text{Rep}_K(G) \rightarrow \text{Bun}(X_v)$ be an exact tensor functor. Then

$$\pi^* F : \text{Rep}_K(G) \rightarrow \text{Bun}(X_v) \rightarrow \text{Bun}(Y_v)$$

is still an exact tensor functor. Since Y is perfectoid, we have $\text{Bun}(Y_v) = \text{Bun}(Y_{\text{ét}})$ by [23, Theorem 3.5.8] (see also [36, Lemma 17.1.8]). We can therefore invoke Proposition 3.3 to see that $\pi^* F$ corresponds to a G -bundle \mathcal{E} on Y . By functoriality of Proposition 3.3 and naturality in the sousperfectoid space, the canonical isomorphism over $Y \times_X Y$

$$\psi : p_1^* \pi^* F \xrightarrow{\sim} p_2^* \pi^* F$$

corresponds to an isomorphism $\psi_{\mathcal{E}} : p_1^* \mathcal{E} \xrightarrow{\sim} p_2^* \mathcal{E}$. Similarly, the cocycle condition for ψ ensures the cocycle condition for $\psi_{\mathcal{E}}$ on $Y \times_X Y \times_X Y$. Hence $\psi_{\mathcal{E}}$ defines a descent datum for a G -torsor along the *v*-cover $Y \rightarrow X$. This defines the desired G -bundle on X_v . ■

As an application, this allows us to answer Question 1.2 for linear algebraic groups: Namely, it shows that Faltings’ *p*-adic Simpson correspondence for proper smooth curves [10] generalizes to linear algebraic groups. More generally, the same is true for the *p*-adic Simpson correspondence of smooth proper rigid spaces of [18].

Theorem 3.9 *Let X be a smooth proper rigid space over K . Let G be any linear algebraic group over K . Then choices of a flat B_{dR}^+/ξ^2 -lift \mathbb{X} of X and of an exponential for K induce an exact tensor equivalence*

$$\{G\text{-bundles on } X_v\} \xrightarrow{\sim} \{G\text{-Higgs bundles on } X_{\text{ét}}\}.$$

Proof The category of G -torsors on X_v is equivalent to the category of G -torsors on $X_{\text{proét}}$ by [15, Corollary 1.2]. For $G = \text{GL}_n$, the result therefore holds by [18, Theorem 5.1], which moreover says that in this case, the correspondence is an exact tensor functor. The case of general G follows by the Tannakian formalism using Proposition 3.8 and Proposition 3.5. ■

We have thus answered Question 1.2 for linear algebraic G , and Question 1.1 for linear algebraic G on abeloid varieties.

4 The small correspondence

In this section, the aspect of the *p*-adic Simpson correspondence for which we explore generalizations to general rigid groups G is the “small” correspondence in the “global” setup. That is, we allow X to be any smooth adic space, not necessarily proper or toric.

The starting point of our discussion is the local *p*-adic Simpson correspondence for small G -bundles constructed in [17, Theorem 6.5]. We first explain in what sense this is functorial with respect to perfectoid abelian Galois covers (c.f. Definition 4.14 below). While it is in general impossible to find such perfectoid covers globally, we

explain at the example of abelian varieties how one can use this to construct instances of a small correspondence for G -bundles in the global setting.

In order to explain the setup, we first recall that in Faltings’ setting [10], the local p -adic Simpson correspondence relies on the following notion.

Definition 4.1 Let U be an affinoid smooth rigid space. A toric chart of U is an étale morphism $h : U \rightarrow \mathbb{T}^d$ to some affinoid torus $\mathbb{T}^d = \text{Spa}(K\langle T_1^{\pm 1}, \dots, T_d^{\pm 1} \rangle)$ which is a composition of rational opens and finite étale maps. We call U toric if a toric chart exists.

The setup of our small correspondence can be divided into a local and a global one.

Setup 4.2 (Local Setup) Let U be a toric smooth rigid space. We assume that

$$f : U_\infty \rightarrow U$$

is a pro-étale Galois cover in Scholze’s pro-étale site $U_{\text{proét}}$ of [31, Section 3] satisfying the following:

- (1) U_∞ is affinoid perfectoid.
- (2) The covering group Δ_U of f is isomorphic as a profinite group to \mathbb{Z}_p^d for $d := \dim U$.
- (3) The following natural Cartan–Leray map associated with f is an isomorphism:

$$\text{Hom}_{\text{cts}}(\Delta_U, \mathcal{O}(U)) \rightarrow H_{\text{cts}}^1(\Delta_U, \mathcal{O}(U_\infty)) \rightarrow H_v^1(U, \mathcal{O}).$$

Here the first morphism is induced by $\text{Hom}_{\text{cts}}(\Delta_U, \mathcal{O}(U)) = H_{\text{cts}}^1(\Delta_U, \mathcal{O}(U))$, and the second map is always an isomorphism due to the assumption that U_∞ is affinoid perfectoid.

Example 4.3 Given a toric chart $h : U \rightarrow \mathbb{T}^d$, pullback along h of the affinoid perfectoid torus $\mathbb{T}_\infty^d \rightarrow \mathbb{T}^d$ induces a pro-étale cover $U_\infty \rightarrow U$. This satisfies Setup 4.2 by [31, Lemmas 4.5 and 5.5]. That being said, we note that we only assume in Setup 4.2 that U is toric, but we do not assume that f is related to any chart.

Setup 4.4 (Global Setup) Let X be a smooth rigid space. We assume that

$$f : X_\infty \rightarrow X$$

is a perfectoid pro-étale Galois cover in $X_{\text{proét}}$ with covering group Δ_X for which there is a basis of $X_{\text{ét}}$ given by affinoids U on which $U_\infty := X_\infty \times_X U \rightarrow U$ satisfies Setup 4.2. We call such morphisms f quasi-toric.

Example 4.5 If X is an abeloid variety of dimension d , then the cover $\tilde{X} = \varprojlim_{[p]} X \rightarrow X$ is a perfectoid $T_p X$ -torsor [3, Theorem 1]. While $T_p X \cong \mathbb{Z}_p^{2d}$ does not have the correct rank to satisfy Setup 4.4, we have the following result in case of ordinary reduction, i.e., when the étale part $H^{\text{ét}}$ of the p -divisible group H of the formal Raynaud extension associated with X (the formal group scheme whose generic fibre is the maximal semi-abelian subgroup of X of good reduction) is $\cong (\mathbb{Q}_p/\mathbb{Z}_p)^d$. In this case, we call any subgroup of $X[p^\infty]$ anticanonical if it is isomorphic to its image in the generic fibre of $H^{\text{ét}}$.

Lemma 4.6 Let X be an ordinary abeloid variety. Fix any anticanonical p -divisible subgroup $D \subseteq X[p^\infty]$ of rank d . Its Tate module $T_p D \subseteq T_p X$ is a submodule which corresponds to a $\Delta_X := T_p X/T_p D$ -torsor $X_\infty \rightarrow X$. This satisfies the conditions of Setup 4.4.

Proof We first explain why X_∞ is perfectoid. If X has good reduction, this can be seen as in [3, Lemma 2.13]. Let \mathfrak{X}_n be the formal model of X/D_n . The transition map $\mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1}$ is the quotient by the canonical subgroup, which reduces to the relative Frobenius mod p . Thus Frobenius is surjective on the mod p fibre of $\varprojlim_n \mathfrak{X}_n$, hence X_∞ is perfectoid.

The general case follows from the one of good reduction via Raynaud uniformisation by the same argument as in [3, Proposition 4.3], replacing $[p]$ in [3, diagram (5)] with the canonical isogeny $X/D_n \rightarrow X/D_{n-1}$. This shows that $X_\infty = \varprojlim_n X/D_n$ is perfectoid.

Since X is ordinary, the Hodge–Tate map admits a factorization

$$\begin{array}{ccc} \mathrm{Hom}(T_p X, K) & \xrightarrow{\mathrm{HT}} & H^0(X, \Omega^1(-1)) \\ & \searrow & \nearrow \text{dotted} \\ & \mathrm{Hom}(T_p C, K) & \end{array}$$

where $T_p C \subseteq T_p X$ is the Tate module of the canonical p -divisible subgroup, the map on the left is the restriction, and the dotted map is an isomorphism. Since D is anticanonical, the composition $T_p C \rightarrow T_p X \rightarrow \Delta_X$ is an isomorphism, so it follows that the map

$$\mathrm{Hom}(\Delta_X, K) \rightarrow \mathrm{Hom}(T_p X, K) \rightarrow H^0(X, \Omega^1(-1))$$

is an isomorphism. Let now $U \subseteq X$ be any affinoid subspace such that $U_\infty = X_\infty \times_X U$ is affinoid perfectoid. Using that Ω_X^1 is a trivial vector bundle in the case of abelian varieties, it then follows by applying $- \otimes_K \mathcal{O}(U)$ that also the following map is an isomorphism

$$\mathrm{Hom}(\Delta_X, \mathcal{O}(U)) \rightarrow \mathrm{Hom}(T_p X, \mathcal{O}(U)) \rightarrow H^0(U, \Omega^1(-1)) = H^1_v(U, \mathcal{O}).$$

■

Remark 4.7 With more work, one can in fact show that there exists a cover satisfying Setup 4.4 for any abelian variety X , not necessarily ordinary. After replacing X by an isogenous abelian variety, we can reduce to the case that X is principally polarized. It then follows from [33, Corollary III.3.12] that X is p -power isogeneous to an abelian variety X' that admits a canonical subgroup (see [33, Definition III.2.7] for the notion of canonical subgroup in this case). For any choice of anti-canonical p -divisible subgroup $(D_n \subseteq X'[p^n])_{n \in \mathbb{N}}$, the tower of canonical isogenies $\varprojlim_{n \in \mathbb{N}} X'/D_n$ then has the desired form. But we will not use this fact in the present paper. It is plausible that a cover satisfying Setup 4.4 exists more generally for abeloid varieties.

In order to construct the small p -adic Simpson correspondence in the setup of Setup 4.4, we now first need to deal with the local situation.

Lemma 4.8 *Let $f : U_\infty \rightarrow U$ be as in Setup 4.2. Then for any $n \geq 1$, there is $\gamma > 0$ such that for any $s \in \mathbb{N}$, the kernel and cokernel of the natural map*

$$\iota_s^n : H^n_{\mathrm{cts}}(\Delta_X, \mathcal{O}^+(U)/p^s) \rightarrow H^n_{\mathrm{cts}}(\Delta_X, \mathcal{O}^+(U_\infty)/p^s)$$

are p^γ -torsion. The same is true for $\iota^n : H^n_{\mathrm{cts}}(\Delta_X, \mathcal{O}^+(U)) \rightarrow H^n_{\mathrm{cts}}(\Delta_X, \mathcal{O}^+(U_\infty))$.

Proof The Cartan–Leray sequence for $U_\infty \rightarrow U$ induces a commutative diagram

$$\begin{array}{ccccc}
 H_{\text{cts}}^n(\Delta_X, \mathcal{O}(U)) & \longrightarrow & H_{\text{cts}}^n(\Delta_X, \mathcal{O}(U_\infty)) & \longrightarrow & H_V^n(U, \mathcal{O}) \\
 \uparrow & & \uparrow & & \uparrow \\
 H_{\text{cts}}^n(\Delta_X, \mathcal{O}^+(U)) & \longrightarrow & H_{\text{cts}}^n(\Delta_X, \mathcal{O}^+(U_\infty)) & \longrightarrow & H_V^n(U, \mathcal{O}^+)
 \end{array}$$

The horizontal maps on the right are (almost) isomorphisms because $R\Gamma_V(U_\infty, \mathcal{O}^+) \stackrel{a}{=} \mathcal{O}^+(U_\infty)$ due to the assumption that U_∞ is affinoid perfectoid. For $n = 1$, the composition of the top row is an isomorphism by assumption. It follows that the top map is an isomorphism for general n . This is because Cartan–Leray is compatible with cup products in group cohomology, respectively sheaf cohomology, and the cup products induce isomorphisms

$$H_{\text{cts}}^n(\Delta_X, \mathcal{O}(U)) = \wedge^n H_{\text{cts}}^1(\Delta_X, \mathcal{O}(U)), \quad H_V^n(U, \mathcal{O}) = \wedge^n H_V^1(U, \mathcal{O}),$$

where the last equality follows from [31, Lemma 5.5] using that U is toric.

To see that the bottom composition is an isomorphism up to bounded p -torsion cokernel, we use again that U admits a toric chart. Considering the Cartan–Leray sequence of the associated toric affinoid cover, this shows that there is an injective map

$$\wedge^n \mathcal{O}^+(U)^d \rightarrow H_V^n(U, \mathcal{O}^+)$$

with p^β -torsion cokernel for some $\beta > 0$. This shows that $p^\beta \cdot \iota^n$ factors through a map $j : H_{\text{cts}}^n(\Delta_X, \mathcal{O}^+(U)) \rightarrow \wedge^n \mathcal{O}^+(U)^d$. As $H_{\text{cts}}^n(\Delta_X, \mathcal{O}^+(U)) = \wedge^n \mathcal{O}^+(U)$, the map j is a morphism between finite free $\mathcal{O}^+(U)$ -modules which is an isomorphism after inverting p . Hence it is injective with p^δ -torsion cokernel for some $\delta > 0$. Setting $\gamma := \beta + \delta$, we get the desired result for ι_s^n . The statement for ι_s^n follows from this by long exact sequences. ■

Let now G be any rigid group over K . Following the notation in Section 2.1, we fix an open subgroup G_0 of G of good reduction and let $k > 0$ be such that the Lie algebra exponential $\exp : \mathfrak{g}_k \rightarrow G_k$ converges and is an isomorphism. In this setting, we can now generalise the technical notion of smallness from [17, Section 6].

Definition 4.9 Let $U_\infty \rightarrow U$ be as in Setup 4.2 and let $\gamma > 0$ be as in Lemma 4.8. Set $c := 10 \cdot \max(\gamma, k)$.

- (1) We call a continuous representation $\rho : \Delta_U \rightarrow G(U)$ an f -small cocycle if it factors through $G_c(U)$. We denote the category of f -small 1-cocycles by $\mathcal{Z}_{\text{cts}}^{1, \text{sm}}(\Delta_U, G(U))$, where the morphisms $\rho_1 \rightarrow \rho_2$ are given by $\{g \in G(U) \mid g\rho_1 g^{-1} = \rho_2\}$.
- (2) A G -bundle on U_v is said to be f -small if it admits a reduction of structure group to G_c . We denote the category of f -small G -bundles on U_v by $\text{Bun}_G^{\text{sm}}(U_v)$.
- (3) A G -Higgs bundle (E, θ) on $U_{\text{ét}}$ is said to be f -small if E is trivial and there is a trivialisation $E \cong G$ with respect to which θ is in the image of the chain of maps

$$\begin{aligned} \text{HTlog: Hom}_{\text{cts}}(\Delta_U, G_c(U)) &\xrightarrow{\log} \text{Hom}_{\text{cts}}(\Delta_U, \mathfrak{g}_c(U)) \xrightarrow{\sim} H^1_{\text{cts}}(\Delta_U, \mathfrak{g}_c(U)) \\ &\rightarrow H^1_{\text{cts}}(\Delta_U, \mathfrak{g}_c(U_\infty)) \rightarrow H^1_v(U, \mathfrak{g}_c) \xrightarrow{\text{HT}} R^1v_*\mathfrak{g}_c(U) \xrightarrow{\sim} H^0(U, \mathfrak{g}_c \otimes \tilde{\Omega}). \end{aligned}$$

We denote the category of *f*-small *G*-Higgs bundles on *U* by $\text{Higgs}_G^{\text{sm}}(U)$.

Remark 4.10

- (1) We emphasize that one way in which these notions depend on *f* is that *c* depends on *f*. Note that we have not characterised the γ in Lemma 4.8 uniquely. This is on purpose, as we need some leeway to increase *c* later on. It would be more precise to call the above notions “*f*-*c*-small”, but we shall drop this for simplicity.
- (2) The reason for the factor 10 in the definition of *c* ultimately lies in the appearance of the same factor in [17, Remark 6.8]. The constant 5*c* appears in [17, Proposition 5.5], and an additional factor of 2 is picked up in the globalization step as explained in detail in [17, Remark 6.8]. That being said, we are not aiming for an optimal value of *c*. For most purposes, the important point is simply that *c* only depends on γ .

The three notions of Definition 4.9 are related as follows.

Lemma 4.11

- (1) *Pushout of the Δ_U -torsor $f : U_\infty \rightarrow U$ along any *f*-small 1-cocycle defines a functor*

$$\mathcal{Z}_{\text{cts}}^{1,\text{sm}}(\Delta_U, G(U)) \rightarrow \text{Bun}_G^{\text{sm}}(U_v).$$

- (2) *Sending an *f*-small 1-cocycle ρ to $(G, \text{HTlog}(\rho))$ defines a functor*

$$\mathcal{Z}_{\text{cts}}^{1,\text{sm}}(\Delta_U, G(U)) \rightarrow \text{Higgs}_G^{\text{sm}}(U).$$

Proof Part (1) is clear from the definitions. Note that any morphism between *f*-small 1-cocycles induces an isomorphism between the attached *v*-*G*-bundles. For part (2), we just need to verify that $\theta := \text{HTlog}(\rho)$ is a Higgs field. Since Δ_U is commutative, it follows from [15, Lemma 3.10] that the image of ρ under log in the above sequence lies in $\text{Hom}_{\text{cts}}(\Delta_U, \mathfrak{g}_c(U))^{[-,-]=0}$. Going through the almost isomorphisms, it is clear from this that its image θ in $H^0(U, \mathfrak{g}_c \otimes \tilde{\Omega})$ is a *G*-Higgs field. ■

Proposition 4.12 *Let $U_\infty \rightarrow U$ be as in Setup 4.2. Let *G* be a rigid group with setup as in Definition 4.9. Then the functors of Lemma 4.11 define equivalences of categories*

$$\text{Bun}_G^{\text{sm}}(U_v) \xleftarrow{\sim} \mathcal{Z}_{\text{cts}}^{1,\text{sm}}(\Delta_U, G(U)) \xrightarrow{\sim} \text{Higgs}_G^{\text{sm}}(U).$$

Proof This is part of [17, Theorem 6.5, Remark 6.8]: More precisely, the axiomatic conditions detailed in [17, Remark 6.8] are satisfied in Setup 4.2 by Lemma 4.8. ■

Definition 4.13 We call the composition

$$\text{LS}_{f,G} : \text{Bun}_G^{\text{sm}}(U_v) \xrightarrow{\sim} \text{Higgs}_G^{\text{sm}}(U)$$

the small *p*-adic Simpson correspondence on *U* attached to *f*.

Next, we explain the naturality of LS_f in *f*. For this we need morphisms of Galois covers.

Definition 4.14 Let $f : X_\infty \rightarrow X$ and $g : Y_\infty \rightarrow Y$ be quasi-toric coverings with Galois groups Δ_X and Δ_Y . A *Galois-equivariant map* $\beta : f \Rightarrow g$ is a triple $\beta = (\varphi, u, \tilde{u})$ consisting of a group homomorphism $\varphi : \Delta_X \rightarrow \Delta_Y$ and a commutative diagram

$$\begin{array}{ccc} X_\infty & \xrightarrow{\tilde{u}} & Y_\infty \\ \downarrow f & & \downarrow g \\ X & \xrightarrow{u} & Y, \end{array}$$

such that \tilde{u} is Δ_X -equivariant with respect to the Δ_X -action on Y_∞ via φ .

Proposition 4.15 Let $f : X_\infty \rightarrow X$, $g : Y_\infty \rightarrow Y$, $h : Z_\infty \rightarrow Z$ be quasi-toric covers and let $\beta = (\varphi, u, \tilde{u}) : f \Rightarrow g$ and $\beta' = (\varphi', u', \tilde{u}') : g \Rightarrow h$ be Galois equivariant morphisms. Let G be a rigid group and let $\gamma > 0$ be large enough such that Lemma 4.8 holds for f, g , and h .

- (1) The equivalence $LS_{f,G}$ is natural in f : There is a natural isomorphism of functors $\alpha_\beta : LS_f \circ u^* \Rightarrow u^* \circ LS_g$ making the following diagram 2-commutative:

$$\begin{array}{ccc} \text{Bun}_G^{\text{sm}}(X_v) & \xrightarrow{LS_f} & \text{Higgs}_G^{\text{sm}}(X) \\ u^* \uparrow & \xrightarrow{\alpha_\beta} & u^* \uparrow \\ \text{Bun}_G^{\text{sm}}(Y_v) & \xrightarrow{LS_g} & \text{Higgs}_G^{\text{sm}}(Y). \end{array}$$

- (2) α is compatible with composition: For the Galois-equivariant map $\beta' \circ \beta$, we have

$$\alpha_{\beta' \circ \beta} = u^* \alpha_{\beta'} \circ \alpha_\beta.$$

Proof For (1), we need to produce natural isomorphisms making the diagram

$$\begin{array}{ccccc} LS_f : \text{Bun}_G^{\text{sm}}(X_v) & \longleftarrow & \mathcal{Z}_{\text{cts}}^{1,\text{sm}}(\Delta_X, G(X)) & \longrightarrow & \text{Higgs}_G^{\text{sm}}(X) \\ u^* \uparrow & & u^* \varphi^* \uparrow & & u^* \uparrow \\ LS_g : \text{Bun}_G^{\text{sm}}(Y_v) & \longleftarrow & \mathcal{Z}_{\text{cts}}^{1,\text{sm}}(\Delta_Y, G(Y)) & \longrightarrow & \text{Higgs}_G^{\text{sm}}(Y) \end{array}$$

2-commutative. For the commutativity of the left square, starting from a 1-cocycle $\rho : \Delta_Y \rightarrow G(Y)$, we need to find a natural isomorphism between the two G -bundles on X_v

$$X_\infty \times^{\Delta_X} G \quad \text{and} \quad X \times_Y (Y_\infty \times^{\Delta_Y} G)$$

where to form the second contracted product we push out $g : Y_\infty \rightarrow Y$ using ρ , and for the first, we push out $f : X_\infty \rightarrow X$ along the composition

$$\Delta_X \xrightarrow{\varphi} \Delta_Y \xrightarrow{\rho} G(Y) \xrightarrow{u} G(X).$$

This is the functoriality of the Cartan–Leray construction. Presenting each of these two G -bundles as quotients, we have by equivariance and universal properties a natural morphism

$$(X_\infty \times G)/\Delta_X \rightarrow (X \times_Y Y_\infty \times G)/\Delta_Y \xrightarrow{\sim} X \times_Y (Y_\infty \times G)/\Delta_Y$$

of G -torsors, which is necessarily an isomorphism.

For commutativity of the right square, it suffices to check commutativity of the diagram

$$\begin{CD} \mathrm{Hom}_{\mathrm{cts}}(\Delta_Y, G_k(Y)) @>\mathrm{HTlog}>> H^0(Y, \mathfrak{g}_c \otimes \tilde{\Omega}) \\ @V u^* VV @VV u^* \varphi^* V \\ \mathrm{Hom}_{\mathrm{cts}}(\Delta_X, G_k(X)) @>\mathrm{HTlog}>> H^0(X, \mathfrak{g}_c \otimes \tilde{\Omega}) \end{CD}$$

where HTlog was defined in Definition 4.9. This is immediate from functoriality of log and the functoriality of the Cartan–Leray map applied to \mathfrak{g}_c . This concludes part (1).

For part (2), it is clear that $(\beta' \circ \beta, u' \circ u, \tilde{u}' \circ \tilde{u})$ is a Galois-equivariant map, and it suffices to prove the composition property for the left and right squares in the above diagram. For the right hand side, this is clear. For the left, by construction, the natural transformation $\alpha_{\beta' \circ \beta}$ comes from

$$(X_\infty \times G)/\Delta_X \xrightarrow{\sim} X \times_Z (Z_\infty \times G)/\Delta_Z$$

whereas $\alpha_{\beta'} \circ \alpha_\beta$ comes from the composition

$$(X_\infty \times G)/\Delta_X \xrightarrow{\sim} X \times_Y (Y_\infty \times G)/\Delta_Y \xrightarrow{\sim} X \times_Z (Z_\infty \times G)/\Delta_Z,$$

It is clear from the universal properties that these two morphisms agree. ■

Given this naturality, we can now glue the local small *p*-adic Simpson correspondences.

Definition 4.16 Let X be a quasi-compact smooth rigid space. Let $f : X_\infty \rightarrow X$ be a quasi-toric cover as in Setup 4.4.

- (1) A G -bundle V on X_v is called (étale) locally small with respect to f if there is an étale cover $h : \tilde{X} \rightarrow X$ by an affinoid \tilde{X} as in Setup 4.2 such that $h_2 : \tilde{X} \times_X \tilde{X} \rightarrow X$ and $h_3 : \tilde{X} \times_X \tilde{X} \times_X \tilde{X} \rightarrow X$ are also as in Setup 4.2, and such that $h^* V$ is small with respect to $\tilde{f} : X_\infty \times_X \tilde{X} \rightarrow \tilde{X}$, and similarly $h_2^* V$ and $h_3^* V$ are also small. We denote the category of locally small G -bundles on X_v by $\mathrm{Bun}_G^{\mathrm{lsm}}(X_v)$.
- (2) A G -Higgs bundle (E, θ) on $X_{\mathrm{ét}}$ is called (étale) locally small with respect to f if there is an étale cover $\tilde{X} \rightarrow X$ by an affinoid as in (1), such that $h^*(E, \theta)$ is small with respect to \tilde{f} and $h_2^*(E, \theta)$ and $h_3^*(E, \theta)$ are also small in this sense. We denote the category of locally small G -Higgs bundles on X by $\mathrm{Higgs}_G^{\mathrm{lsm}}(X)$.

Theorem 4.17 Let X be a smooth rigid space over K with a Galois cover f as in Setup 4.4. Then the local equivalences with respect to f glue to an equivalence of categories

$$S_f^{\mathrm{lsm}} : \mathrm{Bun}_G^{\mathrm{lsm}}(X_v) \rightarrow \mathrm{Higgs}_G^{\mathrm{lsm}}(X).$$

In particular, we obtain such a correspondence for ordinary abeloid varieties as in Lemma 4.6.

Proof This is a formal consequence of Proposition 4.12 and Proposition 4.15. As descent arguments in the context of the *p*-adic Simpson correspondences are subtle in general, we now verify this in detail. Let V be a locally small G -bundle on X_v and

let $h : \tilde{X} \rightarrow X$ be a cover as in Definition 4.16.(1). Then by Proposition 4.12 we have an equivalence

$$\Psi := \text{LS}_{\tilde{\mathcal{F}}} : \text{Bun}_G^{\text{sm}}(\tilde{X}_v) \xrightarrow{\sim} \text{Higgs}_G^{\text{sm}}(\tilde{X}).$$

Let \tilde{f}_2 be the pullback $\tilde{X} \times_X \tilde{X} \times_X X_\infty \rightarrow \tilde{X} \times_X \tilde{X}$ and similarly \tilde{f}_3 the pullback to $\tilde{X} \times_X \tilde{X} \times_X \tilde{X}$, then by assumption we also get $\Psi_2 := \text{LS}_{\tilde{\mathcal{F}}_2}$ on $\tilde{X} \times_X \tilde{X}$ and $\Psi_3 := \text{LS}_{\tilde{\mathcal{F}}_3}$.

Consider the Čech nerve of the cover $\tilde{X} \rightarrow X$. We write the low degree projections as

$$\tilde{X} \times_X \tilde{X} \times_X \tilde{X} \xrightarrow{p_{ij}} \tilde{X} \times_X \tilde{X} \xrightarrow{p_k} \tilde{X}$$

and write π_i for the i -th projection $\tilde{X} \times_X \tilde{X} \times_X \tilde{X} \rightarrow \tilde{X}$. Hence $\pi_1 = p_1 \circ p_{12}$ and so on. The maps \tilde{f}, \tilde{f}_2 and \tilde{f}_3 induce Galois equivariant maps of quasi-toric covers over these. We denote the resulting isomorphisms α from Proposition 4.15 induced by p_i, p_{ij}, π_i by the corresponding subscripts, i.e., α_{p_i} etc.

The pullback $\tilde{V} := h^* V$ carries a natural descent datum $\varphi : p_1^* \tilde{V} \xrightarrow{\sim} p_2^* \tilde{V}$, satisfying the cocycle condition $p_{23}^* \varphi \circ p_{12}^* \varphi = p_{13}^* \varphi$. We need to show that the isomorphism

$$\psi : p_1^* \Psi(\tilde{V}) \xrightarrow{\alpha_{p_1}^{-1}} \Psi_2(p_1^* \tilde{V}) \xrightarrow{\Psi_2(\varphi)} \Psi_2(p_2^* \tilde{V}) \xrightarrow{\alpha_{p_2}} p_2^* \Psi(\tilde{V})$$

satisfies the cocycle condition $p_{23}^* \psi \circ p_{12}^* \psi = p_{13}^* \psi$ and hence provides a descent datum for $\Psi(\tilde{V})$ along $\tilde{X} \rightarrow X$. For this we first claim that the following diagram is commutative

$$\begin{array}{ccc} \Psi_3(\pi_1^* \tilde{V}) & \xrightarrow{\Psi_3(p_{12}^* \varphi)} & \Psi_3(\pi_2^* \tilde{V}) \\ \downarrow \alpha_{\pi_1} & & \downarrow \alpha_{\pi_2} \\ \pi_1^* \Psi(\tilde{V}) & \xrightarrow{p_{12}^* \alpha_{p_1}^{-1}} p_{12}^* \Psi_2(p_1^* \tilde{V}) \xrightarrow{p_{12}^* \Psi_2(\varphi)} p_{12}^* \Psi_2(p_2^* \tilde{V}) \xrightarrow{p_{12}^* \alpha_{p_2}} & \pi_2^* \Psi(\tilde{V}) \end{array}$$

and analogously for p_{12} replaced by p_{23} or p_{13} . If this is the case, note that the bottom map is $p_{12}^* \psi$, so this implies that

$$\begin{aligned} p_{23}^* \psi \circ p_{12}^* \psi &= \alpha_{\pi_3} \circ \Psi_3(p_{23}^* \varphi) \circ \alpha_{\pi_2}^{-1} \circ \alpha_{\pi_2} \circ \Psi_3(p_{12}^* \varphi) \circ \alpha_{\pi_1}^{-1} \\ &= \alpha_{\pi_3} \circ \Psi_3(p_{13}^* \varphi) \circ \alpha_{\pi_1}^{-1} = p_{13}^* \psi \end{aligned}$$

as desired. To see the claim, we use that by Proposition 4.15.(2), the relation $\pi_1 = p_1 \circ p_{12}$ implies that we have $\alpha_{\pi_1} = p_{12}^* \alpha_{p_1} \circ \alpha_{p_{12}}$. Hence the above diagram reduces to

$$\begin{array}{ccc} \Psi_3(\pi_1^* \tilde{V}) & \xrightarrow{\Psi_3(p_{12}^* \varphi)} & \Psi_3(\pi_2^* \tilde{V}) \\ \downarrow \alpha_{p_{12}} & & \downarrow \alpha_{p_{12}} \\ p_{12}^* \Psi_2(p_1^* \tilde{V}) & \xrightarrow{p_{12}^* \Psi_2(\varphi)} & p_{12}^* \Psi_2(p_2^* \tilde{V}). \end{array}$$

This commutes by Proposition 4.15.(1) and hence we are done.

The same argument also works in the other direction for Ψ^{-1} . ■

Example 4.18 Let $X_\infty \rightarrow X$ be an anti-canonical cover of an ordinary abeloid variety as in Lemma 4.6. While it is in general difficult to make the implicit constant c in Definition 4.9 and hence it Definition 4.16 more explicit, we can always find interesting examples of locally small G -bundles on X_v , illustrating the power of Theorem 4.17:

- Any étale-locally free G -bundle V on X_v is clearly locally small. Unravelling the construction, we have $S_f^{\text{lsm}}(V) = (V, 0)$. This example is the reason why we use étale localisation in Definition 4.16 rather than analytic localisation.
- The proof of Lemma 4.6 shows that one can always find an affinoid cover $\tilde{X} \rightarrow X$ such that \tilde{X} and the double and triple fibre products \tilde{X}_2 and \tilde{X}_3 satisfy Setup 4.2. Taking $\gamma > 0$ such that Lemma 4.8 holds for \tilde{X} , \tilde{X}_2 and \tilde{X}_3 , we see that for $c \gg 0$, any continuous representation $\rho : \Delta_X \rightarrow G_c(K)$ defines a locally small G -bundle V_ρ on X by pushout of $X_\infty \rightarrow X$ along ρ . Describing the underlying G -bundle of the associated Higgs bundle $S_f^{\text{lsm}}(V_\rho)$ is a non-trivial task already for $G = \mathbb{G}_m$.

5 Pro-finite-étale torsors

In this section, we move on to Questions 1.1 and 1.3. We therefore study G -torsors for the v -topology which become trivial on the pro-finite-étale cover of Section 2.5.

Let X be a connected smooth proper rigid space over K with a fixed point $x_0 \in X(K)$, and denote by $\tilde{X} \rightarrow X$ the diamantine pro-finite-étale universal cover as in Definition 2.22, which is a $\pi := \pi_1^{\text{ét}}(X, x_0)$ -torsor. The main goal of this section is to prove:

Theorem 5.1 *Let X be a connected smooth proper rigid space over K with a fixed base-point $x_0 \in X(K)$. Let G be a rigid group variety. We consider the natural functor of groupoids*

$$\left\{ \begin{array}{l} \text{continuous homomorphisms} \\ \pi_1^{\text{ét}}(X, x_0) \rightarrow G(K) \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{pro-finite-étale} \\ G\text{-torsors on } X_v \end{array} \right\}$$

$$\rho \mapsto \tilde{X} \times_{\pi_1^{\text{ét}}(X, x_0), \rho} G$$

sending a continuous homomorphisms ρ to the pushout of $\tilde{X} \rightarrow X$ along ρ .

- (1) *If G is connected or commutative, then this functor is essentially surjective.*
- (2) *It is fully faithful if there are no non-constant morphisms $\tilde{X} \rightarrow G$. For example this happens if G is linear analytic, i.e., admits an embedding into GL_n for some n .*

Here as in Proposition 3.4, the morphisms $\rho_1 \rightarrow \rho_2$ between two continuous representations on the left hand side are defined as the set of elements $g \in G(K)$ such that $g \cdot \rho_1(\gamma) \cdot g^{-1} = \rho_2(\gamma)$ for all $\gamma \in \pi_1^{\text{ét}}(X, x_0)$. In the case that $G = \text{GL}_n$, this means that the morphisms are the isomorphisms between representations.

Remark 5.2 While we do not know if (1) holds for any G , the fully faithfulness in (2) can fail in general. We refer to Proposition 5.19 below for a concrete example.

Remark 5.3 Recall that $\pi_1^{\text{ét}}(X, x_0)$ is defined in terms of connected finite étale covers of X . In rigid analytic geometry, there is also a second fundamental group, namely the de Jong fundamental group $\pi_1^{\text{dl}}(X, x_0)$, defined in terms of a different kind of étale covers which are not necessarily finite. While one can show that it is still possible to

attach G -torsors on X_v to continuous representations of $\pi_1^{\text{dJ}}(X, x_0)$, this will typically not result in a fully faithful functor, because the analog of the universal cover for $\pi_1^{\text{dJ}}(X, x_0)$ has many global sections.

Proof of Theorem 5.1.(2) It is clear that the displayed functor is well-defined. If there are no non-constant maps $\tilde{X} \rightarrow G$, this means that $G(\tilde{X}) = G(K)$ and hence

$$H_{\text{cts}}^1(\pi, G(\tilde{X})) = H_{\text{cts}}^1(\pi, G(K)) = \text{Hom}_{\text{cts}}(\pi, G(K))$$

as the π -action on $G(K)$ is trivial. In the case G is linear analytic, since $\text{GL}_n(\tilde{X}) \subseteq \text{M}_n(\tilde{X}) \cong \mathcal{O}(\tilde{X})^{n \times n} = K^{n \times n}$, every map $\tilde{X} \rightarrow \text{GL}_n$ is constant. It follows that $G(\tilde{X}) = G(K)$ as well. ■

Theorem 5.1.(1) is more difficult and will take us the rest of this section.

5.1 Cocycles

In order to describe the G -torsors on X that are trivialized by \tilde{X} , we will study the Čech nerve of the π -torsor $\tilde{X} \rightarrow X$. As a first step, we now reduce this to a computation of group cohomology. In doing so, to simplify notation, let us drop the underline in the notation of the diamond $\underline{\pi}$, i.e., we freely move back and forth between profinite sets and their associated diamonds.

Lemma 5.4 *Let G be any rigid group, and let $n \in \mathbb{N}$ if G is commutative, and $n \in \{0, 1\}$ otherwise. Then there is a natural isomorphism*

$$\check{H}^n(\tilde{X} \rightarrow X, G) = H_{\text{cts}}^n(\pi, G(\tilde{X})),$$

where $G(\tilde{X})$ is endowed with its natural topology via Proposition 2.11.

Example 5.5 Let us clarify our conventions for non-abelian cocycles. Since π acts on \tilde{X} from the left, the induced action on $G(\tilde{X})$ is a right-action. For $n = 1$, the 1-cocycles defining $H_{\text{cts}}^1(\pi, G(\tilde{X}))$ are therefore concretely the morphisms

$$\rho : \pi \times \tilde{X} \rightarrow G$$

for which the natural diagram expressing the relation

$$(6) \quad \rho(\gamma_1 \cdot \gamma_2, x) = \rho(\gamma_1, \gamma_2 x) \cdot \rho(\gamma_2, x)$$

commutes. The equivalence relation defining \check{H}^1 is then defined by declaring that two such cocycles ρ_1 and ρ_2 are equivalent if and only if there is a morphism $\varphi \in G(\tilde{X})$ such that

$$(7) \quad \rho_1(\gamma, x) = \varphi(\gamma x) \rho_2(\gamma, x) \varphi(x)^{-1}.$$

Proof Since $\tilde{X} \rightarrow X$ is a π -torsor, the $(n + 1)$ -fold fibre product of \tilde{X} with itself over X is isomorphic to $\pi^n \times \tilde{X}$. Hence cocycles in the Čech cohomology set with G -coefficients $\check{H}^n(\tilde{X} \rightarrow X, G)$ are given by morphisms of diamonds

$$\rho : \pi^n \times \tilde{X} \rightarrow G.$$

Via Lemma 2.14, such morphisms ρ correspond to continuous morphisms $\pi^n \rightarrow G(\tilde{X})$. By a standard computation, the Čech cocycle relation is identified for Galois

covers like $\tilde{X} \rightarrow X$ with the cocycle condition of continuous group cohomology, and the equivalence relations defining the cohomology on both sides agree via this identification.

Let us give some more details of this identification for $n = 1$ in the non-commutative case, for which we do not know a reference in the literature. We use the isomorphisms

$$(8) \quad \begin{aligned} \pi \times \tilde{X} &\xrightarrow{\sim} \tilde{X} \times_X \tilde{X}, & (\gamma, x) &\mapsto (\gamma x, x) \\ \pi \times \pi \times \tilde{X} &\xrightarrow{\sim} \tilde{X} \times_X \tilde{X} \times_X \tilde{X}, & (\gamma_1, \gamma_2, x) &\mapsto (\gamma_1 \gamma_2 x, \gamma_2 x, x). \end{aligned}$$

It is a straightforward calculation that this identifies the projections $p_{ij} : \tilde{X} \times_X \tilde{X} \times_X \tilde{X} \rightarrow \tilde{X} \times_X \tilde{X}$ with the maps $p_{ij} : \pi \times \pi \times \tilde{X} \rightarrow \pi \times \tilde{X}$ defined as follows

$$p_{12}(\gamma_1, \gamma_2, x) = (\gamma_1, \gamma_2 x), \quad p_{23}(\gamma_1, \gamma_2, x) = (\gamma_2, x), \quad p_{13}(\gamma_1, \gamma_2, x) = (\gamma_1 \gamma_2, x).$$

By definition, 1-cocycles in $\check{H}^1(\tilde{X} \rightarrow X, G)$ are now given by sections $\rho \in G(\tilde{X} \times_X \tilde{X}) = G(\pi \times \tilde{X}) = \text{Map}_{\text{cts}}(\pi, G(\tilde{X}))$ satisfying $p_{12}^*(\rho) \cdot p_{23}^*(\rho) = p_{13}^*(\rho)$. Via the above identifications, this translates to

$$\rho(\gamma_1 \cdot \gamma_2, x) = \rho(\gamma_1, \gamma_2 x) \rho(\gamma_2, x)$$

for all $\gamma_1, \gamma_2 \in \pi, x \in \tilde{X}$, which is precisely the same as in Equation (6).

Finally, the equivalence relation defining $\check{H}^1(\tilde{X} \rightarrow X, G)$ is given by saying that $\rho_1 \sim \rho_2$ if there is $\varphi \in G(\tilde{X})$ such that $\rho_1 = p_1^*(\varphi) \rho_2 p_2^*(\varphi)^{-1}$ where $p_1, p_2 : \tilde{X} \times_X \tilde{X} \rightarrow \tilde{X}$ are the projections. When we identify these projections with maps $\pi \times \tilde{X} \rightarrow \tilde{X}$ via (8), this equivalence relation gets identified with (7). ■

Consider now the map $G(K) \rightarrow G(\tilde{X})$. Since the π -action on $G(K)$ is trivial, this induces a natural map

$$\text{Hom}_{\text{cts}}(\pi, G(K)) \rightarrow H^1_{\text{cts}}(\pi, G(\tilde{X}))$$

defined in terms of geometric 1-cocycles by sending

$$\rho \mapsto (\pi \times \tilde{X} \rightarrow G, \quad \gamma, x \mapsto \rho(\gamma)).$$

The key result that we need for the proof of Theorem 5.1.(1) is the following.

Theorem 5.6 *Let G be any connected rigid group, X any connected smooth proper rigid space. Then the following natural map is surjective:*

$$\text{Hom}_{\text{cts}}(\pi, G(K)) \rightarrow H^1_{\text{cts}}(\pi, G(\tilde{X})).$$

We currently do not know if the connectedness assumption on G is necessary. However, we will see in Theorem 5.14 that it can be removed in the commutative case.

For the proof of Theorem 5.6, we first need some preparations. We start by discussing a variant of the classical Rigidity Lemma from algebraic geometry for \tilde{X} .

5.2 A rigidity lemma for universal covers

The classical rigidity lemma in algebraic geometry says that any morphism between K -schemes $f : X \times Y \rightarrow Z$ where X is irreducible and proper with a K -point x_0 , Y is

irreducible with a K -point y_0 and Z is any scheme, and such that $f(X \times \{y_0\})$ is a single point, factors through the projection $X \times Y \rightarrow Y$. The analogous statement also holds in the rigid analytic setting [26, Lemma 7.1.2].

We need an analog of this statement for \tilde{X} . This will be weaker because \tilde{X} is not in any sense “irreducible”. For example, in the case of the Tate curve X , we easily verify from the explicit description of [3, Theorem 4.6.3] that any open neighbourhood of the identity in \tilde{X} contains an open subspace of the form $\mathbb{Z}_p \times \mathbb{B}^{1,\text{perf}}$, which is manifestly not irreducible as it has many connected components.

We therefore need to prove an analog of a slightly weaker version. If the irreducibility assumptions in the algebraic rigidity lemma are dropped, one can still deduce that f factors through the projection on $X \times U$ where $U \subseteq Y$ is an open neighborhood of y_0 .

Lemma 5.7 (Rigidity Lemma) *Let Y be a diamond and let Z be any adic space over K . Assume that there is a K -point $y_0 \in Y(K)$ and let*

$$f : \tilde{X} \times Y \rightarrow Z$$

be any morphism of diamonds such that $f(\tilde{X} \times y_0) = z_0 := f(x_0, y_0)$ is a single point.

Then there is an open neighbourhood $y_0 \in U \subseteq Y$ such that $f|_{\tilde{X} \times U}$ factors through the projection to U composed with a morphism of diamonds $U \rightarrow Z$.

Proof We first observe that the statement is quasi-pro-étale-local on Y (note that we may lift the base-point $y_0 \in Y(K)$ to quasi-pro-étale covers since K is algebraically closed), so we may replace Y by an affinoid perfectoid space.

Let now $z_0 \in V \subseteq Z$ be an affinoid open neighbourhood of $z_0 = f(x_0, y_0)$. Passing to complements, let $W := (\tilde{X} \times Y) \setminus f^{-1}(V)$. Since $\tilde{X} \rightarrow X$ is a proper morphism of diamonds in the sense of [34, Definition 18.1], the morphism $\tilde{X} \rightarrow \text{Spa}(K)$ is proper, thus so is the projection $q : \tilde{X} \times Y \rightarrow Y$. It follows that $q(W) \subseteq Y$ is closed. Let U be the complement. By assumption, we have $y_0 \in U$, and it follows from the definition that f restricts to a map

$$\tilde{X} \times U \rightarrow V.$$

Since V is affinoid, this is determined by its global sections. But by [19, Proposition 3.9], $\mathcal{O}(\tilde{X} \times U) = \mathcal{O}(U)$. Hence this restriction factors through the projection to U . ■

5.3 Morphism from proper rigid spaces to rigid groups

The goal of this section is to prove the following structural result for rigid groups.

Theorem 5.8 *Any rigid group variety G admits a maximal abeloid closed subgroup $A \subseteq G$. This satisfies the following universal property. Any morphism $\varphi : X \rightarrow G$ from an irreducible proper rigid space X with $0 \in \text{im}(\varphi)$ factors through $A \subseteq G$. Moreover, $A \subseteq Z(G^\circ)$, the center of the identity component of G .*

The most important step in the proof is the following Lemma.

Lemma 5.9 *Any morphism $\varphi : X \rightarrow G$ from an irreducible proper rigid space X with $0 \in \text{im}(\varphi)$ factors through some closed abeloid subgroup $A \subseteq G$.*

The proof relies crucially on the notion of “analytic subsets” of a rigid space Y as introduced in [4, Section 9.5], that we now recall. To avoid confusion with the “analytic topology”, we are instead going to call these “Zariski-closed” subspaces, namely $Z \subseteq Y$ is called Zariski-closed if for any affinoid open $U \subseteq Y$, the subspace $Z \cap U \subseteq U$ is the zero-locus of a set of elements $f_1, \dots, f_d \in \mathcal{O}(U)$. Equivalently, Z is the image of a closed immersion defined by a coherent sheaf of ideals in \mathcal{O}_Y [4, Section 9.5 Corollary 7]. Given a Zariski-closed subspace $Z \subseteq Y$, it is always possible to endow it with the induced-reduced structure by replacing any coherent ideal defining Z with its radical [4, Section 9.5.3, Proposition 4].

The notion of Zariski-closed subspaces has good localisation properties, i.e., whether a subspace is Zariski-closed can be checked analytic-locally, and the pullback of a Zariski-closed subset under a morphism $X \rightarrow Y$ of rigid spaces is again Zariski-closed [4, Section 9.6.3, before Lemma 4]. Moreover, we have the following result due to Kiehl.

Proposition 5.10 ([4, Section 9.6.3, Proposition 3]) *Let $f : X \rightarrow Y$ be a proper morphism of rigid spaces. Then the image $f(M)$ of any Zariski-closed subset $M \subseteq X$ is again Zariski-closed.*

Proposition 5.11 *Let X, Y be rigid spaces over K . Assume that X is proper and Y is separated over K , and let $f : X \rightarrow Y$ be a morphism of rigid spaces. Then f is proper and $f(X)$ endowed with its induced reduced structure is a proper rigid space.*

Proof Since X is quasi-compact, we can without loss of generality assume that Y is quasi-compact. Then by [5, Corollary 5.10(a)], f has a formal model $\mathfrak{f} : \mathfrak{X} \rightarrow \mathfrak{Y}$. Since Y is separated, \mathfrak{Y} is separated (see e.g., [22, Remark 1.3.18(i) and Remark 1.3.19(ii)]). We now use that by [40, Corollaries 4.4 and 4.5], the following are equivalent for a morphism g of formal schemes:

- (1) g is proper,
- (2) the rigid generic fibre of g is proper,
- (3) the special fibre g_0 of g over the residue field k is proper.

This first shows that \mathfrak{X} is proper. Arguing mod p^n for every n , it now follows from [8, Tag 01W6] that \mathfrak{f} is proper, hence in a second step we deduce that f is proper.

To see that $f(X)$ is proper, we may replace Y with $f(X)$ and assume that f is surjective. We can now use the functorial reduction map $\text{red} : Y(K) \rightarrow \mathfrak{Y}(k)$ induced by \mathfrak{Y} , which is surjective by [4, Section 7.1.5 Theorem 4], to see that surjectivity of f implies surjectivity of the special fibre $\mathfrak{f}_0 : \mathfrak{X}_0 \rightarrow \mathfrak{Y}_0$. Then by [8, Tag 03GN], \mathfrak{Y}_0 is proper, hence Y is proper. ■

As studied in [6], there is a good notion of irreducible subspaces of rigid spaces. A non-empty subspace $Z \subseteq X$ of a rigid spaces is called irreducible if it cannot be written as the union $Z = V_1 \cup V_2$ of two Zariski closed subspaces $V_1, V_2 \subsetneq Z$, see [6, Lemma 2.2.3]. One checks as usual (c.f. [8, Tag 0379]) the following fact.

Lemma 5.12 *Let $f : X \rightarrow Y$ be any morphism of rigid spaces. Then the image of any irreducible subspace is irreducible.*

Let now X be any irreducible proper rigid space, G a commutative rigid group, and let us consider a morphism $\varphi : X \rightarrow G$ containing 0 in the image. Then for any $n \in \mathbb{N}$,

we define a morphism

$$\varphi_n : X^n \rightarrow G, \quad (x_1, \dots, x_n) \mapsto \sum_{i=1}^n (-1)^i \varphi(x_i).$$

Let $Z_n \subseteq G$ be the image of φ_n . The fibre product X^n is still proper, and since K is algebraically closed, it is also still irreducible by [6, Lemma 2.2.3]. Since G is separated, φ_n is proper by Proposition 5.11, hence Z_n is Zariski-closed by Proposition 5.10. We can endow it with its induced-reduced structure, so Z_n is a reduced rigid space. The key technical result is now.

Lemma 5.13 *Let $\varphi : X \rightarrow G$ be a morphism from an irreducible proper rigid space to a commutative rigid group containing 0 in the image, and let Z_n be as above.*

Then each Z_n is proper. The Z_n 's form an ascending chain

$$Z_0 \subseteq Z_1 \subseteq Z_2 \subseteq \dots$$

of equidimensional irreducible Zariski-closed subspaces of G . This chain stabilises.

Proof That the Z_n is proper follows from Proposition 5.11. Since X^n is irreducible, Z_n is irreducible by Lemma 5.12. That the Z_n 's form an ascending chain follows from the assumption that $0 \in \text{im}(\varphi)$. Let $e \in X(K)$ be any preimage of 0 under φ , then the diagram

$$\begin{array}{ccc} X^n & \xrightarrow{\varphi_n} & G \\ \downarrow \text{id} \times \{e\} & & \parallel \\ X^{n+1} & \xrightarrow{\varphi_{n+1}} & G \end{array}$$

commutes, and thus $Z_n \subseteq Z_{n+1}$.

By [6, paragraph before Lemma 2.2.3], any irreducible rigid space is equidimensional, meaning that for fixed n , the Krull dimension $\dim \mathcal{O}_{Z_n, z}$ of the stalks is constant for all $z \in Z_n(K)$. Since G is a rigid space, we know that $\dim \mathcal{O}_{G, z} \leq \dim \mathcal{O}_{G, z} < \infty$, from which it follows that there is $n \in \mathbb{N}$ such that the dimension of Z_m is the same for all $m \geq n$. Then $Z_n \subseteq Z_m$ is an analytic subspace of the irreducible rigid space Z_m which is equidimensional of the same dimension as Z_m . By [6, Corollary 2.2.7], it follows that $Z_n = Z_m$. ■

Proof of Lemma 5.9: Since X is irreducible, it is in particular connected, so we can without loss of generality assume that G is connected by replacing G with G° . We can further reduce to the case that G is commutative by the rigid version of a standard argument in classical algebraic geometry. We consider the morphism of rigid spaces

$$f : X \times G \rightarrow G, \quad x, g \mapsto \varphi(x)g\varphi(x)^{-1}g^{-1}.$$

This satisfies $f(x, 1) = 1$ for all $x \in X$. Hence the rigid analytic version of the rigidity lemma [26, Lemma 7.1.2] gives an open neighbourhood $1 \in U \subseteq G$ on which $f|_{X \times U}$ factors through U . Since $f(x_0, g) = 1$ for all $g \in G$ by assumption, we find that $f|_{X \times U} = 1$. It follows that for any K -point $z \in X(K)$, the induced morphism of rigid spaces

$$\psi : G \rightarrow G, \quad g \mapsto \varphi(z)g\varphi(z)^{-1}g^{-1}$$

restricts to $\psi|_U = 1$ on the non-empty open subspace U . As G is irreducible, ψ is uniquely determined by $\psi|_U$. This shows that $\psi = 1$ on all of G . Thus φ factors through $Z(G)$.

Hence we may assume that $\varphi : X \rightarrow G$ is a morphism to a commutative rigid group containing 0 in the image. By Lemma 5.13, it follows from this that there is a proper rigid space $Z = \cup_{m \in \mathbb{N}} Z_m = Z_n$ for $n \gg 0$ with the property that φ factors through a closed immersion $X \rightarrow Z \rightarrow G$. Moreover, by construction, Z contains 0 and the diagram

$$\begin{array}{ccc} Z \times Z & \longrightarrow & Z \\ \downarrow & & \downarrow \\ G \times G & \xrightarrow{m} & G \end{array}$$

commutes, where the top line is the map induced by the natural morphism $Z_{2n} \times Z_{2n} \rightarrow Z_{4n}$.

Due to the alternating sign, Z is also closed under inverses. Namely, the morphisms $Z_n \rightarrow Z_{n+1}$ induced by the map $X^n \rightarrow X^{n+1}$, $(x_1, \dots, x_n) \mapsto (e, x_1, \dots, x_n)$ induce a morphism $Z \rightarrow Z$ over $[-1] : G \rightarrow G$.

All in all, it follows that $Z \subseteq G$ is a rigid subgroup of G . As it is proper and irreducible, in particular connected, it is an abeloid subvariety of G by the structure result of Lütkebohmert [25, Theorem I], as we wanted to see. ■

Proof of Theorem 5.8 Let $A, B \subseteq G$ be any two closed abeloid subgroups. Then $A \times B$ is still smooth proper and connected. By Lemma 5.9 we find that the map

$$A \times B \rightarrow G \times G \xrightarrow{m} G,$$

factors through an abeloid subvariety $C \subseteq G$ that contains A and B . It follows that the union of all abeloid subvarieties of G is an abeloid subvariety. Now the theorem follows from Lemma 5.9. ■

5.4 Cocycles from homomorphisms: The commutative case

With these preparations, we can now prove Theorem 5.6 for commutative G (not necessarily connected). More precisely, we can prove:

Theorem 5.14 *Let G be any commutative rigid group, X any connected smooth proper rigid space. Then the following natural map is surjective:*

$$\text{Hom}_{\text{cts}}(\pi, G(K)) \rightarrow H^1_{\text{cts}}(\pi, G(\tilde{X})).$$

Proof Consider the evaluation morphism

$$G(\tilde{X}) \xrightarrow{ev_{x_0}} G(K)$$

at the fixed base point $x_0 \in \tilde{X}$. This is a surjective group homomorphism, which is split by sending $g \in G(K)$ to the constant map $\tilde{X} \rightarrow \text{Spa}(K) \xrightarrow{g} G(K)$. Denote by $G(\tilde{X})^\circ$ the kernel of this homomorphism ev_{x_0} , this is the subgroup of those morphism $\tilde{X} \rightarrow G$ that send x_0 to 0. Since G is commutative, the splitting of ev_{x_0} now induces a short

exact sequence

$$(9) \quad 0 \rightarrow G(K) \rightarrow G(\tilde{X}) \xrightarrow{\phi} G(\tilde{X})^\circ \rightarrow 0,$$

where ϕ sends $\varphi : \tilde{X} \rightarrow G$ to the map φ° making the following diagram commute:

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\varphi} & G \\ \parallel & & \uparrow \cdot \varphi(x_0) \\ \tilde{X} & \xrightarrow{\varphi^\circ} & G. \end{array}$$

We note that ϕ is a group homomorphism since G is commutative. Observe now that the first morphism $G(K) \rightarrow G(\tilde{X})$ in the exact sequence is clearly π -equivariant. Consequently, $G(\tilde{X})^\circ$ inherits a natural quotient action (note that the action of π does not preserve $G(\tilde{X})^\circ$ as a subspace of $G(\tilde{X})$). Explicitly, the quotient action of π on $G(\tilde{X})^\circ$ is defined by letting $\gamma \in \pi$ send φ to the morphism φ^γ defined by making the following diagram commutative:

$$(10) \quad \begin{array}{ccc} \tilde{X} & \xrightarrow{\varphi} & G \\ \cdot \gamma \uparrow & & \uparrow \cdot \varphi(\gamma x_0) \\ \tilde{X} & \xrightarrow{\varphi^\gamma} & G. \end{array}$$

Recall now from Proposition 2.11 that $G(\tilde{X})$ has a natural topology defined by a system of open neighbourhoods $G_k(\tilde{X})$ where $(G_k)_{k \in \mathbb{N}}$ is a system of subgroups of G . When we endow $G(\tilde{X})^\circ \subseteq G(\tilde{X})$ with the natural subspace topology, then the section $\phi : G(\tilde{X}) \rightarrow G(\tilde{X})^\circ$ is continuous. Indeed, it is immediate from the definition that $\phi^{-1}(G_k(\tilde{X})^\circ) \supseteq G_k(\tilde{X})$.

In summary, this shows that (9) is a short exact sequence of topological π -modules. We deduce that there is an exact sequence

$$(11) \quad \text{Hom}_{\text{cts}}(\pi, G(K)) \rightarrow H^1_{\text{cts}}(\pi, G(\tilde{X})) \rightarrow H^1_{\text{cts}}(\pi, G(\tilde{X})^\circ).$$

More explicitly, this can be interpreted as follows: any geometric 1-cocycle $\pi \times \tilde{X} \rightarrow G$ can be written as a product of a continuous map $\pi \rightarrow G(K)$ obtained by specialising at x_0 (not necessarily a homomorphism) and a morphism $\rho : \pi \times \tilde{X} \rightarrow G$ such that $\rho(\gamma, x_0) = 0$.

We wish to see that the last term in the exact sequence vanishes. The crucial point is:

Lemma 5.15 *The subspace topology of $G(\tilde{X})^\circ \subseteq G(\tilde{X})$ is the discrete one.*

Proof Let $G_k \subseteq G$ be any affinoid open subgroup isomorphic to \mathbb{B}^d as in Proposition 2.11, then the topology of $G(\tilde{X})$ is defined by decreeing that $G_k(\tilde{X})$ is open. But $G_k(\tilde{X}) \cong \mathbb{B}^d(\tilde{X}) = \mathcal{O}_K^d$. Hence $G_k(\tilde{X}) \cap G(\tilde{X})^\circ = \{0\}$ is open, and $G(\tilde{X})^\circ$ is discrete. ■

Lemma 5.16 *Let $\varphi : \tilde{X} \rightarrow G$ be a morphism in $G(\tilde{X})^\circ$, then there is an open subgroup $U \subseteq \pi$ such that the action of U as defined in Equation (10) fixes φ .*

Proof We apply the rigidity lemma 5.7 to the morphism

$$\widetilde{X} \times \underline{\pi} \rightarrow G, \quad (y, \gamma) \mapsto \varphi(\gamma x_0) \varphi(\gamma y)^{-1} \varphi(y)$$

and find an open subgroup U of π such that the restriction to $\widetilde{X} \times U$ vanishes. By the diagram defining the action on $G(\widetilde{X})^\circ$, this implies that $\varphi^\gamma = \varphi$ for all $\gamma \in U$. ■

Lemma 5.17 *For any continuous 1-cocycle $\rho : \pi \rightarrow G(\widetilde{X})^\circ$, there is a connected finite étale cover $X' \rightarrow X$ with Galois group $Q_{X'}$ such that ρ factors through $X' \times Q_{X'}$. Hence the following map is surjective*

$$H_{\text{cts}}^1(\pi, \varinjlim_{X' \rightarrow X} G(X')^\circ) \rightarrow H_{\text{cts}}^1(\pi, G(\widetilde{X})^\circ),$$

where $G(X')^\circ := \ker(G(X') \xrightarrow{\text{ev}_{x_0}} G(K))$ is endowed with the discrete topology.

Proof By Lemma 5.15, ρ is continuous from a profinite set to a discrete set. It follows that it has finite image. Using Lemma 5.16, this shows that ρ factors through $(G(\widetilde{X})^\circ)^U$ for some open subgroup $U \subseteq \pi$. Shrinking U further, we may assume that it moreover factors through π/U . This shows that

$$\varinjlim_{X' \rightarrow X} H^1(Q_{X'}, G(X')^\circ) \rightarrow H_{\text{cts}}^1(\pi, G(\widetilde{X})^\circ)$$

is surjective. By [29, Proposition (1.2.5)], the left hand side equals $H_{\text{cts}}^1(\pi, \varinjlim_{X' \rightarrow X} G(X')^\circ)$. ■

Proposition 5.18 *Let G be any commutative rigid group and X any connected smooth proper rigid space. Then $H_{\text{cts}}^1(\pi, G(\widetilde{X})^\circ) = 0$.*

Proof By Lemma 5.17, it suffices to prove that $H^1(\pi, \varinjlim_{X' \rightarrow X} G(X')^\circ) = 0$. By Theorem 5.8, there is a maximal abeloid subvariety $A \subseteq G$, and we have $G(X')^\circ = A(X')^\circ$. Consider now the short exact sequence of sheaves on $X_{\text{ét}}$

$$0 \rightarrow A[N] \rightarrow A \xrightarrow{[N]} A \rightarrow 0.$$

Taking cohomology over $X' \rightarrow X$, this gives a long exact sequence

$$0 \rightarrow A[N](X') \rightarrow A(X') \rightarrow A(X') \rightarrow H_{\text{ét}}^1(X', A[N]).$$

The first term is equal to $A[N](K)$ since X' is connected. Using that $[N] : A(K) \rightarrow A(K)$ is surjective, it follows that we obtain a left-exact sequence

$$0 \rightarrow A(X')^\circ \xrightarrow{[N]} A(X')^\circ \rightarrow H_{\text{ét}}^1(X', A[N]).$$

Taking the colimit over $X' \rightarrow X$, the last term vanishes since every class in $H^1(X', A[N])$ defines a finite étale torsor over X' and is thus killed by some finite étale cover of X . It follows that

$$[N] : \varinjlim_{X' \rightarrow X} A(X')^\circ \xrightarrow{\sim} \varinjlim_{X' \rightarrow X} A(X')^\circ$$

is an isomorphism. In particular, $\varinjlim_{X' \rightarrow X} A(X')^\circ$ is a \mathbb{Q} -vector space. By [29, Proposition (1.6.2.c)], it follows that $H_{\text{cts}}^1(\pi, \varinjlim_{X' \rightarrow X} A(X')^\circ) = 0$, as desired. ■

Theorem 5.14 now follows from the exact sequence (11) and Proposition 5.18.

5.5 The case of general G

Building on the commutative case, we can now deduce the case of connected rigid groups G .

Proof of Theorem 5.6 The key idea is to use the exact sequence from Lemma 2.9

$$0 \rightarrow Z(G) \rightarrow G \xrightarrow{\text{ad}} \text{Aut}(\mathfrak{g}),$$

where $Z(G)$ is the center as in Definition 2.8.

Working in the category of pro-étale sheaves on Perf_K , we denote by H the sheaf theoretic image of G in $\text{Aut}(\mathfrak{g})$. Then we have a short exact sequence of sheaf of groups on $\text{Perf}_{K,\text{proét}}$

$$0 \rightarrow Z(G) \rightarrow G \rightarrow H \rightarrow 0.$$

Evaluating this over \tilde{X} and over K , we get a commutative diagram of π -equivariant maps

$$\begin{array}{ccccccc} 0 & \longrightarrow & Z(G)(\tilde{X}) & \longrightarrow & G(\tilde{X}) & \longrightarrow & H(\tilde{X}) \longrightarrow 0 \\ & & \uparrow & & \uparrow & & \sim \uparrow \\ 0 & \longrightarrow & Z(G)(K) & \longrightarrow & G(K) & \longrightarrow & H(K) \longrightarrow 0. \end{array}$$

The map $H(K) \rightarrow H(\tilde{X})$ in the right column is an isomorphism. This follows from the fact that $H \subseteq \text{Aut}(\mathfrak{g}) \cong \text{GL}_n$ for $n = \dim G$, and $\text{GL}_n(\tilde{X}) = \text{GL}_n(K)$.

The rows of this diagram are still exact: the bottom row is exact by surjectivity of $G \rightarrow H$ and the fact that $\text{Spa}(K)$ is strictly totally disconnected. The top row is right exact by surjectivity in the bottom row. We now take continuous group cohomology of π with coefficients in the above diagram. Since $Z(G)(\tilde{X}) \subseteq G(\tilde{X})$ is a commutative normal subgroup, by [38, Chapitre I, Proposition 43], we get a commutative diagram of pointed sets

$$\begin{array}{ccccccc} 0 \rightarrow H_{\text{cts}}^1(\pi, Z(G)(\tilde{X})) & \rightarrow & H_{\text{cts}}^1(\pi, G(\tilde{X})) & \rightarrow & H_{\text{cts}}^1(\pi, H(\tilde{X})) & \rightarrow & H_{\text{cts}}^2(\pi, Z(G)(\tilde{X})) \\ & & \uparrow & & \sim \uparrow & & \uparrow \\ 0 \rightarrow H_{\text{cts}}^1(\pi, Z(G)(K)) & \rightarrow & H_{\text{cts}}^1(\pi, G(K)) & \rightarrow & H_{\text{cts}}^1(\pi, H(K)) & \rightarrow & H_{\text{cts}}^2(\pi, Z(G)(K)). \end{array}$$

The leftmost horizontal arrows have trivial kernel since $G(K) \rightarrow H(K)$ is surjective and thus so is $G(\tilde{X})^\pi \rightarrow H(\tilde{X})^\pi = H(K)^\pi = H(K)$.

By Theorem 5.14 the leftmost vertical arrow is surjective. We also have injectivity of the rightmost vertical arrow: Using the sequence (9), its kernel identifies with $H_{\text{cts}}^1(\pi, Z(G)^\circ)$, which is trivial by Proposition 5.18.

Now the crucial point is that $Z(G)(\tilde{X}) \subseteq G(\tilde{X})$ is even a central subgroup. By [38, Chapitre I, Proposition 42], it follows that the maps

$$H_{\text{cts}}^1(\pi, G(K)) \rightarrow H_{\text{cts}}^1(\pi, H(K)) \quad \text{and} \quad H_{\text{cts}}^1(\pi, G(\tilde{X})) \rightarrow H_{\text{cts}}^1(\pi, H(\tilde{X}))$$

are in fact torsors under the group $H^1_{\text{cts}}(\pi, Z(G)(K))$. We can therefore argue as in the proof of the 5-Lemma to see that the second vertical arrow from the left is surjective as well. ■

Combining Theorem 5.6 and Theorem 5.14 completes the proof of Theorem 5.1. ■

5.6 A counter-example

It is not in general true that the functor from Theorem 5.1 is fully faithful. For example, this fails if the natural map $\text{Hom}_{\text{cts}}(\pi, G(K)) \rightarrow H^1_{\text{cts}}(\pi, G(\tilde{X}))$ (which is surjective by Theorem 5.6) is not injective. We now give an example where this happens. Let us assume for simplicity that G is commutative. Then by the long exact sequence of group cohomology for Equation (9), the kernel of this map can be identified with the cokernel of

$$G(X) \rightarrow H^0(\pi, G(\tilde{X})^\circ).$$

Unravelling the explicit description of the π -action on $G(\tilde{X})$ in (10), we see that the second term is the set of $g \in G(\tilde{X})^\circ$ for which $\gamma^*g - g \in G(K)$ for any $\gamma \in \pi$. This can be non-trivial, even in very good cases, as the following example demonstrates.

Proposition 5.19 *Let both $X = A$ and $G = B$ be abelian varieties, considered as rigid analytic spaces. Assume that $\text{Hom}(A, B) = 0$, but that there exists an isomorphism $\tilde{A} \xrightarrow{\sim} \tilde{B}$. Then the composition*

$$\rho : \pi_1^{\text{ét}}(A, 0) \rightarrow \tilde{A} \xrightarrow{\sim} \tilde{B} \rightarrow B = G$$

defines a non-trivial element in $\ker(\text{Hom}_{\text{cts}}(\pi, G(K)) \rightarrow H^1_{\text{cts}}(\pi, G(\tilde{X})))$. In particular, the functor of Theorem 5.1 is not fully faithful in this case.

There are plenty of examples of abelian varieties A and B satisfying the assumptions of Proposition 5.19 by the main result of [14]. For example, we can take A and B to be any non-isogeneous simple abelian varieties of good reduction whose special fibres are isogeneous.

Proof We first note that $\rho = 0$ would imply that $\tilde{A} \rightarrow \tilde{B}$ sends $\pi_1^{\text{ét}}(A, 0)$ to $\pi_1^{\text{ét}}(B, 0)$ and hence would induce an isogeny $A \rightarrow B$, contradicting the assumptions. Hence $\rho \neq 0$.

On the other hand, by [14, Theorem 1.9], we have

$$G(\tilde{X})^\circ = \text{Hom}(\tilde{A}, B) = \text{Hom}(\tilde{A}, \tilde{B}),$$

where the last equality holds by [14, Proposition 3.8, Corollary 3.10]. Observe now that by definition, the π -action on $G(\tilde{X})^\circ$ as described in (10) is such that π acts trivially on homomorphisms. It follows that

$$H^0(\pi, G(\tilde{X})^\circ) = \text{Hom}(\tilde{A}, \tilde{B}) \neq 0$$

which is non-trivial by assumption. All in all, this shows that ρ comes from $G(\tilde{X})^\circ$ via the boundary map. Thus the kernel of $\text{Hom}_{\text{cts}}(\pi, G(K)) \rightarrow H^1_{\text{cts}}(\pi, G(\tilde{X}))$ does not vanish. ■

Remark 5.20 Regarding the setting of Proposition 5.19, we note that already G -torsors on X_{an} are an interesting category when G and X are both abelian varieties. For abelian varieties A and B over K considered as rigid spaces, the Raynaud uniformization

$$0 \rightarrow \Lambda \rightarrow E \rightarrow A \rightarrow 0$$

associated with A induces a homomorphism $\text{Hom}(\Lambda, B) \rightarrow \text{Ext}^1(A, B) \rightarrow H^1(A, B)$. If A has totally degenerate reduction, this map has non-trivial image. Hence, in this case there exist analytic B -torsors over A which are not algebraic, since every Zariski- B -torsor over A is trivial by [37, Section 4, Lemme 4]. It seems an interesting question if there exists a classification of all analytic/étale torsors in this setting which is analogous to the complex analytic characterization of torus bundles over tori [30], and how this changes for the v -topology.

6 The p -adic Lie algebra exponential for commutative rigid groups

6.1 Recollections on the p -adic Lie group logarithm

Let L be any non-archimedean field extension of \mathbb{Q}_p . We will later take $L = K$ algebraically closed, but for now we can work more generally. Recall that in p -adic arithmetic, the p -adic logarithm defines a left-exact sequence of topological groups

$$0 \rightarrow \mu_{p^\infty}(L) \rightarrow 1 + \mathfrak{m}_L \xrightarrow{\log} L.$$

This can be upgraded to a short exact sequence of rigid group varieties in the étale topology [7, Section 7][16, Lemma 2.18]

$$0 \rightarrow \mu_{p^\infty} \rightarrow \mathbb{D}(1) \rightarrow \mathbb{G}_a \rightarrow 0,$$

where for any $a \in L$, we denote by $\mathbb{D}(a)$ the open rigid unit disc around a . There is a partial splitting of \log given by the p -adic exponential

$$\exp : p^\alpha \mathfrak{m}_L \rightarrow 1 + \mathfrak{m}_L, \quad x \mapsto \sum_{n=0}^\infty \frac{x^n}{n!}$$

where $\alpha = \frac{1}{p-1}$ for $p > 2$ and $\alpha = 2$ for $p = 2$. This can again be upgraded to a homomorphism of rigid groups

$$\exp : p^\alpha \mathbb{D}(0) \rightarrow \mathbb{D}(1).$$

But in contrast to the complex case, the exponential series does not converge on all of L . Indeed, the logarithm sequence of rigid groups is never split as any morphism from \mathbb{G}_a to $\mathbb{D}(1)$ is constant. Instead, one can only obtain non-canonical splittings on L -points.

Definition 6.1 By an exponential map for L we mean a continuous group homomorphism

$$\exp : L \rightarrow 1 + \mathfrak{m}_L$$

which splits the p -adic logarithm $\log : 1 + \mathfrak{m}_L \rightarrow L$ and is a continuation of the canonical p -adic exponential on the open subgroup $p^\alpha \mathfrak{m}_L \subseteq L$.

Lemma 6.2 *An exponential map $\exp : L \rightarrow 1 + \mathfrak{m}_L$ exists if $H_{\text{ét}}^1(\text{Spa}(L), \mu_p) = 1$. In particular, it exists if L is algebraically closed.*

Proof It suffices to find a group-theoretic splitting of \log extending \exp , this will automatically be continuous if it extends \exp on $p^\alpha \mathfrak{m}_L$. To find this splitting, we consider the long exact sequence

$$\dots \rightarrow \text{Hom}(L, 1 + \mathfrak{m}_L) \rightarrow \text{Hom}(p^\alpha \mathfrak{m}_L, 1 + \mathfrak{m}_L) \rightarrow \text{Ext}^1(L/p^\alpha \mathfrak{m}_L, 1 + \mathfrak{m}_L)$$

We claim that the last term vanishes. If L is algebraically closed, this is because $1 + \mathfrak{m}_L$ is divisible, in particular injective (c.f. [43, Section A.2.1]). In general, since $L/p^\alpha \mathfrak{m}_L$ is a \mathbb{Z}_p -module, it suffices to prove that $1 + \mathfrak{m}_L$ is p -divisible. For this we consider the sequence on $\text{Spa}(L)_{\text{ét}}$

$$1 \rightarrow \mu_p \rightarrow 1 + \mathfrak{m}_L \mathcal{O}^+ \xrightarrow{x \mapsto x^p} 1 + \mathfrak{m}_L \mathcal{O}^+ \rightarrow 1.$$

Since $H_{\text{ét}}^1(\text{Spa}(L), \mu_p) = 1$, this shows that $1 + \mathfrak{m}_L$ is p -divisible. ■

An exponential is one of the choices that Faltings makes in [10] to define his p -adic Simpson correspondence on curves. In the process, Faltings shows that the datum of \exp induces an exponential map for any commutative algebraic group. The goal of this section is to prove a generalization of this statement for rigid analytic groups, see Theorem 6.12 below.

We now switch back to the algebraically closed field K . From now on, we assume that G is a commutative rigid group over K .

Lemma 6.3 *Let G be a quasi-compact commutative rigid group over K . Then $G[p^n]$ is a finite étale group for each $n \in \mathbb{N}$.*

Proof By [11, Lemme 1], any homomorphism $f : G \rightarrow H$ of rigid groups such that the induced map $\text{Lie}(f) : \text{Lie } G \rightarrow \text{Lie } H$ is bijective is étale. It follows that $[p^n] : G \rightarrow G$ is étale, so $G[p^n]$ is étale. On the other hand, $G[p^n]$ is Zariski-closed in G , because it is the pullback of a Zariski-closed subset $1 \in G$ along $[p^n] : G \rightarrow G$ (see also [11, Exemple 2]). It follows that for any affinoid open $U \subseteq G$, the intersection $G[p^n] \cap U$ is affinoid of dimension 0 in an affinoid, hence finite. As G is quasi-compact, it follows that $G[p^n]$ is finite. ■

For any commutative rigid group G , we have the topologically p -torsion subgroup.

Definition 6.4 ([19, Proposition 2.14]) *The topological p -torsion subgroup $\widehat{G} \subseteq G$ is the image in v -sheaves of the morphism $\underline{\text{Hom}}(\mathbb{Z}_p, G) \xrightarrow{\text{ev}_1} G$.*

We note that this was written as $G(p^\infty)$ in [19]. A priori, $\widehat{G} \subseteq G$ is a v -sheaf, but we can also regard it as a rigid group, which we can use to give a more canonical description of the logarithm map of Lemma 2.5.

Proposition 6.5 ([11, Section 1.6] and [19, Proposition 2.14]) *Let G be a commutative rigid group.*

- (1) *The v -sheaf \widehat{G} is representable by an open subgroup $\widehat{G} \subseteq G$.*
- (2) *There is a unique homomorphism*

$$\log_G : \widehat{G} \rightarrow \mathfrak{g},$$

such that $\text{Lie } \log_G : \text{Lie } \widehat{G} \rightarrow \text{Lie } G$ is the identity. We have $\ker \log_G = G[p^\infty]$.

Assume moreover that there is an admissible formal group \mathfrak{G} over \mathcal{O}_K with adic generic fibre $G = \mathfrak{G}_\eta^{\text{ad}}$ as defined in [35, Section 2.2]. Then:

- (3) The identity component \widehat{G}° is isomorphic to $(\mathfrak{G}^\wedge)_\eta^{\text{ad}}$ where \mathfrak{G}^\wedge is the formal completion at the identity.
- (4) There is a “connected-étale” short exact sequence of rigid groups

$$1 \rightarrow \widehat{G}^\circ \rightarrow \widehat{G} \rightarrow (\mathfrak{G}[p^\infty]^\text{ét})_\eta^{\text{ad}} \rightarrow 0.$$

Proof Parts (1) and (2) are contained in [19, Proposition 2.14.1 and 2]. We reprove (2) to justify the statement about Lie log_G . By Lemma 2.5 and [15, Proposition 3.5], there is an open subgroup $G_0 \subseteq G$ for which there exists an injective homomorphism $\log_0 : G_0 \rightarrow \mathfrak{g}$ with the desired property. For any smooth affinoid algebra R and any $x \in \widehat{G}(R)$, one has $x^{p^n} \in G_0(R)$ for some $n \in \mathbb{N}$. Since \mathfrak{g} is uniquely divisible, we can define $\log(x) = \frac{1}{p^n} \log_0(x^{p^n}) \in \mathfrak{g}(R)$.

Part (4) can be deduced from the connected-étale sequence of the p -divisible group $\mathfrak{G}[p^\infty]$ over \mathcal{O}_K by [11, Corollaire 13]. Alternatively, we can give a v-sheaf-theoretic argument which also shows (3). By the proof of [19, Proposition 2.14], $\widehat{G} \subseteq G$ is the sub-v-sheaf on Perf_K given by sheafification of

$$(R, R^+) \mapsto \{x \in \mathfrak{G}(R^+) \mid x \bmod \mathfrak{m}_K \in \mathfrak{G}(R^+/\mathfrak{m}_K)[p^\infty]\},$$

whereas by [35, Proposition 2.2.2] and the definition of formal completion, $(\mathfrak{G}^\wedge)_\eta^{\text{ad}}$ is the sheafification of

$$(R, R^+) \mapsto \{x \in \mathfrak{G}(R^+) \mid x \equiv 1 \bmod \mathfrak{m}_K \text{ in } \mathfrak{G}(R^+/\mathfrak{m}_K)\}.$$

Since $(\mathfrak{G}^\wedge)_\eta^{\text{ad}}$ is an open disc by [35, Lemma 3.1.2], in particular connected, this shows $(\mathfrak{G}^\wedge)_\eta^{\text{ad}} \subseteq \widehat{G}^\circ$. On the other hand, since R^+/\mathfrak{m}_K is reduced, we have

$$\mathfrak{G}(R^+/\mathfrak{m}_K)[p^\infty] = \mathfrak{G}[p^\infty]^\text{ét}(R^+/\mathfrak{m}_K) = \mathfrak{G}[p^\infty]^\text{ét}(R^+)$$

by Henselian lifting. This exhibits $(\mathfrak{G}^\wedge)_\eta^{\text{ad}}$ as the kernel of a natural map $\widehat{G} \rightarrow (\mathfrak{G}[p^\infty]^\text{ét})_\eta^{\text{ad}}$ from a rigid group to an étale group. Hence $\widehat{G}^\circ \subseteq (\mathfrak{G}^\wedge)_\eta^{\text{ad}}$. ■

There is also a version of \widehat{G} which incorporates torsion coprime to p .

Definition 6.6 ([19, Definition 2.5, Proposition 2.14]) There is a unique open subgroup $G^{\text{tt}} \subseteq G$ that represents the image of the morphism of v-sheaves $\text{Hom}(\widehat{\mathbb{Z}}, G) \xrightarrow{\text{ev}_1} G$. We call it the topological torsion subgroup of G . The natural inclusion $\widehat{G} \rightarrow G^{\text{tt}}$ identifies \widehat{G} with an open and closed subgroup of G^{tt} , and the cokernel is given by the union of all coprime-to- p -torsion points of G .

Lemma 6.7 There is a left exact sequence of rigid groups, functorial in G

$$0 \rightarrow G_{\text{tor}} \rightarrow G^{\text{tt}} \xrightarrow{\log_G} \mathfrak{g},$$

such that \log_G induced the identity on tangent spaces. Here $G_{\text{tor}} = \varinjlim_{N \in \mathbb{N}} G[N]$.

Proof This follows from Proposition 6.5(2). We have $G^{\text{tt}} = \widehat{G} \cdot G_{\text{tor}}$ and define $\log_G = 0$ on G_{tor} . ■

6.2 Criteria for \log_G to be surjective

The goal of this subsection is to prove the following characterisation for when the logarithm morphism is surjective.

Definition 6.8

- (1) Following Fargues [11], we say that G is an analytic p -divisible group if $G = \widehat{G}$ and $[p] : G \rightarrow G$ is finite surjective.
- (2) More generally, we say that a general rigid group G is locally p -divisible if it contains an open subgroup U that is an analytic p -divisible group.

Example 6.9

- (1) Let G be a commutative connected algebraic group. Then by Rosenlicht’s theorem, it is an extension of an abelian variety by a product of \mathbb{G}_a and tori. It follows that $[p] : G \rightarrow G$ is surjective, since this is true for abelian varieties, \mathbb{G}_a and tori. By Proposition 6.10 below, it follows that G is locally p -divisible.
- (2) Any extension of locally p -divisible groups is locally p -divisible.
- (3) The closed unit disc with its additive structure, $G = \mathbb{G}_a^+$, is not locally p -divisible. It is topologically p -torsion, so $G = \widehat{G}$, but clearly $[p] : G \rightarrow G$ is not surjective.

Proposition 6.10 *Let G be a commutative rigid group over K . Consider the statements:*

- (1) $[p] : G \rightarrow G$ is surjective finite étale,
- (2) $[p] : G \rightarrow G$ is surjective,
- (3) $[p] : \widehat{G} \rightarrow \widehat{G}$ is surjective,
- (4) G° is an analytic p -divisible group,
- (5) G is locally p -divisible,
- (6) there exists an open subgroup $U \subseteq G$ on which $[p] : U \rightarrow U$ is surjective,
- (7) $\log_G : \widehat{G} \rightarrow \mathfrak{g}$ is surjective.

Then (1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4) \Leftrightarrow (5) \Leftrightarrow (6) \Leftrightarrow (7). If $\pi_0(G)$ is finite, we have (1) \Leftrightarrow (2). If G is quasi-compact and connected, we have (4) \Leftrightarrow (1), so in this case all statements are equivalent.

Remark 6.11 We suspect that quasi-compactness is not necessary for (4) \Rightarrow (1) to hold.

Proof of Proposition 6.10 (1) \Rightarrow (2) is trivial.

(2) \Rightarrow (3): For any point $x \in \widehat{G}(C, C^+)$ with values in an algebraically closed non-archimedean field extension (C, C^+) of K , there exists some $y \in G(C, C^+)$ such that $[p](y) = x$. But then $[p^n](y) = [p^{n-1}](x) \rightarrow 0$, so $y \in \widehat{G}(C, C^+)$.

(3) \Rightarrow (6): Trivial as $\widehat{G} \subseteq G$ is open.

(4) \Rightarrow (5): We set $U = \widehat{G}^\circ$ which is by assumption analytic p -divisible.

(5) \Rightarrow (6) is trivial.

(6) \Rightarrow (7): Since (2) \Rightarrow (3), we may replace U by \widehat{U} . By Lemma 2.5, the image $M := \log(U)$ is then an open subgroup of \mathfrak{g} . If $[p] : U \rightarrow U$ is surjective, then so is $[p] : M \rightarrow M$. But the only open p -divisible subgroup of \mathfrak{g} is the whole group.

(3) \Rightarrow (4): It is clear that $U := \widehat{G}^\circ$ satisfies $U = \widehat{U}$. By the Snake lemma (applied in the category of v -sheaves), the cokernel of $[p] : \widehat{G}^\circ \rightarrow \widehat{G}^\circ$ is equal to the constant

group given by the cokernel of the map $\widehat{G}[p] \rightarrow \pi_0(\widehat{G})[p]$. But any morphism from \widehat{G}° to a constant group vanishes. Thus the map $[p] : U \rightarrow \mathfrak{g}$ is surjective. It is étale by [11, Lemme 1].

It remains to see that this map is moreover finite. For this, it suffices by [11, Lemme 5] to prove that $U[p]$ is finite. More precisely, as it is a p -torsion abelian group, $U[p]$ is necessarily isomorphic to $(\mathbb{Z}/p\mathbb{Z})^{\oplus I}$ for some index set I , and we need to see that I is finite. We first note that $[p]$ being surjective implies by choosing successive preimages of basis elements under $[p]$ that $U[p^n] = (\mathbb{Z}/p^n\mathbb{Z})^{\oplus I}$, hence

$$U[p^\infty] = (\mathbb{Q}_p/\mathbb{Z}_p)^{\oplus I}.$$

Since we already know that (3) \Rightarrow (7), it follows that we have a short exact sequence of rigid groups

$$0 \rightarrow (\mathbb{Q}_p/\mathbb{Z}_p)^{\oplus I} \rightarrow U \rightarrow \mathfrak{g} \rightarrow 0.$$

We claim that U being connected forces $|I| < \infty$. To see this, let $m := \dim G + 1$ and consider the pushout V of U along any surjective morphism $(\mathbb{Q}_p/\mathbb{Z}_p)^{\oplus I} \rightarrow (\mathbb{Q}_p/\mathbb{Z}_p)^m$, so that we have a commutative diagram of rigid groups

$$\begin{array}{ccccccc} 0 & \longrightarrow & (\mathbb{Q}_p/\mathbb{Z}_p)^{\oplus I} & \longrightarrow & U & \longrightarrow & \mathfrak{g} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & (\mathbb{Q}_p/\mathbb{Z}_p)^m & \longrightarrow & V & \longrightarrow & \mathfrak{g} \longrightarrow 0. \end{array}$$

Then V is still connected, being the image of a surjective map $U \rightarrow V$. Since it is by definition an extension of analytic p -divisible groups, V is clearly itself an analytic p -divisible group. By [11, Théorème 7.2], this implies that $m = \text{rk}(V[p^\infty]) \leq \dim V = \dim G$, as we can discard the first summand in loc. cit. since V is connected. This implies that $|I| \leq \dim G$. Consequently, $U[p]$ is finite, as we wanted to see.

(7) \Rightarrow (4): For this we consider the morphism of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \widehat{G}^\circ[p^\infty] & \longrightarrow & \widehat{G}^\circ & \xrightarrow{\log} & \mathfrak{g} \longrightarrow 0 \\ & & \uparrow [p] & & \uparrow [p] & & \uparrow [p] \\ 0 & \longrightarrow & \widehat{G}^\circ[p^\infty] & \longrightarrow & \widehat{G}^\circ & \xrightarrow{\log} & \mathfrak{g} \longrightarrow 0. \end{array}$$

For each n , denote by Q_n the cokernel of $[p] : \widehat{G}^\circ[p^n] \rightarrow \widehat{G}^\circ[p^n]$ (sheaf-theoretic, e.g., in v -sheaves). Since $\widehat{G}^\circ[p^n] \rightarrow \text{Spa}(K)$ is étale, it is a constant rigid group, and hence so is Q_n . It follows that the cokernel $Q = \varinjlim Q_n$ in the diagram is a constant rigid group. Hence any homomorphism from the connected group \widehat{G}° to Q is trivial. This shows that the middle vertical map is surjective. This proves the result because (3) \Rightarrow (4).

(2) \Rightarrow (1): Assume that $\pi_0(G)$ is finite. Then without loss of generality, we may assume that G is connected. The morphism $[p]$ is étale by [11, Lemme 1]. Since (2) \Rightarrow (4), we know that $G[p]$ is finite. Hence (1) follows from [11, Lemme 5].

Assume now that G is quasi-compact and connected. Then:

(4) \Rightarrow (2): We use Lütkebohmert’s structure result [25, Theorem I]. This says that there is an extension $T \rightarrow E \rightarrow B$ of a quasi-compact group of good reduction B by a rigid torus T such that $G = E/M$ for some lattice $M \subseteq E$. We will use this to reduce to the case of good reduction. Namely, it is clear that $[p] : G \rightarrow G$ is surjective if and only if $[p] : B \rightarrow B$ is surjective. On the other hand, we have

$$\widehat{G} = \widehat{E} \times (M \otimes \mathbb{Q}_p/\mathbb{Z}_p).$$

We deduce that $\widehat{G}^\circ = \widehat{E}^\circ$, and this is itself an extension

$$0 \rightarrow \widehat{T} \rightarrow \widehat{E}^\circ \rightarrow \widehat{B}^\circ \rightarrow 0.$$

We deduce that $[p] : \widehat{B}^\circ \rightarrow \widehat{B}^\circ$ is surjective if $[p] : \widehat{G}^\circ \rightarrow \widehat{G}^\circ$ is surjective.

We have thus reduced to the case that G has good reduction. The reduction \overline{G} is then a commutative connected algebraic group, hence an extension of an abelian variety by a product of tori and additive groups. We need to see that the additive part vanishes, then $[p] : \overline{G} \rightarrow \overline{G}$ is finite surjective, so the same is true for $[p] : G \rightarrow G$ by [4, Section 6.3.5, Theorem 1].

Assume for a contradiction that \overline{G} contains a subgroup isomorphic to \mathbb{G}_a . By Proposition 6.5(4), \widehat{G}° is the analytic p -divisible group associated with the connected part of the p -divisible group of G over \mathcal{O}_K . Consequently, we have a morphism of short exact sequences

$$\begin{array}{ccccccc} 1 & \longrightarrow & \widehat{G}^\circ(K) & \longrightarrow & G(K) & \longrightarrow & \overline{G}(k) \longrightarrow 1 \\ & & [p] \uparrow & & [p] \uparrow & & [p] \uparrow \\ 1 & \longrightarrow & \widehat{G}^\circ(K) & \longrightarrow & G(K) & \longrightarrow & \overline{G}(k) \longrightarrow 1. \end{array}$$

If \overline{G} contains a factor of \mathbb{G}_a , then the kernel of $[p] : \overline{G}(k) \rightarrow \overline{G}(k)$ contains a copy of k . As the kernel of the middle map is finite, this means by the Snake Lemma that the cokernel of the first map is non-trivial. This contradicts the assumption that \widehat{G}° is p -divisible. ■

6.3 Exponentials for G induced by exponentials for \mathbb{G}_m

We can now prove the main result of this section. Namely, a choice of an exponential $\exp : K \rightarrow 1 + \mathfrak{m}_K$ for the group \mathbb{G}_m induces compatible choices of exponentials for all locally p -divisible rigid groups G .

Theorem 6.12 *Let K be a complete algebraically closed extension of \mathbb{Q}_p and let G be a locally p -divisible commutative rigid group over K . Then the choice of an exponential $\exp : K \rightarrow 1 + \mathfrak{m}_K$ for \mathbb{G}_m induces a continuous homomorphism*

$$\exp_G : \text{Lie } G \rightarrow \widehat{G}(K)$$

splitting \log_G from Proposition 6.5(2). The association $G \mapsto \exp_G$ is natural in G .

This generalizes a result of Faltings, who considered the case of commutative algebraic groups [10, p. 856]. The latter are locally p -divisible groups by Example 6.9.

Proof By Proposition 6.10, the group \widehat{G}° is an analytic p -divisible group in the sense of Fargues [11]. By [11, Théorème 3.3], there is then a natural Cartesian diagram of rigid groups

$$\begin{array}{ccc} \widehat{G}^\circ & \longrightarrow & \underline{\mathrm{Hom}}(\Lambda(1), \mathbb{G}_m) \\ \log \downarrow & & \log \downarrow \begin{array}{c} \curvearrowright \\ \exp \end{array} \\ \mathfrak{g} & \longrightarrow & \underline{\mathrm{Hom}}(\Lambda(1), \mathbb{G}_a), \end{array}$$

functorial in G , where $\Lambda := T_p(\widehat{G}^\circ)^\vee$. With this at hand, we can argue as in [43, Proposition A.1.2]: The chosen exponential induces a section of the right map on K -points, thus we obtain a section of the left map on K -points by Cartesianness. ■

7 The Corlette–Simpson correspondence for commutative G

7.1 Recollections on the Hodge–Tate short exact sequence for \mathbb{G}_a and \mathbb{G}_m

Assume that X is a proper, smooth rigid space over K . Set $\widetilde{\Omega} = R^1 v_* \mathcal{O} \cong \Omega_X^1(-1)$. We fix a base point $x_0 \in X(K)$. In this situation, we have by [32, Section 3] the Hodge–Tate sequence

$$(12) \quad 0 \rightarrow H_{\text{ét}}^1(X, \mathbb{G}_a) \rightarrow H_v^1(X, \mathbb{G}_a) \rightarrow H^0(X, \widetilde{\Omega}) \rightarrow 0$$

in which the middle term can be naturally identified as $H_v^1(X, \mathbb{G}_a) = \mathrm{Hom}(\pi_1^{\text{ét}}(X, x_0), K)$ via a Cartan–Leray spectral sequence for the universal cover $\widetilde{X} \rightarrow X$ (c.f. [16, Proposition 2.8.1, Proposition 4.9]). As we now recall, in general, the sequence doesn’t split canonically. However, a splitting can be functorially induced by an additional choice.

Definition 7.1 Let X be a smooth proper rigid space over K . By a (flat) B_{dR}^+/ξ^2 -lift of X we mean a flat morphism of adic spaces $\mathbb{X} \rightarrow \mathrm{Spa}(B_{\text{dR}}^+/\xi^2)$ with an isomorphism

$$\mathbb{X} \times_{\mathrm{Spa}(B_{\text{dR}}^+/\xi^2)} \mathrm{Spa}(K) \xrightarrow{\sim} X.$$

We note that such a datum always exists. There is a canonical choice if X admits a model over a discretely valued subfield $K_0 \subseteq K$ with perfect residue field.

Proposition 7.2 ([12, Proposition 7.2.5]) Any choice of a B_{dR}^+/ξ^2 -lift \mathbb{X} of X induces a splitting of (12). This splitting is functorial in (X, \mathbb{X}) .

7.2 General commutative G

There is an analogous story when we replace \mathbb{G}_a by the group \mathbb{G}_m . By the main result of [16], there is a short exact sequence

$$(13) \quad 0 \rightarrow H_{\text{ét}}^1(X, \mathbb{G}_m) \rightarrow H_v^1(X, \mathbb{G}_m) \rightarrow H^0(X, \widetilde{\Omega}) \rightarrow 0$$

classifying the discrepancy between v -line bundles and étale line bundles in terms of the additional datum of a Higgs field. This time, the splitting requires not only a B_{dR}^+/ξ^2 -lift \mathbb{X} , but also the datum of an exponential \exp as in Definition 6.1, providing a continuation of the Lie group exponential for \mathbb{G}_m .

We now prove an analog of the short exact sequence (13) for general commutative rigid groups, thus giving a generalization of the Hodge–Tate sequence (12) and the multiplicative Hodge–Tate sequence (13) to a large class of commutative rigid groups.

Theorem 7.3 *Let X be a connected smooth proper rigid space with a base point $x_0 \in X(K)$. Let G be a commutative, locally p -divisible rigid group (see Definition 6.8) over K .*

(1) *The Leray spectral sequence for $v : X_v \rightarrow X_{\text{ét}}$ applied to the sheaf G induces a short exact sequence of abelian groups, natural in G ,*

$$(14) \quad 0 \rightarrow H_{\text{ét}}^1(X, G) \rightarrow H_v^1(X, G) \xrightarrow{\text{HTlog}} H^0(X, \tilde{\Omega}) \otimes_K \text{Lie } G \rightarrow 0.$$

(2) *Choices of a B_{dR}^+/ξ^2 -lift of X (Definition 7.1) and an exponential for K (Definition 6.1) induce a natural splitting s_G of HTlog. This is functorial in the following sense: For any $\theta \in H^0(X, \tilde{\Omega}) \otimes_K \text{Lie } G$, these choices induce a v - G -torsor V_θ on X with $\text{HTlog}(V_\theta) = \theta$, and for any homomorphism of commutative locally p -divisible rigid groups $f : G \rightarrow H$ a natural isomorphism*

$$V_\theta \times^G H \xrightarrow{\sim} V_{(\text{Lie } f)(\theta)}.$$

Proof By [17, Corollary 4.3], the Leray spectral sequence for $v : X_v \rightarrow X_{\text{ét}}$ applied to G gives a left-exact sequence

$$0 \rightarrow H_{\text{ét}}^1(X, G) \rightarrow H_v^1(X, G) \rightarrow H^0(X, \tilde{\Omega}) \otimes_K \text{Lie } G.$$

In Theorem 5.1, we have seen that every homomorphism $\rho : \pi_1^{\text{ét}}(X, x_0) \rightarrow G(K)$ induces a class in $H_v^1(X, G)$. Since $\pi_1^{\text{ét}}(X, x_0)$ is profinite, it follows from [19, Proposition 2.14] that the image of ρ is contained in $G^{\text{tt}}(K)$. Composition with the logarithm map $\log : G^{\text{tt}} \rightarrow \mathfrak{g}$ from Lemma 6.7 induces a commutative diagram

$$\begin{array}{ccc} \text{Hom}(\pi_1^{\text{ét}}(X, x_0), G^{\text{tt}}(K)) & \xrightarrow{\log} & \text{Hom}(\pi_1^{\text{ét}}(X, x_0), \text{Lie } G) \\ \downarrow & & \downarrow \text{HT} \\ H_v^1(X, G) & \xrightarrow{\text{HTlog}} & H^0(X, \tilde{\Omega}) \otimes_K \text{Lie } G \end{array}$$

where the right vertical map is induced by applying $- \otimes \text{Lie } G$ to

$$\text{HT} : \text{Hom}(\pi_1^{\text{ét}}(X, x_0), K) \xrightarrow{\sim} H_v^1(X, \mathcal{O}) \rightarrow H^0(X, \tilde{\Omega}).$$

Here the first identification comes from the Cartan–Leray sequence for \mathcal{O} along $\tilde{X} \rightarrow X$.

Now the choice of a B_{dR}^+/ξ^2 -lift of X induces a splitting of the Hodge–Tate sequence (Proposition 7.2) and hence a splitting of the right vertical map. By Theorem 6.12, an exponential for K induces an exponential map $\exp_G : \text{Lie } G \rightarrow \widehat{G}(K)$ that splits the top horizontal map. In particular, both maps are split surjective. Thus HTlog is also split surjective, with a splitting induced by the above choices.

The naturality in G is then immediate from the naturality of \exp_G in Theorem 6.12, and from the fact that the left vertical map is given via Theorem 5.1 by the functorial construction of sending ρ to the pushout V_ρ of ρ along the $\pi_1^{\text{ét}}(X, x_0)$ -torsor $\tilde{X} \rightarrow X$. ■

We can deduce our final main result, the p -adic Simpson correspondence for commutative G , giving answers to each of Questions 1.1, 1.2, 1.3 for commutative G .

Theorem 7.4 *Let X be a connected smooth proper rigid variety over K with a base point $x_0 \in X(K)$, and let G be a commutative locally p -divisible rigid group over K .*

- (1) *Choices of an exponential \exp for K and a B_{dR}^+/ξ^2 -lift \mathbb{X} of X induce an equivalence*

$$\{v\text{-}G\text{-torsors on } X\} \xrightarrow{\sim} \{G\text{-Higgs bundles on } X\}$$

of groupoids that is natural in G and in (X, \mathbb{X}) .

- (2) *In case that every morphism $\tilde{X} \rightarrow G$ is constant (for example if G is linear analytic), this restricts to an equivalence of groupoids*

$$\left\{ \begin{array}{l} \text{continuous homomorphisms} \\ \pi_1^{\text{ét}}(X, x_0) \rightarrow G(K) \end{array} \right\} \xrightarrow{\sim} \left\{ \begin{array}{l} \text{pro-finite-étale} \\ G\text{-Higgs bundles on } X \end{array} \right\}.$$

Proof For simplicity, we write π for $\pi_1^{\text{ét}}(X, x_0)$ below.

For (1), we observe that it suffices to construct a functor from left to right that induces a bijection on isomorphism classes. Indeed, note that since G is commutative, the automorphism group of an object on either side is $G(X) = H_v^0(X, G) = H_{\text{ét}}^0(X, G)$ by Lemma 2.20. By Theorem 7.3.(2), the choices of an exponential for K and a B_{dR}^+/ξ^2 -lift \mathbb{X} of X induce a section s of the map HTlog in (14), and hence an isomorphism

$$H_v^1(X, G) \xrightarrow{\sim} H_{\text{ét}}^1(X, G) \times H^0(X, \tilde{\Omega}^1) \otimes_K \text{Lie } G,$$

which moreover can be upgraded to a functorial construction: For any v - G -torsor V , consider its class in $H_v^1(X, G)$. Applying HTlog , we obtain $\theta \in H^0(X, \tilde{\Omega}^1) \otimes_K \text{Lie } G$. Theorem 7.3.(2) uses the choice of lift and exponential to give us a v - G -bundle V_θ on X with $\text{HTlog}(V_\theta) = \theta = \text{HTlog}(V)$ in a functorial way. Now let E be the v - G bundle $V \otimes^G V_\theta^{-1}$, c.f. Definition 2.21. Since HTlog is additive with respect to contracted products, we see that $\text{HTlog}(E) = 0$. Consequently, E is an étale G -torsor. The desired functor can now be defined by sending V to (E, θ) . Here we note that since G is commutative, the Higgs field condition is vacuous by Lemma 2.20.

This construction is indeed functorial. Any morphism $V \rightarrow V'$ is an isomorphism, hence $\text{HTlog}(V) = \text{HTlog}(V')$ which implies that V_θ agrees for V and V' . So the statement follows from functoriality of $-\otimes^G V_\theta^{-1}$.

The naturality in G follows from naturality in G in Theorem 7.3(2).

It remains to check the naturality in (1) with respect to maps $X \rightarrow Y$ between smooth proper rigid spaces with a lift $f: \mathbb{X} \rightarrow \mathbb{Y}$ over B_{dR}^+/ξ^2 . But this follows immediately from functoriality of the splitting of the Hodge–Tate sequence for \mathbb{G}_a induced by f , see [12, Proposition 7.2.5].

For (2), we use Theorem 5.1 which says that continuous homomorphisms $\rho: \pi \rightarrow G^{\text{tt}}(K)$ are equivalent to pro-finite-étale G -bundles on X_v . Note that by definition, the v - G -torsor V_θ is pro-finite-étale for every $\theta \in H^0(X, \tilde{\Omega}^1) \otimes_K \text{Lie } G$. Hence a v - G -torsor V on X with image θ under HTlog is pro-finite-étale if and only if $V \otimes^G V_\theta^{-1}$ is pro-finite-étale. Applying this to the construction in (1), this implies our claim. ■

Acknowledgements This project originated from discussions the second and the third author had during the workshop *Women in Numbers Europe 4* (Utrecht 2022). We thank the organizers and sponsors of this meeting. Special thanks go to Wiesława Nizioł who coorganized with the second author the research group on “Relative *p*-adic Hodge Theory,” and to the participants of this research group. We would like to thank Lucas Gerth and the anonymous referee for helpful comments on an earlier version. Moreover, we thank Johannes Anschütz, Ian Gleason, Arthur-César Le Bras and Daxin Xu for helpful conversations.

References

- [1] A. Abbes, M. Gros, and T. Tsuji, *The p-adic Simpson correspondence*, 193, Princeton University Press, Princeton, NJ, 2016.
- [2] J. Anschütz, B. Heuer, and A.-C. Le Bras, *Hodge-Tate stacks and non-abelian p-adic Hodge theory of v-perfect complexes on smooth rigid spaces*. Preprint, 2023. [arXiv:2302.12747](https://arxiv.org/abs/2302.12747).
- [3] C. Blakestad, D. Gvirtz, B. Heuer, D. Shchedrina, K. Shimizu, P. Wear, and Z. Yao, *Perfectoid covers of abelian varieties*. *Math. Res. Lett.* 29(2022), no. 3, 631–662.
- [4] S. Bosch, U. Güntzer, and R. Remmert, *Non-Archimedean analysis, Grundlehren der Mathematischen Wissenschaften*, 261, Springer-Verlag, Berlin, 1984.
- [5] S. Bosch and W. Lütkebohmert, *Formal and rigid geometry. II. Flattening techniques*. *Math. Ann.* 296(1993), no. 3, 403–429.
- [6] B. Conrad, *Irreducible components of rigid spaces*. *Ann. Inst. Fourier (Grenoble)* 49(1999), no. 2, 473–541.
- [7] A. J. de Jong, *Étale fundamental groups of non-Archimedean analytic spaces*. *Compos. Math.* 97(1995), nos. 1–2, 89–118, Special issue in honour of Frans Oort.
- [8] A. J. de Jong, *The stacks project*, 2024. Available at <http://stacks.math.columbia.edu>.
- [9] C. Deninger and A. Werner, *Vector bundles on p-adic curves and parallel transport*. *Ann. Sci. École Norm. Sup. (4)* 38(2005), no. 4, 553–597.
- [10] G. Faltings, *A p-adic Simpson correspondence*. *Adv. Math.* 198(2005), no. 2, 847–862.
- [11] L. Fargues, *Groupes analytiques rigides p-divisibles*. *Math. Ann.* 374(2019), no. 1–2, 723–791.
- [12] H. Guo, *Hodge–Tate decomposition for non-smooth spaces*. *J. Eur. Math. Soc. (JEMS)*, 25(2023), no. 4, 1553–1625.
- [13] U. Hackstein, *Principal bundles on p-adic curves and parallel transport*. *Int. Math. Res. Not. IMRN* 050(2008), 39.
- [14] B. Heuer, *Pro-étale uniformisation of abelian varieties*. Preprint, 2021. [arXiv:2105.12604](https://arxiv.org/abs/2105.12604).
- [15] B. Heuer, *G-torsors on perfectoid spaces*. Preprint, 2022. [arXiv:2207.07623](https://arxiv.org/abs/2207.07623).
- [16] B. Heuer, *Line bundles on rigid spaces in the v-topology*. *Forum Math. Sigma* 10(2022), 1–36.
- [17] B. Heuer, *Moduli spaces in p-adic non-abelian Hodge theory*. Preprint, 2022. [arXiv:2207.13819](https://arxiv.org/abs/2207.13819).
- [18] B. Heuer, *A p-adic Simpson correspondence for smooth proper rigid varieties*. Preprint, 2023. [arXiv:2307.01303](https://arxiv.org/abs/2307.01303).
- [19] B. Heuer, *A geometric p-adic Simpson correspondence in rank one*. *Compos. Math.* 160(2024), no. 7, 1433–1466.
- [20] B. Heuer, L. Mann, and A. Werner, *The p-adic Corlette-Simpson correspondence for abeloids*. *Math. Ann.* 385(2023), no. 3–4, 1639–1676.
- [21] R. Huber, *Bewertungsspektrum und rigide Geometrie, Regensburger Mathematische Schriften [Regensburg Mathematical Publications]*, 23, Universität Regensburg, Fachbereich Mathematik, Regensburg, 1993.
- [22] R. Huber, *Étale cohomology of rigid analytic varieties and adic spaces, Aspects of Mathematics*, 30, Springer Fachmedien, Wiesbaden, 1996.
- [23] K. S. Kedlaya and R. Liu, *Relative p-adic Hodge theory, II: Imperfect period rings*. Preprint, 2016. [arXiv:1602.06899](https://arxiv.org/abs/1602.06899).
- [24] R. Liu and X. Zhu, *Rigidity and a Riemann-Hilbert correspondence for p-adic local systems*. *Invent. Math.* 207(2017), no. 1, 291–343.
- [25] W. Lütkebohmert, *The structure of proper rigid groups*. *J. Reine Angew. Math.* 468(1995), 167–219.

- [26] W. Lütkebohmert, *Rigid geometry of curves and their Jacobians*, *Ergebnisse der Mathematik und ihre Grenzgebiete*, 61, Springer, Cham, 2016.
- [27] Y. Min and Y. Wang, *p -adic Simpson correspondence via prismatic crystals*. Preprint, 2022. [arXiv:2201.08030](https://arxiv.org/abs/2201.08030).
- [28] Y. Min and Y. Wang, *Integral p -adic non-abelian hodge theory for small representations*. *Adv. Math.* 458(2024), 109950.
- [29] J. Neukirch, A. Schmidt, and K. Wingberg, *Cohomology of number fields*, Grundlehren der Mathematischen Wissenschaften, 323, Springer-Verlag, Berlin, second edition, 2008.
- [30] R. Palais and T. Stewart, *Torus bundles over a torus*. *Proc. Amer. Math. Soc.* 12(1961), 26–29.
- [31] P. Scholze, *p -adic Hodge theory for rigid-analytic varieties*. *Forum Math. Pi* 1:e1, (2013) 77.
- [32] P. Scholze, *Perfectoid spaces: A survey*. In *Current developments in mathematics 2012*, International Press, Somerville, MA, 2013, pp. 193–227.
- [33] P. Scholze, *On torsion in the cohomology of locally symmetric varieties*. *Ann. of Math.* (2) 182(2015), no. 3, 945–1066.
- [34] P. Scholze, *Étale cohomology of diamonds*. Preprint, 2018. [arXiv:1709.07343](https://arxiv.org/abs/1709.07343).
- [35] P. Scholze and J. Weinstein, *Moduli of p -divisible groups*. *Camb. J. Math.* 1(2013), no. 2, 145–237.
- [36] P. Scholze and J. Weinstein, *Berkeley Lectures on p -adic geometry*, *Annals of Mathematics Studies*, 207, Princeton University Press, Princeton, NJ, 2020.
- [37] J.-P. Serre, *Espaces fibrés algébriques*. *Séminaire Claude Chevalley*. 3(1958), 9–37.
- [38] J.-P. Serre, *Cohomologie galoisienne*, *Lecture Notes in Mathematics*, 5, Springer-Verlag, Berlin, fifth edition, 1994.
- [39] C. T. Simpson, *Higgs bundles and local systems*. *Publ. math. I.H.É.S.* 75(1992), 5–95.
- [40] M. Temkin, *On local properties of non-Archimedean analytic spaces*. *Math. Ann.* 318(2000), no. 3, 585–607.
- [41] T. Tsuji, *Notes on the local p -adic Simpson correspondence*. *Math. Ann.* 371(2018), no. 1-2, 795–881.
- [42] Y. Wang, *A p -adic Simpson correspondence for rigid analytic varieties*. *Algebra Number Theory*, 17(2023), no. 8, 1453–1499.
- [43] D. Xu, *Parallel transport for Higgs bundles over p -adic curves*. Preprint, 2022. [arXiv:2201.06697](https://arxiv.org/abs/2201.06697).

Institut für Mathematik, Goethe-Universität Frankfurt, Robert-Mayer-Str. 6-8, 60325 Frankfurt am Main, Germany

e-mail: werner@math.uni-frankfurt.de heuer@math.uni-frankfurt.de

Princeton University, Fine Hall, Princeton, USA

e-mail: mz9413@princeton.edu