Perhaps it should be stressed that, although it was mathematically correct to refer to S₂ as a concentration, in the absence of a measure of the size of compartment 2, S_2 did not represent the true concentration of the amino acid in this space. S_2 could be considered as a comparative value referenced back to the tryptophan in unit volume of plasma and it was expressed in terms of unit volume.

We thank Sister I. Hockley, Mrs D. Barrett, the nursing staff of East 1 Ward, Whitchurch Hospital, and the medical staffs of Whitchurch Hospital and the Department of Psychological Medicine, Heath Hospital, Cardiff, for their help throughout this study. We would also like to thank the Department of Medical Physics, Heath Hospital, for preparing injections of the radio-isotope and relatives, friends, members of staff of the hospital, the Police Force, the Inland Revenue and local firms for volunteering for the control studies. The Biochemical Psychiatry Laboratory, Whitchurch Hospital, is supported by an MRC Programme Grant.

ERRATA

The Appendix of our previous paper, Shaw et al. (1975),contained the following typographical errors:

(a) Equation (2):

$$\dot{R}_2(t) = S_2 \cdot \dot{a}_2(t) = \lambda_{21} \cdot R_1(t) - \lambda_{02} \cdot R_2(t)$$

This should read:

$$\dot{R}_2(t) = S_2 \cdot \dot{a}_2(t) = \lambda_{21} \cdot R_1(t) - \lambda_{02} \cdot R_2(t) - \lambda_{12} \cdot R_2(t)$$

(b) Equation at bottom of page 213:

$$R_1(0) = S_1 \cdot a_1(0) = S_1(A \div B).$$

This should read:

$$R_1(0) = S_1 \cdot a_1(0) = S_1(A+B)$$
.

N.B. λ_{02} and ρ_{02} in the previous paper are equivalent to λ_{32} and ρ_{32} in this paper.

(c) The specific activity of L-[3H]tryptophan should have been given as 0.5-1 Ci/mMol not 0.5-1 μ Ci/ μ Mol.

REFERENCES

Altschule, M. D., Ascoli, I. & Tillotson, K. J. (1949). Extracellular fluid and plasma volumes in depressed patients given electric shock therapy. Archives of Neurology and Psychiatry (Chicago) 62, 618-623.

Blazek, R., Tidmarsh, S. F., Shaw, D. M. & Riley, G. J.

(1978). In preparation.

Bridges, P. K., Bartlett, J. R., Sepping, P., Kantameneni, B. D. & Curzon, G. (1976). Precursors and metabolites of 5-hydroxytryptamine and dopamine in the ventricular cerebrospinal fluid of psychiatric patients. Psychological Medicine 6, 399-405.

Coppen, A. & Shaw, D. M. (1963). Mineral metabolism in melancholia. British Medical Journal ii, 1439-1444.

Coppen, A., Shaw, D. M. & Farrell, J. P. (1963). Potentiation of the antidepressive effect of a monoamine oxidase inhibitor by tryptophan. Lancet i, 79-81.

Coppen, A., Shaw, D. M., Malleson, A. & Costain, R. (1966). Mineral metabolism in mania. British Medical Journal i,

71-75.

Dawson, J., Hullin, R. P. & Crocket, B. M. (1956). Metabolic variations in manic-depressive psychosis. Journal of Mental Science 102, 168-177.

Greenwood, M., Lader, M., Kantameneni, B. D. & Curzon, G. (1978). The effect of mental arithmetic under stress on tryptophan disposition in psychiatric patients. Submitted for publication.

Hullin, R. P., Bailey, A. D., McDonald, R., Dransfield, G. A. & Milne, H. B. (1967a). Variations in body water during recovery from depression. British Journal of

Psychiatry 113, 573-583.

Hullin, R. P., Bailey, A. D., McDonald, R., Dransfield, G. A. & Milne, H. B. (1967b). Body water variations in manic-depressive psychosis. British Journal of Psychiatry 113, 584-592.

Landis, E. M. & Pappenheimer, J. R. (1963). Exchange of substances though the capillary walls. In Handbook of Physiology, Section 2. Circulation, vol. 11 (ed. W. F. Hamilton), pp. 961-1034. American Physiological Society: Washington.

Riley, G. J. & Shaw, D. M. (1976). Total and non-bound tryptophan in unipolar illness. Lancet ii, 1249.

Riley, G. J. & Shaw, D. M. (1978). In preparation.

Shaw, D. M., Johnson, A. L. & Short, R. (1972). Preliminary studies of the effects of lithium, imipramine and reserpine on the relative sizes of tryptophan pools and flow constants in rabbits. Mathematical Biosciences 15, 137-151.

Shaw, D. M., Johnson, A. L., Tidmarsh, S. F., MacSweeney, D. A., Hewland, H. R. & Woolcock, N. E. (1975). Multicompartmental analysis of amino acids. I. Preliminary data on concentrations, fluxes and flow constants of tryptophan in affective illness. Psychological Medicine 5, 206-213.

Shaw, D. M., Riley, G. J., Tidmarsh, S. & Blazek, R. (1976). Unipolar affective illness. Lancet i, 363-364.

Snedecor, G. W. & Cochran, W. G. (1967). Statistical Methods, 6th edn, pp. 472-503. Iowa State University Press: Ames, Iowa.