



Gastrointestinal helminths of opossums (Mammalia: Didelphidae) from Bolivia

F. Agustín Jiménez¹ , Mariel L. Campbell², Beth Byles¹,
Raymond Philip Scheibel¹ and Scott L. Gardner³ 

Research Article

Cite this article: Jiménez FA, Campbell ML, Byles B, Scheibel RP, Gardner SL (2024). Gastrointestinal helminths of opossums (Mammalia: Didelphidae) from Bolivia. *Parasitology* 1–13. <https://doi.org/10.1017/S0031182024000490>

Received: 14 December 2023

Revised: 9 March 2024

Accepted: 16 April 2024

Keywords:

biological inventories; Bolivia; mouse opossums; parasite collections; taxonomic impediment

Corresponding author:

F. Agustín Jiménez;

Email: agustinjz@zoology.siu.edu

¹School of Biological Sciences, Zoology, Southern Illinois University, Carbondale, IL, USA; ²Museum of Southwestern Biology, 1 University of New Mexico, Albuquerque, NM, USA and ³The Harold W. Manter Laboratory of Parasitology, University of Nebraska, Lincoln, NE, USA

Abstract

A total of 32 taxa of helminths were recovered from 52 individuals corresponding to 17 species of didelphiomorph marsupials collected across Bolivia. From these, 20 taxa are registered for the first time in this landlocked South American country, including the cestode *Mathevotaenia bivittata*, and the nematodes *Moennigia* sp., *Travassostrongylus callis*, *Viannaia didelphis*, *V. hamata*, *V. metachirops*, *V. minispicula*, *V. philanderi*, *V. simplicispicula*, *V. skrjabini*, *V. viannai*, *Cruzia tentaculata*, *Monodelphoxyuris dollmeiri*, *Neohilgertia venusti*, *Pterygodermatites elegans*, *Pterygodermatites jeagerskioldi*, *Spirura guianensis*, *Gongylonemoides marsupialis*, *Turgida turgida* and *Trichuris reesali*. We report for the first time parasites for *Marmosops bishopi*, *Monodelphis emiliae*, *Monodelphis glirina*, *Monodelphis sanctarosae*, *Monodelphis peruviana* and *Thylamys sponsorius* and document 38 new records of parasites infecting marsupials. Twenty-six taxa of helminths infect 2 or more species of didelphiomorph marsupials, with the exception of *Travassostrongylus callis*, *Viannaia didelphis*, *V. hamata*, *V. minispicula* and *V. hamate*, which infected individuals of a single species.

Introduction

Didelphimorphia (opossums) is the most diverse order of marsupials present in the New World with over 125 extant species (Gardner, 2007; Teta *et al.*, 2009; Voss and Jansa, 2009; Gutiérrez *et al.*, 2010; Rossi *et al.*, 2010; Jansa *et al.*, 2014; Voss, 2022). This group includes arboreal, terrestrial, semiaquatic and scansorial representatives (Astúa, 2009; Flores, 2009; Voss and Jansa, 2009). Most of them occur in moderate densities in non-disturbed forests throughout the Neotropics (Lima *et al.*, 2001; Gentile *et al.*, 2004; Püttker *et al.*, 2008; Pires *et al.*, 2010). Several aspects of their ecology remain unknown, perhaps, because a large number of species is arboreal making their collection infrequent (Fontúrbel and Jiménez, 2009; Caceres *et al.*, 2011; José *et al.*, 2019). The territory of Bolivia sits at the juncture of high-altitude deserts, mountainous, temperate, tropical forest and lowland savannah biomes. This results in a large diversity of mammals that includes 24% of the extant species of opossums (Anderson, 1997; Gutiérrez *et al.*, 2010; Rossi *et al.*, 2010; Voss *et al.*, 2012). This diversity includes representatives of all 4 subfamilies and major clades in the Didelphidae Gray and species that inhabit humid and dry forests (Voss and Jansa, 2009; Jansa *et al.*, 2014; Voss, 2022).

From 1984 to 2000, the American Museum of Natural History (AMNH, New York City, USA), the Mammal Division of the Museum of Southwestern Biology (MSB:Mamm, University of New Mexico, Albuquerque New Mexico, USA), the Harold W. Manter Laboratory of Parasitology (HWML, University of Nebraska-Lincoln, Lincoln Nebraska, USA), and the Bolivian National Museum of Natural History in La Paz (Colección Nacional de Fauna Sección Mastozoología, CBF, La Paz, Bolivia) mounted joint US National Science Foundation funded collecting expeditions throughout Bolivia to survey and inventory sylvatic mammals and their parasites. A major part of the work on the mammals has been published by Anderson (1997) and many groups of parasites that were collected from these mammals are still being studied in earnest. From most of the mammals that were collected by these expeditionary research teams in Bolivia, data on habitat, habits and biological associates were also collected and archived in museums. All of the parasites are on deposit in the HWML within the Bolivian Mammal Parasite Collection (BMPC). The BMPC includes all specimens of helminths and parasitic arthropods that were recovered from the more than 16 000 mammals collected and preserved in museums during the course of our work.

Species of helminths that occur in opossums can be grouped in families that include opossum dwelling species, such as Rhopalidae (Radev *et al.*, 2005; Haverkost and Gardner, 2008) or in families that include species occurring in distantly related groups of mammals, i.e. Aspidoderidae, Onchocercidae, Viannidae (Brant and Gardner, 2000; Jiménez *et al.*, 2012; Scheibel *et al.*, 2014). However, the establishment of their specificity or host range is difficult to assess without the existence of a database that relates the distribution of parasite species

© The Author(s), 2024. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

across several species of mammals. Herein, we present the recorded species of gastrointestinal helminths infecting opossums in Bolivia.

Materials and methods

All mammals were collected using Sherman™ live traps baited with a mixture of oatmeal, vanilla, tuna and sardines, or with snap traps baited with peanut butter. Traps were placed in suitable habitat each evening and checked at first daylight the following morning. Details of each mammal collected were recorded in a field-collection catalog book and in the trap data book, copies of which are maintained in the HWML, the originals are in the Department of Mammalogy, AMNH. Mammal voucher specimens are deposited in the AMNH, MSB:Mamm and CBF.

In the field, each organ of the digestive system was examined separately. Platyhelminths found were placed in distilled water until they relaxed and were killed and fixed in either 70% EtOH or hot or cold 10% formalin. Nematodes were either placed directly in 70% ethanol or killed with glacial acetic acid then transferred to either 70% ethanol or 10% formalin solution. Some samples were preserved in 95% ethanol or in liquid nitrogen and then stored in -85°C freezers in the Manter Laboratory Parasite Genomic Research Facility.

Digenetic trematodes, cestodes and acanthocephalans were stained in Semichon's acetocarmine, dehydrated in a graded series of ethanol, cleared in xylene and mounted in Canada balsam or Damar Gum. Nematodes were cleared in lactophenol and mounted on temporary slides. Vouchers for this study were deposited in the HWML. Specimens used for comparison were borrowed from the HWML and 7 additional institutions including:

CHIOC: *Coleção Helminológica do Instituto Oswaldo Cruz*, Oswaldo Cruz Institute, Rio de Janeiro Brazil.

CHLP: Collection of Helminths of the Division of Invertebrates of the Museum of Natural History of the National University of La Plata.

CMNA: Canadian Museum of Nature Parasite Collection, Ottawa, Canada

CNHE: Colección Nacional de Helmintos of the National Autonomous University of Mexico, Mexico City.

CHIAUMSM: Colección de Helmintos e Invertebrados Afines, Museo de Historia Natural, Universidad Nacional Mayor de San Marcos, Lima.

UCDNC: University of California Davis Nematode Collection, University of California, Davis, USA.

USNPC: United States National Parasite Collection of the Smithsonian Institution. Washington D.C. U.S.A.

The list follows current systems of classification (Radev *et al.*, 2005; Beveridge *et al.*, 2014; Mariaux *et al.*, 2017; Hodda, 2022). The helminthological record for most of the marsupials examined is available at http://opensiuc.lib.siu.edu/zool_data/23/. The relation of voucher specimens, numbers and collections is presented below.

Results

A total of 32 taxa of helminths were recovered from 17 species of marsupials collected from 23 localities. The total includes 3 species of digenetic trematodes, 6 species of tapeworms, 22 species of nematodes, 2 of which remain unidentified and an acanthocephalan. The association of parasites and their didelphiomorph hosts is detailed in Tables 1–3. We present this association using the tribe of the mammals.

Phylum Platyhelminthes Gegenbaur, 1859

Class Trematoda Rudolphi, 1808

Order Diplostomida Olson, Cribb, Tkach, Bray and Littlewood, 2003

Superfamily Echinostomoidea Looss, 1902

Family Rhopalidae Looss, 1899

Rhopalias Stiles and Hassall, 1898

1. *Rhopalias caballeroi* Kifune and Uyema, 1982

Site of infection: Small intestine

Type host and locality: *Didelphis marsupialis* L., Huanuco, Peru

Other reported hosts: *Chironectes minimus* (Zimmermann), *Didelphis* sp., *Philander opossum* (L.), *Lutreolina crassicaudata* (Desmarest)

Locality records: Argentina: Buenos Aires, Berisso. Colombia: undetermined. Mexico: Veracruz: Los Tuxtlas. Panama: Panama Canal. Peru: Cusco: Pilcopata; San Martín: Bella Vista; Llamas. Venezuela: Aragua (Tantaleán and Chavez, 2004; Haverkost and Gardner, 2008; Chero *et al.*, 2017)

Records in Bolivia: *Chironectes minimus*: La Paz: La Reserva, $15^{\circ}44'S$, $67^{\circ}31'W$, 850 m, 22 July 1992, HWML70021 (274 specimens) from MSB:MAMM:68330. *Didelphis marsupialis*: Santa Cruz: San Rafael de Amboró, $17^{\circ}21'S$, $63^{\circ}43'W$, 400 m, 24 July 1985, HWML70025 (3 specimens) from MSB:MAMM:55833. *Philander opossum*: Santa Cruz: San Miguel del Rincón, $17^{\circ}23'S$, $63^{\circ}32'W$, 300 m; 13 August 1984, HWML70018 (3 specimens) from MSB:MAMM:55074; Estancia Cachuela Esperanza, $16^{\circ}46'59.99''S$, $63^{\circ}13'59.99''W$, 300 m, 22 August 1984, HWML70021 (18 specimens) from MSB:MAMM:210569.

Additional specimens examined: HWML70014 (1 specimen) from *Lutreolina crassicaudata*, Berisso, Buenos Aires, Argentina; CNHE 4081 (1 specimen) from *Didelphis* sp., Catemaco, Veracruz, Mexico; CNHE965 (1 specimen) from *Didelphis marsupialis*, Aragua, Venezuela.

2. *Rhopalias coronatus* (Rudolphi, 1819) Stiles and Hassall, 1898

Synonyms: *Rhopalias dobbini* Prod'Hon, 1968

Site of infection: Small intestine

Type host and locality: *Didelphis marsupialis*, Brazil.

Other reported hosts: *Didelphis albiventris* Lund, *Didelphis per-nigra* (J.A. Allen), *Lutreolina crassicaudata*, *Metachirus myosurus* (Temminck) and *Philander opossum*.

Locality records: Argentina: Buenos Aires: Berisso. Brazil: Bahía: Igrapiúna; Minas Gerais: Belo Horizonte. Costa Rica: Carii. Mexico: Chiapas: Motozintla; Oaxaca: Cuicatlán; Quintana Roo: La Ceiba; Veracruz: Alvarado and Los Tuxtlas. Panama: Canal Zone. Paraguay: undetermined. Peru: Ancash: Marca; Cajamarca: undetermined; Huánuco: undetermined; Pasco: Villa Rica; San Martín: Llamas. Venezuela: El Tacal (Siebert, 1970; Silva and Costa, 1999; Haverkost and Gardner, 2008; Chero *et al.*, 2017; Polo-Gonzales *et al.*, 2019; Cirino *et al.*, 2020).

Records in Bolivia: *Philander opossum*: Santa Cruz: Estancia Cachuela Esperanza, $16^{\circ}47'$, $63^{\circ}14'$, 300 m, 22 August 1984, HWML70000 (108 specimens) from MSB:MAMM:210569; 15 km S of Santa Cruz, $17^{\circ}53'S$, $63^{\circ}07'W$, 2 August 1987, HWML70002 (39 specimens), from AMNH263965; 3 km SE of Montero, 1 km N of Villa Copacabana, $17^{\circ}23'S$, $63^{\circ}14'W$, 250 m, 26 June 1991, HWML70009 (21 specimens), from AMNH263963.

Additional specimens examined: HWML34950 (1 specimen) from *Didelphis albiventris*, Paraguay. HWML70013 (1 specimen) from *Lutreolina crassicaudata*, Berisso, Buenos Aires, Argentina.

Table 1. Helminthological records in Bolivia for species of marsupials included in the tribe Thylamini

	<i>Marmosops bishopi</i> (n = 1)	<i>Marmosops ocellatus</i> (n = 1)	<i>Marmosops noctivagus</i> (n = 4)	<i>Thylamys pallidior</i> (n = 2)	<i>Thylamys venustus</i> (n = 5)	<i>Thylamys sponsorius</i> (n = 2)	<i>Thylamys</i> sp. (n = 3)
<i>Linstowia schmidtii</i>					USNPC82216		
<i>Atriotaeia</i> sp.			HWML49845 (25%)		HWML70016–18 (20%)	HWML 118760, 118784 (100%)	
<i>Mathevotaenia bivittata</i>							HWML118719 (33%)
<i>Mathevotaenia sanmartini</i>				HWML70016–19 (100%)			
<i>Mathevotaenia</i> sp.	HWML 118786				HWML 118787 (20%)		HWML 118788 (33%)
<i>Pritchardia boliviensis</i>		HWML118789	HWML118790 and 118791 (50%)				
<i>Neohilgertia venusti</i>					HWML118792 (60%)		HWML61553 (66%)
<i>Pterygodermatites elegans</i>	HWML118745						
<i>Gongylonemoides marsupialis</i>					HWML118750 (20%)		
<i>Moennigia</i> sp.			HWML118766 and 118767 (25%)				
<i>Viannaia metachirops</i>			HWML 118724				
<i>Viannaia minispicula</i>			HWML 118726				
<i>Viannaia philanderi</i>					HWML118728		
<i>Viannaia simplicispicula</i>					HWML118761–63	HWML118760	HWML118759 (33%)
<i>Viannaia viannai</i>			HWML 118734–36				HWML118794 (66%)
<i>Turgida turgida</i>	HWML118795		HWML118757 (75%)				
<i>Trichuris reesali</i>			HWML118757 (25%)				

Collection number for a representative and prevalence are provided.

Table 2. Helminthological records in Bolivia for species of marsupials included in the tribe Marmosini

	<i>Monodelphis domestica</i> (n = 9)	<i>Monodelphis sanctarosae</i> (n = 1)	<i>Monodelphis glirine</i> (n = 1)	<i>Monodelphis peruviana</i> (n = 1)	<i>Monodelphis adusta</i> (n = 1)	<i>Marmosa</i> sp. (n = 4)
<i>Rhopalias</i>	HWML118796 and 118797 (22%)					
<i>Paralinstowia schmidti</i>	HWML38846 (11%)					
<i>Mathevotaenia bivittata</i>	HWML118798 (22%)					HWML118799 (50%)
<i>Pritchardia boliviensis</i>	HWML118800 (11%)					HWML118801 (50%)
<i>Didelphoxyuris</i> sp.						HWML118755 (25%)
<i>Monodelphoxyuris dollmeiri</i>	HWML60229, 60130 (22%)					
<i>Aspidodera raillieti</i>	HWML60236 (11%)	HWML118802				HWML118803 (25%)
<i>Spirura guianensis</i>	HWML118756 (33%)					
<i>Pterygodermatites elegans</i>						HWML 118744 (50%)
<i>Pterygodermatites jeagerskioldi</i>	HWML 118746 and 118747 (22%)					HWML 118744 (25%)
<i>Moennigia</i> sp.	HWML 118764 (33%)		HWML 118765			
<i>Viannaia viannai</i>	HWML118805 (33%)			HWML 118737	HWML118806	HWML 118732 and 118733
<i>Trichuris reesali</i>						HWML118807 (25%)

Monodelphis kunsii was infection free. Collection number for a representative and prevalence are provided.

3. *Rhopalias macracanthus* Chandler, 1932

Synonyms: *Rhopalias louisiana* Hearin, 1937

Site of infection: Small intestine

Type host and locality: *Didelphis virginiana* Kerr, Houston, Texas

Other reported hosts: *Didelphis marsupialis*.

Locality Records: Costa Rica: Cariai. Mexico: Colima: Comala, La Esperanza. Chiapas: Jaltengo, Motozintla, and Pueblo Nuevo; Oaxaca: Temazcal; Quintana Roo: Rancho La Ceiba; Veracruz: Alvarado, and Los Tuxtlas. United States: Florida: Tallahassee; Illinois: Jackson Co.; Maryland: Beltsville; Texas: Houston (Siebert, 1970; Alden, 1995; Haverkost and Gardner, 2008).

Records in Bolivia: *Philander opossum*: Santa Cruz: Santa Cruz, 16°28'12"S, 63°08'24"W, HWML70001 (1 specimen); 15 km S of Santa Cruz, 17°53'S, 63°07'W, 2 August 1987, HWML70003 (7 specimens), from AMNH263965 and 263966; 3 km SE of Montero, 1 km N of Villa Copacabana, 17°23'S, 63°14'W, 250 m, 26 June 1991, HWML70010, (3 specimens) from AMNH263963; 10 km N of San Ramón, 16°36'S, 62°42'W, 250 m, 7 August 1985, HWML70028 (1 specimen), from host MSB:MAMM:55857.

Additional specimens examined: USNPC8548 (1 specimen) from *Didelphis marsupialis*, Houston.

Remarks: *Rhopalias macracanthus* and *R. coronatus* cause co-infections in the Gray four-eyed opossum, *Philander opossum*. Haverkost and Gardner (2008) reviewed species in the family across the continent, making observations and identifying reliable characters based on morphometric analyses.

Class Cestoda Rudolphi, 1808

Order Cyclophyllidea van Beneden in Braun, 1900

Family Anoplocephalidae Blanchard, 1891

Subfamily Linstowiinae Fuhrmann, 1907

Mathevotaenia Akhmyan, 1946

4. *Mathevotaenia bivittata* (Janicki, 1904) Yamaguti, 1959

Synonyms: *Oochoristica bivittata* Janicki, 1904; *Linstowia* (*Opossumia*) *bivittata* (Janicki, 1904) Spasskii, 1951; *Opossumia bivittata* Spasskii, 1981.

Site of infection: Small intestine

Type host and locality: *Marmosa* sp., Brazil.

Other reported hosts: *Caluromys derbianus* (Waterhouse), *Didelphis albiventris*, *Didelphis marsupialis*, *Marmosa paraguayana* Tate (as *Micoreus cinereus*), *Marmosa murina* (L.), *Marmosa demerarae* (Thomas), *Metachirus nudicaudatus* (É. Geoffroy), *Monodelphis domestica* (Wagner), *Philander opossum*, and *Thylamys* sp.

Locality records: Argentina: Salta: Orán. Brazil: Pará: Belém, Bassuquara and Bacia de Agua Preta; Mato Grosso do Sul: Bodoquena; Espirito Santo: Santa Teresa; Rio de Janeiro: Angra dos Reis. Panama: Canal Zone. Trinidad and Tobago: Rio Claro, Sangre Grande. French Guiana: Cayenne, Nouragues, Saut Pararé and Saül, Pic Matecho (Foster, 1939; dos Santos, 1968; Campbell et al., 2003; Byles et al., 2013).

Records in Bolivia: *Thylamys* sp.: Tarija: 3 km S of Cuyambuyo, 22°16'S, 64°33'W, 900 m, 3 and 4 August 1991, HWML118719 (207 specimens) from MSB:MAMM:240043.

Additional specimens examined: HWML 17712 from *Marmosa cinerea* (Temminck), Argentina, Salta, Orán. HWML 49769 from *Marmosa murina* French Guiana, Cayenne, Montagne du Tigre.

5. *Mathevotaenia sanmartini* Jiménez, Braun, Campbell and Gardner, 2008

Site of infection: Small intestine

Type host and locality: *Thylamys pallidior* (Thomas), OMNH 34911, Argentina: Jujuy: Susques, 8.2 km south of Sey (by road), 24°00'48.8"S, 66°30'52.8"W, 4167 ± 10 m (31 March and 1 April 2006).

Locality and host records: No additional records available.

Records in Bolivia: *Thylamys pallidior*: Cochabamba: Curubamba, 7.5 km southeast of Rodeo (by road), 17°40'31"S, 65°36'04"W, 4000 m, 24 and 26 July 1993, HWML70016–19 (1 and 3 specimens) from MSB:MAMM:87100 and MSB:MAMM:87102.

Additional specimens examined: CHLP5727 (1 specimen) holotype from *Thylamys pallidior*, Argentina.

Table 3. Helminthological records in Bolivia for species of marsupials included in the tribe Didelphini

	<i>Didelphis albiventris</i> (n = 3)	<i>Didelphis marsupialis</i> (n = 3)	<i>Chironectes minimus</i> (n = 2)	<i>Philander opossum</i> (n = 7)
<i>Rhopalias caballeroi</i>			HWML70021 (50%)	
<i>Rhopalias coronatus</i>				HWML70000, HWML70002; HWML70009 (42%)
<i>Rhopalias macracanthus</i>				HWML70001; HWML70003; HWML70010; HWML70028 (57%)
<i>Mathevotaenia bivittata</i>				HWML118808 (28.6%)
<i>Pritchardia boliviensis</i>				HWML118809 (14.3%)
<i>Didelphoxyuris thylamisis</i>				
<i>Aspidodera railieti</i>	HWML118738 (33%)	HWML61882 and 118810 (100%)		HWML118741 (28.6%)
<i>Turgida turgida</i>	HWML118751 (33%)	HWML118811 (75%)		
<i>Cruzia tentaculata</i>	HWML118741 and 118742 (33%)			HWML118743 (14.3%)
<i>Spirura guianensis</i>			HWML 118748 (50%)	
<i>Travassostrongylus callis</i>			HWML 118721 (50%)	
<i>Viannaia hamata</i>		HWML 118723		
<i>Viannaia simplicispicula</i>		HWML 118758		
<i>Viannaia skrjabini</i>		HWML 118729		
<i>Viannaia viannai</i>		HWML 118730		HWML118812 (43%)
<i>Oligacanthorhynchus microcephala</i>		HWML60388		HWML60600

Collection number for a representative and prevalence are provided.

Remarks: Several of these specimens were reported in the original description of the species (Jiménez *et al.*, 2008).

6. *Mathevotaenia* sp.

Site of infection: Small intestine

Type host and locality: Not yet named.

Records in Bolivia: *Marmosops bishopi* (Pine): La Paz: La Reserva, 15°44'S, 67°31'W, 850 m, 27 July 1992, HWML118786 (1 specimen) from MSB:MAMM:235887. *Marmosops noctivagus* (Tschudi): Cochabamba: 9.5 km by road NE of Tablas Monte, Río Jatún Mayu; 17°2'S, 65°59'W, HWML49845 from MSB:MAMM:70278. *Thylamys pusillus* (Desmarest): Santa Cruz: 53 km E Boyuibe, 20°27'S, 62°50'W, 600 m, 6 July 1991, HWML118788 (6 specimens) from MSB:MAMM:87105. *Thylamys venustus*: Tarija: Tapecua, 21°26'S, 63°55'W, 1500 m, 12 July 1991, HWML 118787 (2 specimens) from AMNH275439.

7. *Paralinstowia schmidti* (Gardner and Campbell, 1992) Beveridge and Spratt, 2003

Site of infection: Small intestine

Type host and locality: *Thylamys elegans venusta* (Thomas), Bolivia: Chuquisaca, El Porvenir 20°45'S, 63°13'W, 675 m, 6 July, 1985, symbiotype: AMNH261257.

Other reported hosts: *Monodelphis domestica* (Wagner)

Locality records: None available.

Records in Bolivia: Chuquisaca: El Porvenir, 20°27'W, 63°07'48"S, 675 m, 15 July 1985, UCDNC2831 (32 specimens) from host MSB:MAMM:211200.

Remarks: Both *Monodelphis domestica* and *Thylamys elegans venusta* -junior synonym of *Thylamys venustus* (Thomas)- were the only marsupials collected in El Porvenir. The species was not found in the other three localities where specimens of *Monodelphis domestica* were collected.

8. *Pritchardia boliviensis* Gardner, Jiménez and Campbell, 2013

Site of infection: Small intestine

Type host and locality: *Marmosops noctivagus*: Cochabamba: 9.5 km by road NE of Tablas Monte, Río Jatun Mayu 17°02'29"S, 65°59'05"W, 1500 m, 14 July 1993, symbiotype MSB: MAMM:70278.

Other reported hosts: *Marmosa paraguayana*, *Metachirus nudicaudatus*, *Gracilinanus* sp., *Marmosops ocellatus* (Tate)

Locality records: Brazil: Paraná: between Corbélia and Cascavel. Paraguay: Alto Paraná: Estación Biológica Limoy (Gardner *et al.*, 2013; Benatti *et al.*, 2023).

Records in Bolivia: *Marmosa* sp.: Santa Cruz: 53 km E of Boyuibe, 20°27'S, 62°50'W, 600 m, 8 July 1991, HWML118801 (139 specimens) from MSB:MAMM:239772. *Marmosops noctivagus*: Cochabamba: 9.5 km by road NE Tablas Monte, 17°02'S, 65°59'W, 14 and 16 July 1993, HWML118790 and HWML118791 (98 and 65 specimens) from MSB:MAMM:70278 and MSB: MAMM: 30279; La Paz: Chijchijpa, 16°09'S, 67°45'W, 1114 m, 8 July 1992, HWML61763 (19 specimens) from host MSB: MAMM:235553. *Marmosops ocellatus*: Santa Cruz: 3.5 km W, Estación El Pailón, 17°39'S, 62°45'W; 300 m, 21 September 1984, HWML118789 from MSB:MAMM:55070. *Metachirus nudicaudatus*: La Paz: La Reserva, 15°44'S, 67°31'W; 840 m. 24 July 1992, CNHE6422, CHIOC37318, USNPC103071, from CBF2310. *Monodelphis domestica*: Santa Cruz: 1 km S and 3 km W of Estancia Isibolos, 19°31'S, 63°36'W, 930 m, 5 July 1991, HWML118800 (52 specimens) from MSB:MAMM:239734. *Philander opossum*: Santa Cruz: 3 km SE Montero, 1 km N Villa Copacabana, 17°23'S, 63°14'W, 250 m, 26 June 1991, HWML118809 (1 specimen) from MSB:MAMM:239685.

Remarks: The holotype for this species was examined and used as a comparative reference.

9. *Atriotaeonia* sp.

Site of infection: Small intestine

Records in Bolivia: *Marmosops noctivagus*: La Paz: La Reserva, 15°44'S, 67°31'W, 850 m, 24 July 1992, HWML118724 (4 specimens) from MSB:MAMM:235815. *Thylamys venustus*: Tarija: 3 km SE Cuyambuyo, 22°16'S, 64°33'W, 900 m, 4 August 1991, HWML118720 (2 specimens) from MSB:MAMM:140296. *Thylamys sponsorius* (Thomas): Tarija: 3 km SE Cuyambuyo, 22°16'S, 64°33'W, 900 m, 4 August 1991, HWML118760 (1 specimen) from MSB:MAMM:67014, HWML118784 (365 specimens from MSB:MAMM:67015).

Remarks: Most of the specimens were contracted, making it difficult to identify to species level.

Phylum Nematoda Cobb, 1932

Class Chromadoria Pearse, 1936

Order Rhabditida Chitwood, 1933

Superfamily Ancylostomatoidea Looss, 1905

Family Ancylostomatidae Looss, 1905

Subfamily Bunostominae Railliet and Henry, 1909

Monodontus Molin, 1860

10. *Monodontus* sp.

Site of infection: Small intestine

Records in Bolivia: *Thylamys venustus*: Tarija: 3 km SE Cuyambuyo, 22°16'S, 64°33'W, 900 m, 4 August 1991, HWML118720 (1 specimen) from MSB:MAMM:140296.

Remarks: This is a single mature female. Species of the genus are known to typically infect rodents.

Superfamily Molineoidea Skrjabin and Shulz, 1937

Family Molineidae Skrjabin and Shulz, 1937

Subfamily Anoplostrongylinae Chandler, 1938

Moennigia Travassos, 1935

11. *Moennigia* sp.

Site of infection: Small intestine

Records in Bolivia: *Monodelphis domestica*: Chuquisaca: Río Limón, 19°33'S, 64°08'W, 1300 m, 3 August 1990 HWML118764 (1 specimen) from MSB:MAMM:63278. *Monodelphis glirina*: Pando: Santa Rosa, 12°07'48"S, 68°14'24"W, 800 m, 1 August 1986, HWML118765 (11 specimens) AMNH M 262399. *Marmosops noctivagus*: Cochabamba: 9.5 km by road NE Tablas Monte, 17°02'S, 65°59'W, 14 and 15 July 1993, HWML118766 and HWML118767 (1 specimen each) from MSB:MAMM:70278 and MSB:MAMM:238453.

Remarks: These individuals belong to a single species which may be new to science. The senior author is attempting to work in the precise identification and description.

Superfamily Heligmosomoidea Cram, 1927

Family Viannaiidae Neveu-Lemaire, 1944

Subfamily Viannaiinae Neveu-Lemaire, 1944

Travassostrongylus Orloff, 1933

12. *Travassostrongylus callis* (Travassos, 1914) Orloff, 1933

Synonyms: *Trichostrongylus callis* Travassos 1914; *Ostertagia callis* (Travassos, 1914) Travassos 1918

Site of infection: Small intestine

Type host and locality: *Didelphis aurita* (Wied-Neuwied), Brazil: Rio de Janeiro: Manguinhos CHIOC 724.

Other reported hosts: *Didelphis marsupialis*, *Philander opossum*.

Locality records: Brazil: Espirito Santo: Sooretama; Rio de Janeiro: Petrópolis. French Guiana: undetermined. Panama: Panama City (Diaw, 1976a, 1976b; Scheibel et al., 2014).

Records in Bolivia: *Chironectes minimus*: La Paz: La Reserva, 15°44'S, 67°31'W, 850 m, 22 July 1992, HWML118721 (6 specimens) from MSB:MAMM:68330.

Additional specimens examined: From *Didelphis aurita*, Brazil: Rio de Janeiro: CHIOC 8426, 8584, 8589, 9608 Manguinhos; CHIOC 9118 Petrópolis; CHIOC 29504, 29505 Espirito Santo: Sooretama.

Viannaia Travassos, 1914

13. *Viannaia didelphis* (Travassos, 1914) Durette-Desset, 1968

Synonyms: *Nematodirus (Mecistocirrus) didelphis* Travassos, 1914

Site of infection: Small intestine

Type host and locality: *Didelphis aurita*, Brazil, Rio de Janeiro, Manguinhos CHIOC 942.

Other reported hosts: *Didelphis marsupialis* and *Didelphis virginiana*.

Locality records: Costa Rica: Guanacaste, Colonia Bolaños. Mexico: Colima: La Esperanza, Madrid. Panama: Panama City. United States: Georgia: Enigma, Bulloch Co.; Louisiana: Jeanerette; Illinois: Urbana, Jackson Co.; North Carolina: undetermined; Tennessee: Reelfoot Lake. Trinidad and Tobago: undetermined; Venezuela: Maracaibo (Guerrero, 1985; Alden, 1995; Monet-Mendoza et al., 2005; Scheibel et al., 2014).

Records in Bolivia: *Marmosa* sp.: La Paz: Chijchijpa, 16°09'S, 67°45'W, 1114 m, 8 July 1992, HWML118722, HWML61763 from host MSB:MAMM:235553.

Remarks: Species in *Viannaia* have been reported in several species of marsupials across the Americas (Dikmans, 1931; Cañeda-Guzmán, 1997; Ellis et al., 1999; Silva and Costa, 1999; Gomes et al., 2003; Antunes, 2005; Byles et al., 2013). A few studies suggest some infections are caused by multiple species (Diaw, 1976a; Guerrero, 1985; Scheibel et al., 2014), thus, individual identification of these nematodes is recommended.

14. *Viannaia hamata* Travassos, 1914

Site of infection: Small intestine

Type host and locality: *Didelphis aurita*, Brazil, Rio de Janeiro, Manguinhos CHIOC 942

Other reported hosts: *Didelphis albiventris*, *Didelphis marsupialis*, *Didelphis virginiana*, *Philander opossum*, *Marmosa cinerea*, *Marmosa murina*.

Locality records: Brazil: Minas Gerais, Belo Horizonte; Pará: Belém; Rio de Janeiro: Glicério, Petrópolis; Paraná: between Corbélia and Cascavel; Rio Grande do Sul: Pelotas; Peru: Pasco; Villa Rica; San Martín: Bella Vista, Lamas. Trinidad and Tobago, undetermined; United States: Georgia, Macintosh Co., Bulloch Co.; North Carolina. Venezuela: Miranda, Guatopo (Wolfgang, 1951; Guerrero, 1985; Ellis et al., 1999; Silva and Costa, 1999; Gomes et al., 2003; Chero et al., 2017; Polo-Gonzales et al., 2019; Benatti et al., 2023).

Records in Bolivia: *Didelphis marsupialis*: La Paz: La Reserva, 15°44'S, 67°31'W, 850 m, 22 July 1992, HWML118723 (2 specimens) from MSB:MAMM:235674.

Additional specimens examined: CHIOC 29289 and 29290 from *Didelphis* sp. Brazil, Rio de Janeiro, Usina da Tijuca.

15. *Viannaia metachirops* Durette-Desset, 1974

Site of infection: Small intestine

Type host and locality: *Philander opossum*, French Guiana

Other reported hosts: None available

Locality records: None available

Records in Bolivia: *Marmosops noctivagus*: La Paz: La Reserva, 15°44'S, 67°31'W, 850 m, 24 July 1992, HWML118724 (5 specimens) from MSB:MAMM:235815. *Marmosa* sp., La Paz:

Chijchijpa, 16°09'S, 67°45'W, 1114 m, 8 July 1992, HWM118725 (3 specimens) from host MSB:MAMM:235553.

Additional specimens examined: None, identification made based on diagnostic traits.

16. *Viannaia minispicula* Guerrero, 1985

Site of infection: Small intestine

Type host and locality: *Marmosa murina*, Venezuela: Amazonas, Caño Yaguá

Other reported hosts: *Marmosa demerarae*, *Philander opossum*

Locality records: French Guiana: Guyanne, Cacao.

Records in Bolivia: *Marmosops noctivagus*: Cochabamba: 9.5 km by road NE Tablas Monte, 17°02'S, 65°59'W, 15 July 1993, HWML118726 (1 specimen) from MSB:MAMM:238453.

Additional specimens examined: None, identification made based on diagnostic traits.

17. *Viannaia philanderi* (Wolfgang, 1951) Durette-Desset, 1968

Site of infection: Small intestine

Type host and locality: *Caluromys philander*, Trinidad.

Other reported hosts: None available

Locality records: None available

Records in Bolivia: *Marmosa* sp.: Santa Cruz: Estancia Cachuela Esperanza, 16°46'59.99"S, 63°13'59.99"W, 300 m, 24 August 1984, HWML118727 (3 specimens) from MSB:MAMM:211050. *Thylamys venustus*: Tarija: 3 km SE Cuyambuyo, 22°16'S, 64°33'W, 900 m, 4 August 1991, HWML118728 (13 specimens) from MSB:MAMM:140297.

Additional specimens examined: None, identification made based on diagnostic traits.

18. *Viannaia simplicispicula* (Navone, Suriano and Pujol, 1991) Jiménez *et al.*, 2024

Synonyms: *Hoineffia simplicispicula* Navone, Suriano and Pujol, 1991

Site of infection: Small intestine

Type host and locality: *Thylamys venustus cinderellus* (Thomas 1902): Argentina: Tucumán: Quebrada Los Sosa, Museo Argentino Bernardino Rivadavia No. 360.

Other reported hosts: *Tlacuatzin canescens* (J.A. Allen).

Locality records: Argentina: Jujuy: Dr Manuel Belgrano, Las Capillas and El Palmar; Salta: Mosconi. Mexico: Oaxaca: Santa Catarina Juquila (Jiménez *et al.*, 2008; Guzmán-Cornejo *et al.*, 2012).

Records in Bolivia: *Didelphis marsupialis*: La Paz: La Reserva, 15°44'S, 67°31'W, 850 m, 22 July 1992, HWML118758 (1 specimen) from MSB:MAMM:235674. *Thylamys* sp.: Tarija: 3 km S of Cuyambuyo, 22°16'S, 64°33'W, 900 m, 4 August 1991, HWML118759 (18 specimens) from MSB:MAMM:240043. *Thylamys sponsorius*: Tarija: 3 km SE Cuyambuyo, 22°16'S, 64°33'W, 900 m, 4 August 1991, HWML118760 (1 specimen) from MSB:MAMM:67015. *Thylamys venustus*: Tarija: 3 km SE Cuyambuyo, 22°16'S, 64°33'W, 900 m, 4 August 1991, HWML118761 (18 specimens) from MSB:MAMM:140296, HWML118762 (10 specimens) from MSB:MAMM:140297; Tapecua, 21°26'S, 63°55'W, 1500 m, 12 July 1991, HWML118763 (4 specimens) from AMNH275439.

Remarks: Guerrero (1985) transferred *Hoineffia cayennensis* Diaw, 1976 to *Viannaia*. This recommendation was based on the observation that the transversally elongated bursa – diagnostic for *Hoineffia* Diaw, 1976 – also occurs in other species featuring a gubernaculum such as *Viannaia venezuelensis* Guerrero, 1985 and *Viannaia barusi* Guerrero, 1985. Furthermore, other species in the genus feature the combination of cordiform bursa and lack

of gubernaculum, such as *Viannaia viannai*. The phenotypic plasticity of the bursa is shown in a subset of species of *Viannaia* collected across Mexico (Ramírez-Cañas *et al.*, 2021). *Hoineffia simplicispicula* Navone, Suriano and Pujol, 1991 was proposed as the second species in the genus; further, the species was recorded in in Mexico and in Argentina (Jiménez *et al.*, 2008; Guzmán-Cornejo *et al.*, 2012). Apparently, Navone *et al.* (1991) were not familiar with the change proposed by Guerrero (1985). We herein consider that the differences in the shape of the caudal bursa, the relative length of the dorsal lobe, dorsal ray and ray 8 are consistent with the intraspecific variability documented by Guerrero (1985). Further this variability is observed in specimens from Argentina, Bolivia and French Guiana. Rather than proposing an amended diagnosis, we refer readers to the diagnosis proposed by Dikmans (1945), who only missed the presence of three ventral ridges proposed by Durette-Desset (1971) in his definition of the genus.

Additional specimens examined: HWML63395 from *Thylamys venustus*, 24.8 km N of Santa Clara (by road), Jujuy, Argentina.

19. *Viannaia skrjabini* Lent and Freitas, 1937

Site of infection: Small intestine

Type host and locality: *Philander opossum*, Brazil: Rio de Janeiro: Petrópolis. CHIOC 7721

Other reported hosts: *Didelphis albiventris*, *Didelphis marsupialis* and *Marmosa robinsoni* Bangs.

Locality records: Brazil: Pernambuco, Exu. Venezuela: Amazonas, Caño Yaguá; Miranda: Río Negro; Distrito Federal: Naiguatá (Guerrero, 1985).

Records in Bolivia: *Didelphis marsupialis*: La Paz: La Reserva, 15°44'S, 67°31'W, 850 m, 22 July 1992, HWML118729 (from HWML61838) (3 specimens) from MSB:MAMM:235674.

Additional specimens examined: Holotype CHIOC7721, from *Philander opossum*, Brazil: Rio de Janeiro, Petrópolis

20. *Viannaia viannai* Travassos, 1914

Site of infection: Small intestine

Type host and locality: *Didelphis aurita*, Brazil, Rio de Janeiro, Manguinhos CHIOC 922.

Other reported hosts: *Didelphis marsupialis*, *Didelphis virginiana*, *Philander opossum*.

Locality records: Brazil: Pernambuco: Exu; Rio de Janeiro: Morro São João, Casimiro de Abreu. Costa Rica: Guanacaste, Colonia Baños. French Guiana: Camp du Tigre. Mexico: Guerrero: Taxco El Viejo. Panama: Panama City. Peru: San Martín: Bella Vista. United States: Illinois, Carbondale; Maryland, Beltsville. Venezuela: Miranda: Caño Yagua, Río Negro and San Antonio (Guerrero, 1985; Monet-Mendoza *et al.*, 2005; Scheibel *et al.*, 2014; Chero *et al.*, 2017).

Records in Bolivia: *Didelphis marsupialis*: La Paz: Chijchijpa, 16°09'S, 67°45'W, 1114 m, 6 July 1992, HWML118730 (1 specimen) from MSB:MAMM:235570; La Reserva, 15°44'S, 67°31'W, 850 m, 22 July 1992, HWML118731 (1 specimen) from MSB:MAMM:235674. *Marmosa* sp.: La Paz: Chijchijpa, 16°09'S, 67°45'W, 1114 m, 8 July 1992, HWML118732 (3 specimens) from host MSB:MAMM:235553; Santa Cruz: Estancia Cachuela Esperanza, 16°46'59.99"S, 63°13'59.99"W, 300 m, 24 August 1984, HWML118733 (7 specimens) from MSB:MAMM:211050. *Marmosops noctivagus*: Cochabamba: 9.5 km by road NE Tablas Monte, 17°02'S, 65°59'W, 14, 15 and 16 July 1993, HWML62620, HWML118734–36 (11, 13 and 10 specimens) from MSB:MAMM:70278, MSB:MAMM:238453 and MSB:MAMM:30279; La Paz: La Reserva, 15°44'S, 67°31'W, 850 m, 24 July 1992, HWML61852 (10 specimens) from MSB:MAMM:235815. *Monodelphis peruviana*: La Paz: La Reserva,

15°44'S, 67°31'W, 850 m, 25 July 1992, HWML118737 (3 specimens) from MSB:MAMM:68336. *Thylamys* sp.: Tarija: Tapehua, 21°26'S, 63°55'W, 1500 m, 1 June 1991, HWML118794 (5 specimens) from MSB:MAMM:238757.

Additional specimens examined: From *Didelphis marsupialis*, HWML67179–81, Panama City, Panama and Colonia Baños, Costa Rica. From *Didelphis virginiana* HWML61798 from Carbondale Illinois, U.S.A.

Order Spirurida Railliet, 1915

Suborder Ascaridina Inglis, 1983

Superfamily Heterakoidea Railliet and Henry, 1912

Family Aspidoderidae Skrjabin and Shikhobalova, 1947

Aspidodera Railliet and Henry, 1912

21. *Aspidodera raillietii* Travassos, 1913

Synonyms: *Aspidodera harwoodi* Chandler, 1932, *Aspidodera vicentei* Pinto, Kohn, Fernandes and Mello, 1981, *Aspidodera diaz-ungriai* Masi-Pallarés and Benítez-Uscher, 1971

Site of infection: Caecum and large intestine

Type host and locality: *Didelphis aurita*, Manguinhos, Brazil

Other reported hosts: *Didelphis marsupialis*, *Didelphis virginiana*, *Didelphis pernigra*, *Gracilinanus agilis* (Burmeister), *Marmosa demerarae*, *Marmosa murina*, *Marmosops ocellatus*, *Metachirus nudicaudatus*, *Metachirus myosurus*, *Philander opossum*, Sigmodontinae: *Nectomys squamipes* (Brants), *Euoryzomys nitidus* (Thomas).

Locality records: Brazil: Bahía: Igrapiúna; São Paulo, Piauí, Formosa. French Guiana: Montagne du Tigre, Nouragues, Saül, Petit Saut, Route de Kaw. Guatemala: Santa Rosa. Mexico: Motozintla; Panama: Panama Canal. Paraguay: Puerto Ibapobó. Suriname: Peru: La Libertad, Bosque del Cachil; San Martín: Bella Vista; Llamas. United States: Texas, Houston; Illinois, Jackson Co., Union Co., (Santos et al., 1990; Alden, 1995; Jiménez-Ruiz et al., 2008; Chero et al., 2017; Polo-Gonzales et al., 2019; Varella et al., 2022)

Records in Bolivia: *Didelphis albiventris*: Tarija: Tapehua, 21° 26'S, 63°55'W, 1500 m, 14 July 1991, 1500 m, HWML118738 (36 specimens) from CBF2379. *Didelphis marsupialis*: La Paz: La Reserva, 15°44'S, 67°31'W, 850 m, 22 and 25 July 1992, HWML118810 (7 specimens) from MSB:MAMM:235674, HWML61882 (2 specimens) from MSB:MAMM:235838. *Marmosops ocellatus*: Santa Cruz: 15 km S of Santa Cruz, 17° 53'S, 63°07'W, 2 August 1984, HWML118739 (2 specimens) from MSB:MAMM:58514. *Monodelphis domestica*: Chuquisaca: El Porvenir, 20°45'S, 63°13'W, 675 m, 7 July 1985, HWML60236 (1 specimen) from MSB:MAMM:55847. *Monodelphis sanctarosae*: Santa Cruz: Santa Rosa de la Roca, 15°30'00"S, 61°16'12"W, 250 m, 6 June 1990, HWML118802 (2 specimens) from MSB:MAMM:237023. *Philander opossum*: Santa Cruz: San Miguel Rincón, 17°22'59"S, 63°31'59"W, 300 m, 14 August 1984, HWML118740 (1 specimen) from MSB: MAMM:210528; 6 km by road W Ascención, 15°25'47"S, 63° 53'59"W, 240 m, 13 August 1985, HWML118741 (6 specimens) from MSB:MAMM:211436.

Additional specimens examined: CHIOC12 (holotype) from *Didelphis aurita*, Rio de Janeiro Brazil.

CHIOC18356, CHIOC19115, from *Didelphis azarae*, Puerto Ibapobó, Paraguay. CHIOC4446 from *Tolypeutes tricinctus* (L.) Tanque, Brazil. CHIOC31879 from *Nectomys squamipes* Formosa, Goiás, Brazil. USNPC8550 from *Didelphis virginiana*, Houston, Texas, U.S. A.CMNA408 from *Didelphis marsupialis*, Saint Vincent, Trinidad, Trinidad and Tobago. CNHE2110 from *Didelphis marsupialis*, Motozintla, Mexico.

Superfamily Cosmocercoidea Railliet, 1916

Family Kathlaniidae Lane, 1914

Cruzia Travassos, 1917

22. *Cruzia tentaculata* (Rudolphi, 1819) Travassos, 1917

Synonyms: *Ascaris tentaculata* Rudolphi, 1819; *Oxysoma tentaculata* Schneider, 1866

Site of infection: Large intestine and caecum

Type host and locality: *Didelphis marsupialis*, Brazil

Other reported hosts: *Didelphis albiventris*, *Didelphis aurita*, *Didelphis pernigra*, *Didelphis virginiana*, *Metachirus nudicaudatus*, *Metachirus myosurus*, *Philander opossum*, *Philander quica* (Temminck).

Locality Records: Brazil: Bahía: Igrapiúna; Minas Gerais: Belo Horizonte, Conceição dos Ouros; Paraíba: Santa Rita; Paraná: Curitiba, Ponta Grossa; São Paulo: São Paulo; Rio Grande do Sul: Porto Alegre; Rio de Janeiro: Glicério, Barra de Marica, Casimiro de Abreu, Serra dos Orgãos, Sumidouro, Petrópolis; Santa Catarina: Santa Catarina Island; Sergipe: Capela, São Cristovão. Colombia: Valle del Cauca: Meléndez. Mexico: Chiapas; Colima; Distrito Federal; Estado de México; Hidalgo: Tasquillo; Guerrero; Jalisco; Morelos; Oaxaca; Veracruz; Yucatán. Peru: Ancash: Marca, Huanchoc; Loreto: Iquitos; Piura: valle del Huancabamba; San Martín: Bella Vista; Cajamarca: Cajamarca. United States: Louisiana, North Carolina, Pennsylvania, Tennessee, Texas, Wisconsin (Alden, 1995; Silva and Costa, 1999; Monet-Mendoza et al., 2005; Chero et al., 2017; Polo-Gonzales et al., 2019; Cirino et al., 2020).

Records in Bolivia: *Didelphis albiventris*: Tarija: Tapehua, 21° 06'S, 63°55'W, 1500 m, 14 July 1991, HWML 118741 (39 specimens) from MSB:MAMM:239823; La Paz: Saynami Rio Zongo, 16°07'39"S, 68°05'59"W, 4 June 1993, HWML118742 (153 specimens) from MSB:MAMM:236299. *Didelphis pernigra*: La Paz: Yanacachi, Valle Aceromarka 16°19'35"S, 67°53'21"W, 3085 (Mollericonna and Nallar, 2014). *Philander opossum*: Santa Cruz: 6 km by road W Ascención, 15°25'47"S, 63°53'59"W, 240 m, 13 August 1985, HWML118743 (99 specimens) from MSB: MAMM:211436.

Remarks: The species has an almost continental distribution and it is known to occur in armadillos and opossums (Ruiz, 1947; Fujita et al., 1995; Adnet et al., 2009; Souza et al., 2022). The material of this species across its putative range needs to be reviewed.

Suborder Spirurina Railliet and Henry, 1915

Superfamily Rictularioidea Hall, 1913

Family Rictulariidae Hall, 1913

Pterygodermatites Wedl, 1861

23. *Pterygodermatites (Paucipectines) elegans* (Travassos, 1928) Quentin, 1969

Synonyms: *Rictularia elegans* Travassos, 1928

Site of infection: Small intestine

Type host and locality: *Eumops perotis* (Schinz): Engenheiro Gomide, São Paulo, Brazil.

Other reported hosts: *Marmosa cinerea*, *Marmosa demerarae*.

Locality Records: Brazil: Cafezal, Belém. French Guiana: Macouria, Montagne du Tigre, Pic Matecho, Saül (Byles et al., 2013).

Records in Bolivia: *Marmosa* sp.: Santa Cruz: Estancia Cachuela Esperanza, 16°46'59.99"S, 63°13'59.99"W, 300 m, 24 August 1984, HWML60078 (15 specimens) from MSB: MAMM:211050; La Paz: Chijchijpa: 16°09'S, 67°45'W, 1114 m, 8 July 1992, HWML118744 (1 specimen) from host MSB: MAMM:235553. *Marmosops bishopi*: La Paz: La Reserva, 15° 44'S, 67°31'W, 850 m, 27 July 1992, HWML118745 (1 specimens) from MSB:MAMM:235887.

Additional specimens examined: HWML67202 from *Marmosa demerarae*, Montagne du Tigre, Cayenne, French Guiana.

24. *Pterygodermatites (Paucipectines) jaegerskioldi* (Lent and Freitas, 1935) Quentin, 1969

Site of infection: Small intestine

Type host and locality: *Caluromys philander* (L.), Rio de Janeiro, Tijuca, Brazil.

Other reported hosts: *Gracilinanus agilis*, *Gracilinanus microtarsus* (Wagner).

Locality records: Brazil: Mato Grosso do Sul, Nhecolândia; Rio de Janeiro: Parque Nacional da Serra dos Orgãos (Lent and Freitas, 1935; Torres *et al.*, 2007, 2009).

Records in Bolivia: *Monodelphis domestica*: Santa Cruz: 27 km S of Santa Cruz, 3 km E and 1 km S Brecha Tres, 18°01'59"S, 63°10'01"W, 20 June 1992, HWML118746 (1 specimen) from MSB:MAMM:67022; 1 km S and 3 km W of Estancia Isibolos, 19°31'S, 63°36', 930 m, 5 July 1991, HWML118747 (5 specimens) from MSB:MAMM:239734.

Superfamily Spiruroidea Oerley, 1885

Family Spiruridae Oerley, 1885

Spirura Blanchard, 1849

25. *Spirura guianensis* (Ortlepp, 1924) Chitwood, 1938

Site of infection: Stomach

Type host and locality: Monki monki (Scientific name not disclosed), Suriname.

Other reported hosts: *Didelphis marsupialis*, *Gracilinanus agilis*, *Marmosa cinerea*, *Marmosa demerarae*, *Marmosa murina*, *Metachirops opossum*, *Philander opossum*, *Saguinus geoffroyi* (Pucheran), *Saguinus nigricollis* (Spix), and *Tamarinus nigricollis* (Spix)

Locality Records: French Guiana: Montagne du Tigre. Brazil: Rio de Janeiro, Itaguaí; Mato Grosso do Sul, Nhecolândia. Panama: Panama Canal (Torres *et al.*, 2009; Byles *et al.*, 2013).

Records in Bolivia: *Chironectes minimus*: La Paz: La Reserva, 15°44'S, 67°31'W, 850 m, 22 July 1992, HWML118748 (34 specimens) from MSB:MAMM:68330. *Monodelphis domestica*: Chuquisaca: El Porvenir, 20°27'W, 63°07'48"S, 675 m, 15 July 1985, HWML 118756 (1 specimen) from host MSB:MAMM:211199

Family Gongylonematidae Hall, 1916

Gongylonemoides Lent and Freitas, 1937

26. *Gongylonemoides marsupialis* (Vaz and Pereira, 1934) Freitas and Lent, 1937

Synonym: *Gongylonema marsupialis* Vaz and Pereira, 1934

Site of infection: Esophagus

Type host and locality: *Didelphis aurita*, São Paulo, Brazil

Other reported hosts: *Didelphis aurita*, *Metachirops opossum*.

Locality Records: Brazil: Rio de Janeiro; São Paulo: undetermined. Peru: San Martín: Llamas (Gomes *et al.*, 2003; Chero *et al.*, 2017).

Records in Bolivia: *Marmosa* sp.: La Paz: Chijchijpa, 16°09'S, 67°45'W, 1114 m, 8 July 1992, HWML118749 (55 specimens) from host MSB:MAMM:235553; *Thylamys venustus*: Tarija: 3 km SE Cuyambuyo, 22°16'S, 64°33'W, 900 m, 4 August 1991, HWML118750 (3 specimens) from MSB:MAMM:140296.

Superfamily Physalopteroidea Railliet, 1893

Family Physalopteridae Railliet, 1893

Turgida Travassos, 1920

27. *Turgida turgida* (Rudolphi, 1819) Travassos, 1919

Synonyms: *Physaloptera turgida* Rudolphi, 1819; *Spiroptera turgida* Dujardin, 1845; *Physaloptera didelphidis* Leidy, 1851.

Site of infection: Stomach

Type host and locality: Brazil

Other reported hosts: *Caluromys derbianus*, *Didelphis aurita*, *Didelphis albiventris*, *Didelphis marsupialis* and *Didelphis virginiana*, *Metachirus nudicaudatus*, and *Philander opossum*.

Locality Records: Argentina: Santiago del Estero. Brazil: Goiás: Nerópolis; Rio de Janeiro: Casimiro de Abreu, Angra dos Reis, Sumidouro; Santa Catarina: Santa Catarina Island. Mexico: Chiapas: Motozintla, Tonalá; Colima: Colima, Comala, Dos Amates, La Esperanza, Madrid; Distrito Federal; Estado de México: Tequesquahuac; Guerrero: Coyuquilla, Taxco El Viejo; Hidalgo: Tasquillo; Jalisco: Chamela; Michoacán, El Hortigal; Morelos; Oaxaca: Temazcal; Veracruz, Los Tuxtlas. Panama: Panama Canal. Peru: Loreto: Iquitos; Piura: San Felipe de Vichayal; San Martín: Bella Vista. United States: California, Connecticut, Colorado, Florida, Georgia, Illinois, Kansas, Louisiana, Oklahoma, New York, North Carolina, Pennsylvania, Tennessee, Texas, Virginia, Wisconsin. Trinidad and Tobago. Venezuela: Maracaibo (Alden, 1995; Monet-Mendoza *et al.*, 2005; Chero *et al.*, 2017; Polo-Gonzales *et al.*, 2019).

Records in Bolivia: *Didelphis albiventris*: Tarija: Tapecua, 21°06'S, 63°55'W, 1500 m, 14 July 1991, HWML118751 (69 specimens) from MSB:MAMM:239823; *Philander opossum*: Pando: Bella Vista, 11°13'48"W, 67°07'12"W, 170 m, 26 July 1986, HWML118752 (2 specimens) from MSB:MAMM:211891.

Suborder Oxyurinae Railliet, 1895

Superfamily Oxyuroidea Cobbold, 1864

Family Oxyuridae Cobbold, 1864

Didelphoxyuris Gardner and Hugot, 1995

28. *Didelphoxyuris thylamisis* Gardner and Hugot, 1995

Site of infection: Large intestine and caecum

Type host and locality: *Thylamys venustus*: Santa Cruz, 5 km NE Quiñe, 18°03'S, 64°19'W, 1900 m, 27 May 1991.

Other reported hosts: Other than symbiotype, none available

Locality records: Other than type locality, none available

Records in Bolivia: *Marmosa* sp.: Santa Cruz: 53 km E Boyuibe, 20°27'S, 62°50'W, 600 m, 6 July 1991, HWML118755 (12 specimens) from MSB:MAMM:239772. *Thylamys venustus*: Tarija: 3 km SE Cuyambuyo, 22°16'S, 64°33'W, 900 m, 4 August 1991, HWML118753 from MSB:MAMM:140296, HWML118754 (256 specimens) from MSB:MAMM:140297; Tarija: Tapecua, 21°26'S, 63°55'W, 1500 m, 12 July 1991, HWML61315 (17 specimens) from AMNH275439; Santa Cruz: 5 km NE Quiñe, 18°03'S, 64°19'W, 1900 m, 27 May 1991, HWML61086 NK22813 MSB:MAMM:87107 (360 specimens). *Thylamys pusillus*: Santa Cruz: 53 km E Boyuibe, 20°27'S, 62°50'W, 600 m, 6 July 1991, HWML61267 (28 specimens) from MSB:MAMM:87105

Additional specimens examined: HWML39072 holotype

Remarks: The symbiotype was originally identified as *Thylamys elegans* Waterhouse. However, a systematic review of the genus revealed that this species is restricted to the western slope of the Andean Cordillera; furthermore, fat-tailed opossums infected with pinworms used in the species description belong to *Thylamys venustus* (Giarla *et al.*, 2010).

Monodelphoxyuris Guerrero and Hugot, 2003

29. *Monodelphoxyuris dollmeiri* Guerrero and Hugot, 2003

Site of infection: Large intestine and caecum

Type host and locality: *Monodelphis emiliae* (Thomas): San Martín, Rio Camisea, Cusco, Peru, 11°47'10"S, 72°42'5"W, 474 m; May 08, 1997. Accession number CHIAUMSM1175. Symbiotype 14149 Mammal Collection of the Museo de Historia Natural, Universidad Nacional Mayor de San Marcos, Lima.

Other reported hosts: None available

Locality records: Other than type locality, none available

Records in Bolivia: Monodelphis domestica: Chuquisaca: El Porvenir, 20°27'W, 63°07'48"S, 675 m, 15 July 1985, HWML 118756 (158 specimens) from host MSB:MAMM:211199; HWML 60229 (88 specimens) from host AMNH M 261233. *Thylamys venustus*: Chuquisaca: El Porvenir, 20°27'W, 63°07'48"S, 675 m, 13 July 1985, HWML 60130 (2 specimens) from host MSB:MAMM:211181

Neohilgertia Navone, Suriano and Pujol, 1990

30. *Neohilgertia venusti* Navone, Suriano and Pujol, 1990

Site of infection: Large intestine and caecum

Type host and locality: *Thylamys venustus*, Tucuman, Burruyacu. Bernardino Rivadavia Helminthological Collection 362.

Other reported hosts: None available

Locality records: Other than type locality, none available

Records in Bolivia: Thylamys sp.: Tarija: 3 km S of Cuyambuyo, 22°16'S, 64°33'W, 900 m, 4 and 5 August 1991, HWML61553 (36 specimens) from MSB:MAMM:240043; HWML118792 (42 specimens) from MSB:MAMM:240056

Class Dorylaimea Hodda, 2007

Order Trichocephalida Spasski, 1954

Family Trichuridae Ransom, 1911

Trichuris Roederer, 1761

31. *Trichuris reesali* Wolfgang, 1951

Site of infection: Large intestine and caecum

Type host and locality: *Didelphis marsupialis*, Trinidad

Other reported hosts: *Didelphis marsupialis*, *Marmosa demerarae*, *Marmosa murina*, *Philander opossum* (Wolfgang, 1951; Byles *et al.*, 2013).

Locality Records: French Guiana: Camp du Tigre, Saül, Macouria.

Records in Bolivia: Marmosa sp.: Santa Cruz: Estancia Cachuela Esperanza, 16°46'59.99"S, 63°13'59.99"W, 300 m, 24 August 1984, HWML118807 (2 specimens) from MSB:MAMM:211050; *Marmosops noctivagus*: La Paz: La Reserva, 15°44'S, 67°31'W, 850 m, 24 July 1992, HWML118757 (7 specimens) from MSB:MAMM:235815.

Phylum Acanthocephala Rudolphi, 1808

Class Archiacanthocephala Meyer, 1931

Order Moniliformida Schmidt, 1972

Family Oligoacanthorhynchidae Southwell and McPhie, 1925

Oligoacanthorhynchus Travassos, 1915

32. *Oligoacanthorhynchus microcephalus* (Rudolphi, 1819)

Schmidt, 1972

Synonyms: *Echinorhynchus microcephalus* Rudolphi, 1819; *Echinorhynchus tortuosa* Leidy, 1850; *Hamanniella microcephalus* (Rudolphi, 1819) Travassos, 1915; *Hamanniella tortuosa* (Leidy, 1850) Van Cleave, 1924; *Hamanniella tumida* (Van Cleave, 1947) Van Cleave, 1953; *Oligoacanthorhynchus tortuosa* (Leidy, 1850) Schmidt, 1972; *Oligoacanthorhynchus tumida* (Van Cleave, 1947) Schmidt, 1972; *Travassosia tumida* Van Cleave, 1947

Site of infection: Small intestine

Type host and locality: *Caluromys philander*, Brazil

Other reported hosts: *Didelphis albiventris*, *Didelphis marsupialis*, *Didelphis virginiana*, *Marmosa demerarae*, *Marmosa murina*, *Metachirus myosurus*, *Philander opossum*. *Dasyurus novemcinctus* L., *Euphractus sexcinctus* (L.).

Locality Records: Brazil: Bahía: Igrapiúna; Rio de Janeiro: Rio de Janeiro; São Paulo: São Paulo. Colombia: Chocó. Mexico: Campeche, Colima, Michoacán, Guanajuato, Morelos, Oaxaca,

Tabasco, Veracruz, Yucatán. French Guiana: Montagne du Tigre, Macouria, Pic Matecho. Paraguay: Chaco Boreal. Suriname. United States: Alabama, Arkansas, Florida, Georgia, Illinois, Louisiana, Mississippi, Oklahoma, South Carolina, Texas. Venezuela (Richardson *et al.*, 2014; Acosta-Virgen *et al.*, 2015; Cirino *et al.*, 2020).

Remarks: *Oligoacanthorhynchus microcephalus* was collected from *Metachirus opossum* and *Philander opossum* from Santa Cruz, Bolivia and used for a redescription of the species (Richardson *et al.*, 2014).

Discussion

Bolivia contains a large variety of biomes and it is rich in natural resources (Ergueta and Salazar, 1991; Auty, 1994; Anderson, 1997; Hancock *et al.*, 2018). Among these resources, some minerals are used in high demand (Finer *et al.*, 2008; Hancock *et al.*, 2018), and vast areas of the country have been converted to agricultural use (Cuellar and Noss, 2014). The exploitation of these natural resources wipes out natural habitats and causes extinction of native biodiversity (Finer *et al.*, 2008; Cuellar and Noss, 2014; Gardner *et al.*, 2021). These abrupt modifications change the interactions among species occurring in the biome, which includes the dynamics that regulate the interactions between parasites and hosts (Gardner and Campbell, 1992b). The parasite checklist herein presented is a historical document that summarizes the marsupial parasite association present in Bolivia by the end of the Twentieth Century.

Of the 35 species of marsupials recorded for Bolivia we document the helminthological record for 17. Most of the specimens were collected in localities across the Chaco, and Yungas, with few individuals collected from the Amazon basin. The majority of the specimens representing both parasites and hosts were preserved and archived in relevant repositories for biodiversity, which include the Harold W. Manter Laboratory of Parasitology (Lincoln, Nebraska), the Museum of Southwestern Biology (Albuquerque, New Mexico), and the American Museum of Natural History (New York, New York). The records resulted from the synergistic effort of mammalogists and parasitologists participating in the Bolivian Faunistic Inventories completed by the end of the previous century (Anderson, 1997). The present checklist includes the helminths collected from didelphiomorphs and it expands on other published checklists that document the diversity of parasites in mammals present in Bolivia (Dick *et al.*, 2007; Pucu *et al.*, 2014; Sanchez *et al.*, 2018).

A greater effort is necessary to complete the inventory of the parasites of marsupials present in Bolivia, since 17 species of didelphiomorphs, plus *Lestoros inca* (Thomas), representative of Coenolestidae, have not been surveyed for helminths. Furthermore, the assessment of the helminth fauna associated with each species may be hindered by the relatively large sample size necessary to survey the parasite species richness in marsupials (Jiménez *et al.*, 2011; Byles *et al.*, 2013).

Nematodes represent the most diverse group in this checklist; it includes 22 species of which 9 are included in Viannaiidae and only occur in didelphiomorph marsupials. The second largest group includes cyclophyllidean tapeworms (6 species) followed by rhopaliid trematodes (3 species) and 1 species of acanthocephalan. From the total of parasite species, 16 are monoxenous, and include nematodes of the Viannaiidae, Oxyuridae, Trichuridae, Molineidae and Ancylostomatidae. The other 16 species are heteroxenous, and they depend on molluscs or insects to infect their definitive hosts. Heteroxenous species may be used as indicators of the local biodiversity by revealing the taxonomic and trophic levels that still function in any given locality (Gardner and Campbell, 1992b). A modest fraction of the organisms used in this checklist have served

as the foundation for systematic reviews and species descriptions for trematodes, cestodes and nematodes (Gardner and Campbell, 1992a; Haverkost and Gardner, 2008; Jiménez *et al.*, 2008; Gardner *et al.*, 2013), underscoring the relevance of species descriptions as the sole records documenting biodiversity.

As a consequence, the present checklist incorporates additional localities that better represent the distribution of parasites and their host spectrum. The results listed in this checklist should act as a starting point to build upon the diversity of mutualists, micropredators or parasites associated with marsupials. As stated elsewhere (Gardner and Campbell, 1992b; Wood *et al.*, 2023), the changes in the quality of the habitat and the diversity of organisms will determine the likelihood of heteroxenous species to complete their life cycle.

The identification of helminths herein presented documents their distribution in nearly 50% of the marsupial biodiversity of Bolivia. None of the species included in this checklist are known to have a zoonotic potential. Nevertheless this list should complement the efforts to screen these organisms for microparasites, which may be of zoonotic relevance and have been documented across Bolivia (Messenger *et al.*, 2015). This is important because several species of marsupials are synanthropic and thrive in human altered environments (Bezerra-Santos *et al.*, 2021; Voss, 2022). This innate ability makes it important to continue the surveillance of parasites in these mammals, since they can expose human populations to unanticipated outbreaks. The surveillance is an important component of proposed novel protocols to prevent outbreaks resulting from anthropogenic alterations (Hoberg *et al.*, 2022; Gardner *et al.*, 2023).

Data availability statement. The helminthological record for most of the marsupials examined is available at http://opensiuc.lib.siu.edu/zool_data/23/.

Acknowledgements. The following organizations provided logistic support in the field: Museo Nacional de Historia Natural, La Paz; The Museum of Southwestern Biology, University of New Mexico and Instituto Boliviano de Biología de la Altura, La Paz. We also thank the students and staff of Museums in Bolivia who are working hard to learn and conserve their biodiversity heritage. Dr Gabor Racz of the H.W. Manter Laboratory assisted in work in the laboratory, loans and deposition of specimens. Cristina Damborenea (CHLP), Luis García Prieto (CNHE), Eric Hoberg (USNPC), Luis Muniz (CHIOC), Graciela Navone, Judith Price (CMNA), and Lidia Sánchez (CHIAUMSM) made part of the material they have under their care available for comparisons.

Author contributions. S. L. G. conceived and designed the logistics to secure specimens in the field. S. L. G. M. L. C. and F. A. J. conducted data gathering. F. A. J., M. L. C., B. B. and R. P. S. performed analyses. F. A. J. and S. L. G. wrote the article.

Financial support. This work was supported by United States National Science Foundation Grants BSR8612329, BSR8612329, BSR9024816, DEB9496263, and DEB9631295 to S. L. G. BSR8408923 to T. L. Yates; BSR8316740 to S. Anderson. Additional support was provided directly by the American Museum of Natural History, The Museum of Southwestern Biology, the Tinker Foundation, SIU-Carbondale and the Harold W. Manter Laboratory of Parasitology.

Competing interests. The authors declare that there is no competing interest.

Ethical standards. Most recent version of protocols describing the use of vertebrates includes protocol 21-017 approved by IACUC of SIUC.

References

Acosta-Virgen K, Lopez-Caballero J, Garcia-Prieto L and Mata-Lopez R (2015) Helminths of three species of opossums (Mammalia, Didelphidae) from Mexico. *Zookeys* **511**, 131–152.

- Adnet FAO, Anjos DHS, Menezes-Oliveira A and Lanfredi RM (2009) Further description of *Cruzia tentaculata* (Rudolphi, 1819) Travassos, 1917 (Nematoda: Cruzidae) by light and scanning electron microscopy. *Parasitology Research* **104**, 1207–1211.
- Alden KJ (1995) Helminths of the opossum *Didelphis virginiana*, in southern Illinois, with a compilation of all helminths reported from this host in North America. *Journal of the Helminthological Society of Washington* **62**, 197–208.
- Anderson S (1997) Mammals of Bolivia, taxonomy and distribution. *Bulletin of the American Museum of Natural History* **231**, 1–652.
- Antunes GM (2005) *Diversidade e potencial zoonótico de parasitos de Didelphis albiventris Lund, 1841 (Marsupialia: Didelphidae)**, PhD thesis. Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil.
- Astúa D (2009) Evolution of scapula size and shape in Didelphid marsupials (Didelphimorphia: Didelphidae). *Evolution* **63**, 2438–2456.
- Auty RM (1994) The resource course thesis – minerals in Bolivian development, 1970–90. *Singapore Journal of Tropical Geography* **15**, 95–111.
- Benatti D, Moraes MFD, Pacheco CAA, Machado DMR, Oliveira WJ, Perin PP, Andrietti LF, Cândido Júnior JF, Vogliotti A, Tebaldi JH and Hoppe EGL (2023) Endoparasites of marsupials in fragments of the Atlantic rain-forest, western Paraná State, Brazil. *Revista Brasileira de Parasitologia Veterinária* **32**, e005823.
- Beveridge A, Spratt DM and Durette-Desset MC (2014) Order Strongylida (Railliet & Henry, 1913). In Schmidt-Rhaesa A (ed.), *Handbook of Zoology. Gastrotricha, Cycloneuralia and Gnathifera*, Vol. 2: *Nematoda*. Boston, Massachusetts: de Gruyter, pp. 557–612.
- Bezerra-Santos MA, Ramos RAN, Campos AK, Dantas-Torres F and Otranto D (2021) *Didelphis* spp. opossums and their parasites in the Americas: a one health perspective. *Parasitology Research* **120**, 4091–4411.
- Brant SV and Gardner SL (2000) Phylogeny of species of the genus *Litomosoides* (Nematoda: Onchocercidae): evidence of rampant host switching. *Journal of Parasitology* **86**, 545–554.
- Byles B, Catzeffis F, Scheibel RP and Jiménez FA (2013) Gastrointestinal helminths of two species of mouse opossums (*Marmosa demerarae* and *Marmosa murina*) from French Guiana. *Comparative Parasitology* **80**, 210–216.
- Caceres NC, Nápoli RP and Hannibal W (2011) Differential trapping success for small mammals using pitfall and standard cage traps in a woodland savannah region of southwestern Brazil. *Mammalia* **75**, 45–52.
- Campbell ML, Gardner SL and Navone GT (2003) A new species of *Mathevotaenia* (Cestoda: Anoplocephalidae) and other tapeworms from marsupials in Argentina. *Journal of Parasitology* **89**, 1181–1185.
- Cañeda-Guzmán IC (1997) *Parásitos de tres especies de marsupiales de la estación 'Los Tuxtles' y algunas zonas cercanas, Veracruz, México*, Thesis. Universidad Nacional Autónoma de México, Mexico City, Mexico.
- Chero JD, Sáez G, Mendoza-Vidaurre C, Iannacone J and Cruces CL (2017) Helminths of the common opossum *Didelphis marsupialis* (Didelphimorphia: Didelphidae), with a checklist of helminths parasitizing marsupials from Peru. *Revista Mexicana de Biodiversidad* **88**, 560–571.
- Cirino BS, Neto SFC, Maldonado AJ and Gentile R (2020) First study on the helminth community structure of the neotropical marsupial *Metachirus myosuros* (Didelphimorphia, Didelphidae). *Revista Brasileira de Parasitologia* **29**, e005420.
- Cuellar E and Noss AJ (2014) Mammal diversity and local participation in the conservation of the Bolivian Gran Chaco. *Therya* **5**, 39–60.
- Diaw OT (1976a) Coexistence des nématodes trichostrongyloides chez des marsupiaux de Guyane. *Annales de parasitologie humaine et comparée* **51**, 355–363.
- Diaw OT (1976b) Contribution à l'étude de nématodes Trichostrongyloidea parasites de Xenarthre, Marsupiaux et Rongeurs néotropicaux. *Bulletin du Muséum National d'Histoire Naturelle* **282**, 1065–1089.
- Dick CW, Gettinger D and Gardner SL (2007) Bolivian ectoparasites: a survey of bats (Mammalia Chiroptera). *Comparative Parasitology* **74**, 372–377.
- Dikmans G (1931) A new nematode worm *Viannaia bursobscura*, from the opossum with a note on the other parasites of the opossum. *Proceedings of the United States National Museum* **79**, 1–4.
- Dikmans G (1945) The occurrence of *Viannaia viannai* Travassos (Nematoda: Heligmosomidae) in opossums in North America. *Proceedings of the Helminthological Society of Washington* **10**, 6–8.
- dos Santos E (1968) Redescricao de *Mathevotaenia bivittata* (Janicki, 1904), parasito de marsupial (Cestoda, Linstowiinae). *Atas da Sociedade de Biologia do Rio de Janeiro* **11**, 193–194.
- Durette-Desset MC (1971) Essai de classification des nématodes Héligmosomes. Corrélations avec la paléobiogéographie des hôtes.

- Mémoires du Muséum National d'Histoire Naturelle. Série A. Zoologie* **69**, 1–126.
- Ellis RD, Pung OJ and Richardson DJ (1999) Site selection by intestinal helminths of the Virginia opossum (*Didelphis virginiana*). *Journal of Parasitology* **85**, 1–5.
- Ergueta P and Salazar J (1991) *Fauna silvestre de Bolivia*. La Paz, Bolivia: El Instituto Geográfico Militar and Centro de Datos Para la Conservación.
- Finer M, Jenkins CN, Pimm SL, Keane B and Ross C (2008) Oil and gas projects in the western Amazon: threats to wilderness, biodiversity, and indigenous peoples. *PLoS One* **3**, e2932.
- Flores DA (2009) Phylogenetic analyses of postcranial skeletal morphology in didelphid marsupials. *Bulletin of the American Museum of Natural History* **320**, 1–81.
- Fontúrbel FE and Jiménez JE (2009) Underestimation of abundances of the monito del monte (*Dromiciops gliroides*) due to a sampling artifact. *Journal of Mammalogy* **90**, 1357–1362.
- Foster AO (1939) Some helminths of the woolly opossum in Panama. *Transactions of the American Microscopical Society* **58**, 185–198.
- Fujita O, Abe N, Oku Y, Sanabria L, Inchausti A and Kamiya M (1995) Nematodes of armadillos in Paraguay: a description of a new species *Aspidodera esperanzae* (Nematoda: Aspidoderidae). *Journal of Parasitology* **81**, 936–941.
- Gardner AL (ed.) (2007) *Mammals of South America*. Chicago, IL: Chicago University Press.
- Gardner SL and Campbell ML (1992a) A new species of *Linstowia* (Cestoda: Anoplocephalidae) from marsupials in Bolivia. *Journal of Parasitology* **78**, 795–799.
- Gardner SL and Campbell ML (1992b) Parasites as probes for biodiversity. *Journal of Parasitology* **78**, 596–600.
- Gardner SL, Jiménez FA and Campbell ML (2013) *Pritchardia boliviensis* n. gen., n. sp. (Anoplocephalidae: Linstowinae), a tapeworm from opossums (Didelphidae) in the Yungas and lowlands of Bolivia and Atlantic forest of Paraguay. *Occasional Papers of the Texas Tech University Museum* **319**, 1–8.
- Gardner SL, Botero-Cañola S, Aliaga-Rossel E, Dursahinhan AT and Salazar-Bravo J (2021) Conservation status and natural history of *Ctenomys*, tuco-tucos in Bolivia. *Therya* **12**, 15–36.
- Gardner SL, Brooks DR, Boeger WA and Hoberg E (2023) *An Evolutionary Pathway for Coping with Emerging Infectious Disease*. Lincoln, NE: Zea Books.
- Gentile R, Finotti R, Rademaker V and Cerqueira R (2004) Population dynamics of four marsupials and its relation to resource production in the Atlantic forest in southeastern Brazil. *Mammalia* **68**, 109–119.
- Giarla TC, Voss RS and Jansa SA (2010) Species limits and phylogenetic relationships in the didelphid marsupial genus *Thylamys* based on mitochondrial DNA sequences and morphology. *Bulletin of the American Museum of Natural History* **346**, 1–67.
- Gomes DC, Cruz RPD, Vicente JJ and Pinto RM (2003) Nematode parasites of marsupials and small rodents from the Brazilian Atlantic rain forest in the State of Rio de Janeiro, Brazil. *Revista Brasileira de Biologia* **20**, 699–707.
- Guerrero R (1985) Nematoda: Trichostrongyloidea parásitos de mamíferos silvestres de Venezuela. II. Revisión del género *Viannaia* Travassos, 1914. *Memoria de la Sociedad de Ciencias Naturales La Salle* **45**, 9–47.
- Gutiérrez EE, Jansa SA and Voss RS (2010) Molecular systematics of mouse opossums (Didelphidae: *Marmosa*): assessing species limits using mitochondrial DNA sequences, with comments on phylogenetic relationships and biogeography. *American Museum Novitates* **3692**, 1–22.
- Guzmán-Cornejo C, García-Prieto L, Acosta-Gutiérrez R, Falcón-Ordaz J and León-Paniagua L (2012) Metazoarios parásitos de *Tlacuatzin canescens* y *Marmosa mexicana* (Mammalia: Didelphimorphia) de México. *Revista Mexicana de Biodiversidad* **83**, 557–561.
- Hancock L, Ralph N and Ali SH (2018) Bolivia's lithium frontier: can public private partnerships deliver a minerals boom for sustainable development? *Journal of Cleaner Production* **178**, 551–560.
- Haverkost TR and Gardner SL (2008) A review of species in the genus *Rhopalias* (Rudolphi, 1819). *Journal of Parasitology* **94**, 716–726.
- Hoberg EP, Boeger WA, Molnár O, Földvári G, Gardner SL, Juarrero A, Kharchenko VA, Ortiz E, Preiser W, Trivellone V and Brooks DR (2022) The DAMA protocol, an introduction: finding pathogens before they find us. *MANTER: Journal of Parasite Biodiversity* **21**, 1–20.
- Hodda M (2022) Phylum Nematoda: a classification, catalogue and index of valid genera, with a census of valid species. *Zootaxa* **5114**, 1–289.
- Jansa SA, Barker FK and Voss RS (2014) The early diversification history of didelphid marsupials: a window into South America's 'splendid isolation'. *Evolution* **68**, 684–695.
- Jiménez FA, Braun JK, Campbell ML and Gardner SL (2008) Endoparasites of fat-tailed opossums (*Thylamys*: Didelphidae) from northwestern Argentina and southwestern Bolivia, with the description of a new species of tapeworm. *Journal of Parasitology* **94**, 1098–1102.
- Jiménez FA, Gardner SL and Catzeffis F (2011) Structure of parasite component communities of didelphid marsupials: insight from a comparative study. *Journal of Parasitology* **97**, 779–787.
- Jiménez FA, Gardner SL, Navone GT and Ortí G (2012) Four events of host-switching in Aspidoderidae (Nematoda) involve convergent lineages of mammals. *Journal of Parasitology* **98**, 1166–1175.
- Jiménez-Ruiz FA, Gardner SL, Noronha D and Pinto RM (2008) The systematic position of Lauroinae Skrjabin and Schikhobalova, 1951 (Nematoda: Heterakoidea: Aspidoderidae), as revealed by the analysis of traits used in its diagnosis. *Cladistics* **24**, 459–476.
- José H, Macedo I and Loss MC (2019) A new and simple method to capture small arboreal mammals: the suspended pitfall. *Revista Brasileira de Zoociencias* **20**, 1–14.
- Lent H and Freitas JFTD (1935) Sobre dois novos nematodeos parasitos da quica: *Caluromys philander* (L.). *Memorias do Instituto Oswaldo Cruz* **30**, 535–542.
- Lima M, Stenseth NC, Yoccoz NG and Jaksic FM (2001) Demography and population dynamics of the mouse opossum (*Thylamys elegans*) in semi-arid Chile: seasonality, feedback structure and climate. *Proceedings of the Royal Society of London Series B-Biological Sciences* **268**, 2053–2064.
- Mariaux J, Tkach VV, Vasileva GP, Waeschenbach A, Beveridge I, Dimitrova YD, Haukismäki V, Greiman SE, Littlewood DTJ, Makarikov AA, Phillips AJ, Razafiarisolo T, Widmer V and Georgiev BB (2017) Cyclophyllidea van Beneden in Braun, 1900. In Caira JN and Jensen K (eds), *Planetary Biodiversity Inventory (2008–2017): Tapeworms from Vertebrate Bowels of the Earth*. Lawrence, KS: University of Kansas, Natural History Museum, pp. 77–148.
- Messenger LA, Garcia L, Vanhove M, Huaranca C, Bustamante M, Torrico M, Torrico F, Miles MA and Llewellyn MS (2015) Ecological host fitting of *Trypanosoma cruzi* TcI in Bolivia: mosaic population structure, hybridization and a role for humans in Andean parasite dispersal. *Molecular Ecology* **24**, 2406–2422.
- Mollericonna JL and Nallar R (2014) *Cruzia tentaculata* (Rudolphi, 1819) Travassos, 1917 in *Didelphis pernigra* (Allen, 1900) in the Acero Marka Valley of the Yungas of La Paz, Bolivia. *Neotropical Helminthology* **8**, 487–492.
- Monet-Mendoza A, Osorio-Sarabia D and García-Prieto L (2005) Helminths of the Virginia opossum *Didelphis virginiana* (Mammalia: Didelphidae) in Mexico. *Journal of Parasitology* **91**, 213–219.
- Navone GT, Suriano DM and Pujol CA (1991) *Travassostrongylus yungaensis* n. sp. and *Hoineffia simplicispicula* n. sp. (Nematoda: Trychostrongyloidea) from *Thylamys venustus* and *Lutreolina crassicaudata* (Marsupialia: Didelphidae) in the Northwest Argentina. *Systematic Parasitology* **19**, 187–193.
- Pires MP, Martins EG, Silva MNF and Reis SFD (2010) *Gracilinanus microtarsus* (Didelphimorphia: Didelphidae). *Mammalian Species* **42**, 33–40.
- Polo-Gonzales A, Sánchez L and Pacheco V (2019) Helminths of the genus *Didelphis* (Didelphimorphia: Didelphidae) of four regions in Peru. *Neotropical Helminthology* **13**, 273–286.
- Pucu E, Lareschi M and Gardner SL (2014) Bolivian ectoparasites: a survey of the fleas of *Ctenomys* (Rodentia: Ctenomyidae). *Comparative Parasitology* **81**, 114–118.
- Püttker T, Meyer-Lucht Y and Sommer S (2008) Effects of fragmentation on parasite burden (nematodes) of generalist and specialist small mammal species in secondary forest fragments of the coastal Atlantic Forest, Brazil. *Ecological Research* **23**, 207–215.
- Radev V, Gardner AL and Kanev I (2005) Family Rhopalidae Looss, 1899. In Jones A, Bray RA and Gibson DI (eds), *Keys to the Trematoda*, Vol. 2. Oxfordshire, UK: CABI Publishing, pp. 119–121.
- Ramírez-Cañas SA, López-Caballero JD and Mata-López R (2021) Morphological and molecular data reveal two new species of *Viannaia* (Nematoda: Viannidae), parasitizing opossums (Mammalia: Didelphidae) in Mexico. *Journal of Parasitology* **107**, 388–403.
- Richardson DJ, Gardner SL and Allen JW (2014) Redescription of *Oligacanthorhynchus microcephalus* (Rudolphi, 1819) Schmidt 1972 (syn.

- Oligacanthorhynchus tortuosa* (Leidy, 1850) Schmidt 1972 (Acanthocephala: Oligacanthorhynchidae). *Comparative Parasitology* **81**, 53–60.
- Rossi RV, Voss RS and Lunde DP** (2010) A revision of the didelphid marsupial genus *Marmosa* part 1. The species in Tate's 'mexicana' and 'Mitts' sections and other closely related forms. *Bulletin of the American Museum of Natural History* **334**, 3–83.
- Ruiz JM** (1947) *Revisão gênero Cruzia (Nematoda: Oxiuroidea) e estudo das espécies Brasileiras*. Thesis. Universidade de São Paulo, São Paulo, Brazil.
- Sanchez J, Lareschi M, Salazar-Bravo J and Gardner SL** (2018) Fleas of the genus *Neotyphloceras* associated with rodents from Bolivia: new host and distributional records, description of a new species and remarks on the morphology of *Neotyphloceras rosenbergi*. *Medical and Veterinary Entomology* **32**, 462–472.
- Santos CP, Lent H and Gomes DC** (1990) The genus *Aspidodera* Railliet and Henry, 1912 (Nematoda: Heterakoidea): revision, new synonyms and key for species. *Revista Brasileira de Biologia* **50**, 1017–1031.
- Scheibel RP, Catzeflis F and Jiménez FA** (2014) The relationships of marsupial-dwelling Viannaiidae and description of *Travassostrongylus scheibelorum* n. sp. (Trichostrongylina: Heligmosomoidea), from mouse opossums (Didelphidae) from French Guiana. *Folia Parasitologica* **61**, 242–254.
- Siebert AE** (1970) A new record for the trematods *Rhopalias macracanthus* Chandler, 1932, and a key to the genus *Rhopalias* Stiles and Hassel, 1989. *Proceedings of the Louisiana Academy of Sciences* **33**, 35–37.
- Silva MGQE and Costa HMA** (1999) Helminths of white-bellied opossum from Brazil. *Journal of Wildlife Diseases* **35**, 371–374.
- Souza R, Vilela RDV, Gentile R, Lopes-Torres EJ, Cordeiro-Estrela P, Moratelli R, da Costa-Neto SF, Cardoso TDS, Varella K and Maldonado Júnior A** (2022) Population genetic structure and morphological diversity of *Cruzia tentaculata* (Nematoda: Ascaridida), a parasite of marsupials (Didelphinae), along the Atlantic Forest on the eastern coast of South America. *Parasitology* **149**, 1487–1504.
- Tantaleán M and Chavez J** (2004) Endoparasitos (Nemathelminthes y Platyhelminthes) de animales de vida silvestre de la Reserva de Biósfera del Manu, Perú. *Revista peruana de biología* **11**, 219–222.
- Teta P, D'Elia G, Flores DA and de la Sancha NU** (2009) Diversity and distribution of the mouse opossums of the genus *Thylamys* (Didelphimorphia, Didelphidae) in northeastern and central Argentina. *Gayana* **73**, 180–199.
- Torres EL, Maldonado A and Lanfredi RM** (2007) *Pterygodermatites (Paucipectines) jagerskioldi* (Nematoda: Rictulariidae) from *Gracilinanus agilis* and *G. microtarsus* (Marsupialia: Didelphidae) in Brazilian pantanal and Atlantic forest by light and scanning electron microscopy. *Journal of Parasitology* **93**, 274–279.
- Torres E JL, Maldonado A and Lanfredi RM** (2009) Spirurids from *Gracilinanus agilis* (Marsupialia: Didelphidae) in Brazilian Pantanal wetlands with a new species of *Physaloptera* (Nematoda: Spirurida). *Veterinary Parasitology* **163**, 87–92.
- Varella K, Vilela RDV, Gentile R, Cardoso TDS, da Costa-Neto SF and Maldonado Júnior A** (2022) Population genetic structure and phenotypic diversity of *Aspidodera raillieti* (Nematoda: Heterakoidea), a parasite of Didelphini marsupials in Brazil's South and Southeast Atlantic Forest. *Parasites & Vectors* **15**, 203.
- Voss RS** (2022) An annotated checklist of recent opossums (Mammalia: Didelphidae). In Knight M (ed.), *Bulletin of the American Museum of Natural History*, Vol. **455**. New York, NY: American Museum of Natural History, p. 74.
- Voss RS and Jansa SA** (2009) Phylogenetic relationships and classification of didelphid marsupials, and extant radiation of New World metatherian mammals. *Bulletin of the American Museum of Natural History* **322**, 1–177.
- Voss RS, Pine RH and Solari S** (2012) A new species of the Didelphid marsupial genus *Monodelphis* from eastern Bolivia. *American Museum Novitates* **3740**, 1–14.
- Wolfgang RW** (1951) Studies on the endoparasitic fauna of Trinidad Mammals. VIII. Parasites of marsupials. *Canadian Journal of Zoology* **29**, 352–373.
- Wood CL, Welicky RL, Preisser WC, Leslie KL, Mastick N, Greene C, Maslenikov KP, Tornabene L, Kinsella JM and Essington TE** (2023) A reconstruction of parasite burden reveals one century of climate-associated parasite decline. *Proceedings of the National Academy of Sciences* **120**, e2211903120.