



# Immunological differences between heart- and kidney-transplanted children: a cross-sectional study – CORRIGENDUM

## Corrigendum

**Cite this article:** Ekman-Joelsson B-M, Brandström P, Allén M, Andersson B, Wåhlander H, Mellgren K, and Ekwall O (2023) Immunological differences between heart- and kidney-transplanted children: a cross-sectional study – CORRIGENDUM. *Cardiology in the Young* 33: 842. doi: [10.1017/S1047951122002347](https://doi.org/10.1017/S1047951122002347)

First published online: 22 July 2022

Britt-Marie Ekman-Joelsson, Per Brandström, Maria Allén, Bengt Andersson, Håkan Wåhlander, Karin Mellgren and Olov Ekwall

DOI: <https://doi.org/10.1017/S1047951122001743> Published online by Cambridge University Press: 24 June 2022

The authors apologise that upon publication of Ekman-Joelsson, B., Brandström, P., Allén, M., Andersson, B., Wåhlander, H., Mellgren, K., & Ekwall, O. (2022), the first line of the abstract was incomplete.

The full abstract is as below

### Abstract

Post-transplantation lymphoproliferative disorder is a potentially mortal complication after heart transplantation in children. As the immune system plays a crucial role in the development of lymphoma, we explored the influence of thymus function in relation to immunosuppressive treatment in organ-transplanted children and healthy control subjects. A prospective case-control study was performed at a single centre, in which 36 children who had undergone heart transplantation were compared to two control groups: 34 kidney-transplanted children and 33 healthy age- and sex-matched children. T- and B-lymphocyte subtypes and monocytes were analysed by flow cytometry, and T-cell receptor excision circles were assessed using quantitative polymerase chain reaction. Heart-transplanted children had a lymphocyte profile characterised by reduced or absent thymic function with low numbers of T-cell receptor excision circles and total and naïve T cells, together with immune activation against the allograft. Despite similar immunosuppressive treatment, the kidney-transplanted group showed an activated T-lymphocyte compartment.

The online version of this article has been updated.

### Reference

Ekman-Joelsson, B., Brandström, P., Allén, M., Andersson, B., Wåhlander, H., Mellgren, K., & Ekwall, O. (2022). Immunological differences between heart- and kidney-transplanted children: A cross-sectional study. *Cardiology in the Young*, 1-6. doi: [10.1017/S1047951122001743](https://doi.org/10.1017/S1047951122001743)

© The Author(s), 2022. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

