

Cité, Paris, France; ⁴INSPECT-LB (Institut National de Santé Publique, d'Épidémiologie Clinique et de Toxicologie-Liban), Beirut; ⁵School of Medicine, Lebanese American University, Byblos, Lebanon; ⁶Department of Primary Care and Population Health, University of Nicosia Medical School, Nicosia, Cyprus; ⁷Faculty of Pharmacy, Lebanese University, Hadat, Lebanon; ⁸Faculty of Pharmacy, Université Laval and ⁹Oncology Division, CHU de Québec-Université Laval Research Center, Quebec, Canada

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.305

Introduction: Alcohol and sedative substance use disorders are escalating global public health challenges. Lebanon has grappled with multiple crises, including economic, healthcare, and social issues.

Objectives: This study aimed to assess the correlates of the alcohol and sedative substance use risk scores with sociodemographic and clinical factors, including sleep disorders, chronotype, anxiety, and depression.

Methods: A cross-sectional study was conducted among the Lebanese population using several validated scales to assess the risk of alcohol and sedative substance use, including the Alcohol, Smoking, and Substance Involvement Screening Test (ASSIST). Other tools evaluated chronotype, sleep, and mood disturbances. Bivariate and multivariable analyses were then performed, taking the alcohol and sedative scores as dependent variables.

Results: A total of 646 participants were included. Multivariate analysis revealed positive and significant correlations between higher ASSIST-alcohol scores and personal history of alcohol abuse ($B=4.61$), family history of prescription substance abuse ($B=1.763$), psychiatric disorders ($B=2.898$), and worse Insomnia Severity Index scores ($Beta=0.14$). Conversely, ASSIST-alcohol scores negatively correlated with weight ($B=-0.39$) and morning chronotype ($B=-0.084$). Positive correlations were identified between higher ASSIST-alcohol scores and personal history of illicit substance abuse ($B=2.834$), prescription substance abuse ($B=2.252$), sleep quality ($B=0.130$), and sleep severity ($B=0.082$), while negatively correlating with cigarette smoking ($B=-0.038$).

Conclusions: This study elucidates the role of several predisposing factors to alcohol and sedative use disorders in Lebanon, including history of substance abuse, psychiatric disorders, sleep disorders, and chronotype. These findings advocate, in particular, for the integration of sleep disorder assessment and management into addiction rehabilitation programs.

Disclosure of Interest: None Declared

EPP0077

Changing drinking patterns among Italians: 7 out of 10 students experience Binge Drinking

F. Marcolini*, D. De Ronchi and A. R. Atti

Department of Biomedical and Neuromotor Sciences, University of Bologna, Bologna, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.306

Introduction: The expression Binge Drinking (BD) refers to dysregulated alcohol consumption, characterized by the intake of large

quantities of alcohol, regardless of their nature, consecutively in a limited period of time. BD is a significant public health problem in many European countries, including Italy. According to data from the *Istituto Superiore di Sanità*, dated 2020, over 4 million Italians exhibit episodic excessive alcohol consumption (compared to 2019 data, there was an increase of approximately 5,3%).

Objectives: This study aims to examine alcohol consumption habits in the Italian population, evaluating psychopathological correlations that can explain its diffusion.

Methods: Between January and May 2023, an anonymous online questionnaire was randomly sent to the general population. Alongside with tests to evaluate psycho-social features, to estimate the presence of alcohol abuse or dependence the AUDIT scale (Saunders *et al.* *Addict Abingdon Engl.* 1993; 88:791–804) was used. It included two specific questions to frame the phenomenon of BD (Cranford *et al.* *Alcohol Clin Exp Res.* 2006; 30:1896–905). No other study conducted in Italy has so far used the aforementioned validated questions.

Results: The sample consists of 308 people (189 F, 119 M), with an average age of 32 years (sd 14). The AUDIT indicates a state of chronic alcohol consumption in 11,7% (95% confidence interval 8,5%-15,7%), of the recruited sample, positively correlating with the element of impulsivity ($p<0,005$) confirming what has already been reported in literature. BD prevalence reaches 56% (M 57%, F 55%) without any significant correlation with impulsivity, personality disorders, emotional dysregulation, or sensitivity to rejection. Among university students the prevalence of BD exceeds 70% (95% confidence interval 60%-76%), with a number of drinks reported for a single occasion reaching up to 25 units and a reported number of binge episodes, in a two-week span, ranging from 2 to 10.

Conclusions: Despite possible *biases*, this study raises the relevant issue of the extremely high prevalence of BD disorder, which is particularly alarming in light of the numerous issues related to the behavior itself. A direct correlation with reduced school performance, an increase in risky sexual behavior, and an increase in cases of drunk driving have been evaluated. Considering these consequences, it is of primary importance on a medical, but even more social level, to best characterize this phenomenon in such a way as to be able to implement awareness-raising and prevention interventions.

Disclosure of Interest: None Declared

EPP0079

A preliminary analysis of clinical characteristics of patient with alcohol use disorder and suicidal ideation

R. F. Palma-Alvarez*, A. Rios-Landeo, G. Ortega-Hernandez, E. Ros-Cucurull, C. Daigre, M. Perea-Ortueta, M. Sorribes, L. Grau-López and J. A. Ramos-Quiroga

Hospital Universitari Vall d'Hebron, Barcelona, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.307

Introduction: Suicidal behaviors are frequently observed among patients with substance use disorder, including suicidal ideation (SI) (1). Alcohol use disorder (AUD) is one of the most prevalent addictions and may be related to suicidal behaviors (2,3). However,