

VOL. III.—1889.

c

THE JOURNAL
OF
LARYNGOLOGY
AND
RHINOLOGY;

*AN ANALYTICAL RECORD OF CURRENT LITERATURE RELATING TO
THE THROAT AND NOSE.*

EDITED BY

SIR MORELL MACKENZIE, M.D. Lond.

AND

R. NORRIS WOLFENDEN, M.D. Cantab.

WITH THE CO-OPERATION OF

DR. JOHN N. MACKENZIE (BALTIMORE)
DR. BRYSON DELAVAN (NEW YORK)
DR. W. PORTER (ST. LOUIS)
DR. JOAL (MONT DORE)
PROF. MASSEI (NAPLES)
PROF. GUYE (AMSTERDAM)
DR. CAPART (BRUSSELS)
DR. CONSTANTIN KARWOWSKI (WARSAW)
DR. BARCLAY J. BARON (BRISTOL)

DR. VALERIUS IDELSON (BERNE)
DR. MICHAEL (HAMBURG)
PROF. RAMON DE LA SOTA Y LASTRA
(SEVILLE)
DR. HOLGER MYGIND (COPENHAGEN)
DR. SMYLY (DUBLIN)
DR. DAVID COLLINGWOOD (SYDNEY)
DR. G. W. MAJOR (MONTREAL)
DR. MAXWELL ROSS (EDINBURGH)

AND

Dr. G. HUNTER MACKENZIE (Edinburgh).

Published Monthly.

LONDON:
R. ANDERSON & CO.,
14, COCKSPUR STREET,
CHARING CROSS, S.W.

PHILADELPHIA:
P. BLAKISTON, SON & CO.,
1012, WALNUT STREET.

c

CONTENTS.

Original Article, p. 403.—Notes on a Case of Primary Sarcoma of the Tonsil, by Dr. R. NORRIS WOLFENDEN.

Therapeutics, Diphtheria, &c., p. 405.—SUCHANNEK: Sozoiiodol. GOUGUENHEIM: Salol in Throat Inflammations. FRITSCHÉ, SCHNIBER, MUGDAN: Whooping Cough. MEIG'S Opium in Spasmodic Croup. NELSON, RACHFORD: Etiology of Diphtheria. NEW YORK MEDICAL JOURNAL: Transmissibility of Diphtheria. CHANTMESSE and VIDAL, JACOBI, CALDWELL: Treatment of Diphtheria. JACKSON, MCCOLLOM: Notes on Diphtheria in Boston. BOKAI: Pharyngeal Diphtheria. HAILES, DILLON BROWN: Intubation in Diphtheria.

Mouth, Pharynx, Œsophagus, &c, p. 411.—WYETH: Extirpation of both Parotids for Sarcoma. MILONAS: Double Parotitis of Reflex Origin. LIMBECK: Hemiatrophic Tongue. KIRCHHOFF: Neoplasms of the Tongue, WARREN, KRAUSE: Cancer of Tongue. VERGELY: Syncope and Convulsions from lowering of the Tongue. FOURNIER: Gummata of Palatine Arches. MACCOY: Sarcoma of the Tonsil. JUHEL-RÉNOY: Gummata of Tonsil. ZISMAN: Excision of Enlarged Tonsils. MCBURNEY: Perforation of Hard Palate. CLAIBORNE: Papilloma of Fauces. LAVRAND: Œdema of Eyelids and Facial Erysipelas. MAURIN: Primary Gangrenous Angina. DELAVAN: Adenoid Hypertrophy of Pharynx. RAULIN: Tuberculosis of Pharynx. FRAENKEL: Syphilitic Stricture of Pharynx. VANDERPOEL: Pharyngeal Mycosis. JOAL, OSGOOD: Œsophagismus. ALSBERG: Carcinoma of Œsophagus. LUTZ: Gastrostomy. SCHULZ, IVANHOFF, WAHL: Foreign Bodies in Œsophagus.

Nose and Naso-Pharynx, p. 421.—RICE: Antiseptic Nasal Surgery. POTTER: Membranous Rhinitis. SCHUCHARDT: Nature of Ozena. NOQUET: Papilloma of Nasal Fossa. THORNER: Hematoma of Septum) CHEATHAM: Diseases of Eye, Ear, Throat, and Nose. WAGNIER: Modified Employment of Galvanic Loop for Polypi. WRIGHT: Œdema of Lip and Intranasal Pressure. FOSTER: Obstruction of Nose and Cause of Nervousness. SOKOLOFF: Foreign Bodies in the Nose. GAREL: Rhinolith. LUBET-BARBON, COUSTOUX: Adenoid Vegetations of the Naso-Pharynx. ROLLAND: Dilatation of Sphenoidal Sinus.

Larynx, &c., p. 425.—FRENCH: Photography of Larynx during formation of Singing Registers. VINEBERG: Laryngismus Stridulus. NATIER: Lupus. VOLTOLINI: Operations under Electric Light. HEYMANN, LUBLINSKI, SCHADEWALT, FRAENKEL, MARTIUS: Cases at the Berlin Laryngological Society. RUAULT: Catamenial Laryngeal Hemorrhages. MOURE: Laryngeal Hæmorrhage. GAREL: Laryngeal Ictus. NATIER: Œdema of Vocal Cords, &c. AUDUBERT: Dry Laryngitis. TCHERNOFF: Traumatic Laryngitis. RICARDO-BOTEY: Laryngeal Polypus. GOUGUENHEIM: Dilatation in Laryngeal Stricture. KRAUSE: New Methods of Treating Laryngeal Tuberculosis. LOCOARRET: Syphilis of Ventricular Bands. CARTAZ: Late Hereditary Syphilis. TCHERNOFF: Laryngitis Syphilitica. BESCHORNER: Carcinoma of Larynx. SCHMIDT: Laryngeal Surgical Casuistics. WHARTON, VIERHUFF: Foreign Bodies in Air Passages. GANGHOFNER: Fatal Hæmorrhage after Tracheotomy. ALEXANDROFF: Foreign Body in the Trachea.

Neck, Thyroid, &c, p. 433.—HÜBER: Cervico-Tracheal Fistula. FOWLER: Hernia of the Pleura. SCHOETZ: Erythema Multiforme. WEIR: Thyroid Tumours. WELCH: Cystic Goitre. LACAÏLLE, AUDRY: Exophthalmic Goitre. WÖFLER: Struma Migrans.

Association Meetings, p. 434.—Report of the Fortieth Annual Meeting of the American Medical Association.

Notes, p. 444.—A New Collaborator. The American Public Health Association.

SOME GOOD POINTS.

LONDON.

OCTOBER.

1889.

SULPHONAL-BAYER AS AN HYPNOTIC.

From a very extensive use of Sulphonal in private and hospital practice, covering cases of insanity [mania and melancholia], delirium tremens, morphine habit, sleeplessness from mitral insufficiency, from syphilis, from malarial poisoning, from anaemia, and in Bright's disease, the following conclusions have been arrived at, viz. :—

1. It is a safe hypnotic, of remarkable intrinsic value, especially in cases of insanity. In all cases reported, except one, refreshing slumber of from two to eight hours' duration was produced by fifteen grains, dry or in hot soup, repeated in one hour if necessary.

2. It is effective in those cases where bromides, chloral, paraldehyde and hypnone have been tried in vain.

3. It has a secondary effect, *i.e.*, its calmative power seems to extend over one day, and exert a modified hypnotic influence the second night after its exhibition.

4. Its advantages over other hypnotic remedies are that it does not constipate as do the preparations of opium; the sleep produced by it seems to more closely resemble natural sleep than does that of any other drug; having neither taste nor odour, it is easily given in cases where patients object to take medicines.

5. In morphine habit it is regarded as the most valuable remedy yet discovered to control insomnia, and with a 1/75 grain Tabloid of Hydrobromate of Hyoscine hypodermically for delirium, these usually troublesome cases are being treated with marked success.

Sulphonal Tabloids (5 gr.)—the most eligible form of the drug—are supplied in bottles containing 25 and 100 at 1s. and 3s. each (subject to fluctuation of the market).

Sulphonal (Bayer) supplied to the medical profession at 1s. 6d. per oz. (subject to the fluctuation of the market).

LANOLINE FOR THE SKIN AND HAIR.

Lanoline is the purified fat contained in the wool of sheep. Dr. Oscar Liebreich, of the University of Berlin, the discoverer of chloral, says that "wool-fat" was used in the time of Herodotus under the name of *Oesypus*, and his description is reproduced in most of the pharmacopœias published up to the eighteenth century. It was also used by the ancients as a cosmetic, and Ovid mentions in his "*Ars Amatoria*," that the Roman ladies anointed their faces and skin with *Oesypus* brought from Athens. Certain soaps, puffed up by means of cunningly-devised advertisements, often cause pimples, blotches and sores on the

face and prevent eruptions already formed from healing. They remove the fat and dry the skin, so that this tissue becomes hard, inelastic and rough, and loses its natural colour. Lanoline Soap, on the other hand, containing as it does a fat which is natural to, and absorbed by, the skin is acknowledged by dermatologists to be the very best preservative of this tissue, restoring its soft, pliable and elastic nature, and healthful surface.

Lanoline has also been acknowledged by the highest authorities as a hair preservative. "Hair can no longer be regarded as an effete and dead tissue, but in a certain though limited sense an organ capable of elaboration. During its growth it can elaborate a cholesterine fat compatible with water, and it is this which maintains the hair lustrous and soft without greasiness." Lanoline (Liebreich) contains this cholesterine principle and does not undergo decomposition like other fats. Lanoline Pomade since its introduction has been recommended by the profession as the natural unguent for preserving the hair. "The fatty acids produced by the decomposition of oils and fats are peculiarly injurious, and it is to be feared that the existence of such in many of the ordinary pomades is only concealed by the perfume."—JAMESON.

Lanoline (Liebreich) supplied to the medical profession in 1 lb. and 7 lb. tins at 2s. 8d. per lb. Anhydrous Lanoline (Lanolinum Anhydricum), at 3s. 4d. per lb. Lanoline Base (Ung. Lanolini), 2s. 2d. per lb. Anhydrous-Lanoline Base, 2s. 10d. per lb. Corecin, 1s. 6d. per lb.; Paraffinum Liquidum, 1s. per lb. (packages extra). Lanoline Cold Cream supplied in pots at 1s. 2d. each. Lanoline Hazeline Ointment, in pots at 1s. 2d. each. Lanoline Pomade, in pots at 1s. 2d. each. Lanoline Toilet Soap, 2s. per box of 3 Tablets. Lanoline Eucalyptine Soap, 2s. per box of 3 Tablets. Lanoline Pinol Soap, 2s. per box of 3 Tablets. Toilet Lanoline, in tubes at 5d. each. Lano-Creolin supplied in Collapsible Tubes.

A Pamphlet on the "Place of Lanoline in Pharmacy" will be sent post free to any Physician on request.

THE VERDICT FOR THE KEPLER EXTRACT OF MALT.

The *Lancet* reports :—"Kepler's is the best known, and in this country the largest used extract of Malt. It is as distinct an advance in therapeutics as was the introduction of cod liver oil. It is one of the best remedies for atonic dys-pepsia, and is undoubtedly useful in consumption and other wasting diseases."

The *British Medical Journal* reports :—"We can recommend the Kepler Malt Extract.—Being prepared at a very low temperature, its qualities are not deteriorated or its flavour spoiled.—It is very favourably spoken of by physicians both in respect to its nutritive and digestive properties and as being of very agreeable flavour."

The *Medical Record* reports :—"Kepler Extract of Malt is undoubtedly the best, the richest in diastase, and the most largely used . . . It is the Extract of Malt which every physician prescribes."

Kepler Extract of Malt and Kepler Extract of Malt with Cod Liver Oil (Kepler Solution) supplied to the medical profession in bottles, containing $\frac{1}{4}$ lb., and $\frac{1}{2}$ lb., at 1s. 10d. and 3s. 2d. Retail prices, 2s. 6d. and 4s. Special terms to Hospitals. Supplied by all Chemists throughout the World.

BURROUGHS, WELLCOME & Co., Snow Hill Buildings, London, E.C.

Will be happy to supply trial Specimens to Medical Men free on request.