

## Riboflavin supplementation in a resettlement village in north-east Thailand

BY D. I. THURNHAM,\* P. MIGASENA, NIYOMSRI VUDHIVAI  
AND VENUS SUPAWAN

*Department of Tropical Nutrition, Faculty of Tropical Medicine,  
Mahidol University, Bangkok, Thailand*

(Received 1 September 1971 – Accepted 13 December 1971)

1. A high prevalence of biochemical ariboflavinosis was shown to be present in the preschool children and a small group of adults in a village in north-east Thailand using an *in vitro* test based on erythrocyte glutathione reductase (EGR) activity.
2. A riboflavin supplement of 10 mg/d was given to all subjects for 7 d.
3. It was demonstrated that the riboflavin supplement cured the biochemical ariboflavinosis and the results supported the previous suggestion that a stimulation  $\geq 20\%$  of the *in vitro* EGR activity was a good indicator of ariboflavinosis.
4. A dietary survey carried out on twenty preschool children during the experimental period showed riboflavin intake to be low,  $0.18 \pm 0.06$  mg/Mcal.
5. No glucose-6-phosphate dehydrogenase (G-6-PD)-deficient subjects were shown to be deficient in riboflavin by the *in vitro* test. The possibility of using *in vivo* stimulation of EGR activity by riboflavin is discussed as a possible way of demonstrating ariboflavinosis in G-6-PD-deficient subjects.

The diet of rural Thai people has been shown to be deficient in riboflavin, and clinical signs commonly associated with ariboflavinosis, glossitis and angular stomatitis have frequently been found (Chandrapond & Ritchie, 1952; Chandrapond, 1955; Hauck & Sudsaneh, 1959; Interdepartmental Committee on Nutrition for National Defense, 1962). In this department studies were undertaken to investigate the use of the erythrocyte glutathione reductase (EGR) test of Glatzle, Körner, Christeller & Wiss (1970) to assess biochemically the prevalence of ariboflavinosis. These studies were carried out in two villages in north-east Thailand, and the prevalence of ariboflavinosis, although fluctuating with the seasons, was generally found to be high (Thurnham, Migasena & Pavapootanon, 1970; Thurnham, Migasena & Vudhivai, 1971). A good correlation has been found between the dietary intake of riboflavin and the prevalence of biochemical ariboflavinosis (Thurnham, Migasena, Vudhivai & Supawan, 1971), but it was considered that additional evidence, that the EGR test measured ariboflavinosis, would be gained by a supplementation experiment. This was done and the results are reported in this paper.

\* Present address: Department of Human Nutrition, London School of Hygiene and Tropical Medicine, Keppel Street (Gower Street), London WC1E 7HT.

## METHODS

*Subjects.* The experiment was carried out with preschool children and their parents in a resettlement village in north-east Thailand, in the dry season between February and March 1971.

*Blood samples.* Blood samples were taken from sixty children and twenty-five parents (Table 1). Seven of the children were found to be glucose-6-phosphate dehydrogenase (G-6-PD) deficient, but unfortunately the true number of G-6-PD-deficient parents in the initial group is not known as the reagents were not working satisfactorily on the day the parents were examined. Three parents in the sample group, however, were later found to be G-6-PD deficient.

*Collection and preservation of haemolysates*

Capillary blood was collected in Unibore heparinized capillary tubes (Harshaw Chemicals Ltd, London) and the packed cells were used to prepare haemolysates as described by Thurnham *et al.* (1970). EGR activity was measured in 100  $\mu$ l samples of haemolysate by the method of Glatzle *et al.* (1970) except that the concentration of reduced nicotinamide adenine-dinucleotide phosphate (NADP) was doubled to 0.4  $\mu$ mol/cuvette. EGR activity in each haemolysate with and without flavin-adenine dinucleotide (FAD) was measured by observing the decrease of reduced NADP at 334 nm, and the ratio of the stimulated over the basic activity was used to calculate an activation coefficient ( $\alpha$ ). Enzyme units are expressed as  $\mu$ mol reduced NADP oxidized by EGR in 1 ml erythrocytes in 10 min at 35°.

*Dietary studies.* Nine families containing twenty children less than 8 years old were selected randomly from the participants who began the supplementation study. Each family was visited for 1 full day and the food intake of each child was measured by methods described elsewhere (Thurnham, Pongpaew, Migasena, Supawan and Changbumrung, in preparation).

*Riboflavin supplement.* Ten tablets, each containing 10 mg riboflavin (Atlantic Trading Co. Ltd, Bangkok), were given to each person with the instructions to take one tablet each day for 10 d. Each family was visited on the 5th day and then daily until the 10th day to check that the instructions were being followed. At least seven tablets were taken during the treatment period by each person included in the results.

*G-6-PD deficiency.* G-6-PD deficiency was detected by the method of Bernstein (1962).

## RESULTS

*Biochemical findings*

It was reported by Flatz (1970) that G-6-PD deficiency may alter the binding properties of EGR for FAD. In view of this, the results obtained for those persons with G-6-PD deficiency are treated separately in this report, but discussion concerning the G-6-PD-deficient subjects will be brief as it is dealt with more fully elsewhere (Thurnham, 1972).

Table 1. *Basic erythrocyte glutathione reductase (EGR) activity, determined in vitro, of haemolysates of preschool children and of their parents before and after receiving a riboflavin supplement*

(Mean values and standard deviations; size of group shown in parentheses)

G-6-PD status	EGR activity (units†)			
	Children		Parents	
	Before	After	Before	After
Normal				
Whole group	12.1 ± 3.6 (53)	—	11.3 ± 4.3	—
Sample	12.3 ± 3.9 (22)	17.4 ± 3.0 (22)***	10.1 ± 2.9 (10)	16.1 ± 3.8 (10)***
Deficient				
Whole group	21.6 ± 4.0 (7)	—	—	—
Sample	20.8 ± 3.5 (3)	22.4 ± 1.0 (3)	20.0 ± 2.9 (3)	22.7 ± 4.1 (3)

'Sample' refers to those subjects who completed the supplementation experiment, i.e. took at least 10 mg riboflavin/d for 7 d. Differences between the 'sample' and 'whole' groups were not significant.

\*\*\* Increase in EGR activity was significant ( $P < 0.001$ ) in both children and parents when tested by a test for paired values.

†  $\mu\text{mol}$  reduced NADP oxidized by EGR in 1 ml erythrocytes in 10 min at 35°.

The results in Table 1 show the mean values of the basic EGR activity before and after riboflavin supplementation. The 'sample' group refers to those children and parents who returned to give a second sample of blood 15 d later and who had taken the supplementary riboflavin as directed (see page 92). The values for basic EGR activities of all the sample groups before taking the supplement shown in Table 1 did not differ significantly from those for the respective whole groups.

After the riboflavin supplement had been taken, the mean values of the basic EGR activities of all sample groups increased (Table 1). The increase in EGR activity was inversely proportional to the EGR activity before supplementation and only three individuals of those who completed the experiment showed results that differed from those given in Table 2. Two of these were children with high basic EGR activities and low activation coefficients before supplementation, and it was probable that they were not deficient in riboflavin at the start. The third person was one of the parents whose initial basic EGR activity was low (9.6 units) but whose activation coefficient indicated no ariboflavinosis ( $\alpha = 1.05$ ). On supplementation, however, the basic EGR activity only increased by 0.4 activity units but the activation coefficient increased to 1.24. It is possible that there may have been a technical error in the assay and this would partly explain the results; alternatively he may not have taken the tablets as he reported. The mean stimulation of the basic EGR activity in response to the riboflavin supplement of the G-6-PD-normal parents was  $67 \pm 51\%$  and for the children in the same group it was  $56 \pm 31\%$ .

The prevalence of ariboflavinosis in February 1971, as measured by the activation

Table 2. Correlation between the degree of ariboflavinosis as measured by activation coefficient and the resultant increase in erythrocyte glutathione reductase (EGR) activity, determined *in vitro*, caused by riboflavin supplementation in preschool children

(Mean values and standard deviations; no. of subjects in parentheses)

G-6-PD status	Before supplementation		After supplementation	
	Activation coefficient	Basic EGR activity (units†)	Increase in basic EGR activity (units†)	Stimulation in basic EGR activity (%)
Normal	≥ 1.20	9.3 ± 1.9 (12)	7.2 ± 1.8 (12)***	77 ± 19 (12)***
	1.10-1.19	13.0 ± 1.5 (5)	3.4 ± 1.6 (5)**	27 ± 14 (5)*
	< 1.10	18.1 ± 1.4 (5)	2.9 ± 2.2 (4)‡	16 ± 14 (4)‡
Deficient	< 1.10	20.8 ± 3.5 (3)	2.1 (2)‡	12 (2)‡

Activation coefficients were calculated by dividing the stimulated EGR (i.e. determined in the presence of added FAD) by the basic EGR.

\*  $P < 0.01$ , \*\*  $P < 0.005$ , \*\*\*  $P < 0.001$ , by a test for paired comparisons.

†  $\mu\text{mol}$  reduced NADP oxidized by EGR in 1 ml erythrocytes in 10 min at 35°.

‡ No stimulation occurred in one child in each of these groups.

coefficient method of Glatzle *et al.* (1970), was 43% for the children and 39% for the parents. That is, twenty-six of the sixty children and eleven of the twenty-eight parents had activation coefficients greater than 1.20 or, alternatively expressed, had a basic EGR activity that showed an *in vitro* stimulation greater than 20% when FAD was added before incubation. None of the G-6-PD-deficient persons was deficient in riboflavin as assessed by this method. After the riboflavin supplementation the activation coefficients of all persons were reduced, with the exception of that of the adult mentioned above, and all fell below 1.20.

The degree of stimulation of the basic EGR values produced by the riboflavin supplement was in direct proportion to the activation coefficient of the pre-supplement samples (Table 2). Generally speaking, it was found, for both children and parents before supplementation, that those whose activation coefficient was greater than 1.20 usually had the lowest *in vitro* basic EGR activities and that these subjects after receiving the riboflavin showed the greatest stimulation of the *in vitro* EGR activity.

#### *Dietary intake over the experimental period*

No control group was included in this study as it was considered that each group would act as its own control. However, it was found that dietary riboflavin intake had been higher in the early part of the previous year (i.e. 1970) (Thurnham, Migasena, Vudhivai *et al.* 1971) and that this increase in riboflavin intake corresponded to a fall in the prevalence of ariboflavinosis which occurred in March of that year in this village (Thurnham, Migasena & Vudhivai, 1971; Thurnham, Migasena, Vudhivai *et al.* 1971). In view of the possibility that dietary riboflavin intake might increase during the experimental period, dietary studies on preschool children were carried out in the first 5 d of riboflavin supplementation. The results showed that, of the twenty children

studied, only two received more than 0.3 mg/Mcal, the highest intake being 0.35 mg. The mean dietary riboflavin intake of the group was  $0.18 \pm 0.06$  mg/Mcal. Eleven of the children who took part in the dietary studies were in fact present in the sample group who completed the supplementation experiment. Ten of these children had normal G-6-PD values and the mean basic EGR activity of these ten children, measured before riboflavin supplementation, was compared with that of the other twelve children in the survey group whose diet was not checked during the period of supplementation. Mean values for both groups were identical ( $12.3 \pm 4.8$  and  $12.3 \pm 3.2$  respectively), indicating that the dietary habits of the two groups of children were probably very similar.

#### DISCUSSION

The village where this study was made contained forty-one families and about eighty children under 8 years of age. It was expected from earlier results of Migasena, Thurnham, Jintakanon & Pongpaew (1972), who worked in the same village on an earlier (1969) supplementation experiment with iron, that not more than 60% would take part in the experiment satisfactorily. It was decided therefore to change previous policies (Migasena, Thurnham, Pongpaew, Jintakanon & Harinasuta, 1971) and include parents as well as preschool children in order to obtain as many subjects as possible. The subjects were not divided into control and test groups for this would have reduced the number in the experimental group. Nor was it considered practical to include another village, where the riboflavin status and dietary intakes might be different and where the villagers might not co-operate when it came to taking the second sample of blood. The study on the dietary intakes of twenty children in the village was therefore the only controlling aspect in the study to indicate what the riboflavin nutrition was like during the experimental period and what the biochemical situation may have been without the supplement. The fact that the dietary intake of riboflavin was below marginal levels (Horwitt, Harvey, Hills & Liebert, 1950) and that the subjects were in an environment exposed to the normal stresses of life would suggest that the biochemical changes observed were due to the supplement.

It is interesting to note how low the dietary riboflavin intake of preschool children was during the supplementation period in 1971 in comparison with the values found in the previous year (Thurnham, Migasena, Vudhivai *et al.* 1971). In January and April 1970 it was found that the riboflavin intakes were significantly higher  $0.26 \pm 0.10$  (14) ( $P < 0.01$ ) and  $0.28 \pm 0.16$  (13) mg/Mcal ( $P < 0.02$ ) respectively than the intake in February 1971, which was  $0.18 \pm 0.06$  (20) mg/Mcal. One observation that might account for this difference was that, in 1970, many families had made extensive use of the recently installed irrigation system for household vegetable cultivation and vegetable intake was higher (Thurnham, Migasena, Vudhivai *et al.* 1971), whereas at the time of the supplementation study in 1971 the emphasis was more on the cultivation of tobacco and 'kitchen' gardens were not as numerous.

The dose of riboflavin taken by the subjects who completed the experiment was at least 70 mg, but in most instances 90 mg, and it was taken over 7-10 d, according to the information collected. Various supplements of riboflavin have been used to

stimulate EGR activity by different workers, varying from 5 mg for 8 d (Beutler, 1969*a*) to 10 mg for 15 d (Bamji & Sharada, 1971). Horwitt *et al.* (1950), in one of the more intensive studies of this subject, considered that a daily supplement of 6 mg over 15 d was required to saturate riboflavin-depleted subjects. The arrangement adopted in this experiment was a compromise in which the period of supervision was reduced to 10 d but the total amount taken was in most instances 90 mg, i.e. the amount recommended by Horwitt *et al.* (1950).

There are various modifications of the *in vitro* glutathione reductase test to determine riboflavin status, but they are all basically the same. The test determines two things – the basic EGR activity and the degree of stimulation of this activity that can be produced by adding FAD to haemolysate before incubation. Both measurements are related to the riboflavin status of the host, but the degree of stimulation produced by the *in vitro* addition of FAD is considered the more useful index of riboflavin status. Beutler (1969*a*) and Bamji (1969) both assessed riboflavin status in terms of 'FAD effect' (or percentage stimulation) whereas Glatzle (1970) introduced the term activation coefficient and proposed the value 1.20 to separate the normal from the riboflavin-deficient subjects. Results of studies carried out in Thailand using the latter method supported this suggestion of Glatzle (Thurnham *et al.* 1970).

Beutler (1969*b*) has pointed out that alterations in the order of addition of FAD to the reagents in the *in vitro* test influence greatly the stimulation produced by FAD. This has been confirmed in this laboratory using the reagent concentrations of Glatzle *et al.* (1970), and it was found that the stimulated EGR activity could be increased such that activation coefficients were less frequently below 1.00 (unpublished observations). It has also been noticed in this department that approximately 15% of the haemolysates obtained from rural children show activation coefficients below 1.00 (cf. Thurnham *et al.* 1970). That is, FAD causes some inhibition of *in vitro* EGR activity using the method of Glatzle *et al.* (1970). Approximately half of these samples with activation coefficients below 1.00 are from G-6-PD-deficient subjects (unpublished results). The effect of altering the order of addition of reagents to that recommended by Beutler (1969*b*) was to increase the activation coefficient of all samples. To have used this modification would have necessitated establishing a new threshold to separate the nutritionally sufficient from the deficient and therefore it was not adopted in these studies. The only change made was to increase the reduced NADP concentration. It was previously found that often with a haemolysate of high EGR activity the oxidation of reduced NADP was not linear for the full 10 min of the test. Increasing the reduced NADP concentration rectified this.

The riboflavin supplement was given to thirty-eight subjects and, with the exception of one adult male whose consumption was suspect, the results supported the generally held belief (Bamji, 1969; Beutler, 1969*b*; Glatzle *et al.* 1970) that the EGR test is a good indication of riboflavin status. In those haemolysates in which the *in vitro* stimulation of EGR activity produced by FAD was less than 20% (i.e.  $\alpha < 1.20$ ), the increase in basic EGR activity produced by the riboflavin supplement was significantly lower than in those subjects whose initial activation coefficients were greater than 1.20. That is, these results suggest that when FAD, added *in vitro*, produces a percentage

stimulation of the basic EGR activity that is greater than 20% ( $\alpha \geq 1.20$ ) the subject is probably riboflavin-deficient.

It is interesting to note that no ariboflavinosis could be demonstrated in the G-6-PD-deficient subjects, either male or female, by this method. Flatz (1970) described an experiment in northern Thailand on G-6-PD-deficient and normal Thai people in which he gave the same riboflavin supplement and obtained results similar to those reported here. He suggested that the FAD might be bound more tightly to the glutathione reductase enzyme in G-6-PD-deficient persons. His values cannot be compared direct with those reported here as the conditions of assay of EGR activity were different, but the basic EGR activity of his G-6-PD-deficient subjects appeared to be stimulated by the in vivo supplement to a smaller extent than was observed in the study reported here. This difference might reflect a difference in riboflavin status, and work is in progress to determine whether the response of the basic EGR activity of G-6-PD-deficient persons to in vivo supplementation with riboflavin can be used as a method of assessing the riboflavin status of G-6-PD-deficient persons (Thurnham, 1972)

These studies were partly supported by research grants T.69-01 and T.70-05 from SEAMEO-TROPED project. The authors are grateful to Professor B. G. Maegraith for criticizing the manuscript and to Mr P. Hongtong for technical assistance. The senior author (D. I. T.) was provided under Colombo Plan Technical Assistance to Thailand by the United Kingdom.

## REFERENCES

- Bamji, M. S. (1969). *Clinica chim. Acta* **26**, 263.  
 Bamji, M. S. & Sharada, D. (1971). *Clinica chim. Acta* **31**, 409.  
 Bernstein, R. E. (1962). *Nature, Lond.* **194**, 192.  
 Beutler, E. (1969a). *Science, N. Y.* **165**, 613.  
 Beutler, E. (1969b). *J. clin. Invest.* **48**, 1957.  
 Chandrapond, A. (1955). *J. med. Ass. Thailand* **38**, 111. (In Thai with English abstract.)  
 Chandrapond, A. & Ritchie, J. (1952). *J. med. Ass. Thailand* **35**, 11.  
 Flatz, G. (1970). *Nature, Lond.* **226**, 755.  
 Glatzle, D. (1970). In *Proceedings of the Seventh International Congress of Clinical Chemistry* Vol. 2 *Clinical Enzymology* p. 89 [J. Frei and M. Jemelin, editors]. Basel: Karger.  
 Glatzle, D., Körner, W. F., Christeller, S. & Wiss, O. (1970). *Int. Z. VitamForsch.* **40**, 166.  
 Hauck, H. M. & Sudsaneh, S. (1959). *J. Am. diet. Ass.* **35**, 1149.  
 Horwitt, M. K., Harvey, C. C., Hills, O. W. & Liebert, E. (1950). *J. Nutr.* **41**, 247.  
 Interdepartmental Committee on Nutrition for National Defense (1962). Nutrition Survey: The Kingdom of Thailand. Washington, DC: US Government Printing Office.  
 Migasena, P., Thurnham, D. I., Pongpaew, P., Jintakanon, K. & Harinasuta, C. (1971). In *1st Southeast Asian Regional Seminar on Nutrition*, 1969, Djakarta. Bangkok: SEAMEO Office. (In the Press.)  
 Migasena, P., Thurnham, D. I., Jintakanon, K. & Pongpaew, P. (1972). *Southeast Asian J. trop. Med. publ. Hlth.* (In the Press.)  
 Thurnham, D. I. (1972). *Ann. trop. Med. Parasit.* (In the Press.)  
 Thurnham, D. I., Migasena, P. & Pavapootanon, N. (1970). *Mikrochim. Acta* **5**, 988.  
 Thurnham, D. I., Migasena, P. & Vudhivai, N. (1971). *Southeast Asian J. trop. Med. publ. Hlth* **2**, 259.  
 Thurnham, D. I., Migasena, P., Vudhivai, N. & Supawan, V. (1971). *Southeast Asian J. trop. Med. publ. Hlth* **2**, 552.