

DISCRETE CONVEX FUNCTIONS AND PROOF OF THE SIX CIRCLE CONJECTURE OF FEJES TÓTH

IMRE BÁRÁNY, ZOLTÁN FÜREDI AND JÁNOS PACH

1. Introduction, results. A system \mathcal{C} of openly disjoint discs in the plane is said to form a *6-neighbourued circle packing* if every $C \in \mathcal{C}$ is tangent to at least 6 other elements of \mathcal{C} . (It is evident that such a system consists of infinitely many discs.) The simplest example is the regular circle packing all of whose circles are of the same size and have exactly 6 neighbours. L. Fejes Tóth conjectured that the regular circle packing has the interesting extremal property that, if we slightly “perturb” it, then there will necessarily occur either arbitrarily small or arbitrarily large circles. More precisely, he asked whether or not the following “zero or one law” (cf. [3], [6]) is valid: If \mathcal{C} is a 6-neighbourued circle packing, then

$$\frac{\inf r(C)}{\sup r(C)} = \begin{cases} 1 & \text{if } \mathcal{C} \text{ is regular} \\ 0 & \text{otherwise} \end{cases}$$

where $r(C)$ denotes the radius of circle C , \inf and \sup are taken over all $C \in \mathcal{C}$.

We shall prove this conjecture in the following stronger form:

THEOREM 1. *If \mathcal{C} is a 6-neighbourued circle packing which is not regular, then $\inf_{C \in \mathcal{C}} r(C) = 0$.*

Given a circle packing \mathcal{C} , we define a graph $G_{\mathcal{C}} = (V_{\mathcal{C}}, E_{\mathcal{C}})$ on the vertex set $V_{\mathcal{C}} := \mathcal{C}$ as follows: $C, C' \in \mathcal{C}$ are joined by an edge (i.e., $CC' \in E_{\mathcal{C}}$) if and only if C and C' are tangent.

Now let $G = (V, E)$ be an arbitrary graph. For any $x \in V$, let $\Gamma_G(x)$ denote the set of all neighbours of x . That is,

$$\Gamma_G(x) = \{y \in V \mid xy \in E\}.$$

A function $f: V \rightarrow \mathbf{R}$ is called *convex* (or *subharmonic*) on G if

$$\frac{1}{|\Gamma_G(x)|} \sum_{y \in \Gamma_G(x)} f(y) \geq f(x) \quad \text{for every } x \in V.$$

Any convex function defined on a connected finite graph is obviously constant. It is somewhat more surprising (but still easily seen) that every convex function on e.g. the rectangular lattice in the plane has either arbitrarily large positive values, or it is again constant (see [2]). This

Received June 7, 1983.

observation inspired our proof of Theorem 1, though we will not rely upon it in what follows.

Our next theorem states that if \mathcal{C} is a 6-neighbourd circle packing then the function $r^{-1}(C)$, ($C \in \mathcal{C}$) is convex on the graph $G_{\mathcal{C}}$ defined above.

THEOREM 2. *Let C be a disc of radius $r(C)$ which is tangent to at least 6 non-overlapping discs C_1, \dots, C_k , ($k \geq 6$). Then*

$$\frac{1}{k} \sum_{i=1}^k \frac{1}{r(C_i)} \geq \frac{1}{r(C)}$$

with equality if, and only if $k = 6, r(C) = r(C_1) = \dots = r(C_k)$.

A graph $G = (V, E)$ is called *locally finite* if every $x \in V$ has only finitely many neighbours in G , i.e., $|\Gamma_G(x)| < \infty$. We need the following general result on convex functions:

THEOREM 3. *Let $G = (V, E)$ be a locally finite graph, $f: V \rightarrow \mathbf{R}$ a convex function on G . Assume further that $x_0 y_0 \in E, f(x_0) = 0, f(y_0) = 1$; and let V_i denote the set of those points in G which can be reached from y_0 by a monotone increasing path of length at most i , i.e.,*

$$V_i = \left\{ y \in V \mid \begin{array}{l} \exists y_1, \dots, y_{j-1} \in V \text{ such that } j \leq i, \\ y_0 y_1, y_1 y_2, \dots, y_{j-1} y \in E \text{ and} \\ f(y_0) \leq f(y_1) \leq \dots \leq f(y) \end{array} \right\}.$$

Then, for every natural number n , we have

$$(1) \quad \max_{x \in V_n} f(x) \geq 1 + \sum_{i=1}^n \frac{1}{|E_i|},$$

where

$$E_i = \{ uv \in E \mid u \in V_{i-1}, v \in V_i \setminus V_{i-1}, f(u) \leq f(v) \}, \quad 1 \leq i \leq n.$$

We close this section by showing how Theorem 1 follows from the last two results.

Let \mathcal{C} be a 6-neighbourd circle packing, $G_{\mathcal{C}} = (V_{\mathcal{C}}, E_{\mathcal{C}})$ the corresponding graph defined above. If \mathcal{C} is the regular circle packing then there is nothing to prove. Hence, we may suppose without loss of generality that there are two tangent discs $C', C'' \in \mathcal{C} = V_{\mathcal{C}}$ such that $r(C') > r(C'')$. Our Theorem 2 implies that

$$f(C): = \left(\frac{1}{r(C)} - \frac{1}{r(C')} \right) / \left(\frac{1}{r(C'')} - \frac{1}{r(C')} \right), \quad C \in \mathcal{C}$$

is a convex function on $G_{\mathcal{C}}$ which meets the conditions of Theorem 3 with $x_0 = C', y_0 = C''$. Thus, for every natural n , we get (1).

Suppose now, in order to obtain a contradiction, that

$$\inf_{C \in \mathcal{C}} r(C) = \epsilon > 0.$$

Since all elements of V_i are covered by a circle of radius $r(C'')i$ around the centre of C'' , we have

$$|V_i| \leq \left(\frac{r(C'')}{\epsilon} \right)^2 i^2, \text{ for every } i.$$

On the other hand, $G_{\mathcal{C}}$ is a planar graph, thus any subgraph of $G_{\mathcal{C}}$ contains at most 3 times more edges than vertices. This yields

$$(2) \quad \sum_{j=1}^i |E_j| \leq 3 |V_i| \leq 3 \left(\frac{r(C'')}{\epsilon} \right)^2 i^2 \quad (\forall i).$$

The following easy technical lemma is a simple special case of Karamata's inequality (cf. [1], [5]).

LEMMA. Let e_1, \dots, e_n and A be positive numbers satisfying

$$\sum_{j \leq i} e_j \leq Ai^2 \text{ for every } i = 1, \dots, n.$$

Then

$$\sum_{i \leq n} \frac{1}{e_i} > (\log n)/(2A).$$

Hence (2) implies

$$(2') \quad \sum_{i=1}^n \frac{1}{|E_i|} \geq \frac{\epsilon^2}{6r^2(C'')} \log n.$$

Putting (1) and (2') together we obtain

$$\lim_{n \rightarrow \infty} \max_{x \in V_n} f(x) = \infty$$

which is tantamount to $\inf_{C \in \mathcal{C}} r(C) = 0$, a contradiction.

2. Proof of theorem 2. We may assume without loss of generality that C is *exactly surrounded* by the discs C_1, \dots, C_k : i.e., each C_i is tangent to C , C_{i-1} and C_{i+1} (where the subscripts are taken modulo k). Let O and O_i denote the centers of C and C_i ($i = 1, \dots, k$), respectively, and let $\angle O_i O O_{i+1} = \varphi_i$. We are going to show that

$$(3) \quad \frac{1}{r(C_i)} + \frac{1}{r(C_{i+1})} \geq \frac{1}{r(C)} \left(\tan \frac{\varphi_i}{4} + \cot \frac{\varphi_i}{4} - 2 \right), \quad i = 1, \dots, k.$$

Let us have a closer look e.g. at the discs C, C_1 and C_2 . Let P denote the centre of the circle D inscribed in $\Delta O O_1 O_2$, and put $\angle O O_1 P = \alpha$, $\angle O O_2 P = \beta$. Then we have




Figure 1

$$\begin{aligned} \frac{r(D)}{r(C_1)} + \frac{r(D)}{r(C_2)} &= \tan\alpha + \tan\beta \geq 2 \tan\frac{\alpha + \beta}{2} \\ &= 2 \tan\frac{\pi - \varphi_1}{4} = \tan\frac{\varphi_1}{2} \cdot \frac{(1 - \tan\frac{\varphi_1}{4})^2}{\tan\frac{\varphi_1}{4}} \\ &= \frac{r(D)}{r(C)} \left(\tan\frac{\varphi_1}{4} + \cot\frac{\varphi_1}{4} - 2 \right), \end{aligned}$$

as required.

Summing up (3) for $i = 1, \dots, k$ and using Jensen’s inequality, we obtain

$$\begin{aligned} 2 \sum_{i=1}^k \frac{1}{r(C_i)} &\geq \frac{1}{r(C)} \sum_{i=1}^k \left(\tan\frac{\varphi_i}{4} + \cot\frac{\varphi_i}{4} - 2 \right) \\ &\geq \frac{k}{r(C)} \left(\tan\frac{\pi}{2k} + \cot\frac{\pi}{2k} - 2 \right) \geq \frac{2k}{r(C)}, \end{aligned}$$

for $k \geq 6$. This completes the proof.

3. Proof of theorem 3. Throughout this section n will be kept fixed,

$$M := \max_{x \in V_n} f(x).$$

Define a subgraph $G' = (V', E') \subseteq G$ by

$$V' := V_n \cup \{x_0\}$$

$$E' := \{uv \in E \mid u, v \in V_n\} \cup \{x_0y_0\}.$$

We turn G' into a digraph $\vec{G}' = (V', \vec{E}')$ by supplying its edges with the following orientation: (a) If $uv \in E'$ and $f(u) < f(v)$, then let $\vec{uv} \in \vec{E}'$. (b) Every edge belonging to $E_i \subseteq E'$ will be directed from V_{i-1} to V_i , $i = 1, \dots, n$. (c) Any further edge $uv \in E'$ (with $f(u) = f(v)$) can be directed arbitrarily.

Note that if $uv \in E'$ for some $u \in V_{i-2}$ and $v \in V_i \setminus V_{i-1}$, then $f(u)$ is necessarily larger than $f(v)$, consequently $uv \notin E_i$, and $\vec{vu} \in \vec{E}'$.

Let \mathcal{F} denote the class of all functions $\varphi: V' \rightarrow \mathbf{R}$ satisfying the following three conditions:

- (i) $\varphi(x_0) = 0$.
- (ii) $f(x) \leq \varphi(x) \leq M$ for every $x \in V'$.
- (iii) There exists a digraph $\vec{G}'' = (V', \vec{E}'') \subseteq \vec{G}'$ such that $\vec{x_0y_0} \in \vec{E}'' \subseteq \vec{E}'$, $\varphi(u) \leq \varphi(v)$ for every $\vec{uv} \in \vec{E}''$, and

$$(4) \quad \frac{1}{|\Gamma_{G''}(x)|} \sum_{y \in \Gamma_{G''}(x)} \varphi(y) \geq \varphi(x) \text{ for every } x \in V_{n-1}.$$

Here G'' denotes the graph obtained from \vec{G}'' by ignoring the orientation of the edges, and $\Gamma_{G''}(x)$ is, as usual, the set of neighbours of x in G'' .

We clearly have $f \in \mathcal{F}$. To see this, it suffices to show that $\varphi = f$ satisfies (iii) with $\vec{G}'' := \vec{G}'$. But this follows immediately from the fact that, if $x \in V_{n-1}$, then $f(y) < f(x)$ for every $y \in \Gamma_G(x) \setminus \Gamma_{G'}(x)$.

Let f_0 be a maximal element of \mathcal{F} , and let $\vec{G}_0'' = (V', \vec{E}_0'')$ denote the corresponding digraph in (iii). (The existence of a maximal element is obvious, e.g. by compactness arguments.) Next we prove that f_0 satisfies (4) with equality at every (non-isolated) vertex $x \in V_{n-1}$. That is,

$$(5) \quad \frac{1}{|\Gamma_{G_0''}(x)|} \sum_{y \in \Gamma_{G_0''}(x)} f_0(y) = f_0(x) \quad (x \in V_{n-1}, \Gamma_{G_0''}(x) \neq \emptyset).$$

Assume indirectly that, for some $x \in V_{n-1}$, the left-hand side of (5) is strictly larger than the right-hand side. Set

$$\begin{aligned} \vec{F} &:= \{ \vec{xz} \in \vec{E}_0'' \mid f_0(z) = f_0(x) \}, \\ f_\epsilon(t) &:= \begin{cases} f_0(t) & \text{if } t \neq x \\ f_0(x) + \epsilon & \text{if } t = x. \end{cases} \end{aligned}$$

If $\epsilon > 0$ is sufficiently small, then f_ϵ and $\vec{G}_\epsilon'' := (V', \vec{E}_0'' \setminus \vec{F})$ meet all the conditions (i), (ii) and (iii), contradicting the maximum property of f_0 .

Now let $\vec{f}_0(\vec{xy}) := f_0(y) - f_0(x)$ for every $\vec{xy} \in \vec{E}_0''$, and observe that (5) is equivalent to

$$(5') \quad \sum_{\vec{xy} \in \vec{E}_0''} \vec{f}_0(\vec{xy}) = \sum_{\vec{vx} \in \vec{E}_0''} \vec{f}_0(\vec{vx}) \quad (\forall x \in V_{n-1}).$$

Hence, in another terminology we can say that \vec{G}_0'' is a network (with one source x_0 and some sinks in $V_n \setminus V_{n-1}$), and assigning the value $\vec{f}_0(\vec{uv}) \geq 0$ to each edge $\vec{uv} \in \vec{E}_0''$ we obtain a flow. (Note that there can not be any source in $V_n \setminus V_{n-1}$, because $f_0(x) = M$ for every $x \in V_n \setminus V_{n-1}$.)

Since the flow value disappearing at the sinks is equal to the value entering our network, we have

$$(6) \quad \sum_{y \in V_n \setminus V_{n-1}} \sum_{\vec{xy} \in \vec{E}_0''} \vec{f}_0(\vec{xy}) = \vec{f}_0(\vec{x_0y_0}) = f_0(y_0).$$

Further, it is obvious that every edge of positive flow value leaving V_{i-1} belongs to E_i , thus

$$(7) \quad \sum_{\substack{\vec{xy} \in \vec{E}_0'' \\ \vec{xy} \in E_i}} \vec{f}_0(\vec{x}) \geq \vec{f}_0(\vec{x_0y_0}) = f_0(y_0), \quad i = 1, \dots, n.$$

Hence it follows by (6) that

$$\begin{aligned} \sum_{xy \in E_0''} (f_0(y) - f_0(x))^2 &= \sum_{y \in V''} f_0(y) \sum_{x \in \Gamma_{G_0}(y)} (f_0(y) - f_0(x)) \\ &= \sum_{y \in V_n \setminus V_{n-1}} f_0(y) \sum_{x \in \Gamma_{G_0}(y)} (f_0(y) - f_0(x)) \\ &= \sum_{y \in V_n \setminus V_{n-1}} M \sum_{\substack{\bar{xy} \in \tilde{E}_0'' \\ \bar{xy} \in E_i}} \tilde{f}_0(\bar{xy}) = Mf_0(y_0). \end{aligned}$$

On the other hand, applying (7) and Jensen’s inequality, we obtain

$$\begin{aligned} \sum_{xy \in E_0''} (f_0(y) - f_0(x))^2 &\geq (f_0(y_0) - f_0(x_0))^2 + \sum_{i=1}^n \sum_{\substack{\bar{xy} \in \tilde{E}_0'' \\ \bar{xy} \in E_i}} \tilde{f}_0^2(\bar{xy}) \\ &\geq f_0^2(y_0) + \sum_{i=1}^n \frac{f_0^2(y_0)}{|E_0'' \cap E_i|} \\ &\geq f_0^2(y_0) \left(1 + \sum_{i=1}^n \frac{1}{|E_i|} \right). \end{aligned}$$

Now using the fact that $f_0(y_0) \geq f(y_0) = 1$, (1) immediately follows. This completes the proof of Theorem 3.

4. Remarks and acknowledgement. We would like to express our gratitude to L. Fejes Tóth and N. Dolbilin for their valuable comments and suggestions.

Special thanks are due to I. Vincze who has considerably simplified our first proof of Theorem 2. Our original argument, however, established also the following result:

THEOREM 4. *Let C, C_1, \dots, C_k ($k \geq 3$) be non-overlapping discs in the plane, and suppose that C is exactly surrounded by C_1, \dots, C_k (see Section 2). Further, let $|n| \geq 100$. Then*

$$\frac{1}{k} \sum_{i=1}^k r^n(C_i) \geq \left(\frac{\sin(\pi/k)}{1 - \sin(\pi/k)} r(C) \right)^n,$$

with equality if and only if $r(C_1) = \dots = r(C_k)$.

The proof can be omitted here, since L. Fejes Tóth [4] has recently managed to prove the same assertion for all $n \geq 1$.

Added in proof. We thank Adrian Bondy for calling our attention to a result of C. St. J. A. Nash-Williams: Random walk and electric current in networks (Proc. Cambridge Philos. Soc. 55 (1959), 181-199) which is closely related to our Theorem 3.

REFERENCES

1. E. F. Beckenbach and R. Bellman, *Inequalities* (Springer, Berlin-Göttingen-Heidelberg, 1961).
2. E. B. Dynkin and A. A. Juschkevitsch, *Sätze und Aufgaben über Markoffsche Prozesse* (Springer, Berlin-Heidelberg-New York, 1969).
3. L. Fejes Tóth, *Research problem*, Periodica Math. Hung. 8 (1977), 103-104.
4. ——— *Compact packing of circles*, Studia Sci. Math. Hungar. (to appear).
5. J. Karamata, *Sur une inégalité relative aux fonctions convexes*, Publ. Math. Univ. Belgrade 1 (1932), 145-148.
6. K. A. Rybnikov, *Problems of combinatorial analysis* (in Russian) (MIR Publ. Co., Moscow, 1980).

*Hungarian Academy of Sciences,
Budapest, Hungary*