

# High net accumulation rates at Campo de Hielo Patagónico Sur, South America, revealed by analysis of a 45.97 m long ice core

TAKAYUKI SHIRAIWA,<sup>1</sup> SHIRO KOHSHIMA,<sup>2</sup> RYU UEMURA,<sup>3</sup> NAOHIRO YOSHIDA,<sup>3</sup>  
SUMITO MATOBA,<sup>1</sup> JUN UETAKE,<sup>2</sup> MARIA ANGELICA GODOI<sup>4</sup>

<sup>1</sup>*Institute of Low Temperature Science, Hokkaido University, Sapporo 060-0819, Japan*

*E-mail: shiraiwa@pop.lowtem.hokudai.ac.jp*

<sup>2</sup>*Faculty of Bioscience and Biotechnology, Tokyo Institute of Technology, Okayama 2-12-1, Meguro-ku, Tokyo 152-8551, Japan*

<sup>3</sup>*Department of Environmental Science and Technology, Tokyo Institute of Technology, Yokohama 226-8502, Japan*

<sup>4</sup>*Instituto de la Patagonia, Universidad de Magallanes, Casilla 113-D, Punta Arenas, Chile*

**ABSTRACT.** A 45.97 m long ice core was recovered in the accumulation area of Glaciar Tyndall (50°59'05" S, 73°31'12" W; 1756 m a.s.l.), Campo de Hielo Patagónico Sur (southern Patagonia icefield), during December 1999. The firn core was subjected to visual stratigraphic observation and bulk density measurements in the field, and later to analyses of water isotopes ( $\delta^{18}\text{O}$ ,  $\delta\text{D}$ ), major dissolved ions and snow algal biomass. The drillhole remained dry down to about 43 m depth, where a water-soaked layer appeared. Seasonal cycles were found for  $\delta^{18}\text{O}$ ,  $\delta\text{D}$  and the D-excess, although the amplitudes of the cycles decreased with depth. Major dissolved ions ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Cl}^-$ ,  $\text{SO}_4^{2-}$ ) and algal biomass exhibit rapid decreases in the upper 3 m, probably due to meltwater elution. Annual increments defined by the  $\delta^{18}\text{O}$  and D-excess peaks suggest that the minimum net accumulation rates at this location were  $17.8 \text{ m a}^{-1}$  in 1997/98–1998/99 and  $> 11.0 \text{ m a}^{-1}$  in 1998/99–1999/2000. These are much higher values than those previously obtained from past ice-core studies in Patagonia, but are of the same order of magnitude as those predicted from various observations in ablation areas of Patagonian glaciers.

## INTRODUCTION

Since the pioneering work of Meier (1984), the contribution of mountain glaciers to sea-level rise has been a key issue for understanding the effects of global warming. Of particular importance is the mass balance of the huge temperate glaciers in Alaska and Patagonia (Houghton and others, 1996) because the sensitivity of equilibrium-line altitude to temperature change is more pronounced at high precipitation rates (Oerlemans and Fortuin, 1992). The glaciers in Patagonia are maritime; furthermore, Campo de Hielo Patagónico Norte (HPN; northern Patagonia icefield) at  $4200 \text{ km}^2$  and Campo de Hielo Patagónico Sur (HPS; southern Patagonia icefield) at  $13\,000 \text{ km}^2$  together constitute the world's largest ice mass outside the polar regions (Warren and Sugden, 1993; Aniya and others, 1996). Accordingly, changes in their mass balances are closely related to world sea level.

Mass-balance estimates have been made using ice cores recovered from the icefields. Three ice-core drillings have been performed so far; these yielded estimated net accumulation rates of  $3.45 \text{ m a}^{-1}$  at the Glaciar San Rafael drilling site (HPN,  $46^\circ 44' \text{ S}$ ,  $73^\circ 32' \text{ W}$ ; 1296 m a.s.l.) (Yamada, 1987),  $1.2 \text{ m a}^{-1}$  at Glaciar Perito Moreno (HPS,  $50^\circ 38' \text{ S}$ ,  $73^\circ 15' \text{ W}$ ; 2680 m a.s.l.) (Aristarain and Delmas, 1993) and  $2.2 \text{ m a}^{-1}$  at Glaciar Nef (HPN,  $45^\circ 56' \text{ S}$ ,  $73^\circ 19' \text{ W}$ ; 1500 m a.s.l.) (Matsuoka and Naruse, 1999). Aristarain and Delmas (1993) suggested that the  $1.2 \text{ m a}^{-1}$  value might have been influenced by wind scouring and surface melting. Surface melting could be more critical for ice cores at lower altitudes. It is therefore necessary to obtain ice cores from a site where there is minimal

influence on mass balance from meltwater discharge and wind scouring.

This paper presents a new estimate of net accumulation rates at HPS, based on analyses of  $\delta^{18}\text{O}$  and  $\delta\text{D}$  as well as major ions and algal biomass along a 45.97 m long ice core. The high values calculated are considered to be typical of this glacier in the light of synoptic-scale climate analyses of the region. Glaciar Tyndall can therefore be classed as one of the most extreme maritime glaciers in the world.

## GEOGRAPHICAL SETTING

The drilling site is located at 500 m east of a divide between Glaciar Tyndall to the east and Glaciar Amaria to the west in the southernmost part of HPS ( $50^\circ 59' 05'' \text{ S}$ ,  $73^\circ 31' 12'' \text{ W}$ ; 1756 m a.s.l.). A map of the region is shown in Figure 1. Glaciar Tyndall flows southwards for 32 km, covering a total area of about  $331 \text{ km}^2$ , and eventually calves into a proglacial lake (Aniya and others, 1996).

The glacier surface at the drilling site is inclined by  $1^\circ$  to the east. There are nunataks approximately 2 km south of the drilling site (Fig. 2), but the underlying surface is otherwise flat for at least 2 km in the other directions. Preliminary radio-echo soundings suggest that the glacier might be  $> 600 \text{ m}$  thick (personal communication from A. Rivera and G. Casassa, 1999), although its exact thickness at the drilling site is unknown.

According to our meteorological observations during the 24 day drilling campaign, weather conditions in this region are extreme: we recorded an average hourly wind




Fig. 1. Location of the drilling site. Previous drilling sites and the location of Punta Arenas are also shown.

speed of  $13.6 \text{ m s}^{-1}$  and a snow accumulation of approximately 3.5 m (about 1.75 m w.e.) from 30 November to 23 December 1999. Visibility exceeded 50 m on only 4 days out of 24. Air temperature varied from  $3.3^\circ$  to  $-8.2^\circ\text{C}$ , with an average of  $-3.5^\circ\text{C}$  during the campaign.

### ICE-CORE DRILLING AND ANALYSIS

Because of the harsh environment, we drilled the ice core from a snow cave located below the glacier surface. The tents (see Fig. 2) were finally abandoned due to difficulties in maintaining them in the heavy snowfall. Details of the campaign are given in Godoi and others (in press). Because the air temperature inside the snow cave was nearly  $0^\circ\text{C}$ , the recovered ice cores had to be processed immediately. First, the core stratigraphy was observed with the aid of a fluorescent light; here special attention was paid to a distinction between firn and melt layers. Then the bulk density of the core was determined. For analysis, the core was sliced into 25 cm sections, each of which was vertically cut into three pieces (one half and two quarters). The half-section was packed in a sealed plastic bag and left untouched. External parts of the two other pieces were removed using a pre-cleaned ceramic knife, and the samples were packed in separate, sealed plastic bags. Unfortunately, weather conditions enforced an emergency evacuation with limited samples; hence, only one set of samples (i.e. one-quarter of the core) could be saved for laboratory analyses. These samples were kept frozen until their arrival at the Institute of Low Temperature Science in Japan, where they were analyzed.

Measurement of the borehole temperature after drilling was planned, but was prevented by meltwater inflow into the borehole from the top while the last sections were being drilled. The borehole was dry down to 42.55 m depth, where the firn became soaked with meltwater. The glacier is thus



Fig. 2. Landscape near the drilling site on Glacier Tyndall.

classified as temperate. Further details of the drilling procedure are given in Kohshima and others (2002).

The 25 cm long samples were melted, heated to  $+3^\circ\text{C}$ , and immediately analyzed for major ions, water isotopes and algal biomass. Major ions were analyzed with an ion chromatography system (Dionex model 500x with AS-12A and CS12-A separate columns and  $500 \mu\text{L}$  sample loops) at the clean laboratory of the Institute of Low Temperature Science. Blank values were measured and the detection limit for the major ions was found to be 10 ppb or less. Stable oxygen and hydrogen isotopes were measured by a mass spectrometer (Finnigan model MATdelta 252) at the Department of Environmental Science and Technology, Tokyo Institute of Technology. The measurement errors for  $\delta^{18}\text{O}$  and  $\delta\text{D}$  are  $\pm 0.05\text{‰}$  and  $\pm 1\text{‰}$ , respectively. From the errors measured in this analysis, we can determine D-excess to an accuracy of  $\pm 1.4\text{‰}$ .

Algal biomass in each sample was observed and measured at the Faculty of Bioscience and Biotechnology, Tokyo Institute of Technology, and they are expressed as total algal volume ( $\mu\text{m}^3 \text{ mL}^{-1}$ ). Several algal species were found in the ice-core samples, but *Chloromonas sp.* of unicellular green algae was the most abundant. The volume was determined by quantifying the number of cells and their volume with the aid of a microscope. Further details of the method are given in Yoshimura and others (2000).

### PHYSICAL PROPERTIES OF THE ICE CORE

The core consisted of layers of firn and melt features that varied in thickness between 1 and 50 mm down to 42.55 m depth, below which water-soaked firn–ice layers continued to 45.97 m depth. Neither dirt layers nor ash layers were found throughout the profile. Snow grains were mostly coarse granules.

The bulk density  $\rho$  gradually increases with depth  $d$  and attains the firn/ice transition ( $\sim 830 \text{ kg m}^{-3}$ ) at 42.5 m depth (Fig. 3). The best fit by a quadratic equation is  $\rho_d = -0.147d^2 + 14.66d + 459.91$  ( $r^2 = 0.94$ ). The depth of pore close-off, 42.5 m, is greater than the depths of 20–30 m recorded in other temperate glaciers (Kawashima and Yamada, 1997). This may be due to the abnormally high accumulation rate mentioned below.

Figure 3 also compares the depth–density curve of Glacier




Fig. 3. Profile of the ice-core bulk density (solid dots). Ordinate is the measured core depth  $d$  (m). For comparison, profiles of bulk densities from other glaciers on the Patagonia icefields are given: Glacières San Rafael (Yamada, 1987), Perito Moreno (Aristarain and Delmas, 1993) and Nef (Matsuoka and Naruse, 1999). Arrows indicate depths at which water-saturated firn first appeared.

Tyndall with those from previous measurements on other Patagonian glaciers. The curve of Glacières Tyndall (1756 m) is similar to that from Glacières Perito Moreno (2680 m at HPS; Aristarain and Delmas, 1993), despite the 924-m difference in elevation. Altitudes of the drilling sites at Glacières San Rafael (1296 m at HPN; Yamada, 1987) and Nef (1500 m at HPN; Matsuoka and Naruse, 1999) are much closer to that of Glacières Tyndall (1756 m). At a given depth, however, the firn densities of Glacières San Rafael and Nef are much higher than those of Glacières Tyndall and Perito Moreno.

Another important point is the occurrence of water-saturated firn-ice layers at 42.5 m on Glacières Tyndall, 19.7 m on Glacières San Rafael and 13.3 m on Glacières Nef. This suggests that below these depths any chemical and isotopic signals in the ice cores may have been eliminated by the meltwater, as already found at the two glaciers in HPN.

### WATER-ISOTOPE, ION-CHEMISTRY AND BIOMASS ANALYSES

#### $\delta^{18}\text{O}$ , $\delta\text{D}$ and D-excess

Average values and the standard deviations of all 183 samples of  $\delta^{18}\text{O}$ ,  $\delta\text{D}$  and D-excess ( $d = \delta\text{D} - 8\delta^{18}\text{O}$ ; Dansgaard, 1964) measured from the surface to the bottom of the ice core are  $-11.9 \pm 1.5\text{‰}$ ,  $-84 \pm 12\text{‰}$  and  $10 \pm 2\text{‰}$ , respectively. In order to check the dependence of water isotopes on altitude, the averaged value of the  $\delta^{18}\text{O}$  is compared with those obtained for Glacières San Rafael (Yamada, 1987), Perito Moreno (Aristarain and Delmas, 1993) and Nef (Matsuoka and Naruse, 1999) and for Punta Arenas (53.00°S, 70.51°W; 37 m a.s.l.) (Fig. 4). Average values were obtained for Glacières San Rafael and Nef by averaging all of the  $\delta^{18}\text{O}$  values appearing in each report. The averaged  $\delta^{18}\text{O}$  of Glacières Perito Moreno was calculated and converted from the reported  $\delta\text{D}$




Fig. 4. Relationship between averaged values of  $\delta^{18}\text{O}$  and altitudes in the two Patagonia icefields and the nearest meteorological station. Original data for Glacières San Rafael are from Yamada (1987); for Glacières Perito Moreno from Aristarain and Delmas (1993); and for Glacières Nef from Matsuoka and Naruse (1999). Data for Punta Arenas were obtained from the GNIP database (<http://www.iaea.org/programs/ri/gnip/gnipmain.htm>). Solid dots and bars are the average value and the standard deviation of each dataset used for analyses. See text for further explanation.

values by assuming a meteoric water relationship. The averaged  $\delta^{18}\text{O}$  for Punta Arenas was obtained from the International Atomic Energy Agency/World Meteorological Organization global network of isotopes in precipitation (GNIP) database, Release 3 October 1999 (<http://www.iaea.org/programs/ri/gnip/gnipmain.htm>) for the period 1996–97.

The four average values of  $\delta^{18}\text{O}$  are linearly related to altitude  $z$  (m a.s.l.); the best-fit equation is  $\delta^{18}\text{O} = -0.002z - 8.49$ . The  $\delta^{18}\text{O}$ /altitude effect is  $-0.2\text{‰}$  for each 100 m, which is the same as for the Swiss Alps (Schotterer and others, 1997). Although the data used are variable in both quality and quantity and may lack statistical significance, it seems that the water isotopes over the Patagonian icefields are controlled mainly by altitude.

Profiles of  $\delta^{18}\text{O}$  and D-excess in the ice core are given in Figure 5. Hereafter, all depths are given by their water equivalent. The amplitudes of the two signals are relatively large in the upper 10 m, ranging from  $-8\text{‰}$  to  $-17\text{‰}$  in the case of  $\delta^{18}\text{O}$ , but smaller at greater depths, ranging from  $-9\text{‰}$  to  $-14\text{‰}$ . It is, however, possible to find oscillations whose strongest  $\delta^{18}\text{O}$  peaks are at 11.0 and 28.8 m depths if the upper 10 m is smoothed by an 11-data running mean (thick line in Fig. 5). D-excess has peaks at 8.6 and 23.3 m and shows fluctuations similar to those of  $\delta^{18}\text{O}$ . The difference between the two profiles is that they are slightly shifted in their peaks: D-excess peaks postdate those of  $\delta^{18}\text{O}$ .

To explain why the  $\delta^{18}\text{O}$  peaks predate D-excess peaks in the ice core, we examined seasonal fluctuations in the two signals observed at the nearest meteorological station, Punta Arenas (Fig. 1), where water-isotope data are currently available at the GNIP database mentioned previously. These are plotted in Figure 6 by calculating monthly averages from the dataset during 1990–97. Although the measurement




Fig. 5. Profiles of  $\delta^{18}\text{O}$  (‰), D-excess (‰), major ions (ppb or  $\mu\text{g L}^{-1}$ ) and algal biomass ( $\mu\text{m}^3\text{mL}^{-1}$ ). Depth is given by the water equivalent value. Thick lines in  $\delta^{18}\text{O}$  and D-excess are 11-data running mean.




Fig. 6. Seasonal changes in  $\delta^{18}\text{O}$  (‰) and D-excess parameter at Punta Arenas ( $53.00^\circ\text{S}$ ,  $70.51^\circ\text{W}$ ; 37 m a.s.l.). The original data were obtained from the GNIP database, and the monthly average data from 1990 to 1997 were used for this analysis.

period is not long enough to determine definite statistical relationships, the monthly averages of  $\delta^{18}\text{O}$  and D-excess show slightly shifted seasonal patterns too: the strongest  $\delta^{18}\text{O}$  and D-excess peaks appear in February and April, respectively. This pattern is very similar to what we found in the ice core, where  $\delta^{18}\text{O}$  peaks appeared approximately several meters lower, i.e. earlier, than D-excess peaks. D-excess's time lag behind  $\delta^{18}\text{O}$  remains unexplained, but it suggests that the fluctuations of both signals found in the ice core would correspond to the annual cycles in water isotopes.

**Major ions and biomass**

Major dissolved ions ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Cl}^-$ ,  $\text{SO}_4^{2-}$ ) exhibit rapid decreases in the upper 4 m (Fig. 5).  $\text{NO}_3^-$  was below the detection limit throughout the profile. Several additional peaks are, however, detectable above the upper level of the water-soaked layer at 28.94 m w.e. These peaks are located at 5–7, 20–21 and 24–25 m. The peaks for the dissolved ions correspond to the peaks for the snow algal biomass.

Concentrations of each ion in this ice core are one order of magnitude higher than those reported from Glaciar Perito Moreno (Aristarain and Delmas, 1993) as far as the samples from the surface 4 m are concerned. This is not due to contamination during processing of the samples and the analy-




Fig. 7. Profiles of  $\delta^{18}\text{O}$  (‰), D-excess (‰),  $\text{Cl}^-/\text{Na}^+$  ( $\mu\text{eq L}^{-1}/\mu\text{eq L}^{-1}$ ) and melt-layer percentage. Thick lines in each profile are 11-data running mean. The two horizontal bands are possible summer (February) horizons; the two arrows in the D-excess profile are possible April horizons.

ses: if this were the case, a high concentration of ions would also be found in the lower part, where samples are processed and analyzed in the same way as samples from the upper 4 m.

To check whether the major ions are eluted by meltwater, a ratio of  $\text{Cl}^-$  ( $\mu\text{eq L}^{-1}$ ) and  $\text{Na}^+$  ( $\mu\text{eq L}^{-1}$ ) was compared with a typical sea-salt ratio of 1.16 (Wilson, 1975, cited in Keene and others, 1986) (Fig. 7). Original compositions of  $\text{Cl}^-$  and  $\text{Na}^+$  in precipitation supplied to HPS are known to be close to the marine reference value (Aristarain and Delmas, 1993), and the original ratios can be disturbed by post-depositional elution if ice-core melting is significant (Davies and others, 1982). The  $\text{Cl}^-/\text{Na}^+$  ratio is close to the typical sea-salt ratios of 1.16 at depths of 5–10 m and 12–15 m where the  $\delta^{18}\text{O}$  is relatively low (Fig. 7). At other depths, the ratios are significantly lower than the typical sea-salt ratios. Because the  $\delta^{18}\text{O}$  is supposed to be a seasonal signal having maxima in summer and minima in winter in this ice core, the original sea-water ratio is considered to be maintained in winter depths of the ice core. Therefore, the lower values found for the  $\text{Cl}^-/\text{Na}^+$  ratios are probably due to the influence of elution after the deposition of snow.

The profile of algal biomass also shows a rapid decrease in the surface 3 m (Fig. 5), and several minor peaks appear at depths where  $\delta^{18}\text{O}$  is relatively low. Yoshimura and others (2000) proposed that high peaks of the algal biomass are markers of summer depths in the Himalayan ice core, but this seems not to be the case for this ice core.

### ESTIMATE OF ANNUAL ACCUMULATION RATES

Our multi-signal analyses on the ice core suggest that fluctuations of  $\delta^{18}\text{O}$  and D-excess along the ice core are similar to those obtained at the nearest meteorological station, Punta Arenas; thus we may use these signals in the ice core as seasonal markers. However, the fluctuations of the signals for the major ions and algal biomass are difficult to interpret since they are considered to have been substantially disturbed by meltwater after deposition. Meltwater can also destroy the isotopic signals, as occurred on Glaciares San Rafael and Nef in HPN; nevertheless, the original isotopic profiles can be maintained for at least a few years when the amount of meltwater is not significant (e.g. Krouse and others, 1977). Therefore, it may be possible to determine net accumulation rates at the drilling site from the  $\delta^{18}\text{O}$  and D-excess in the depth profiles.

The highest peaks in the  $\delta^{18}\text{O}$  profile are located at depths of 11.0 and 28.8 m (the two horizontal bands in Fig. 7). The profile suggests that another peak is forming at the surface. Because the  $\delta^{18}\text{O}$  peak appears in February at Punta Arenas, we tentatively assume that  $\delta^{18}\text{O}$  in the ice core also peaks in February. D-excess peaks are located at depths of 8.6 and 23.3 m, which could represent April snowfall (arrows in Fig. 7). This seems reasonable since the topmost level of D-excess is still relatively low; thus another peak probably appears much later than our drilling period in December.

Occurrences of melt layers in the ice core support the above-mentioned speculations (Fig. 7). The melt layers are usually formed in summer, when meltwater at the surface percolates to a certain depth of the glacier.\* In this ice core, however, one can recognize two horizons where melt layers are almost absent: at 4–7 and 20–25 m. These two depths are also characterized by lower  $\delta^{18}\text{O}$  values, so they would correspond to winter horizons.

As a result, we calculate that the net accumulation rate from the austral summer (February) of 1997/98–1998/99 was  $17.8 \text{ m a}^{-1}$  w.e., whereas that of 1998/99–1999/2000 is expected to be  $11.0 \text{ m a}^{-1}$  plus the amount that was supplied until February 2000, as indicated by the two horizontal bands in Figure 7. These values do not include the meltwater discharged from the site, so the calculated values are lower bounds of the actual net balances. However, the amount of meltwater from firn layers is likely insignificant at this altitude since the isotopic profiles would have washed out if the amount were high.

Mass balances of the Patagonian icefields are poorly understood to date. Besides the ice-core data mentioned above, specific mass balances were estimated, for a part of HPS only, by glacial dynamical methods (Skvarca and Naruse, 1997; Rott and others, 1998). Annual precipitation over both the HPS and HPN accumulation areas has been estimated either by the hydrological method (Escobar and others, 1992) or by meteorological observations (Fujiyoshi and others, 1987). All of the studies estimated a high annual mass input of 5–10 m w.e. on average in the accumulation area. Although our estimates are almost double the above-mentioned estimates, we believe that they are not far from the realistic value if we remember the actual accumulation rate we observed during our drilling operation (3.5 m in snow height per 24 days in December 1999).

Finally, we ask whether the calculated high net accumulation rates are normal or abnormal at this location. K. Kubota (unpublished information) analyzed the synoptic-scale weather conditions over Patagonia by using data on geopotential height, wind speed and moisture flux at 850 hPa from 1985 to 1999. He compared the December monthly averages of these parameters from 1999 with those from other years. December 1999 was found to be nearly the average of the 15 year period, so this was not an unusual year. Hence, the extremely high snow accumulation observed for December 1999 is likely typical for this glacier. This means that Glaciar Tyndall must rank as one of the most maritime glaciers in the world.

## CONCLUSIONS

From our ice coring at Glaciar Tyndall on HPS we found that

- (1) field campaigns here are very strenuous due to high winds and heavy snowfall;
- (2) the glacier is temperate even at its highest saddle at 1750 m a.s.l.;
- (3) the seasonal cycles in  $\delta^{18}\text{O}$ ,  $\delta\text{D}$  and D-excess were more or less preserved for at least 2 years;

(4) major ions and algal biomass were substantially eluted by meltwater;

(5) the seasonal cycles of  $\delta^{18}\text{O}$  and D-excess together with the melt-layer distribution helped to reconstruct the lower bounds of net accumulation rates for 1997/98–1998/99 ( $17.8 \text{ m a}^{-1}$ ) and 1998/99–1999/2000 ( $>11.0 \text{ m a}^{-1}$ ).

## ACKNOWLEDGEMENTS

The authors would like to thank K. Kubota, M. Arevalo, J. Quinteros, N. Takeuchi, R. Naruse and M. Aniya for valuable support during the field campaign. The evacuation operation headed by G. Cassasa and A. Rivera is highly appreciated. The workshop staff at the Institute of Low Temperature Science provided technical support for developing the drilling system. Comments from B. Stauffer, D. Wagenbach, an anonymous reviewer and K. Matsuoka helped the authors to revise the manuscript. This research was funded by grants-in-aid for Basic Research A (No. 10041105; principal investigator (PI): M. Aniya) and Basic Research C (No. 11640422; PI: R. Naruse), both from Ministry of Education, Science, Sports and Culture of Japan, and a grant from Fondo Nacional de Ciencias y Tecnología (FONDECYT 1980293) of Chile.

## REFERENCES

- Aniya, M., H. Sato, R. Naruse, P. Skvarca and G. Casassa. 1996. The use of satellite and airborne imagery to inventory outlet glaciers of the Southern Patagonia Icefield, South America. *Photogramm. Eng. Remote Sensing*, **62**(12), 1361–1369.
- Aristarain, A. J. and R. J. Delmas. 1993. Firn-core study from the southern Patagonia ice cap, South America. *J. Glaciol.*, **39**(132), 249–254.
- Dansgaard, W. 1964. Stable isotopes in precipitation. *Tellus*, **16**(4), 436–468.
- Davies, T. D., C. E. Vincent and P. Brimblecombe. 1982. Preferential elution of strong acids from a Norwegian ice cap. *Nature*, **300**(5888), 161–163.
- Escobar, F., F. Vidal, C. Garin and R. Naruse. 1992. Water balance in the Patagonia icefield. In Naruse, R. and M. Aniya, eds. *Glaciological researches in Patagonia, 1990*. Nagoya, Japanese Society of Snow and Ice. Data Center for Glacier Research, 109–119.
- Fujiyoshi, Y., H. Kondo, J. Inoue and T. Yamada. 1987. Characteristics of precipitation and vertical structure of air temperature in the northern Patagonia. *Bull. Glacier Res.* **4**, 15–23.
- Godoi, M., T. Shiraiwa, S. Kohshima and K. Kubota. In press. Firn-core drilling operation at Tyndall Glacier, Southern Patagonia Icefield. In Sepulveda, F., G. Casassa and R. Sinclair, eds. *The Patagonian icefields: a unique natural laboratory for environmental and climate change studies*. New York, Kluwer/Plenum.
- Houghton, J. T., L. G. M. Filho, B. A. Callander, N. Harris, A. Kattenberg and K. Maskell. 1996. *Climate change 1995: the science of climate change*. Cambridge, etc., Cambridge University Press.
- Kawashima, K. and T. Yamada. 1997. Experimental studies on the transformation from firn to ice in the wet-snow zone of temperate glaciers. *Ann. Glaciol.*, **24**, 181–185.
- Keene, W. C., A. A. P. Pszenny, J. N. Galloway and M. E. Hawley. 1986. Sea-salt corrections and interpretation of constituent ratios in marine precipitation. *J. Geophys. Res.*, **91**(D6), 6647–6658.
- Kohshima, S., T. Shiraiwa, M. Godoi, K. Kubota, N. Takeuchi and K. Shinbori. 2002. Ice core drilling at Southern Patagonia Icefield — development of a new portable drill and the field campaign in 1999. *Natl. Inst. Polar Res. Mem., Special Issue* **56**, 49–58.
- Krouse, H. R., K. West, R. Hislop, H. M. Brown and J. L. Smith. 1977. Climatic and spatial dependence of the retention of D/H and  $\text{O}^{18}/\text{O}^{16}$  abundances in snow and ice of North America. *International Association of Hydrological Sciences Publication* 118 (Symposium at Grenoble 1975 — *Isotopes and Impurities in Snow and Ice*), 242–247.
- Matsuoka, K. and R. Naruse. 1999. Mass balance features derived from a firn core at Hielo Patagónico Norte, South America. *Arct. Antarct. Alp. Res.*, **31**(4), 333–340.
- Meier, M. F. 1984. Contribution of small glaciers to global sea level. *Science*, **226**(4681), 1418–1421.
- Oerlemans, J. and J. P. F. Fortuin. 1992. Sensitivity of glaciers and small ice caps to greenhouse warming. *Science*, **258**(5079), 115–117.

\* Due to the high value of the water percolation at this site, melt layers could also indicate winter (and not summer) layers, meltwater refreezing not just below the surface, but deeper, in the preceding winter layer.

- Rott, H., M. Stuefer, A. Siegel, P. Skvarca and A. Eckstaller. 1998. Mass fluxes and dynamics of Moreno Glacier, Southern Patagonia Icefield. *Geophys. Res. Lett.*, **25**(9), 1407–1410.
- Schotterer, U., K. Fröhlich, H.W. Gäggeler, S. Sandjordj and W. Stichler. 1997. Isotope records from Mongolian and Alpine ice cores as climate indicators. *Climatic Change*, **36**(3–4), 519–530.
- Skvarca, P. and R. Naruse. 1997. Dynamic behavior of Glaciar Perito Moreno, southern Patagonia. *Ann. Glaciol.*, **24**, 268–271.
- Warren, C. R. and D. E. Sugden. 1993. The Patagonian icefields: a glaciological review. *Arct. Alp. Res.*, **25**(4), 316–331.
- Wilson, T. R. S. 1975. Salinity and the major elements of sea water. In Riley, J. P. and G. Skittow, eds. *Chemical oceanography Vol. 1*. London, Academic Press, 365–413.
- Yamada, T. 1987. Glaciological characteristics revealed by 376-m deep core drilled at the accumulation area of San Rafael Glacier, the Northern Patagonia Icefield. *Bull. Glacier Res.* **4**, 59–67.
- Yoshimura, Y., S. Kohshima, N. Takeuchi, K. Seko and K. Fujita. 2000. Himalayan ice-core dating with snow algae. *J. Glaciol.*, **46**(153), 335–340.