

Variable dust formation by the double-line spectroscopic binary WR 70 (HD 137603, WC9vd+B0I)

Peredur M. Williams

*Institute for Astronomy, University of Edinburgh,
Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, Scotland, UK*

Karel A. van der Hucht

*SRON Nationaal Instituut voor Ruimteonderzoek,
Sorbonnelaan 2, NL-3584 CA Utrecht, Nederland*

Fred Marang

*South African Astronomical Observatory,
P.O. Box 9, Observatory, 7935, South Africa*

Abstract. Infrared photometry of the WC9vd+B0I spectroscopic binary WR 70 over two decades shows variable dust emission. The variations are relatively slow and show evidence for an underlying process with a period $P = 1045 \pm 60$ d, together with modulation on shorter and longer time-scales.

Variability of WR 70 in the near-IR was reported by Williams, van der Hucht & Thé (1987), based on comparison of their 1983-86 observations from the SAAO and ESO, with earlier observations by Cohen (1975) and Pitault *et al.* (1983). We have continued IR-monitoring WR 70, to search for systematic variations. The synoptic light curves (Figure 1) show amplitude increasing with wavelength, consistent with the spectral energy distribution (SED) which showed dust emission, deviating from the stellar wind SED only longward of $1.5 \mu\text{m}$ (Williams *et al.* 1987). The *J*-photometry suggests, that the stellar wind flux from WR 70 is constant to within a few per cent, and that variations at longer wavelengths must, therefore, be due to variable dust emission; which is present even at minimum. The light curves delineate clear features, including a sharp maximum in 1989.25 and a well defined fading and recovery in 1992-93.

The WC9 and B0I components of WR 70 display RV variations, showing it to be an SB2 (Golombek 1983; Niemela 1995) and, therefore, a colliding-wind binary. This prompted a search for periods in the variation of the dust emission, which might be modulated by orbit-related effects. A CLEAN power spectrum (using PERIOD, Dhillon & Privett 1997) of 128 *K*-magnitudes gave a strong, broad maximum at 0.356 y^{-1} (1026 d). On the basis that the (*K-L*) colour variations, attributable to dust formation starting and stopping, may be a better indicator of underlying processes occurring in a colliding-wind system, 126 (*K-L*) colours were also searched, giving maximum power at 0.348 y^{-1} (1050 d). The string-length method, useful when the form of the light curve is




Figure 1. *JHK* light curves of WR 70 from observations at ESO (\times ; mostly 1983-93) and SAAO (\bullet ; mostly 1989-2001), combined by applying shifts determined from comparison of contemporaneous ($|\Delta T| \leq 6$ d) observations.

not known, gave a broad minimum at 0.36 y^{-1} (1045 d) and clear minima at 2P and 3P. These tests suggest that there is an underlying process in WR 70, modulating dust formation with a period of about 1045 ± 50 d, but light curves phased to this period show significant scatter, suggesting that other processes on both shorter and longer time-scales are at work, such as might be caused by high-density structures in one of the stellar winds, and a third stellar component in a longer period orbit, respectively.

References

- Cohen, M. 1975, *MNRAS* 173, 489
- Dhillon, V.S., Privett, G.J. 1997, *PERIOD* Time-Series Analysis Package, Starlink User Note 167.5, Rutherford Appleton Laboratory
- Golombek, D.A. 1983, *Estudio Espectrografico de Una Estrella Supergigante con Líneas de Emisión*, PhD thesis, Universidad de Buenos Aires, Argentina
- Niemela, V.S. 1995, in: K.A. van der Hucht & P.M. Williams (eds), *Wolf-Rayet Stars: Binaries, Colliding Winds, Evolution*, Proc. IAU Symposium No.163 (Dordrecht: Kluwer), p. 223
- Pitault, A., Epchtein, N., Gómez, A.E., Lortet, M.C. 1983 *A&A* 120, 53
- Williams, P.M., van der Hucht, K.A., Thé, P.S. 1987, *A&A* 182, 91