

- patients (CGI-BPD): a scale sensible to detect changes. *Actas Esp Psiquiatr* 2007; **35**: 229–35.
- 51** Zanarini MC, Vujanovic AA, Parachini EA, Boulanger JL, Frankenbreg FR, Hennen J. Zanarini Rating Scale for Borderline Personality Disorder (ZAN-BPD): a continuous measure of DSM-IV borderline psychopathology. *J Pers Disord* 2003; **17**: 233–42.
- 52** Zanarini MC, Frankenbreg FR, Hennen J, Reich DB, Silk KR. Axis I comorbidity in patients with borderline personality disorder: 6-year follow-up and prediction of time to remission. *Am J Psychiatry* 2004; **161**: 2108–14.
- 53** National Collaborating Centre for Mental Health. *Borderline Personality Disorder: Treatment and Management. Full Guideline*. Clinical Guideline 78. NICE, 2009 (<http://www.nice.org.uk/nicemedia/pdf/CG78fullGuideline.pdf>)
- 54** Oldham JM. *Guideline Watch: Practice Guideline for the Treatment of Patients with Borderline Personality Disorder*. American Psychiatric Association, 2005.
- 55** Herpertz SC, Zanarini M, Schulz CS, Siever L, Lieb K, Moller HJ. World Federation of Societies of Biological Psychiatry (WFSBP) guidelines for biological treatment of personality disorders. *World J Biol Psychiatry* 2007; **8**: 212–44.



psychiatry  
in pictures

## The Kiss

by Ismond Rosen (1924–1996)

The sculptor Ismond Rosen was a distinguished psychoanalyst and consultant psychiatrist who became a fellow to both the Royal College of Psychiatrists and the Society of Portrait Sculptors. He presented this 1947 white marble carving to the Royal Society of Medicine in 1987, where it can now be seen framed by a window on the south façade. The stillness of its simple curves symbolises the importance of early attachments as a template for all subsequent relationships. In Dr Rosen's own words, 'The two heads relate above and fuse below in a manner expressive of human loving, both in its earliest mother-child relationship and in adult sexual congress.'

Text by Dr Alexandra Pitman. Image courtesy of the Royal Society of Medicine.

The British Journal of Psychiatry (2010)  
196, 12. doi: 10.1192/bjp.196.1.12

