

THE LOGIC OF HYPERLOGIC. PART B: EXTENSIONS AND RESTRICTIONS

ALEXANDER W. KOCUREK 

Sage School of Philosophy, Cornell University

Abstract. This is the second part of a two-part series on the logic of hyperlogic, a formal system for regimenting metalogical claims in the object language (even within embedded environments). Part A provided a minimal logic for hyperlogic that is sound and complete over the class of all models. In this part, we extend these completeness results to stronger logics that are sound and complete over restricted classes of models. We also investigate the logic of hyperlogic when the language is enriched with hyperintensional operators such as counterfactual conditionals and belief operators.

§B1. Introduction. This is the second part of a two-part series on the logic of *hyperlogic*, a hyperintensional semantics designed to regiment metalogical claims (e.g., “Intuitionistic logic is correct” or “The law of excluded middle holds”) in the object language. To recap, this regimentation is achieved using:

- a multigrade entailment operator \triangleright ;
- propositional quantifiers $\forall p$ and $\exists p$;
- interpretation terms i that double as atomic formulas (“ i is correct”);
- hybrid operators $@_i$ (“according to i ”) and $\downarrow i$ (“where i is the current interpretation”).

The semantics of hyperlogic introduces the notion of a “hyperconvention,” i.e., a complete interpretation of the propositional variables, Boolean connectives, and \triangleright over some space of possible worlds propositions. Interpretation terms denote “conventions,” modeled as sets of hyperconventions. Propositional quantifiers range over (special kinds of) index propositions, i.e., sets of world-hyperconvention pairs. Models in this semantics determine (i) a set of worlds W ; (ii) a domain of admissible conventions D_C ; (iii) a domain of admissible index propositions D_P ; and (iv) a valuation V . Truth-in-a-model is evaluated relative to worlds and hyperconventions. Operators like $@_i$ can shift the hyperconvention parameter. This allows formulas to be assessed on alternative interpretations of the connectives and entailment. Hyperintensionality is thus captured through shifting these interpretations.

In Part A [34], a complete axiomatization for this semantics was given. The axiomatization in Part A captures consequence over the class of *all* models. Almost no constraints are placed on either a model’s convention or proposition domain. The

Received: April 25, 2022.

2020 *Mathematics Subject Classification*: Primary 03A05, 03B16, 03B42, 03B45, 03B60.

Key words and phrases: hyperlogic, hyperintensionality, hybrid logic, propositional quantifiers, counterfactuals, counterpossibles, counterlogicals, belief, logical omniscience.

© The Author(s), 2022. Published by Cambridge University Press on behalf of The Association for Symbolic Logic. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

resulting logic for hyperlogic is, therefore, fairly minimal. For example, no constraints are placed on the interpretations the entailment operator \triangleright can receive. Yet, intuitively, it would be a stretch to say \triangleright really represents a notion of “entailment” if, say, it wasn’t factive (i.e., if $\triangleright\phi$ did not imply ϕ), or if it wasn’t reflexive or transitive. It would then be natural to inquire into how imposing such constraints affects the underlying logic of hyperlogic.

Furthermore, hyperlogic was initially motivated by concerns with the interaction between metalogical claims and hyperintensional operators such as attitude verbs, counterfactuals, and so on. Yet the language of hyperlogic introduced in Part A does not contain any of such operators.

In Part B of this series, we take initial steps to filling these gaps. We start by studying stronger logics for hyperlogic that can be obtained by adding additional rules and axioms in Section B2. These stronger logics can be shown to be sound and complete over classes of models whose convention and proposition domains satisfy certain natural constraints. In Section B3, we examine how the completeness results from Part A carry over to languages with hyperintensional operators. We conclude in Section B4 with some questions left open by this two-part investigation into the logic of hyperlogic. Section B5 is a technical appendix containing proofs of completeness for various classes of hypermodels.

Note: as this is a continuation of a two-part series, I will freely refer back to the definitions, notation, and results from Part A [34], rather than repeat them. Labels for sections, definitions, theorems, and tables are prefixed with the part that they refer to (e.g., “Section A3” refers to Section 3 of Part A).

§B2. Restrictions on hypermodels. Let us start by exploring constraints we may impose on the class of hypermodels and how that affects the logic of hyperlogic. In Section B2.1, we look at general constraints on the convention domain and present axiomatizations in the quantifier-free fragment over those hypermodels. In Section B2.2, we extend some of these results to languages with propositional quantifiers. Finally, in Section B2.3, we examine constraints on the proposition domain.

B2.1. Quantifier-free fragment. Table B1 contains a sample of constraints one may want to impose on the convention domain. For the analyticity constraint, we write $c \approx c'$ to mean c and c' are exactly alike except possibly in how they interpret propositional variables (i.e., $c(\Delta) = c'(\Delta)$ for all Δ). The intersection of each class in Table B1 is denoted by concatenation (e.g., the class of analytic and full hypermodels is AnF). Where X is a class of hypermodels, we define classical and universal entailment over X , written $\Gamma \models_X \phi$ and $\Gamma \Vdash_X \phi$ respectively, as in Definition A2.16 except restricting to hypermodels in X .

Table B2 contains axiomatizations of consequence over various classes. Some of the axioms make use of the following abbreviations:

$$\widehat{@}_i \phi := \sim @_i \sim \phi, \quad (\phi)^t = (\psi)^\kappa := (@_i \phi = @_\kappa \psi).$$

Here are their truth conditions (where $\llbracket \phi \rrbracket^C = \bigcap_{c \in C} \llbracket \phi \rrbracket^c$):

$$\begin{aligned} w, c \Vdash \widehat{@}_i \phi & \Leftrightarrow \text{for some } c' \in V(i) : w, c \Vdash \phi, \\ w, c \Vdash (\phi)^t = (\psi)^\kappa & \Leftrightarrow \llbracket \phi \rrbracket^{V(i)} = \llbracket \psi \rrbracket^{V(\kappa)}. \end{aligned}$$

Table B1. *Some constraints on convention domains.*

Name	Class	Constraint (on all $C \in D_C$)
full	F	$\pi_c = \emptyset$ for each $c \in C$
atomic	At	$\{w\} \in \pi_c$ for each $c \in C$ and $w \in W$
boolean	B	π_c is closed under complement and finite intersection for each $c \in C$
quantification uniform	U_q	$\pi_c = \pi_{c'}$ for each $c, c' \in C$
operation uniform	U_o	$c(\Delta) = c'(\Delta)$ for each $c, c' \in C$ and each Δ
singular	Si	$ C = 1$
analytic	An	for any $c, c' \in D_{\mathbb{H}}$, if $c \in C$ and $c \approx c'$, then $c' \in C$
S5-universal	S_5	each $c \in C$ is classical
classically complete	Co_{cl}	$V(cl) = \{c \in \mathbb{H}_W \mid c \text{ is classical}\}$

Table B2. *Axiomatizations in \mathcal{L}^H for various classes from Table B1. Axiomatizations in \mathcal{L}^{HE} (except those appealing to RAN, which becomes infinitary when add \triangleright) are obtained by replacing \mathbf{H} with $\mathbf{H}_{\triangleright}$ and generalizing the corresponding axioms accordingly.*

Name	Axiom/Rule
Uni _o	$\kappa \in \iota, \lambda \in \iota, \vec{\phi}^\kappa = \vec{\psi}^\lambda \Vdash (\Delta(\vec{\phi}))^\kappa = (\Delta(\vec{\psi}))^\lambda$
Sing	$\Vdash I _1$
Self-Dual _@	$@_i \phi \Vdash \widehat{@}_i \phi$
Bool	$\star \phi \Vdash \star \phi$ $(\phi \circ \psi) \Vdash (\phi \bullet \psi)$
RAN	if $\vec{\alpha}, \kappa _1, \lambda _1, (\vec{p})^\kappa = (\vec{q})^\lambda \Vdash (\Delta(\vec{p}))^\kappa = (\Delta(\vec{q}))^\lambda$ for each Δ where none of \vec{p}, \vec{q} are in $\vec{\alpha}$, then $\vec{\alpha}, \kappa _1, \lambda _1 \Vdash (\kappa \in \iota) \equiv (\lambda \in \iota)$
Class	Axiomatization
F, U_q , At, B	\mathbf{H}
U_o	$\mathbf{H} + \text{Uni}_o$
Si	$\mathbf{H} + \text{Sing} = \mathbf{H} + \text{Self-Dual}_@$
AnF, An U_q	$\mathbf{H} + \text{RAN}$
S_5	$\mathbf{H} + \text{Bool}_ $

In addition, we write $\vec{\phi}$ for ϕ_1, \dots, ϕ_n , and $\vec{\phi}^\kappa = \vec{\psi}^\lambda$ for $\bigwedge_{i=1}^n (\phi_i^\kappa = \psi_i^\lambda)$. Where \mathbf{L} is a logic and \mathbf{A} is an axiom, $\mathbf{L} + \mathbf{A}$ is the result of extending \mathbf{L} with \mathbf{A} (i.e., the rules still apply to formulas derived using \mathbf{A}). If \mathbf{R} is a rule, $\mathbf{L} + \mathbf{R}$ is the result of closing \mathbf{L} under \mathbf{R} along with the other rules. Given this, we have the following:

THEOREM B2.1 (Relative completeness in \mathcal{L}^H and \mathcal{L}^{HE}). *The axiomatic systems in Table B2 are sound and complete for (consequence over) the relevant class of hypermodels.¹ (See Section B5.1 for the proof.)*

¹ It is an open question whether consequence in \mathcal{L}^H over An or Co_{cl} can be axiomatized. An axiomatization for An in \mathcal{L}^{QH} is given in Section B2.2. (Interestingly, the key axiom invokes $\forall\exists$ -quantification, which cannot be directly expressed in \mathcal{L}^H .) By contrast, consequence for Co_{cl} in \mathcal{L}^{QH} is provably unaxiomatizable (Corollary B2.5).

Table B3. *Some constraints on the interpretation of \triangleright .*

Name	Constraint (on all $c, c' \in D_{\mathbb{H}}$)
reflexive	$(X \triangleright_c X) = W$
transitive	$(\vec{X} \triangleright_c \vec{Y}) \cap (\vec{Y} \triangleright_c \vec{Z}) \subseteq (\vec{X} \triangleright_c \vec{Z})$
monotonic	$(\vec{X}, \vec{Y} \triangleright_c \vec{Z}) \subseteq (\vec{X}, U, \vec{Y} \triangleright_c \vec{Z})$
contractive	$(\vec{X}, U, U, \vec{Y} \triangleright_c \vec{Z}) \subseteq (\vec{X}, U, \vec{Y} \triangleright_c \vec{Z})$
commutative	$(\vec{X}, U_1, U_2, \vec{Y} \triangleright_c \vec{Z}) = (\vec{X}, U_2, U_1, \vec{Y} \triangleright_c \vec{Z})$
congruential	$((\vec{X} \triangleright_c \vec{Y}) \cap (\vec{Y} \triangleright_c \vec{X}) \cap (\vec{X} \triangleright_c \vec{Z})) \subseteq (\vec{Y} \triangleright_c \vec{Z})$
self-aware	$(\vec{X} \triangleright_c (\vec{Y} \triangleright_c \vec{Z})) = (\vec{Y} \triangleright_c \vec{Z})$
fully aware	$(\vec{X} \triangleright_c (\vec{Y} \triangleright_{c'} \vec{Z})) = (\vec{Y} \triangleright_{c'} \vec{Z})$
import-export	$(\vec{X} \triangleright_c (\vec{Y} \triangleright_c \vec{Z})) = (\vec{X}, \vec{Y} \triangleright_c \vec{Z})$
\supset -residuation	$(\vec{X}, Y \triangleright_c \vec{Z}) = (\vec{X} \triangleright_c (\vec{Y} \cup \vec{Z}))$
\rightarrow -residuation	$(\vec{X}, Y \triangleright_c \vec{Z}) = (\vec{X} \triangleright_c (Y \rightarrow_c \vec{Z}))$
&-fusion	$(\vec{X}, U_1, U_2, \vec{Y} \triangleright_c \vec{Z}) = (\vec{X}, U_1 \cap U_2, \vec{Y} \triangleright_c \vec{Z})$
\wedge -fusion	$(\vec{X}, U_1, U_2, \vec{Y} \triangleright_c \vec{Z}) = (\vec{X}, U_1 \wedge_c U_2, \vec{Y} \triangleright_c \vec{Z})$
factive	$((\vec{X} \triangleright_c Y) \cap X_1 \cap \dots \cap X_n) \subseteq Y$
noncontingent	either $(\vec{X} \triangleright_c Y) = W$ or $(\vec{X} \triangleright_c Y) = \emptyset$
strict	$(\vec{X} \triangleright_c Y) = \{w \in W \mid X_1 \cap \dots \cap X_n \subseteq Y\}$

In addition, we can consider imposing restrictions specifically on the interpretation of \triangleright . Usual suspects include reflexivity, transitivity, monotonicity, etc. But there are also “unusual” suspects to consider (e.g., factivity) since \triangleright is an object language operator. Table B3 contains examples of such constraints, with their corresponding axioms stated in Table B4. Following our earlier convention, we write \vec{X} for X_1, \dots, X_n . (If $n = 0$, then $\vec{X} = \langle \rangle$.) We also write $(\vec{X} \triangleright_c \vec{Y})$ for $\bigcap_i (\vec{X} \triangleright_c Y_i)$.

THEOREM B2.2 (Relative completeness for \triangleright). *The axiomatic systems resulting from adding the relevant axioms in Table B4 to $\mathbf{H}_{\triangleright}$ are sound and complete for the relevant class of hypermodels.*

Proof (Sketch). We revise the definition of the proposition space for canonical hyperconventions (Definition A3.30) so that $\pi_{c_\kappa} = \{X \mid [X]_\kappa \neq \emptyset\}$.² The completeness proof in Section §A3.2 remains in tact. We just need to verify that if we impose an axiom, the canonical model satisfies the corresponding constraint. The proof is more-or-less the same for each case. We illustrate with the transitivity case. Suppose $\Delta \in (\vec{X} \triangleright_{c_\kappa} \vec{Y}) \cap (\vec{Y} \triangleright_{c_\kappa} \vec{Z})$. Since $\Delta \in (\vec{X} \triangleright_{c_\kappa} \vec{Y})$, there are some $\vec{\phi} \in [\vec{X}]_\kappa$ and $\vec{\psi} \in [\vec{Y}]_\kappa$ such that $@_{\kappa}(\vec{\phi} \triangleright \psi_i) \in \Delta$ for each i (note: we can let $\vec{\phi}$ be the same for each ψ_i by Lemma A3.29 and Rep $_{\triangleright}$). Since $\Delta \in (\vec{Y} \triangleright_{c_\kappa} \vec{Z})$, there is a $\chi \in [Z]_\kappa$ such that $@_{\kappa}(\vec{\psi} \triangleright \chi) \in \Delta$. By Tr, $@_{\kappa}(\vec{\phi} \triangleright \chi) \in \Delta$. Hence, $\Delta \in (\vec{X} \triangleright_{c_\kappa} \vec{Z})$. \square

B2.2. Adding quantifiers. Adding propositional quantifiers to the language allows us the ability to distinguish between classes of models that previously generated the

² Observe that this revised definition of π_{c_κ} is not guaranteed to be full or atomic, so this proof does not automatically carry over when these constraints are also imposed.

Table B4. Axiomatizations in \mathcal{L}^{HE} for various classes from Table B3.

Name	Axiom	Corresponding Constraint
Id	$\Vdash(\phi \triangleright \phi)$	reflexive
Tr	$(\vec{\phi} \triangleright \vec{\psi}), (\vec{\psi} \triangleright \chi) \Vdash (\vec{\phi} \triangleright \chi)$	transitive
Weak	$(\vec{\alpha}, \vec{\beta} \triangleright \chi) \Vdash (\vec{\alpha}, \phi, \vec{\beta} \triangleright \chi)$	monotonic
Contr	$(\vec{\alpha}, \phi, \phi, \vec{\beta} \triangleright \chi) \Vdash (\vec{\alpha}, \phi, \vec{\beta} \triangleright \chi)$	contractive
Perm	$(\vec{\alpha}, \phi, \psi, \vec{\beta} \triangleright \chi) \Vdash (\vec{\alpha}, \psi, \phi, \vec{\beta} \triangleright \chi)$	commutative
Cong	$(\vec{\phi} \triangleright \vec{\psi}), (\vec{\psi} \triangleright \vec{\phi}), (\vec{\phi} \triangleright \chi) \Vdash (\vec{\psi} \triangleright \chi)$	congruential
Self-Aware	$(\vec{\phi} \triangleright (\vec{\psi} \triangleright \chi)) \Vdash (\vec{\psi} \triangleright \chi)$	self-aware
Aware	$(\vec{\phi} \triangleright (\vec{\psi} \triangleright_i \chi)) \Vdash (\vec{\psi} \triangleright_i \chi)$	fully aware
IE	$(\vec{\phi} \triangleright (\vec{\psi} \triangleright \chi)) \Vdash (\vec{\phi}, \vec{\psi} \triangleright \chi)$	import-export
Res \supset	$(\vec{\phi} \triangleright (\psi \supset \chi)) \Vdash (\vec{\phi}, \psi \triangleright \chi)$	\supset -residuation
Res \rightarrow	$(\vec{\phi} \triangleright (\psi \rightarrow \chi)) \Vdash (\vec{\phi}, \psi \triangleright \chi)$	\rightarrow -residuation
Fus \cap	$(\vec{\alpha}, \phi, \psi, \vec{\beta} \triangleright \chi) \Vdash (\vec{\alpha}, \phi \& \psi, \vec{\beta} \triangleright \chi)$	$\&$ -fusion
Fus \wedge	$(\vec{\alpha}, \phi, \psi, \vec{\beta} \triangleright \chi) \Vdash (\vec{\alpha}, \phi \wedge \psi, \vec{\beta} \triangleright \chi)$	\wedge -fusion
T \triangleright	$(\vec{\phi} \triangleright \psi), \vec{\phi} \Vdash \psi$	factive
Rigid \triangleright	$(\vec{\phi} \triangleright \psi) \Vdash \blacksquare (\vec{\phi} \triangleright \psi)$	noncontingent
Strict \triangleright	$(\vec{\phi} \triangleright \psi) \Vdash \blacksquare (\vec{\phi} \supset \psi)$	strict

same logic. Notably, the consequence relations over F, U_q, At, and B are now all distinguishable. In addition, we can now present an axiomatization for An, which was absent from Section B2.1 (see footnote 1).

Axiomatizations for some of those classes are given in Table B5. Where Σ is a set of axioms of the form $\Vdash\sigma$, we let $\mathbf{L} \cup \Sigma$ be the proof system defined as follows: $\Gamma \Vdash_{\mathbf{L} \cup \Sigma} \phi$ iff $\Gamma \cup \{\sigma \mid (\Vdash\sigma) \in \Sigma\} \Vdash_{\mathbf{L}} \phi$ (in other words, Σ are treated as premises, not axioms; this means, among other things, that one cannot necessarily derive the universal generalization of members of Σ). The axiomatizations in Table B5 make use of the following abbreviations:

$$\begin{aligned} \pi_\kappa \subseteq \pi_\lambda &:= \forall p \exists q (p^\kappa = q^\lambda), & \pi_\kappa = \pi_\lambda &:= (\pi_\kappa \subseteq \pi_\lambda) \& (\pi_\lambda \subseteq \pi_\kappa), \\ |\pi_\kappa|_1 &:= \forall p \forall q (p =_\kappa q), & \kappa \approx \lambda &:= \& \{ \Delta_\kappa = \Delta_\lambda \} \Delta \& (\pi_\kappa = \pi_\lambda). \end{aligned}$$

These have the obvious truth conditions assuming $|V(\kappa)| = |V(\lambda)| = 1$ (which is the only relevant case for the axiomatizations below).

THEOREM B2.3 (Relative completeness in \mathcal{L}^{QH}). *The proof systems in Table B5 are sound and complete for the relevant class of hypermodels. (See Section B5.2.)*

Notice that no axiomatization for F is stated. This is because consequence over F is unaxiomatizable.

THEOREM B2.4 (Unaxiomatizability of full consequence in \mathcal{L}^{QH}). \models_F in \mathcal{L}^{QH} is unaxiomatizable. Moreover, where X is the intersection of any of the classes mentioned in Tables B1 and B6, if $F_X \neq \emptyset$, then \models_{F_X} in \mathcal{L}^{QH} is unaxiomatizable.

Table B5. Axiomatizations in \mathcal{L}^{QH} for various classes from Table B1.

Name	Axiom/Rule
Atom	$\Vdash \exists p(p \& \forall q(\blacksquare(p \supset @_i q) + \blacksquare(p \supset \sim @_i q)))$
BoolEx	$\Vdash E \sim p$ $\Vdash E(p \& q)$
Ex	$\Vdash E \phi$
An	$\sim i _1, \kappa \in i, \lambda _1, \kappa \approx \lambda \Vdash \lambda \in i$
Many _I Nom	$\Vdash (i _1 \& i _1 \& i \approx i) \supset (i = i)$ where $i \in \text{INom}$
Uni _q	$\kappa \in i, \lambda \in i \Vdash \pi_\kappa = \pi_\lambda$
Uni _o \forall	$\kappa \in i, \lambda \in i \Vdash \kappa \approx \lambda$
Class	Axiomatization
At	QH + Atom
B	QH + BoolEx
U _q	QH + Uni _q
U _o	QH + Uni _o \forall
An	(QH + An) \cup Many _I Nom
Si	QH + Sing = QH + Self-Dual _@
S ₅	QH + Bool _{\Vdash} + Ex

Table B6. Some constraints on proposition domains.

Name	Class	Constraint (on all $P \in D_{\mathbb{P}}$)
complete	Cp	$D_{\mathbb{P}} = \mathbb{P}_{D_{\mathbb{H}}}$
correlated	Cr	$P(c) = P(c')$ whenever $c, c' \in C$
closed under Φ	Cl _{Φ}	$\llbracket \phi \rrbracket^{\mathcal{M}} \in D_{\mathbb{P}}$ for all $\phi \in \Phi$
strongly closed under Φ	Cl _{Φ} ⁺	$\llbracket \phi \rrbracket^{\mathcal{M}'} \in D_{\mathbb{P}}$ for all $\phi \in \Phi$ and all $\mathcal{M}' \approx \mathcal{M}$
definable in Φ	Df _{Φ}	if $P \in D_{\mathbb{P}}$, then there is a $\phi \in \Phi$ such that $\llbracket \phi \rrbracket^{\mathcal{M}} = P$
discerning	Di	for all $c, c' \in D_{\mathbb{H}}$, if $c \neq c'$, then for some $P \in D_{\mathbb{P}}$, $P(c) \neq P(c')$
combinatorial	Cb	if $X_1 \in \pi_{c_1}, \dots, X_n \in \pi_{c_n}$ for some distinct $c_1, \dots, c_n \in D_{\mathbb{H}}$, then for some $P \in D_{\mathbb{P}}$ such that $P(c_i) = X_i$ for $i \leq n$

Proof. Let $\text{At}(p) := \blacklozenge p \& \forall q(\blacksquare(p \supset q) + \blacksquare(p \supset \sim q))$. It is easy to verify that if c is full, then $\mathcal{M}, w, c \Vdash \text{At}(p)$ iff $|V(p)(c)| = 1$. Let Δ consist of the following formulas:

$$\begin{aligned}
 \forall p @_k (\neg p = \sim p), & & \forall p @_k (\lozenge p = \neg \square \neg p), \\
 \forall p \forall q @_k ((p \wedge q) = (p \& q)), & & \forall p \forall q @_k \blacksquare(\square(p \rightarrow q) \rightarrow (\square p \rightarrow \square q)), \\
 \forall p \forall q @_k ((p \vee q) = (p + q)), & & \forall p @_k (\blacksquare p \supset \blacksquare \square p), \\
 \forall p \forall q @_k ((p \rightarrow q) = (p \supset q)), & & \forall p @_k \blacksquare(\forall q(\text{At}(q) \supset \square(q \supset p)) \supset \square p).
 \end{aligned}$$

Since the propositionally quantified modal logic $\mathbf{K}\pi+$ is unaxiomatizable [19], it suffices to show that for any $\phi \in \mathcal{L}^Q$ (the language of propositionally quantified modal

logic), $\models_{\mathbf{K}\pi+} \phi$ iff $\Delta, |k|_1 \models_{\mathbf{F}} @_k \phi$. We do this by constructing, for each $\mathbf{K}\pi+$ -model, an equivalent full hypermodel of Δ and vice versa.

First, let $\mathcal{K} = \langle W, R, V \rangle$ be a $\mathbf{K}\pi+$ -model. Let c_k be defined as follows:

$$\begin{aligned} \pi_{c_k} &= \emptyset W, & \neg_{c_k} X &= \overline{X}, \\ c_k(p) &= V(p), & X \wedge_{c_k} Y &= X \cap Y, \\ \Box_{c_k} X &= \{w \in W \mid R[w] \subseteq X\}, & X \vee_{c_k} Y &= X \cup Y, \\ \Diamond_{c_k} X &= \{w \in W \mid R[w] \cap X \neq \emptyset\}, & X \rightarrow_{c_k} Y &= \overline{X} \cup Y. \end{aligned}$$

Define $\mathcal{M}^{\mathcal{K}} = \langle W, D_{\mathbb{C}}, D_{\mathbb{P}}, V^{\mathcal{K}} \rangle$ so that (i) $c_k \in D_{\mathbb{H}}$, (ii) each $c \in D_{\mathbb{H}}$ is full, (iii) $V^{\mathcal{K}}(p) = P_p$, and (iv) $V^{\mathcal{K}}(k) = \{c_k\}$. Clearly, $\mathcal{M}^{\mathcal{K}}, w, c_k \Vdash \Delta \cup \{|k|_1\}$. Moreover, by induction, for all $\phi \in \mathcal{L}^{\mathcal{Q}}$ and all Q_1, \dots, Q_n where $Q_i(c_k) = X_i$, we have $\mathcal{K}_{X_1, \dots, X_n}^{q_1, \dots, q_n}, w \Vdash \phi$ iff $(\mathcal{M}^{\mathcal{K}})_{Q_1, \dots, Q_n}^{q_1, \dots, q_n}, w, c_k \Vdash \phi$. Hence, $\mathcal{K}, w \Vdash \phi$ iff $\mathcal{M}^{\mathcal{K}}, w, c_k \Vdash @_k \phi$.

Next, let $\mathcal{M} = \langle W, D_{\mathbb{C}}, D_{\mathbb{P}}, V \rangle$ be a full hypermodel satisfying $\Delta \cup \{|k|_1\}$. Let c_k be such that $V(k) = \{c_k\}$. Define $\mathcal{K}^{\mathcal{M}} = \langle W, R, V^{\mathcal{M}} \rangle$ so that (i) wRv iff for all $X \subseteq W$, if $w \in \Box_{c_k} X$, then $v \in X$, and (ii) $V^{\mathcal{M}}(p) = c_k(p)$. We establish by induction that for all $\phi \in \mathcal{L}^{\mathcal{Q}}$ and all Q_1, \dots, Q_n where $Q_i(c_k) = X_i$, we have $\mathcal{M}_{Q_1, \dots, Q_n}^{q_1, \dots, q_n}, w, c_k \Vdash \phi$ iff $(\mathcal{K}^{\mathcal{M}})_{X_1, \dots, X_n}^{q_1, \dots, q_n}, w \Vdash \phi$. The only interesting case is the \Box -clause. Observe that $R[w] = \{v \in W \mid w \in \Diamond_{c_k} \{v\}\}$.³ For notational ease, let $\mathcal{M}^* = \mathcal{M}_{Q_1, \dots, Q_n}^{q_1, \dots, q_n}$ and $\mathcal{K}^* = (\mathcal{K}^{\mathcal{M}})_{X_1, \dots, X_n}^{q_1, \dots, q_n}$.

(\Rightarrow) Suppose $\mathcal{M}^*, w, c_k \Vdash \Box \phi$. Thus, $w \in \Box_{c_k} \llbracket \phi \rrbracket^{\mathcal{M}^*, c_k}$. Let $v \in R[w]$. Then for all $X \subseteq W$, if $w \in \Box_{c_k} X$, then $v \in X$. Hence, $v \in \llbracket \phi \rrbracket^{\mathcal{M}^*, c_k}$, which by IH means $v \in \llbracket \phi \rrbracket^{\mathcal{K}^*}$. Hence, $\mathcal{K}^*, w \Vdash \Box \phi$.

(\Leftarrow) Suppose $\mathcal{M}^*, w, c_k \not\Vdash \Box \phi$. Thus, $w \notin \Box_{c_k} \llbracket \phi \rrbracket^{\mathcal{M}^*, c_k}$. Since c_k is full, by Definition A2.11 (constraint (ii) on $D_{\mathbb{P}}$), there exists a P such that $P(c_k) = \llbracket \phi \rrbracket^{\mathcal{M}^*, c_k}$. By the definition of Δ , $(\mathcal{M}^*)_P^p, w, c_k \Vdash \exists q(\text{At}(q) \wedge \Diamond(q \wedge \neg p))$. Let Q be such that $(\mathcal{M}^*)_{P, Q}^{p, q}, w, c_k \Vdash \text{At}(q) \wedge \Diamond(q \wedge \neg p)$. Thus, $Q(c_k) = \{v\}$ for some $v \notin P(c_k) = \llbracket \phi \rrbracket^{\mathcal{M}^*, c_k}$. By IH, $v \notin \llbracket \phi \rrbracket^{\mathcal{K}^*}$, i.e., $\mathcal{K}^*, v \not\Vdash \phi$. And since $w \in \Diamond_{c_k} (Q(c_k) \cap \overline{P(c_k)}) = \Diamond_{c_k} \{v\}$, that means $v \in R[w]$, and so $\mathcal{K}^*, w \not\Vdash \Box \phi$. \square

COROLLARY B2.5 (Unaxiomatizability of classically complete consequence in $\mathcal{L}^{\mathcal{QH}}$). $\models_{\text{Co}_{cl}}$ in $\mathcal{L}^{\mathcal{QH}}$ is unaxiomatizable, as is $\models_{\text{Co}_{cl}X}$ for any X that is the intersection of any of the classes mentioned in Tables B1 and B6 where $\text{Co}_{cl}X \neq \emptyset$.

Proof. Since $V(cl) = \{c \in \mathbb{H}_W \mid c \text{ is classical}\}$, there is a $c \in V(cl)$ such that c is full. So adding $@_{cl} \forall p \exists q(p = @_k q)$ to Δ is enough to ensure that c_k is full. \square

B2.3. Constraints on propositions. Let's now turn to constraints on the proposition domain. A sample of such constraints is given in Table B6. For strong closure, we write $\mathcal{M} \approx \mathcal{M}'$ to mean \mathcal{M} and \mathcal{M}' are based on the same hyperframe (i.e., only differ in

³ For the \subseteq -direction: If $v \in R[w] = \{u \in W \mid \forall X \subseteq W : w \in \Box_{c_k} X \Rightarrow u \in X\}$, then $w \notin \Box_{c_k} \overline{\{v\}} = \Box_{c_k} \neg_{c_k} \{v\}$, and so $w \in \Diamond_{c_k} \{v\}$. For the \supseteq -direction: If $w \in \Diamond_{c_k} \{v\}$, then $w \notin \Box_{c_k} \overline{\{v\}}$. So let $X \subseteq W$ where $w \in \Box_{c_k} X$. If $v \notin X$, then $X \subseteq \overline{\{v\}}$. Thus, $(X \rightarrow_{c_k} \overline{\{v\}}) = W$. By the necessitation formula, $\Box_{c_k} W = W$. Hence, $w \in \Box_{c_k} (X \rightarrow_{c_k} \overline{\{v\}})$. By the K axiom formula, $w \in \Box_{c_k} \overline{\{v\}}$, $\not\perp$. Hence, $v \in X$.

Table B7. Axiomatizations in \mathcal{L}^{QH} for various classes from Table B6.

Name	Axiom/Rule
Corr	$\kappa \in \iota, \lambda \in \iota \Vdash \forall p(p^\kappa = p^\lambda)$
Elim $_{\forall\Phi}$	$\forall p\phi \Vdash \phi[\chi/p]$ where $\chi \in \Phi$ and χ is free for p
Ex $_{\Phi}$	$\Vdash \exists p \&_{i=1}^n (p =_{\iota_i} \chi)$ where $\chi \in \Phi$ and p does not occur free in χ
Ex $_{\Phi}^-$	$\Vdash \text{E}\chi$ where $\chi \in \Phi$
PII $^+$	$ \iota _1, \kappa _1, \forall p(p^\iota = p^\kappa) \Vdash (\iota = \kappa)$
PII $_1^+$	$ \iota _1, \kappa _1, \forall p(p^\iota = p^\kappa), \pi_\iota _1 \Vdash (\iota = \kappa)$
Split	$\{ \iota_i _1\}_{i=1}^n, \{(\iota_i \neq \iota_j)\}_{i \neq j} \Vdash \exists p \&_{i=1}^n (p =_{\iota_i} q_i)$ where $p \notin \{q_1, \dots, q_n\}$
Hom $_{\Phi}$	$\Vdash \forall p(p =_{\iota} \chi \supset p =_{\kappa} \chi)$ where $\chi \in \Phi$ and p does not occur free in χ
Gen $_{\forall\Phi}$	if each $\chi \in \Phi$ is free for p in ψ and $\vec{\alpha} \Vdash \psi[\chi/p]$ for each $\chi \in \Phi$, then $\vec{\alpha} \Vdash \forall p\psi$
Class	Axiomatization
Cr	QH + Corr
Cl $_{\Phi}$	QH \cup Ex $_{\Phi}$
Cl $_{\Phi}^+$	QH + Ex $_{\Phi}$ = QH + Elim $_{\forall\Phi}$
Df $_{\Phi}$	QH + Gen $_{\forall\Phi}$ (only weakly complete if Φ is infinite)
Cl $_{\Phi}$ Df $_{\Phi}$	(QH \cup Ex $_{\Phi}$) \cup Hom $_{\Phi}$ = (QH \cup Ex $_{\Phi}^-$) \cup Hom $_{\Phi}$
Cl $_{\Phi}^+$ Df $_{\Phi}$	(QH + Ex $_{\Phi}$) \cup Hom $_{\Phi}$
Di	QH + PII $^+$
Cb	QH + PII $^+$ + Split = QH + PII $_1^+$ + Split
CpSi	QH + Split + Sing = QH + Split + Self-Dual $_{@}$

valuation). Axiomatizations for consequence over some of these classes are stated in Table B7. Some of the axioms use the following abbreviation: $(\phi =_{\iota} \psi) := @_{\iota}(\phi = \psi)$. Completeness for the intersections of these classes can be gotten from combining the relevant axiomatizations, with the exception of Cl $_{\Phi}$ Df $_{\Phi}$ and Cl $_{\Phi}^+$ Df $_{\Phi}$, which are mentioned explicitly in Table B7.

THEOREM B2.6 (Relative completeness in \mathcal{L}^{QH}). *The proof systems in Table B7 are sound and complete over the relevant class. (See Section B5.3.)*

§B3. Hyperintensional operators. In this section, we enrich the language of hyperlogic with hyperintensional operators and explore their logic(s). We start by adding a counterfactual conditional and then show how a similar approach can apply to belief and knowledge operators. In Section B3.1, we expand the syntax and semantics from Section §A2 to include a counterfactual conditional (following Kocurek [33]). In Section B3.2, we axiomatize the minimal counterfactual hyperlogic on this semantics. In Section B3.3, we extend this axiomatization to include an entailment operator/propositional quantifiers. In Section B3.4, we explore stronger counterfactual hyperlogics obtained by imposing restrictions on the selection function. Finally, in Section B3.5, we show how a similar approach can address the hyperlogic of belief/knowledge.

B3.1. Selection semantics. For any language \mathcal{L}^* mentioned in Part A, we can consider the language $\mathcal{L}_{\square\rightarrow}^*$, that results from extending \mathcal{L}^* with a counterfactual conditional $\square\rightarrow$. For instance, $\mathcal{L}_{\square\rightarrow}^0$ is the result of extending \mathcal{L}^0 with $\square\rightarrow$, $\mathcal{L}_{\square\rightarrow}^H$ the result of extending \mathcal{L}^H with $\square\rightarrow$, and so on. To extend hyperlogic with a counterfactual conditional, Kocurek [33] proposes we allow counterfactuals to shift the hyperconvention parameter of an index. This can be achieved by simply replacing worlds in the standard (intensional) selection semantics for counterfactuals [39, 48] with world-hyperconvention pairs. Thus, we revise Definitions A2.11 and A2.12 as follows:

DEFINITION B3.7 (Selection hypermodel). A selection hyperframe is a tuple $\mathcal{F} = \langle W, D_C, D_P, f \rangle$ where $\langle W, D_C, D_P \rangle$ is a hyperframe and $f : \wp \mathbb{I}_{D_H} \times \mathbb{I}_{D_H} \rightarrow \wp \mathbb{I}_{D_H}$ is a selection function. A selection hypermodel over \mathcal{F} is a selection hyperframe paired with a valuation for \mathcal{F} . Satisfaction is defined as in Definition A2.12 with the following addition, where $\llbracket \phi \rrbracket^{\mathcal{M}} = \{ \langle v, d \rangle \in \mathbb{I}_{D_H} \mid \mathcal{M}, v, d \Vdash \phi \}$:

$$\mathcal{M}, w, c \Vdash \phi \square\rightarrow \psi \iff f(\llbracket \phi \rrbracket^{\mathcal{M}}, w, c) \subseteq \llbracket \psi \rrbracket^{\mathcal{M}}.$$

At the outset, we impose no restrictions on the selection function. Some theorists (e.g., Cohen [15] and Nolan [43]) argue that if counter(meta)logicals are nonvacuous, then the logic of counterfactuals is trivial. For example, it is nearly universally accepted that $\phi \square\rightarrow (\psi \wedge \chi) \Vdash \phi \square\rightarrow \psi$. Yet, here is an alleged counterexample:

- (1) a. If every instance of conjunction elimination had failed, Alice and Beth would be sad.
- b. $\not\Rightarrow$ If every instance of conjunction elimination had failed, Alice would be sad.

Similar “counterexamples” can be constructed to nearly every principle of counterfactual reasoning.⁴ Even principles such as $\Vdash \phi \square\rightarrow \phi$ have been called into question [43, p. 555].⁵

Hyperlogic offers refuge to those who find this disheartening. As we’ll see, even though counter(meta)logicals are nonvacuous in hyperlogic, its counterfactual logic is

⁴ Nolan [43] makes an exception for modus ponens ($\phi \square\rightarrow \psi, \phi \Vdash \psi$), which is immune to counterexamples of this sort.

⁵ We might try to save the standard logic for counterfactuals with possible antecedents [6, 12]. It is not obvious this will work, though. Imagine Alice endorses a strange logic on which every instance of conjunction elimination fails. Then (i) is as problematic as (1) despite only having counterfactuals with possible antecedents (Alice could have had the right views about logic).

- (i) a. If Alice were right about logic, every instance of conjunction elimination would fail.
- b. If Alice were right about logic, Beth and Cher would be sad.
- c. Therefore, if Alice were right about logic, Beth would be sad.

One may try to block this counterexample by denying the first premise on the grounds that the antecedent is possible and “nothing impossible would obtain were something that’s possible to obtain.” This reasoning appeals to what Nolan [43] calls the “Strangeness of Impossibility Condition”: no impossible world can occur closer to the actual world than any possible world. But this principle has been called into question [4, 13, 43, 56]. Hyperlogic, by contrast, can explain what’s going on in examples like (1) and (i) without taking a stand on this issue.

nontrivial: the standard counterfactual principles can be salvaged when the connectives used to state those principles are classically rigidified. This means, among other things, that imposing constraints on the selection function is not incompatible with the nonvacuity of counter(meta)logicals, such as those in (1).

B3.2. Completeness. Let’s turn to the logic of counterfactual hyperlogic. Given that we are not placing any constraints on the selection function, what counterfactual principles, if any, are valid?

Kocurek and Jerzak [35, Appendix] show that the logic of classical consequence in $\mathcal{L}_{\Box \rightarrow}^0$ is the same as the logic of the standard “impossible worlds” semantics for counterfactuals, where we can model an impossible world as an arbitrary set of formulas. But this is only because (as Cohen [15] and Nolan [43] suggest) there are no valid principles of counterfactual reasoning that aren’t already substitution instances of **S5**-theorems. Thus, without further constraints, (1) is invalid in the hyperconvention semantics when regimented so:

$$(\forall p \forall q \sim((p \wedge q) \triangleright p) \Box \rightarrow (a \wedge b)) \therefore (\forall p \forall q \sim((p \wedge q) \triangleright p) \Box \rightarrow a).$$

Fortunately, counterfactual hyperlogic in $\mathcal{L}_{\Box \rightarrow}^H$ is more interesting, since it has the expressive resources to “hold fixed” the interpretation of a certain connective within the scope of a counterfactual [35, p. 21]. If we require a certain formula within a counterfactual to be interpreted according to, say, a classical hyperconvention, then any entailments that formula generates in classical logic must be preserved. For example, the reason (1) seems to invalidate conjunction elimination in the consequent is that the word “and” in the consequent is being interpreted relative to a logic where conjunction elimination fails. If we force that “and” to be interpreted classically, however, then the argument is valid. That is, (1) is valid when regimented so:⁶

$$(\forall p \forall q \sim((p \wedge q) \triangleright p) \Box \rightarrow (a \& b)) \therefore (\forall p \forall q \sim((p \wedge q) \triangleright p) \Box \rightarrow a).$$

This could explain why (1) has a ring of plausibility to it. Even though the counterlogical supposition is asking us to interpret conjunction so that conjunction

⁶ As an anonymous referee points out, hyperlogic predicts the following inference is still (universally) valid:

$$(\forall p \forall q \sim((p \& q) \triangleright p) \Box \rightarrow (a \& b)) \therefore (\forall p \forall q \sim((p \& q) \triangleright p) \Box \rightarrow a).$$

Here, “(the law of) conjunction elimination” is regimented using $\&$ rather than \wedge . I am unsure whether this is an unwelcome result (we are, after all, still using our actual notion of entailment to reason about these counterfactuals, not the notion of entailment denoted by \triangleright in the antecedent). However, if we want to avoid this result, we could revise the semantics of hyperlogic, following a suggestion from Kocurek [33, p. 13683], so that counterfactuals can shift the denotation of interpretation nominals (though not interpretation variables). Since $\&$ is defined in terms of cl , this revision would allow that $\&$ no longer has its classical meaning in the consequent. The resulting counterfactual logic would still be nontrivial, since the inference would hold if we regiment the premise as follows (given interpretation variables have rigid denotation):

$$\downarrow i.@_{cl} \downarrow k.@_i (\forall p \forall q \sim((p \wedge q) \triangleright p) \Box \rightarrow \downarrow j.@_k (@_j a \wedge @_j b)).$$

It is an open question how this revision would affect the resulting logic of hyperlogic.

Table B8. *Axioms and rules for provability in $\mathcal{L}_{\Box\rightarrow}^H$ (with some derivable rules). The rules for \Vdash can be converted into rules for \vdash (given κ isn't cl) by applying C2U, U2C, and Cl.*

H $_{\Box\rightarrow}$
 All the axioms and rules in **H**, plus:

K$_{\Box\rightarrow}$	$\phi\Box\rightarrow(\psi\supset\chi), \phi\Box\rightarrow\psi \Vdash \phi\Box\rightarrow\chi$
Nec$_{\Box\rightarrow}$	$@_i \blacksquare \psi \Vdash \phi\Box\rightarrow(\iota\supset\psi)$
Gen$_{\Box\rightarrow}$	if $\vec{\alpha}, \kappa _1 \Vdash \phi\Box\rightarrow(\kappa\supset\psi)$ where κ does not occur free in $\vec{\alpha}, \phi$, or ψ , then $\vec{\alpha} \Vdash \phi\Box\rightarrow\psi$
REA	if $\vec{\alpha} \Vdash \phi =_{\kappa} \phi'$ where κ does not occur free in $\vec{\alpha}, \phi$, or ϕ' , then $\vec{\alpha} \Vdash (\phi\Box\rightarrow\psi) = (\phi'\Box\rightarrow\psi)$

Derivable rules:

Gen$_{\Box\rightarrow, \iota}$	if $\vec{\alpha}, \kappa _1 \Vdash \phi\Box\rightarrow_{\iota}(\kappa\supset\psi)$ where κ does not occur free in $\iota, \vec{\alpha}, \phi$, or ψ , then $\vec{\alpha} \Vdash \phi\Box\rightarrow_{\iota}\psi$
RK$_{\Box\rightarrow(\iota)}$	if $\psi_1, \dots, \psi_n \Vdash \chi$, then $\phi\Box\rightarrow_{(\iota)}\psi_1, \dots, \phi\Box\rightarrow_{(\iota)}\psi_n \Vdash \phi\Box\rightarrow_{(\iota)}\chi$

elimination fails, it's nevertheless tempting to hold on to our “standard” classical way of interpreting “and” when evaluating the consequent.⁷

We can generalize this observation by axiomatizing consequence in $\mathcal{L}_{\Box\rightarrow}^H$. The axiomatic system **H $_{\Box\rightarrow}$** is given in Table B8. Some notation:

$$\begin{array}{ll}
 \phi\Box\rightarrow_{\iota}\psi & := @_{\iota}(\phi\Box\rightarrow\psi), & \phi\Diamond\rightarrow_{\iota}\psi & := \sim @_{\iota}(\phi\Box\rightarrow\sim\psi), \\
 \Box_{\phi, \iota}\psi & := \phi\Box\rightarrow_{\iota}\psi, & \Diamond_{\phi, \iota}\psi & := \phi\Diamond\rightarrow_{\iota}\psi, \\
 \Box_{\phi, \iota}^{\alpha}\psi & := \alpha\supset\Box_{\phi, \iota}\psi, & \Diamond_{\phi, \iota}^{\alpha}\psi & := \alpha\&\Diamond_{\phi, \iota}\psi, \\
 \Box_{\phi, \iota}^{\alpha}\psi & := \Box_{\phi_1, \iota_1}^{\alpha_1} \dots \Box_{\phi_n, \iota_n}^{\alpha_n} \psi, & \Diamond_{\phi, \iota}^{\alpha}\psi & := \Diamond_{\phi_1, \iota_1}^{\alpha_1} \dots \Diamond_{\phi_n, \iota_n}^{\alpha_n} \psi.
 \end{array}$$

As before, let $\mathcal{L}_{\Box\rightarrow}^{H+}$ be the expansion of \mathcal{L}^H with Prop⁺ and INom⁺.

DEFINITION B3.8 (Lindenbaum set). *A set $\Gamma \subseteq \mathcal{L}_{\Box\rightarrow}^{H+}$ is Lindenbaum if it is a $\mathcal{L}_{\Box\rightarrow}^{H+}$ -maximal consistent set that satisfies constraints (i)–(iii) from Definition A3.23 (nominalized, witnesses $\sim@s$, differentiates terms) as well as the following:*

- iv. Γ^+ **differentiates antecedents**: if $(\Box_{\phi, \iota}\psi \neq \Box_{\phi', \iota}\psi) \in \Gamma^+$, then $|l^+|_1 \in \Gamma^+$ and $(\phi \neq_{l^+} \phi') \in \Gamma^+$ for some fresh $l^+ \in \text{INom}^+$.
- v. Γ^+ **witnesses actual \Diamond s**: if $\Diamond_{\phi, \iota}^{\alpha}\psi \in \Gamma^+$, then $|l^+|_1 \in \Gamma^+$ and $\Diamond_{\phi, \iota}^{\alpha}(l^+\&\psi) \in \Gamma^+$ for some fresh $l^+ \in \text{INom}^+$.
- vi. Γ^+ **witnesses possible \Diamond s**: if $\Diamond(\alpha_0 \wedge \Diamond_{\phi, \iota}^{\alpha}\psi) \in \Gamma^+$, then $|l^+|_1 \in \Gamma^+$ and $\Diamond(\alpha_0 \wedge \Diamond_{\phi, \iota}^{\alpha}(l^+\&\psi)) \in \Gamma^+$ for some fresh $l^+ \in \text{INom}^+$.

⁷ This strategy requires we interpret “and” in the consequent of (1a) in terms of & even though we interpret “(the law of) conjunction elimination” in terms of \wedge . We see a similar phenomenon with in-scope de re readings of counterfactuals. Consider:

- (i) If I hadn't gone to college, my professor would find the class easier to teach.

Here, “my professor” in the consequent picks out the speaker's professor in the actual world even though we are entertaining the speaker never going to college. The claim that “and” in the consequent of (1a) can be interpreted according to our actual (classical) conventions even though we are entertaining an alternative convention is similar.

LEMMA B3.9 (Counterfactual Lindenbaum). *If $\Gamma \subseteq \mathcal{L}_{\square \rightarrow}^H$ is consistent, then there is a Lindenbaum set $\Gamma^+ \subseteq \mathcal{L}_{\square \rightarrow}^{H+}$ such that $\Gamma \subseteq \Gamma^+$.*

Proof. The construction is the same as that in Lemma A3.24 except for how we define Γ_{k+1} from Γ'_k (both l^+ and p^+ are unused throughout):

$$\Gamma_{k+1} = \begin{cases} \Gamma'_k \cup \{l^+ \in i, \neg @_{l^+} \psi\}, & \text{if } \phi_k \in \Gamma'_k \text{ where } \phi_k = \neg @_i \psi, \\ \Gamma'_k \cup \{@_i p^+ \neq @_{\kappa} p^+\}, & \text{if } \phi_k \in \Gamma'_k \text{ where } \phi_k = (i \neq \kappa) \wedge |i|_1 \wedge |\kappa|_1, \\ \Gamma'_k \cup \{|l^+|_1, \phi \neq_{l^+} \phi'\}, & \text{if } \phi_k \in \Gamma'_k \text{ where } \phi_k = (\Box_{\phi, i} \psi \neq \Box_{\phi', i} \psi), \\ \Gamma'_k \cup \{\diamond_{\phi, i}^\alpha (l^+ \& \psi), |l^+|_1\}, & \text{if } \phi_k \in \Gamma'_k \text{ where } \phi_k = \diamond_{\phi, i}^\alpha \psi, \\ \Gamma'_k \cup \{\diamond(\alpha_0 \wedge \diamond_{\phi, i}^\alpha (l^+ \& \psi)), |l^+|_1\}, & \text{if } \phi_k \in \Gamma'_k \text{ where } \phi_k = \diamond(\alpha_0 \wedge \diamond_{\phi, i}^\alpha \psi), \\ \Gamma'_k, & \text{otherwise.} \end{cases}$$

Suppose for reductio that Γ_{k+1} is inconsistent. The only cases we need to check are where $\phi_k = (\Box_{\phi, i} \psi \neq \Box_{\phi', i} \psi)$, where $\phi_k = \diamond_{\phi, i}^\alpha \psi$, and where $\phi_k = \diamond(\alpha_0 \wedge \diamond_{\phi, i}^\alpha \psi)$. Assume throughout the contradiction is derivable from $\gamma_1, \dots, \gamma_n \in \Gamma_k$.

Suppose $\phi_k = (\Box_{\phi, i} \psi \neq \Box_{\phi', i} \psi)$. Thus:

$$\begin{array}{ll} \widehat{\gamma}, \Box_{\phi, i} \psi \neq \Box_{\phi', i} \psi, |l^+|_1 \vdash \phi =_{l^+} \phi', & \\ cl, l_\Gamma, \widehat{\gamma}, \Box_{\phi, i} \psi \neq \Box_{\phi', i} \psi, |i|_1 \Vdash \phi =_i \phi' & \text{Lemma 3.19, C2U,} \\ @_{l_\Gamma} cl, @_{l_\Gamma} \widehat{\gamma}, \Box_{\phi, i} \psi \neq \Box_{\phi', i} \psi, |i|_1 \Vdash \phi =_i \phi' & \text{Gen@, Red, Red,} \\ @_{l_\Gamma} cl, @_{l_\Gamma} \widehat{\gamma}, \Box_{\phi, i} \psi \neq \Box_{\phi', i} \psi \Vdash \phi = \phi' & \text{Gen}_\downarrow, \text{Vac}_\downarrow, \text{Idle}_\downarrow, \\ @_{l_\Gamma} cl, @_{l_\Gamma} \widehat{\gamma}, \Box_{\phi, i} \psi \neq \Box_{\phi', i} \psi \Vdash \phi =_{l^+} \phi' & \text{Gen@, Red,} \\ l_\Gamma, |l_\Gamma|_1, \widehat{\gamma}, \Box_{\phi, i} \psi \neq \Box_{\phi', i} \psi \Vdash \phi =_{l^+} \phi' & \text{Intro@, Cl,} \\ l_\Gamma, |l_\Gamma|_1, \widehat{\gamma}, \Box_{\phi, i} \psi \neq \Box_{\phi', i} \psi \vdash \phi =_{l^+} \phi' & \text{U2C,} \\ l_\Gamma, |l_\Gamma|_1, \widehat{\gamma}, \Box_{\phi, i} \psi \neq \Box_{\phi', i} \psi \vdash \Box_{\phi, i} \psi = \Box_{\phi', i} \psi & \text{REA, } \not\downarrow \text{ (} l_\Gamma, |l_\Gamma|_1 \in \Gamma_k \text{).} \end{array}$$

Suppose $\phi_k = \diamond_{\phi, i}^\alpha \psi$. Thus:

$$\begin{array}{ll} \widehat{\gamma}, \diamond_{\phi, i}^\alpha \psi, |l^+|_1 \vdash \neg \diamond_{\phi, i}^\alpha (l^+ \& \psi), & \\ \widehat{\gamma}, \diamond_{\phi, i}^\alpha \psi, |l^+|_1 \vdash \Box_{\phi, i}^\alpha \sim (l^+ \& \psi) & \text{def. of } \diamond_{\phi, i}^\alpha, \\ \widehat{\gamma}, \diamond_{\phi, i}^\alpha \psi, |l^+|_1 \vdash \Box_{\phi, i}^\alpha (l^+ \supset \sim \psi) & \text{RK}_{\square \rightarrow i}, \\ \widehat{\gamma}, \diamond_{\phi, i}^\alpha \psi \vdash \Box_{\phi, i}^\alpha \sim \psi & \text{Gen}_{\square \rightarrow i}, \\ \widehat{\gamma} \vdash \neg \diamond_{\phi, i}^\alpha \psi & \text{def. of } \diamond_{\phi, i}^\alpha, \not\downarrow. \end{array}$$

Suppose $\phi_k = \diamond(\alpha_0 \wedge \diamond_{\phi, i}^\alpha \psi)$. Thus:

$$\begin{array}{ll} \widehat{\gamma}, \diamond(\alpha_0 \wedge \diamond_{\phi, i}^\alpha \psi), |l^+|_1 \vdash \Box(\alpha_0 \rightarrow \Box_{\phi, i}^\alpha (l^+ \supset \sim \psi)), & \\ \diamond \widehat{\gamma}, \diamond(\alpha_0 \wedge \diamond_{\phi, i}^\alpha \psi), |l^+|_1, \alpha_0 \vdash \Box_{\phi, i}^\alpha (l^+ \supset \sim \psi) & \text{S5, Rigid,} \\ \diamond \widehat{\gamma}, \diamond(\alpha_0 \wedge \diamond_{\phi, i}^\alpha \psi), \alpha_0 \vdash \Box_{\phi, i}^\alpha \sim \psi & \text{Gen}_{\square \rightarrow i}, \\ \diamond \widehat{\gamma}, \diamond(\alpha_0 \wedge \diamond_{\phi, i}^\alpha \psi) \vdash \Box(\alpha_0 \rightarrow \Box_{\phi, i}^\alpha \sim \psi) & \text{RK, S5,} \\ \widehat{\gamma} \vdash \Box(\alpha_0 \rightarrow \diamond_{\phi, i}^\alpha \sim \psi) & \text{S5, } \not\downarrow. \end{array}$$

□

LEMMA B3.10 (Counterfactual existence). *Where $\Delta \in W_\Gamma$:*

- a. *If $\Box \phi \notin \Delta$, then there is a $\Delta' \in W_\Gamma$ such that $\phi \notin \Delta'$.*

b. If $\diamond_{\phi,i}(l \& \theta) \in \Delta$ where $|l|_1 \in \Delta$, then there is a $\Delta' \in W_\Gamma$ extending $\{l_\Delta, |l_\Delta|_1, |l|_1, @_i \theta\} \cup \{@_i \chi \mid \square_{\phi,i}(l \supset \chi) \in \Delta\}$.

Proof. Start with (a). By the proof of Lemma A3.26, $\Delta^\square \cup \{\neg\phi\}$ is consistent. Enumerate all formulas of the form $\neg @_i \psi$, of the form $\diamond_{\phi,i}^\alpha \psi$, or of the form $\diamond(\alpha_0 \wedge \diamond_{\phi,i}^\alpha \psi)$ as $\chi_1, \chi_2, \chi_3, \dots$. We define a sequence of formulas $\delta_0, \delta_1, \delta_2, \dots$ depending on the form of χ_{n+1} . As before, $\delta_0 := \neg\phi$. If $\chi_{n+1} = \neg @_i \psi$, then define δ_{n+1} as in Lemma A3.26. If $\chi_{n+1} = \diamond_{\phi,i}^\alpha \psi$, define $\delta_{n+1} := \chi_{n+1} \rightarrow (|l^+|_1 \wedge \diamond_{\phi,i}^\alpha(l^+ \& \psi))$, where l^+ is the first nominal such that $\Delta^\square, \delta_0, \dots, \delta_n, \chi_{n+1} \rightarrow (|l^+|_1 \wedge \diamond_{\phi,i}^\alpha(l^+ \& \psi)) \not\vdash \perp$. Suppose for reductio there were no such l^+ . Reasoning as in Lemma A3.26, we can conclude that $\square(\widehat{\delta} \rightarrow \chi_{n+1}) \in \Delta$ and $\square(\widehat{\delta} \rightarrow \neg(|l^+|_1 \wedge \diamond_{\phi,i}^\alpha(l^+ \& \psi))) \in \Delta$ for all l^+ , and that $\diamond(\widehat{\delta} \wedge \chi_{n+1}) \in \Delta$. Since Δ witnesses possible \diamond s, there is an l^+ such that $\diamond(\widehat{\delta} \wedge \diamond_{\phi,i}^\alpha(l^+ \& \psi)) \in \Delta, \not\vdash$.

If $\chi_{n+1} = \diamond(\alpha_0 \wedge \diamond_{\phi,i}^\alpha \psi)$, define $\delta_{n+1} := \chi_{n+1} \rightarrow (|l^+|_1 \wedge \diamond(\alpha_0 \wedge \diamond_{\phi,i}^\alpha(l^+ \& \psi)))$, where l^+ is the first such that $\Delta^\square, \delta_0, \dots, \delta_n, \chi_{n+1} \rightarrow (|l^+|_1 \wedge \diamond(\alpha_0 \wedge \diamond_{\phi,i}^\alpha(l^+ \& \psi))) \not\vdash \perp$. Suppose there is no such l^+ . Then $\square(\widehat{\delta} \rightarrow \chi_{n+1}) \in \Delta$ and $\square(\widehat{\delta} \rightarrow \neg(|l^+|_1 \wedge \diamond(\alpha_0 \wedge \diamond_{\phi,i}^\alpha(l^+ \& \psi)))) \in \Delta$ for all l^+ . As before, $\diamond(\widehat{\delta} \wedge \chi_{n+1}) \in \Delta$, i.e., $\diamond(\widehat{\delta} \wedge \diamond(\alpha_0 \wedge \diamond_{\phi,i}^\alpha \psi)) \in \Delta$. By S5, $\diamond \widehat{\delta} \wedge \diamond(\alpha_0 \wedge \diamond_{\phi,i}^\alpha \psi) \in \Delta$. Since Δ witnesses possible \diamond s, there is an l^+ such that $\diamond(\alpha_0 \wedge \diamond_{\phi,i}^\alpha(l^+ \& \psi)) \in \Delta$. By S5 again, $\diamond(\widehat{\delta} \wedge \diamond(\alpha_0 \wedge \diamond_{\phi,i}^\alpha(l^+ \& \psi))) \in \Delta, \not\vdash$.

Now for (b). Let $\Delta^{\square_{\phi,i}} = \{@_i \chi \mid \square_{\phi,i}(l \supset \chi) \in \Delta\}$. Then $\Delta^{\square_{\phi,i}} \cup \{l_\Delta, |l_\Delta|_1, |l|_1, @_i \theta\}$ is consistent. For suppose not. Then for some $@_i \chi_1, \dots, @_i \chi_n \in \Delta^{\square_{\phi,i}}$, we have

$l_\Delta, l_\Delta _1, l _1, @_i \widehat{\chi} \vdash \sim @_i \theta$	Bool,
$l_\Delta, l_\Delta _1, l _1, @_i \widehat{\chi} \vdash @_i \sim \theta$	Dist _@ ,
$cl, l_\Delta, l_\Delta _1, l _1, @_i \widehat{\chi} \Vdash @_i \sim \theta$	C2U,
$@_{l_\Delta} cl, l_\Delta _1, l _1, @_i \widehat{\chi} \Vdash @_i \sim \theta$	Gen _@ , Ref, Red,
$l_\Delta \in cl, l _1, @_i \widehat{\chi} \Vdash @_i \sim \theta$	Elim _{&} ,
$l_\Delta \in cl, l _1, l, \widehat{\chi} \Vdash \sim \theta$	Intro _@ , Elim _@ ,
$l_\Delta \in cl, l _1, l \supset \widehat{\chi} \Vdash l \supset \sim \theta$	S5,
$\square_{\phi,i}(l_\Delta \in cl), \square_{\phi,i} l _1, \square_{\phi,i} \widehat{\chi} \Vdash \square_{\phi,i} \sim \theta$	RK _{\square_{\rightarrow i}} ,
$l_\Delta \in cl, l _1, \square_{\phi,i} \widehat{\chi} \Vdash \square_{\phi,i} \sim \theta$	Rigid, Nec _{\square_{\rightarrow i}} , Gen _{\square_{\rightarrow i}} .

Since $\square_{\phi,i} \widehat{\chi}, (l_\Delta \in cl), |l|_1 \in \Delta$, that means $\square_{\phi,i} \sim \theta \in \Delta$, contrary to our initial assumption that $\diamond_{\phi,i} \theta \in \Delta, \not\vdash$.

Now, suppose $\square \chi \in \Delta$. Thus, $@_i \blacksquare @_{l_\Delta} \chi \in \Delta$ (by Rigid, Intro_@, Red, Bool, and Dist_@). By Nec_{\square_{\rightarrow i}}, $\square_{\phi,i}(l \supset @_{l_\Delta} \chi) \in \Delta$. Hence, $@_i @_{l_\Delta} \chi \in \Delta^{\square_{\phi,i}}$. Since $l_\Delta, |l_\Delta|_1 \vdash @_i @_{l_\Delta} \chi \leftrightarrow \chi$, the set $\Sigma := \Delta^{\square_{\phi,i}} \cup \{l_\Delta, |l_\Delta|_1, |l|_1, @_i \theta\} \cup \{\chi \mid \square \chi \in \Delta\}$ is consistent. The proof strategy from here is essentially the same as in part (a), though some changes need to be made to ensure the steps go through. The main change is that we need to replace $\square(\widehat{\delta} \rightarrow \dots)$ and $\diamond(\widehat{\delta} \wedge \dots)$ with $\square_{\phi,i}(\widehat{\delta} \supset \dots)$ and $\diamond_{\phi,i}(\widehat{\delta} \& \dots)$. To illustrate, I'll use the case where $\chi_{n+1} = \diamond(\alpha_0 \wedge \diamond_{\beta,\kappa}^\alpha \psi)$. As before, we define $\delta_{n+1} := \chi_{n+1} \rightarrow (|l^+|_1 \wedge \diamond(\alpha_0 \wedge \diamond_{\beta,\kappa}^\alpha(l^+ \& \psi)))$, where l^+ is the first such that: $\Sigma, \delta_0, \dots, \delta_n, \chi_{n+1} \rightarrow (|l^+|_1 \wedge \diamond(\alpha_0 \wedge \diamond_{\beta,\kappa}^\alpha(l^+ \& \psi))) \not\vdash \perp$. By Bool and the fact that $l_\Delta, |l_\Delta|_1 \in \Sigma$, this

condition is equivalent to $\Sigma, \delta_0, \dots, \delta_n, \chi_{n+1} \supset (|l^+|_1 \& @_{l_\Delta} \blacklozenge (\alpha_0 \& \blacklozenge_{\beta, \kappa}^\alpha (l^+ \& \psi))) \not\vdash \perp$. Suppose, for reductio, there's no such l^+ . So for all l^+ , there are some $\gamma_1, \dots, \gamma_n \in \Sigma$ such that $\widehat{\gamma} \vdash \widehat{\delta} \supset \sim (\chi_{n+1} \supset @_{l_\Delta} \blacklozenge (\alpha_0 \& \blacklozenge_{\beta, \kappa}^\alpha (l^+ \& \psi)))$. By $\text{RK}_{\square \rightarrow, i}$, $\square_{\phi, i} \widehat{\gamma} \vdash \square_{\phi, i} (\widehat{\delta} \supset \sim (\chi_{n+1} \supset @_{l_\Delta} \blacklozenge (\alpha_0 \& \blacklozenge_{\beta, \kappa}^\alpha (l^+ \& \psi))))$. Since $\square_{\phi, i} \widehat{\gamma} \in \Delta$,⁸ this means $\square_{\phi, i} (\widehat{\delta} \supset (\chi_{n+1} \& \sim (|l^+|_1 \& @_{l_\Delta} \blacklozenge (\alpha_0 \& \blacklozenge_{\beta, \kappa}^\alpha (l^+ \& \psi)))) \in \Delta$ for all l^+ . As before, $\blacklozenge_{\phi, i} (\widehat{\delta} \& \chi_{n+1}) \in \Delta$, i.e., $\blacklozenge_{\phi, i} (\widehat{\delta} \& @_{l_\Delta} \blacklozenge (\alpha_0 \& \blacklozenge_{\beta, \kappa}^\alpha \psi)) \in \Delta$. By $\text{RK}_{\square \rightarrow, i}$, $\blacklozenge_{\phi, i} \widehat{\delta} \in \Delta$ and $\blacklozenge_{\phi, i} @_{l_\Delta} \blacklozenge (\alpha_0 \& \blacklozenge_{\beta, \kappa}^\alpha \psi) \in \Delta$. By $\text{Nec}_{\square \rightarrow}$ and $\text{Gen}_{\square \rightarrow, i}$, $@_{l_\Delta} \blacksquare (\alpha_0 \& \blacklozenge_{\beta, \kappa}^\alpha \psi) \Vdash \square_{\phi, i} @_{l_\Delta} \blacksquare (\alpha_0 \& \blacklozenge_{\beta, \kappa}^\alpha \psi)$. Since $|l_\Delta|_1 \in \Delta$, that means $@_{l_\Delta} \blacklozenge (\alpha_0 \& \blacklozenge_{\beta, \kappa}^\alpha \psi) \in \Delta$. Since $l_\Delta \in \Delta$, that means $\blacklozenge (\alpha_0 \wedge \blacklozenge_{\beta, \kappa}^\alpha \psi)$ by Bool . Since Δ witnesses possible \blacklozenge s, there is an l^+ such that $|l^+|_1 \& @_{l_\Delta} \blacklozenge (\alpha_0 \& \blacklozenge_{\beta, \kappa}^\alpha (l^+ \& \psi)) \in \Delta$. By $\text{Nec}_{\square \rightarrow}$, $\square_{\phi, i} (|l^+|_1 \& @_{l_\Delta} \blacklozenge (\alpha_0 \& \blacklozenge_{\beta, \kappa}^\alpha (l^+ \& \psi))) \in \Delta$. By $\text{RK}_{\square \rightarrow, i}$, $\blacklozenge_{\phi, i} (\widehat{\delta} \& (|l^+|_1 \& @_{l_\Delta} \blacklozenge (\alpha_0 \& \blacklozenge_{\beta, \kappa}^\alpha (l^+ \& \psi)))) \in \Delta$, $\not\zeta$. \square

The proofs of the other intermediate lemmas are all as before. To finish the proof, we need to define the selection function for our canonical model.

DEFINITION B3.11 (Defining formula). *Where $A \subseteq \mathbb{I}_{W_\Gamma}$, we define the set $[A] := \{\phi \in \mathcal{L}_{\square \rightarrow}^{\text{H}^+} \mid A = \{\langle \Delta, c_\kappa \rangle \mid @_\kappa \phi \in \Delta\}\}$.*

LEMMA B3.12 (Replacement of definitions). *For all $A \subseteq \mathbb{I}_{W_\Gamma}$, all c_κ , all $\phi, \phi' \in [A]$, and all ψ , we have $((\phi \square \rightarrow_\kappa \psi) = (\phi' \square \rightarrow_\kappa \psi)) \in \Gamma$.*

Proof. Suppose for reductio that $((\phi \square \rightarrow_\kappa \psi) = (\phi' \square \rightarrow_\kappa \psi)) \notin \Gamma$. Since Γ differentiates antecedents, there are some l^+ such that (by $\text{Dist}_{@}$) $(@_{l^+} \phi \neq @_{l^+} \phi') \in \Gamma$. Since $\phi, \phi' \in [A]$, $(@_{l^+} \phi = @_{l^+} \phi') \in \Gamma_n$, $\not\zeta$. \square

DEFINITION B3.13 (Canonical selection function). *We define f_Γ , the canonical selection function for Γ , as follows for all $A \subseteq \mathbb{I}_{W_\Gamma}$, all $\Delta \in W_\Gamma$, and all c_κ . First, if $[A] = \emptyset$, then $f_\Gamma(A, \Delta, c_\kappa) = \{\langle \Delta, c_\kappa \rangle\} \cap A$. Second, if $\phi \in [A]$, then $\langle \Delta', c_\lambda \rangle \in f_\Gamma(A, \Delta, c_\kappa)$ iff for all $\psi \in \mathcal{L}_{\square \rightarrow}^{\text{H}^+}$, if $(\phi \square \rightarrow_\kappa (\lambda \supset \psi)) \in \Delta$, then $@_\lambda \psi \in \Delta'$.*

By Lemma B3.12, if $\phi, \phi' \in [A]$, then $(\phi \square \rightarrow_\kappa (\lambda \supset \psi)) \in \Delta$ iff $(\phi' \square \rightarrow_\kappa (\lambda \supset \psi)) \in \Delta$, so this definition for f_Γ is well-defined.

DEFINITION B3.14 (Canonical model). *The canonical selection hypermodel of Γ is the selection hypermodel $\mathcal{M}_\Gamma = \langle W_\Gamma, D_{\square \Gamma}, D_{\mathbb{P}\Gamma}, f_\Gamma, V_\Gamma \rangle$ where $\langle W_\Gamma, D_{\square \Gamma}, D_{\mathbb{P}\Gamma}, V_\Gamma \rangle$ is defined as in Definition A3.32 and f_Γ is defined as in Definition B3.13.*

LEMMA B3.15 (Truth). $\mathcal{M}_\Gamma, \Delta, c_\kappa \Vdash \phi$ iff $@_\kappa \phi \in \Delta$.

Proof. The inductive steps are all the same as before. We just need to check the $\square \rightarrow$ step goes through. First, $\Delta, c_\kappa \Vdash \phi \square \rightarrow \psi$ iff $f_\Gamma(\llbracket \phi \rrbracket, \Delta, c_\kappa) \subseteq \llbracket \psi \rrbracket$. By Lemma B3.12 and by IH ($\phi \in \llbracket \phi \rrbracket$), this holds iff the following holds for all Δ' and c_λ :

$$\text{if } \forall \chi \in \mathcal{L}_{\square \rightarrow}^{\text{H}^+} : (\phi \square \rightarrow_\kappa (\lambda \supset \chi)) \in \Delta \Rightarrow @_\lambda \chi \in \Delta', \text{ then } @_\lambda \psi \in \Delta'.$$

We now show this condition holds for all Δ' and c_λ iff $@_\kappa (\phi \square \rightarrow \psi) \in \Delta$.

(\Leftarrow) Suppose $@_\kappa (\phi \square \rightarrow \psi) \in \Delta$. Let Δ' and c_λ be such that for all $\chi \in \mathcal{L}_{\square \rightarrow}^{\text{H}^+}$, if $(\phi \square \rightarrow_\kappa (\lambda \supset \chi)) \in \Delta$, then $@_\lambda \chi \in \Delta'$. Since $@_\kappa (\phi \square \rightarrow \psi) \in \Delta$, we have by $\text{RK}_{\square \rightarrow, \kappa}$ that $@_\kappa (\phi \square \rightarrow (\lambda \supset \psi)) \in \Delta$. Hence, $@_\lambda \psi \in \Delta'$.

⁸ This is the step that would not have gone through without the relevant change, since we do not have $\square \widehat{\gamma} \in \Delta$.

Table B9. *Axioms and rules for provability in $\mathcal{L}_{\square\rightarrow}^{QH}$*

$\mathbf{QH}_{\square\rightarrow}$	All the axioms and rules in \mathbf{QH} and $\mathbf{H}_{\square\rightarrow}$, plus:
$\mathbf{BF}_{\square\rightarrow}$	$\forall p(\phi \square\rightarrow \psi) \Vdash \phi \square\rightarrow \forall p\psi$ where p does not occur free in ϕ

Table B10. *Some constraints on selection functions.*

Name	Class	Constraint (on all $A, B \subseteq \mathbb{I}_{D_{\text{III}}}$)
success	Suc	$f(A, w, c) \subseteq A$
weak centering	W	if $\langle w, c \rangle \in A$, then $\langle w, c \rangle \in f(A, w, c)$
strong centering	C	if $\langle w, c \rangle \in A$, then $f(A, w, c) = \{\langle w, c \rangle\}$
Stalnaker's assumption	Stal	$ f(A, w, c) \leq 1$
vacuism	Vac	if $A(c) = \emptyset$, then $f(A, w, c) = \emptyset$
necessary consequent	NC	$f(A, w, c) \subseteq W \times \{c\}$
necessary entailment	NEC	$f(A, w, c) \subseteq A(c) \times \{c\}$
strangeness of impossibility	SIC	if $A(c) \neq \emptyset$, then $f(A, w, c) \subseteq W \times \{c\}$
operational rigidity	R_{\circ}	$f(A, w, c) \subseteq W \times \{c' \in D_{\text{III}} \mid c \approx c'\}$

(\Rightarrow) Suppose $@_{\kappa}(\phi \square\rightarrow \psi) \notin \Delta$. Thus, $\phi \diamond\rightarrow_{\kappa} \sim\psi \in \Delta$. Since Δ witnesses actual \diamond s, there is an l^+ such that $\phi \diamond\rightarrow_{\kappa}(l^+ \& \sim\psi)$. By Lemma B3.10, there is a $\Delta' \in W_{\Gamma}$ such that $\Delta' \supseteq \{\neg @_{l^+} \psi\} \cup \{ @_{l^+} \chi \mid \phi \square\rightarrow_{\kappa}(l^+ \supset \chi) \in \Delta \}$. Hence, $\langle \Delta', l^+ \rangle$ is the counterexample we need. \square

B3.3. Adding \triangleright and quantifiers. What changes if we add an entailment operator or propositional quantifiers to $\mathcal{L}_{\square\rightarrow}^H$? In the former case, no additional axioms are required apart from those in $\mathbf{H}_{\triangleright}$ and $\mathbf{H}_{\square\rightarrow}$: all the proofs in Section B3.2 go through in the presence of \triangleright . In the latter case, we do need one additional axiom. Observe that the Barcan formula and its converse are universally valid for counterfactuals (where p does not occur free in ϕ):

$$\forall p(\phi \square\rightarrow \psi) \Vdash \phi \square\rightarrow \forall p\psi.$$

The converse Barcan formula is easily derived just by combining \mathbf{QH} and $\mathbf{H}_{\square\rightarrow}$:

$$\begin{array}{ll} \forall p\psi \Vdash \psi & \text{Elim}_{\forall}, \\ \phi \square\rightarrow \forall p\psi \Vdash \phi \square\rightarrow \psi & \text{RK}_{\square\rightarrow}, \\ \forall p(\phi \square\rightarrow \forall p\psi) \Vdash \forall p(\phi \square\rightarrow \psi) & \text{RK}_{\forall}, \\ \phi \square\rightarrow \forall p\psi \Vdash \forall p(\phi \square\rightarrow \psi) & \text{Vac}_{\forall}. \end{array}$$

However, the Barcan formula, which is needed to prove the analogue of Lemma A4.41, must be added separately. Other than that, the proofs of completeness for \mathcal{L}^{QH} and $\mathcal{L}_{\square\rightarrow}^H$ can be straightforwardly combined to yield a proof of completeness for $\mathcal{L}_{\square\rightarrow}^{QH}$.

B3.4. Constraints on selection function. Let's now look at some constraints on the selection function. Table B10 contains several such constraints. We can extend the completeness result to include such constraints by adding the relevant axioms from Table B11.

Table B11. *Axiomatizations in $\mathcal{L}_{\Box\rightarrow}^0$ for various classes from Table B10.*

<i>Name</i>	<i>Axiom(s)</i>	<i>Corresponding Constraint</i>
$\text{Id}_{\Box\rightarrow}$	$\Vdash \phi \Box \rightarrow \phi$	success
$\text{MP}_{\Box\rightarrow}$	$\phi, \phi \Box \rightarrow \psi \Vdash \psi$	weak centering
Cen	$\phi \Vdash (\phi \Box \rightarrow \psi) \equiv \psi$	strong centering
CEM	$\Vdash (\phi \Box \rightarrow \psi) + (\phi \Box \rightarrow \sim \psi)$	Stalnaker’s assumption
Vac	$\sim \blacklozenge \phi \Vdash \phi \Box \rightarrow \psi$	vacuism
NC	$\blacksquare \psi \Vdash \phi \Box \rightarrow \psi$ $\Vdash \downarrow i. (\phi \Box \rightarrow i)$	necessary consequent
NEC	$\blacksquare (\phi \supset \psi) \Vdash \phi \Box \rightarrow \psi$ $\Vdash \downarrow i. (\phi \Box \rightarrow (i \& \phi))$	necessary entailment
SIC	$\blacklozenge \phi, \blacksquare \psi \Vdash \phi \Box \rightarrow \psi$ $\blacklozenge \phi \Vdash \downarrow i. (\phi \Box \rightarrow i)$	strangeness of impossibility
R_o	$\Vdash \downarrow i. \Box \phi \downarrow j. [\Delta(\vec{\phi}) = @_i \Delta(@_j \vec{\phi})]$	operational rigidity

THEOREM B3.16 (Relative completeness in $\mathcal{L}_{\Box\rightarrow}^H$). *The proof systems in Table B11 are sound and complete for the relevant class of selection hypermodels. (See Section B5.4.)*

Let me briefly explain the motivation behind some of these constraints. Vacuism is the view that all counterpossibles (counterfactuals with impossible antecedents) are vacuously true.⁹ Often, vacuists also endorse the necessary consequent and necessary entailment principles, which are all coderivable given success (the labels come from [20]). Some of these principles have equivalent “hybrid” formulations. In the hyperconvention semantics (with success), counterpossibles are vacuous when we hold fixed the interpretation of the antecedent. This goes back to one of the main motivations for considering hyperlogic as a semantics for metalogical claims, viz., it can formalize “conventionalist” approaches to hyperintensionality, which explain hyperintensionality in terms of convention-shifting (Section §A1). We can regiment this idea of “holding fixed” an interpretation using the hybrid binder \downarrow , which is what allows alternative axiomatizations for some of these principles.

Second, the “strangeness of impossibility condition” was introduced by Nolan [43, p. 550]. If we think of selection functions as selecting the “closest” or “most similar” antecedent-worlds, then the condition effectively says that impossible worlds are always “far out” in that they’re less similar than any possible world.¹⁰ French et al. [20] present an impossible worlds semantics where this corresponds to the axiom $(\blacklozenge \phi \wedge \Box \psi) \rightarrow (\phi \Box \rightarrow \psi)$, which has an analogue in Table B11. Again, this has an equivalent formulation in terms of convention-shifting: counterconventional readings only arise when the antecedent in question is impossible (on its actual interpretation).

Finally, operational rigidity, in effect, states *counterlogical* vacuism, i.e., the view that all counterlogicals (counterfactuals with logically impossible antecedents) are vacuously true. Some nonvacuists have held that even if counterpossibles are generally nonvacuous, counterlogicals are a special exception, while others have argued there’s

⁹ For a defense of vacuism, see [3, 18, 37, 39, 48, 52, 57, 58]. For criticism, see [4, 6, 12, 14, 15, 22, 29–31, 36, 40, 41, 43, 53, 56, 60]. See [7, 32] for an overview.

¹⁰ See [4, 13, 27, 31, 36, 40, 43, 56] for discussion of this principle. See [32] for an overview.

no relevant difference between counterlogicals and other counterpossibles.¹¹ In hyperlogic, this turns on whether counterfactuals are allowed to shift the interpretation of the connectives. Thus, those who maintain that counterlogicals are a special exception can hold that counterfactuals are only allowed to shift the interpretation of nonlogical vocabulary.

B3.5. Belief and knowledge. Thus far, we have focused on counterfactual hyperlogic. But the selection semantics (or something like it) is also often employed as a semantics for dyadic belief and knowledge, where $B^\phi\psi$ says the agent believes that ψ given ϕ and likewise for $K^\phi\psi$.¹² It is standard to define the monadic belief operator as $B\phi := B^\top\phi$ (here, we can define $\top := (p + \sim p)$). Letting $R(w, c) := f(\llbracket \top \rrbracket, w, c)$, we then have the following semantics for monadic belief:

$$\mathcal{M}, w, c \Vdash B\phi \iff \text{for all } \langle v, d \rangle \in R(w, c) : \mathcal{M}, v, d \Vdash \phi.$$

Thus, we can import the results in Section B3.2 to give a logic of belief and knowledge within hyperlogic. As in Section B3.4, one could consider imposing any of the usual restrictions on R to obtain stronger logics.

One application of doxastic/epistemic hyperlogic is to the problem of logical omniscience. It is well known that on the standard intensional semantics, belief is closed under classical entailment: if $\phi \models \psi$, then $B\phi \models B\psi$.¹³ Attempts to avoid this result generally often appeal to limitations or defects in cognitive states, e.g., lack of computational resources, awareness, or informational access. However, another (less discussed) way logical omniscience can fail is via different views on logic. If Inej believes intuitionistic logic is correct, then her beliefs will not generally be closed under classical consequence even if she is a perfect reasoner.

Doxastic hyperlogic is well-equipped to handle such cases. While it does not require that beliefs are closed under classical consequence, it does validate a more modest closure principle: $@_i \blacksquare (\phi \supset \psi), B_i, B\phi \Vdash B\psi$. Restricting to hypermodels where \triangleright_c is factive and noncontingent (Table B3), we can simplify this principle: $@_i (\phi \triangleright \psi), B_i, B\phi \Vdash B\psi$. In other words, beliefs are closed under whatever logic the agent adopts (if there is one, assuming it's reasonable). We obtain the "classical" picture only when we assume BcI holds.¹⁴

Of course, doxastic hyperlogic is not a complete solution to the problem of logical omniscience. For one, it still assumes agents are perfect reasoners within their own logic, and is thus not a good model of logical error. Moreover, beliefs are still assumed to be closed under universal consequence: if $\phi \Vdash \psi$, then $B\phi \Vdash B\psi$. The moral, rather, is that there are several different problems of logical omniscience that likely need to be addressed with different tools. Appeals to computation, awareness, fragmentation, etc.

¹¹ For defenses of counterlogical vacuism, see [22, 31]. For defenses of counterlogical nonvacuism, see [6, 12, 15, 30, 36, 40, 43, 56]. Kocurek and Jerzak [35] defend an intermediate position, viz., counterlogicals are only nonvacuous on counterconventional readings.

¹² See, e.g., [2, 11, 21, 38, 42, 54, 55].

¹³ For discussion of this problem, see [1, 5, 8–10, 16, 17, 23–28, 44, 46, 47, 49–51, 59].

¹⁴ Sedlár [45] likewise explores a doxastic logic where the belief operator is nonclassical, though the base logic is classical. In some ways, Sedlár's system is similar to doxastic hyperlogic, although the latter is more flexible in the range of logics an agent's beliefs may be sensitive to. Thanks to an anonymous referee for noting this parallel.

are better equipped for modeling logical error, whereas doxastic hyperlogic is better equipped for modeling ideal yet nonclassical agents.

§B4. Conclusion. This concludes the two-part series exploring the logic of hyperlogic. In Part A of this series, we axiomatized a minimal logic of hyperlogic. In Part B, we extended these results to stronger logics over a restricted class of models as well as to languages with hyperintensional operators. In this final section, I wish to sketch a few possible directions for future research that would be worth pursuing.

First, it is an open question how best to extend hyperlogic with first-order quantifiers. We could have hyperconventions specify a domain of individuals, but this might bring technical complications with tracking the denotations of variables across shifts in hyperconvention. Another option would be to have hypermodels directly specify a single domain across all hyperconventions. This might be more manageable, though it builds in substantive metaontological assumptions. Second, there are many questions remaining for the model theory of hyperlogic, especially concerning “finite” hypermodels. For example, it is easy to see that any satisfiable \mathcal{L}^{QH} -formula is satisfiable in a convention-finite model (i.e., one where D_{C} is finite): just reduce the hypermodel to the denotations of the free terms in the formula. Yet, there are satisfiable (quantified) \mathcal{L}^{QH} -formulas that not satisfiable in a *hyperconvention-finite* model (i.e., one where D_{H} is finite). What about any satisfiable \mathcal{L}^{H} -formula, though? Does **H** satisfy the finite model property?

Third, we made a number of choice points regarding the initial setup of the hyperconvention semantics that could be revised. One is that we required the “classical” hyperconventions to all interpret \Box and \Diamond as universal modals. It would be natural to weaken this requirement so that \Box and \Diamond only obey weaker normal modal logics. Another choice point concerned whether to treat iterated “according to” operators as redundant. I suspect there is more than one way to naturally generalize the semantics for “according to” so that iteration matters.

Finally, the hyperconvention semantics only concerns claims about logics for the propositional modal language. It does not have a way of capturing metalogical claims concerning alternative logics for *hyperlogic*—specifically, for the propositional quantifiers, hybrid operators, or counterfactuals (except insofar as they also concern alternative logics for the connectives). While Kocurek [33, Section 6] sketches a possible extension to such a language, it is unclear what the resulting logic of this proposed solution is or whether there might be more elegant solutions waiting to be explored.

§B5. Appendix: Proofs of relative completeness. In this appendix, we give the proofs of various completeness theorems relative to restricted classes of models (Theorems B2.1, B2.3, B2.6, and B3.16). First, we state a helpful lemma, which follows straightforwardly from Corollary A3.27 and Definition A3.30:

LEMMA B5.17 (Canonical operations). *Let $|\kappa|_1, |\lambda|_1 \in \Gamma$. Where $\phi_i \in [X_i]_{\kappa}$ and $\psi_i \in [Y_i]_{\lambda}$, $\Delta_{c_{\kappa}}(\vec{X}) = \Delta_{c_{\lambda}}(\vec{Y})$ iff $(\Delta(\vec{\phi})^{\kappa} = \Delta(\vec{\psi})^{\lambda}) \in \Gamma$.*


In each case, the proof of soundness is straightforward and left to the reader. Completeness requires showing the canonical model is in the relevant class. In some

cases, we must revise the canonical model construction and/or the Lindenbaum construction.

B5.1. Theorem B2.1. The proofs of completeness for $F, U_q, At,$ and B are immediate since the canonical hypermodel (Definition A3.32) is full (and thus, quantification uniform, atomic, and boolean).

U_o . We need to make a slight revision to the definition of a canonical hyperconvention. In particular, we need to revise the third clause to say that c_κ interprets the connectives classically if the following is in Γ for some ι_1, \dots, ι_n and $\lambda_1, \dots, \lambda_n$:

$$(\kappa \in \iota_1) \wedge (\lambda_1 \in \iota_1) \wedge (\lambda_1 \in \iota_2) \wedge (\lambda_2 \in \iota_2) \wedge \dots \wedge (\lambda_n \in \iota_n) \wedge (\lambda_n \in cl)$$



So unlike Definition A3.30, c_κ can be classical even if $@_\kappa cl \notin \Gamma$, so long as it satisfies this “zigzag” condition. Now, Lemma A3.33 needs to be restated as the following:

CLAIM. *If $(\kappa \in \iota), (\lambda \in \iota) \in \Gamma$ and c_κ is classical, then c_λ is classical.*

Proof. Suppose first that κ satisfies the zigzag condition. Then the zigzag can be extended to λ via ι , and thus c_λ is classical. Suppose instead that κ does not satisfy the zigzag condition. Then $c_\kappa(\neg)(X) = \{\Delta \in W_\Gamma \mid \exists \phi \in [X]_\kappa : @_\kappa \neg \phi \in \Delta\}$. Suppose $[X]_\kappa = \emptyset$. Then $c_\kappa(\neg)(X) = \emptyset$. But since c_κ is classical, $c_\kappa(\neg)(X) = \bar{X}$. So $X = W$, even though $(p + \sim p) \in [W]_\kappa, \not\in$. Hence, there is no X where $[X]_\kappa = \emptyset$. This can only happen if W_Γ is finite. List the members of W_Γ as $\Delta_1, \dots, \Delta_n$. Since these are all distinct maximal consistent sets, there must be some $\delta_1, \dots, \delta_n$ such that $\delta_i \in \Delta_j$ iff $i = j$. Each $X \subseteq W_\Gamma$ is then definable by a disjunction of these δ_i s (if $X = \emptyset$, then it’s definable by \perp). Now, let $\iota_\Gamma \in \Gamma$ and for each $X \subseteq W_\Gamma$, let $\delta_X = @_{\iota_\Gamma} \delta_{i_1} + \dots + @_{\iota_\Gamma} \delta_{i_k}$, where the disjunction of $\delta_{i_1}, \dots, \delta_{i_k}$ defines X . By Red and $Dist_{@}$, $((\delta_X)^\kappa = (\delta_X)^\lambda) \in \Gamma$ for every X . By $Unio.$, $((\star \delta_X)^\kappa = (\star \delta_X)^\lambda) \in \Gamma$. Thus, $c_\kappa(\star) = c_\lambda(\star)$ by Lemma B5.17, and so c_λ is classical. \square

Using this claim in place of Lemma A3.33 in the inductive step for the connectives in Lemma A3.34, the completeness proof goes through as before. We just need to check that D_{C_Γ} is operationally uniform. Let $c_\kappa, c_\lambda \in C_\Gamma$. By the above claim, c_κ is classical iff c_λ is classical. If both are classical, then we’re done. So suppose otherwise. I just prove the \star -case for illustration. If $[X]_\kappa = \emptyset$, then $[X]_\lambda = \emptyset$ (otherwise, if $\phi \in [X]_\lambda$, then $@_\lambda \phi \in [X]_\kappa$). If $[X]_\kappa = [X]_\lambda = \emptyset$, then $c_\kappa(\star)(X) = c_\lambda(\star)(X) = \emptyset$. So suppose $\phi \in [X]_\kappa$ and $\psi \in [X]_\lambda$. Then $@_\kappa \phi \in \Delta$ iff $@_\lambda \psi \in \Delta$. By Corollary A3.27 and Bool, $@_\kappa \phi = @_\lambda \psi \in \Gamma$. By $Unio.$, $@_\kappa \star \phi = @_\lambda \star \psi \in \Gamma$. Hence, $c_\kappa(\star)(X) = c_\lambda(\star)(X)$ by Lemma B5.17.

Si Completeness is straightforward. To establish that $\mathbf{H} + Sing = \mathbf{H} + Self-Dual^+$, we just need to show that Sing is coderivable with $Self-Dual_{@}$ in \mathbf{H} . $Self-Dual_{@}$ trivially follows from $Dist_{@}$ and Sing. Here’s the other direction:

$$\iota, i \Vdash \sim @_\iota \sim i \qquad \text{Elim}_{@},$$

$\iota, i \Vdash @_\iota i$	Self-Dual _@ ,
$\iota \Vdash \downarrow i. @_\iota i$	Gen _{\downarrow} , Vac _{\downarrow} ,
$\Vdash \iota _1$	Gen _@ , Ref, def. of $ \iota _1$.

AnF, AnU_q. We revise the Lindenbaum construction, specifically the definition of Γ_{k+1} . Let $\kappa \approx \lambda$ abbreviate $((\vec{p}^+)^{\kappa} = (\vec{q}^+)^{\lambda}) \wedge \bigvee \left\{ (\Delta(\vec{p}^+))^{\kappa} \neq (\Delta(\vec{q}^+))^{\lambda} \right\}_{\Delta}$, where \vec{p}^+ and \vec{q}^+ are unused at this point in the construction. Then we revise the definition of Γ_{k+1} so that $\Gamma_{k+1} = \Gamma'_k \cup \{ \kappa \approx \lambda \}$ if $\phi_k \in \Gamma'_k$ where $\phi_k = (\kappa \in \iota) \wedge \neg(\lambda \in \iota) \wedge | \lambda |_1$. Suppose Γ_{k+1} is inconsistent in this case. Then for some $\gamma_1, \dots, \gamma_n \in \Gamma'_k$, we have $\widehat{\gamma}, \kappa \in \iota, \lambda \notin \iota, | \lambda |_1, (\vec{p}^+)^{\kappa} = (\vec{q}^+)^{\lambda} \Vdash \Delta(\vec{p}^+)^{\kappa} = \Delta(\vec{q}^+)^{\lambda}$ for each Δ . By RAn, $\widehat{\gamma}, \kappa \in \iota, \lambda \notin \iota, | \lambda |_1 \Vdash (\kappa \in \iota) \equiv (\lambda \in \iota)$. Hence, Γ'_k is inconsistent, $\not\zeta$.

It suffices to show that the canonical hypermodel is analytic. Suppose $c_{\kappa} \in C_{\iota}$ and $c_{\kappa} \approx c_{\lambda}$. So $(\kappa \in \iota) \in \Gamma$ and $| \lambda |_1 \in \Gamma$. Moreover, if $(\lambda \in \iota) \notin \Gamma$, then by the revised Lindenbaum construction, $\kappa \approx \lambda \in \Gamma$, contrary to $c_{\kappa} \approx c_{\lambda}, \not\zeta$. Hence, $(\lambda \in \iota) \in \Gamma$.

S₅. We revise Definition A3.30 so that $c(\Delta)$ is always defined classically. The only revision needed to the proofs is to verify the connective case in the truth lemma (Lemma A3.34). This follows from the fact that $| \kappa |_1 \Vdash @_{\kappa} \Delta(\vec{\phi}) \equiv \Delta(@_{\kappa} \vec{\phi})$ is $(\mathbf{H} + \text{Bool}_{\Vdash})$ -derivable (by Bool_{\Vdash} , Gen_@, and Dist_@ (for \Vdash)).

B5.2. Theorem B2.3. For some of these proofs, we use the lemma below, which follows from Definition A4.42 and \exists -witnessing.

LEMMA B5.18 (Canonical proposition space). *Let $| \kappa |_1, | \lambda |_1 \in \Gamma$. Then $\pi_{c_{\kappa}} \subseteq \pi_{c_{\lambda}}$ iff $(\pi_{\kappa} \subseteq \pi_{\lambda}) \in \Gamma$, and $| \pi_{c_{\kappa}} | = 1$ iff $| \pi_{\kappa} |_1 \in \Gamma$.*

We omit the proofs for B, U_q, U_o, Si, and S₅, which are routine.

At. Let $c_{\kappa} \in D_{\text{H}\Gamma}$ and $\Delta \in W_{\Gamma}$. First, observe that $\phi \rightarrow \Box(@_{\kappa} p^+ \rightarrow \phi) \in \Delta$. For by Atom, Bool, and Dist_@, $\exists p(@_{\kappa} p \wedge \forall q(\Box(@_{\kappa} p \rightarrow @_{\iota_{\Delta}} q) \vee \Box(@_{\kappa} p \rightarrow \neg @_{\iota_{\Delta}} q))) \in \Delta$. Since $I_{\Delta} \in \Delta$, we have $\exists p(@_{\kappa} p \wedge \forall q(\Box(@_{\kappa} p \rightarrow q) \vee \Box(@_{\kappa} p \rightarrow \neg q))) \in \Delta$. By \exists -witnessing, $@_{\kappa} p^+ \wedge \forall q(\Box(@_{\kappa} p^+ \rightarrow q) \vee \Box(@_{\kappa} p^+ \rightarrow \neg q)) \in \Delta$ for some p^+ . By Elim_{\forall}, CLEx, and \exists -witnessing, $\Box(@_{\kappa} p^+ \rightarrow \phi) \vee \Box(@_{\kappa} p^+ \rightarrow \neg \phi) \in \Delta$. By S5, $\phi \rightarrow \Box(@_{\kappa} p^+ \rightarrow \phi) \in \Delta$.

So suppose $@_{\kappa} p^+ \in \Delta'$ and suppose $\phi \in \Delta$. Thus, $\Box(@_{\kappa} p^+ \rightarrow \phi) \in \Delta$. By Corollary A3.27, $\phi \in \Delta'$. Hence, $\Delta' = \Delta$. So $p^+ \in \{ \{ \Delta \} \}_{\kappa}$, i.e., $\{ \Delta \} \in \pi_{c_{\kappa}}$.

An. Since members of $\text{INom}^+ = \{ I_1^+, I_2^+, I_3^+, \dots \}$ might not be allowed to denote singletons (since conventions must be closed under \approx), the Henkin construction needs to be revised so that INom^+ is replaced with $\text{IVar}^+ = \{ i_1^+, i_2^+, i_3^+, \dots \}$ (though we don't allow formulas in $\mathcal{L}^{\text{QH}^+}$ to bind members of IVar^+). We also need to make the following amendments to the definition of the canonical model:

$$D_{C\Gamma} = \{ C_{\iota} \mid \neg | \iota |_1 \in \Gamma \} \cup \{ C_{\iota} \mid | \iota |_1 \in \Gamma \} \cup \{ \{ c_{\kappa} \} \mid c_{\kappa} \approx c_{\iota} \mid | \iota |_1 \in \Gamma \},$$

$$V_{\Gamma}(\iota) = \begin{cases} \{ c_{\iota} \}, & \text{if } | \iota |_1 \in \Gamma \text{ and } \iota \in \text{IVar} \cup \text{IVar}^+, \\ C_{\iota}, & \text{otherwise.} \end{cases}$$

The proof of the truth lemma (Lemma A3.34) remains intact (the only difference is the $@_i$ -case where ι is i and $|i|_1 \in \Gamma$; in that case, $\Delta, c_\kappa \Vdash @_i \phi$ iff $\Delta, c_i \Vdash \phi$ iff $@_i \phi \in \Delta$ iff $@_\kappa @_i \phi \in \Delta$.) Trivially, $\{c_\kappa \mid c \approx c_i\}$ is analytic. So we need to show that C_l is analytic if $\neg |l|_1 \in \Gamma$, and that C_l is analytic if $|l|_1 \in \Gamma$.

First, suppose $\neg |l|_1 \in \Gamma$. Let $c_\kappa \in C_l$ and let $c_\lambda \neq c_\kappa$ be such that $c_\kappa \approx c_\lambda$. Since $(\kappa \in \iota), |\lambda|_1 \in \Gamma$, it suffices to show that $(\kappa \approx \lambda) \in \Gamma$; for then by An, $(\lambda \in \iota) \in \Gamma$, and so $c_\lambda \in C_l$. By Lemma B5.18, $(\pi_\kappa = \pi_\lambda) \in \Gamma$ since $\pi_{c_\kappa} = \pi_{c_\lambda}$. Moreover, if $((p^+)^{\kappa} = (q^+)^{\kappa}) \in \Gamma$, then $\{\Delta \in W_\Gamma \mid @_\kappa p^+ \in \Delta\} = \{\Delta \in W_\Gamma \mid @_\lambda q^+ \in \Delta\}$. Since $c_\kappa \approx c_\lambda$, that means $c_\kappa(\star)(X) = c_\lambda(\star)(X)$. So by Lemma B5.17, $((\star p^+)^{\kappa} = (\star q^+)^{\lambda}) \in \Gamma$. Therefore, $((p^+)^{\kappa} = (q^+)^{\lambda}) \supset ((\star p^+)^{\kappa} = (\star q^+)^{\lambda}) \in \Gamma$. Since Γ witnesses $\exists s, \forall p \forall q ((p^\kappa = q^\lambda) \supset ((\star p)^\kappa = (\star q)^\lambda)) \in \Gamma$, i.e., $(\star_\kappa = \star_\lambda) \in \Gamma$. Similarly, $(\circ_\kappa = \circ_\lambda) \in \Gamma$. Hence, $(\kappa \approx \lambda) \in \Gamma$.

Next, suppose $|l|_1 \in \Gamma$. Let $c_\kappa \approx c_l$. By the reasoning above, $(\kappa \approx l) \in \Gamma$. Since $|\kappa|_1, |l|_1 \in \Gamma$, it follows by Many_{INom} that $(\kappa = l) \in \Gamma$. Hence, by Lemma A4.44, $c_\kappa = c_l$, and thus $c_\kappa \in C_l$.

B5.3. Theorem B2.6. We omit the proofs for Cr and Di, which are routine.

$C_{l\Phi}$. We revise the Henkin construction. Let $\text{Prop}^\circ = \{p_1^\circ, p_2^\circ, p_3^\circ, \dots\}$ be new propositional variables, and let $\mathcal{L}^{\text{QH}+\circ}$ be the result of expanding $\mathcal{L}^{\text{QH}+}$ with Prop° (again, not including formulas with quantifiers binding these variables). Enumerate the members of Φ as $\chi_1, \chi_2, \chi_3, \dots$. Let Δ be the set of all formulas of the form $p_k^\circ =_l \chi_k$, where $l \in \text{INom}^+$ and $\chi_k \in \Phi$. The Henkin construction is the same except we redefine Γ'_k so that $\Gamma'_k = \Gamma_k \cup \{\phi_k\}$ if $\Gamma_k, \Delta, \phi_k \not\vdash_{\text{QH} \cup \text{Ex}_\Phi} \perp$ (and $= \Gamma_k$ otherwise). Clearly, if $\Gamma_k \cup \Delta$ is $(\text{QH} \cup \text{Ex}_\Phi)$ -consistent, then so is $\Gamma'_k \cup \Delta$. The proof that $\Gamma_{k+1} \cup \Delta$ is consistent if $\Gamma'_k \cup \Delta$ is consistent is essentially the same. Thus, we just need to show that $\Gamma_1 \cup \Delta$ is $(\text{QH} \cup \text{Ex}_\Phi)$ -consistent. Suppose it's not. Since l_Γ occurs nowhere in Δ , we can eliminate l_Γ by the same reasoning as in Lemma A3.24. Thus, there are some $\alpha_1, \dots, \alpha_k$ that are instances of Ex_Φ , some $\delta_1, \dots, \delta_n \in \Delta$ where δ_i is of the form $q_i^\circ =_{k_i} \psi_i$ for some $\psi_i \in \Phi$ and $k_i \in \text{INom}^+$, and some $\gamma_1, \dots, \gamma_m \in \Gamma$ such that $\widehat{\alpha}, \widehat{\delta} \vdash \neg \widehat{\gamma}$ (throughout, I'll use \vdash for provability in QH and \vdash_{Ex_Φ} for provability in $\text{QH} \cup \text{Ex}_\Phi$). Now, it may be that $q_i^\circ = q_j^\circ$ for some i and j . So let $q_i^\circ \approx \psi_i$ be the conjunction all δ_j s such that $q_j^\circ = q_i^\circ$ —that is, $q_i^\circ \approx \psi_i$ has the form $(q_i^\circ =_{k_{i_1}} \psi_{i_1}) \wedge \dots \wedge (q_i^\circ =_{k_{i_j}} \psi_{i_j})$. (Given how Δ is defined and how Φ is enumerated, it is never the case that $q_i^\circ = q_j^\circ$ but $\psi_i \neq \psi_j$; so this definition is well-defined.) Thus, $\widehat{\alpha}, q_1^\circ \approx \psi_1, \dots, q_n^\circ \approx \psi_n \vdash \neg \widehat{\gamma}$. By Lemma A4.38, $\widehat{\alpha}, r_1 \approx \psi_1, \dots, r_n \approx \psi_n \vdash \neg \widehat{\gamma}$ where $r_1, \dots, r_n \in \text{Prop}$ are fresh. By RK_\exists , Vac_\exists , and VDist_\exists , $\widehat{\alpha}, \exists r_1 (r_1 \approx \psi_1), \dots, \exists r_n (r_n \approx \psi_n) \vdash \neg \widehat{\gamma}$. So by $\text{Ex}_\Phi, \vdash_{\text{Ex}_\Phi} \neg \widehat{\gamma}, \not\vdash$.

The rest of the proof of the Henkin lemma (Lemma A4.40) goes through as before. And since $\Gamma_k \cup \Delta$ is $(\text{QH} \cup \text{Ex}_\Phi)$ -consistent for each k , $\Gamma^+ \cup \Delta$ is $(\text{QH} \cup \text{Ex}_\Phi)$ -consistent, which by maximality means $\Delta \subseteq \Gamma^+$. Hence, Γ^+ has the following property: for each $\chi \in \Phi$, there is a p° such that for all $\iota \in \text{ITerm}^+$, $(p^\circ =_\iota \chi) \in \Gamma^+$.

To complete the proof, we revise the definition of π_{c_κ} (when $@_\kappa cl \notin \Gamma$) and $D_{\mathbb{P}\Gamma}$:

$$\pi_{c_\kappa} = \{X \mid \exists p \in \text{Prop}^+ \cup \text{Prop}^\circ : p \in [X]_\kappa\},$$

$$D_{\mathbb{P}\Gamma} = \{P \in \mathbb{P}_{D_{\mathbb{H}\Gamma}} \mid \exists p \in \text{Prop}^* \cup \text{Prop}^\circ \forall c_\kappa \in D_{\mathbb{H}\Gamma} : p \in [P(c_\kappa)]_\kappa\}.$$

The rest of the proof goes through as before. So by Lemma A4.47, $\llbracket \chi_i \rrbracket^{\mathcal{M}_{\Gamma, c_\kappa}} = \{\Delta \in W_\Gamma \mid @_\kappa \chi_i \in \Delta\} = \{\Delta \in W_\Gamma \mid @_\kappa p_i^\circ \in \Delta\}$, so $P_{p_i^\circ}$ can be our witness for $\chi_i \in \Phi$. Hence, $D_{\mathbb{P}\Gamma}$ is closed under Φ .

Cl_{Φ}^+ . The proof is roughly the same as Cl_{Φ} , but we need to make some revisions. Let $\Phi' = \{\chi[q'_1/q_1, \dots, q'_n/q_n] \mid q'_1, \dots, q'_n \in Prop^* \cup Prop^{\circ}\}$. Enumerate the members of Φ' as $\chi_1, \chi_2, \chi_3, \dots$ in such a way that p'_k never occurs in χ_1, \dots, χ_k . Proceed with the Henkin construction in the same manner as before, replacing Φ throughout with Φ' . To establish that $\Gamma_1 \cup \Delta$ is $(\mathbf{QH} + Ex_{\Phi})$ -consistent, we use the same reasoning, except the last step needs further justification, since ψ_i may not be in Φ but rather of the form $\psi_i = \chi[q'_1/q_1, \dots, q'_n/q_n]$ for some $\chi \in \Phi$. However, since Ex_{Φ} is now an axiom, that means if $\chi \in \Phi$, then $\vdash \exists r(r \approx \chi)$. So by $Gen_{\forall}, \vdash \forall_{q_1} \dots \forall_{q_n} \exists r(r \approx \chi)$. Hence, by $Elim_{\forall}, \vdash \exists r(r \approx \psi_i)$.

Making the same revisions as before, the rest of the completeness proof goes through. So we just need to show now that $D_{P_{\Gamma}}$ is strongly closed under Φ . Let $\mathcal{M} = \langle W_{\Gamma}, D_{C_{\Gamma}}, D_{P_{\Gamma}}, V \rangle$. Then $V(q_i) = P_{q'_i}$ for some q'_i . Hence, by Lemma A4.36, $\llbracket \phi \rrbracket^{\mathcal{M}} = \llbracket \phi[q'_1/q_1, \dots, q'_n/q_n] \rrbracket^{\mathcal{M}_{\Gamma}}$. By how Γ was constructed, there is a p° such that for all i , $p^{\circ} =_i \phi[q'_1/q_1, \dots, q'_n/q_n] \in \Gamma$. By Lemma A4.47, $\llbracket \phi[q'_1/q_1, \dots, q'_n/q_n] \rrbracket^{\mathcal{M}_{\Gamma}} = \llbracket p^{\circ} \rrbracket^{\mathcal{M}_{\Gamma}} \in D_{P_{\Gamma}}$. Hence, $D_{P_{\Gamma}}$ is strongly closed under Φ .

To establish that $\mathbf{QH} + Ex_{\Phi} = \mathbf{QH} + Elim_{\forall\Phi}$, it suffices to show that Ex_{Φ} is coderivable with $Elim_{\forall\Phi}$. Deriving Ex_{Φ} from $Elim_{\forall\Phi}$ is straightforward by S5 and $Dual_{\forall}$. For the other direction, it follows by induction (or completeness over the class of all models) that if χ is free for p , and i_1, \dots, i_n are the free interpretation terms in ϕ , then $p = \chi, p =_{i_1} \chi, \dots, p =_{i_n} \chi \Vdash \phi = \phi[\chi/p]$. Hence:

$\forall p\phi, p = \chi, \dots, p =_{i_n} \chi \Vdash \phi[\chi/p]$	above, $Elim_{\forall}$,
$\downarrow i. \forall p\phi, \downarrow i.(p =_i \chi \& \dots \& p =_{i_n} \chi) \Vdash \downarrow i. \phi[\chi/p]$	$Gen_{\downarrow}, Idle_{\downarrow}, Dist_{\downarrow}$,
$\forall p\phi, \downarrow i.(p =_i \chi \& \dots \& p =_{i_n} \chi) \Vdash \phi[\chi/p]$	Vac_{\downarrow} ,
$\forall p\phi, \exists p \downarrow i.(p =_i \chi \& \dots \& p =_{i_n} \chi) \Vdash \phi[\chi/p]$	$RK_{\exists}, VDist_{\exists}, Vac_{\exists}$,
$\forall p\phi, \downarrow i. \exists p(p =_i \chi \& \dots \& p =_{i_n} \chi) \Vdash \phi[\chi/p]$	BF_{\downarrow} ,
$\forall p\phi \Vdash \phi[\chi/p]$	$Ex_{\Phi}, Gen_{\downarrow}$.

Df_{Φ} . I will only prove weak completeness here; it's easy to check that if Φ is finite, then strong completeness can be established via the same method. Suppose ϕ is $(\mathbf{QH} + Gen_{\forall\Phi})$ -consistent. Enumerate the members of Φ as $\chi_1, \chi_2, \chi_3, \dots$. Parallel to $\Gamma_1, \Gamma_2, \Gamma_3, \dots$, we construct a new sequence of sets $\Delta_1, \Delta_2, \Delta_3, \dots$. First, $\Gamma_1 = \{\phi\} \cup \{l^+, |l^+|_1\}$ and $\Delta_1 = \emptyset$. Next, define Γ_{k+1} as in the proof of Lemma A4.40. Finally, define Δ_{k+1} as follows:

$$\Delta_{k+1} = \begin{cases} \Delta_k \cup \{q^+ =_{l^+} \chi \mid (q^+ =_{l^+} \chi) \in \Delta_k\}, & \text{if } (*) \text{ holds,} \\ \Delta_k \cup \{p^+ =_{l^+} \chi \mid l^+ \in INom^+ \text{ occurs in } \Gamma_{k+1}\}, & \text{if } (**) \text{ holds,} \\ \Delta_k, & \text{otherwise,} \end{cases}$$

- (*) $\phi_k = \neg @_i \psi$ and l^+ is the witness introduced in Γ_{k+1} ,
- (**) $\phi_k = \exists p\psi$, where p^+ is the witness introduced in Γ_{k+1} and χ is the first of Φ such that $\Gamma_{k+1}, \Delta_k, \{p^+ =_{l^+} \chi \mid l^+ \in INom^+ \text{ occurs in } \Gamma_{k+1}\} \not\vdash \perp$.

Finally, $\Gamma^+ = \bigcup_{k \geq 1} \Gamma_k$. We first show that for each k , $\Gamma_k \cup \Delta_k$ is $(\mathbf{QH} + Gen_{\forall\Phi})$ -consistent. Clearly this holds for $k = 1$. And clearly if $\Gamma_k \cup \Delta_k$ is $(\mathbf{QH} + Gen_{\forall\Phi})$ -consistent, then so is $\Gamma'_k \cup \Delta_k$ and $\Gamma_{k+1} \cup \Delta_k$. So we just need to show that if $\Gamma_{k+1} \cup \Delta_k$ is $(\mathbf{QH} + Gen_{\forall\Phi})$ -consistent, then so is $\Gamma_{k+1} \cup \Delta_{k+1}$. If $\phi_k = \exists p\psi$, then $\Gamma_{k+1} \cup \Delta_{k+1}$

is $(\mathbf{QH} + \text{Gen}_{\forall\Phi})$ -consistent by construction of Δ_{k+1} , assuming it's defined. Here's the proof that it is always defined, i.e., there always is such a χ in this case. Suppose otherwise. That means for all $\chi \in \Phi$, where $\gamma := \bigwedge \Gamma_k$, $\delta := \bigwedge \Delta_k$, and l_1^+, \dots, l_n^+ are the nominals occurring in some formula of Γ_{k+1} , $\gamma, \delta \vdash \neg(p^+ =_{l_1^+} \chi \wedge \dots \wedge p^+ =_{l_n^+} \chi)$. By $\text{Gen}_{\forall\Phi}$, $\gamma, \delta \vdash \forall p \neg(p^+ =_{l_1^+} p \wedge \dots \wedge p^+ =_{l_n^+} p)$. By Elim_{\forall} , $\gamma, \delta \vdash \neg(p^+ =_{l_1^+} p^+ \wedge \dots \wedge p^+ =_{l_n^+} p^+)$. Hence, by S5, $\gamma, \delta \vdash \perp$, ζ .

Now suppose $\phi_k = \neg @_{l^+} \psi$ and suppose for reductio that $\Gamma_{k+1} \cup \Delta_{k+1}$ is $(\mathbf{QH} + \text{Gen}_{\forall\Phi})$ -inconsistent. Then for some formula of the form $q_i^+ =_{l^+} \chi_i$, we have $\neg @_{l^+} \psi, l^+ \in l, \gamma, \delta, (q_1^+ =_{l^+} \chi_1), \dots, (q_n^+ =_{l^+} \chi_n) \vdash @_{l^+} \psi$. Repeating the reasoning in Lemma A3.24, $\gamma, \delta, \downarrow i. (q_1^+ =_{l^+} \chi_1), \dots, \downarrow i. (q_n^+ =_{l^+} \chi_n) \vdash @_{l^+} \psi$. Since $l_1^+, |l_1^+|_1 \in \Gamma_k$: $\gamma, \delta, (q_1^+ =_{l_1^+} \chi_1), \dots, (q_n^+ =_{l_1^+} \chi_n) \vdash @_{l^+} \psi$. But $(q_1^+ =_{l_1^+} \chi_1), \dots, (q_n^+ =_{l_1^+} \chi_n) \in \Delta_k$. Thus, $\gamma, \delta \vdash @_{l^+} \psi$. So $\Gamma'_k \cup \Delta_k$ is already $(\mathbf{QH} + \text{Gen}_{\forall\Phi})$ -inconsistent, ζ . Hence, $\Gamma_{k+1} \cup \Delta_{k+1}$ is $(\mathbf{QH} + \text{Gen}_{\forall\Phi})$ -consistent. Therefore, $\Gamma^+ \cup \bigcup_k \Delta_k$ is $(\mathbf{QH} + \text{Gen}_{\forall\Phi})$ -consistent, and so by maximality, $\Delta_k \subseteq \Gamma^+$ for all k .

By construction, for each $p^+ \in \text{Prop}^+$, there is a $\chi \in \Phi$ such that $(p^+ =_{l^+} \chi) \in \Gamma^+$ for all $l^+ \in \text{INom}^+$. From here, the completeness proof proceeds as before. To complete the proof, we show $D_{\mathbb{P}\Gamma}$ is definable in Φ . Where $P = P_{p^+} \in D_{\mathbb{P}\Gamma}$, let $\chi \in \Phi$ be such that $p^+ =_{l^+} \chi \in \Gamma^+$ for all $l^+ \in \text{INom}^+$. Then by Lemma A4.47, $P_{p^+}(c_\kappa) = \llbracket p^+ \rrbracket^{c_\kappa} = \llbracket \chi \rrbracket^{c_\kappa}$. Hence, $P_{p^+} = \llbracket \chi \rrbracket$.

$Cl_\Phi Df_\Phi$. We use the same construction as in Df_Φ . We need to show (i) that we can dispense with the $\text{Gen}_{\forall\Phi}$ rule in the proof above, and (ii) $D_{\mathbb{P}\Gamma}$ is closed under Φ . (To establish that $\mathbf{QH} \cup \text{Hom}_\Phi \cup \text{Ex}_\Phi = \mathbf{QH} \cup \text{Hom}_\Phi \cup \text{Ex}_\Phi^-$, simply observe that $\{\forall p((p = \chi) \supset (p =_{l^+} \chi)) \mid i \leq n\}, \text{E}\chi \Vdash \exists p \&_{i=1}^n (p =_{l^+} \chi)$.)

For (i), note that $\text{Gen}_{\forall\Phi}$ was only used to establish that in the Henkin construction, if $\phi_k = \exists p \psi$ is added to Γ'_k and p^+ is the witness introduced into Γ_{k+1} , then there is a $\chi \in \Phi$ such that $\Gamma_{k+1}, \Delta_k, \{p^+ =_{l^+} \chi \mid l^+ \in \text{INom}^+\}$ occurs in $\Gamma_{k+1} \not\vdash \perp$. For all $\chi \in \Phi$, there is an l^+ such that $(p^+ =_{l^+} \chi) \notin \Gamma_{k+1}$. Then for all $\chi \in \Phi$, there exist some $\gamma_1, \dots, \gamma_n \in \Gamma_{k+1}$, some $\delta_1, \dots, \delta_m \in \Delta_k$, some $\alpha_1, \dots, \alpha_k \in \text{Hom}_\Phi$, and some $\beta_1, \dots, \beta_j \in \text{Ex}_\Phi^-$ such that $\widehat{\alpha}, \widehat{\beta}, \widehat{\gamma}, \widehat{\delta}, p^+ =_{l_1^+} \chi, \dots, p^+ =_{l_n^+} \chi \vdash \perp$. Since $\forall p(p = \chi \supset p =_{l^+} \chi) \in \text{Hom}_\Phi$ for each l_i^+ , we can assume these are included in $\widehat{\alpha}$. Hence, by Elim_{\forall} , $\widehat{\alpha}, \widehat{\beta}, \widehat{\gamma}, \widehat{\delta}, p^+ = \chi \vdash \perp$. So by Lemma A4.38, where r is fresh, $\widehat{\alpha}, \widehat{\beta}, \widehat{\gamma}, \widehat{\delta}, r = \chi \vdash \perp$. By Intro_{\exists} , VDist_{\exists} and Vac_{\exists} , $\widehat{\alpha}, \widehat{\beta}, \widehat{\gamma}, \widehat{\delta}, \text{E}\chi \vdash \perp$. So, $\widehat{\gamma}, \widehat{\delta} \vdash_{\text{Hom}_\Phi \cup \text{Ex}_\Phi^-} \perp$, ζ . (Notice we did not rely on Γ_{k+1} being finite, so the same strategy establishes strong completeness.) For (ii), let $\chi \in \Phi$. By Ex_Φ , $(p^+ = \chi) \in \Gamma$ for some $p^+ \in \text{Prop}^+$. By Hom_Φ and Elim_{\forall} , $(p^+ =_{l^+} \chi) \in \Gamma$ for all $l^+ \in \text{ITerm}^+$. Hence, $p^+ \in \llbracket \chi \rrbracket^{c_\kappa}$ for all c_κ , i.e., $\llbracket \chi \rrbracket \in D_{\mathbb{P}\Gamma}$.

$Cl_\Phi^+ Df_\Phi$. Similar to $Cl_\Phi Df_\Phi$, except using Ex_Φ as an axiom to show that $D_{\mathbb{P}\Gamma}$ is strongly closed (as in the proof of completeness over Cl_Φ^+).

Cb. Let $X_1 \in \pi_{c_{\kappa_1}}, \dots, X_n \in \pi_{c_{\kappa_n}}$ where $c_{\kappa_1}, \dots, c_{\kappa_n}$ are distinct. Let $p_1^+, \dots, p_n^+ \in \text{Prop}^+$ be such that $p_i^+ \in [X_i]_{\kappa_i}$. Since $c_{\kappa_1}, \dots, c_{\kappa_n}$ are distinct, by the same reasoning as in Di, $(\kappa_i \neq \kappa_j) \in \Gamma$ for $i \neq j$. By Split and Bool , $\exists p \bigwedge_{i=1}^n (p =_{\kappa_i} p_i^+) \in \Gamma$. By witnessing \exists s, there is a $p^+ \in \text{Prop}^+$ such that $p^+ =_{\kappa_i} p_i^+ \in \Gamma$ for $1 \leq i \leq n$. Hence, $P_{p^+}(c_{\kappa_i}) = X_i$.

To establish that $\mathbf{QH} + \mathbf{PII}^+ + \mathbf{Split} = \mathbf{QH} + \mathbf{PII}_1^+ + \mathbf{Split}$, we just need to show that \mathbf{PII}^+ is $(\mathbf{QH} + \mathbf{PII}_1^+ + \mathbf{Split})$ -derivable. By \mathbf{PII}_1^+ , it suffices to show that $\forall p(p' = p^\kappa), |i|_1, |\kappa|_1, (i \neq \kappa) \Vdash |\pi_i|_1$ is derivable using \mathbf{Split} :

$\forall p(p' = p^\kappa) \Vdash p' = p^\kappa$	Elim $_{\forall}$,
$\forall p(p' = p^\kappa) \Vdash r' = r^\kappa$	Elim $_{\forall}$,
$\forall p(p' = p^\kappa), p' = q', p^\kappa = r^\kappa \Vdash q' = r'$	S5,
$\forall p(p' = p^\kappa), \exists p(p' = q' \& p^\kappa = r^\kappa) \Vdash q' = r'$	RK $_{\exists}$, VDist $_{\exists}$, Vac $_{\exists}$,
$\forall p(p' = p^\kappa), i _1, \kappa _1, (i \neq \kappa) \Vdash q' = r'$	Split,
$\forall p(p' = p^\kappa), i _1, \kappa _1, (i \neq \kappa) \Vdash \pi_i _1$	RK $_{\forall}$, Vac $_{\forall}$, Dist $_{@}$.

CpSi. We must revise the definition of the canonical model so that $D_{\mathbb{P}\Gamma} = \mathbb{P}D_{\mathbb{H}\Gamma}$. The only thing that needs to be redone is the \forall inductive step of Lemma A4.47. The argument that if $\mathcal{M}_\Gamma, \Delta, c_\kappa \Vdash \forall p\phi$, then $@_\kappa \forall p\phi \in \Delta$ is the same. For the other direction, we first need the following intermediate result:

CLAIM. For all ϕ and all $\lambda \in \text{ITerm}^+$ such that $|\lambda|_1 \in \Gamma$, there is a formula ϕ^\uparrow such that where \vec{p} are the free propositional variables in ϕ :

- i. ϕ^\uparrow contains no interpretation binders $\downarrow i$.
- ii. If i and κ occur in ϕ^\uparrow and i isn't κ , then $(i \neq \kappa) \in \Gamma$.
- iii. For all $\Delta \in W_\Gamma, \mathcal{M}_\Gamma, \Delta, c_\lambda \Vdash \forall \vec{p}(\phi = \phi^\uparrow)$.
- iv. For all $\Delta \in W_\Gamma, \forall \vec{p}(\phi = \phi^\uparrow) \in \Delta$.

Proof. First, since Γ witnesses $\neg@_s$, for each free i , there is an $l_i^+ \in \text{INom}^+$ such that $(l_i^+ \in i) \in \Gamma$. By Sing and Intro $_{=}$, $(l_i^+ = i) \in \Gamma$. Let l_i^+ be the first in INom^+ with this property. By SubId, we can replace each i that occurs free in ϕ with l_i^+ . Call the result ϕ' . Now proceed as follows:

- a. If $\downarrow i$ does not occur in the scope of any $@_\kappa$ or any $\downarrow j$, replace each free i in its scope with l_i^+ . Then delete this $\downarrow i$.
- b. Repeat (a) on the result until there are no more $\downarrow i$ s that do not occur in the scope of any $@_\kappa$ or any $\downarrow j$.
- c. For each subformula of the form $@_{l^+} \psi$ that does not occur in the scope of any $@_\kappa$ operator, repeat (a) and (b) on ψ except with l^+ in place of l_i^+ . Continue until there are no more binders $\downarrow i$ left. Call the result ϕ^\uparrow .

It is now easy to verify that ϕ^\uparrow satisfies (i)–(iv). □

So suppose $\mathcal{M}_\Gamma, \Delta, c_\kappa \not\Vdash \forall p\phi$. By the claim above, $\mathcal{M}_\Gamma, \Delta, c_\kappa \not\Vdash \forall p\phi^\uparrow$. Thus, there is a $P \in \mathbb{P}D_{\mathbb{H}\Gamma}$ such that $(\mathcal{M}_\Gamma)_P, \Delta, c_\kappa \not\Vdash \phi^\uparrow$. Let l_1^+, \dots, l_n^+ be the interpretation terms in ϕ^\uparrow . By CIE x and $\Vdash \text{Ep}$ (by Intro $_{\exists}$), for each l_i^+ , there is a p_i^+ such that $p_i^+ \in [P(c_{l_i^+})]_{l_i^+}$, i.e., $\Delta' \in P(c_{l_i^+})$ iff $@_i p_i^+ \in \Delta'$. Since $(l_i^+ \neq l_j^+) \in \Gamma$ when $i \neq j$, it follows by Split that $\exists p \&_{i=1}^n (p =_{l_i^+} p_i^+) \in \Gamma$. By witnessing $\exists s$, there is a p^+ such that $\&_{i=1}^n (p^+ =_{l_i^+} p_i^+) \in \Gamma$. Thus, for each i and Δ' : $@_{l_i^+} p^+ \in \Delta'$ iff $@_{l_i^+} p_i^+ \in \Delta'$. Hence, $\Delta' \in P(c_{l_i^+})$ iff $@_i p^+ \in \Delta'$. By Lemma A4.36, $(\mathcal{M}_\Gamma)_P, \Delta, c_\kappa \Vdash \phi^\uparrow$ iff $\mathcal{M}_\Gamma, \Delta, c_\kappa \Vdash \phi^\uparrow [p^+ / p]$. Hence, $\mathcal{M}_\Gamma, \Delta, c_\kappa \not\Vdash \phi^\uparrow [p^+ / p]$. By IH, $@_\kappa \phi^\uparrow [p^+ / p] \notin \Delta$. By Elim $_{\forall}$, $\forall p @_\kappa \phi^\uparrow \notin \Delta$. By the claim above, $\forall p @_\kappa \phi \notin \Delta$. By CBF $_{@}$, $@_\kappa \forall p\phi \notin \Delta$.

B5.4. Theorem B3.16. In each case, it suffices to show that f_Γ satisfies the corresponding constraint given the axiom. Moreover, f_Γ is already defined to satisfy the relevant constraint when $[A] = \emptyset$. So assume throughout that $[A] \neq \emptyset$.

Suc. Suppose $\langle \Delta', c_\lambda \rangle \in f_\Gamma(A, \Delta, c_\kappa)$. Let $\phi \in [A]$. By Lemma B3.12 and Definition B3.13, if $(\phi \Box \rightarrow_\kappa (\lambda \supset \psi)) \in \Delta$ where $\psi \in \mathcal{L}_{\Box \rightarrow}^{H+}$, then $@_\lambda \psi \in \Delta'$. By $\text{Id}_{\Box \rightarrow}$ and $\text{RK}_{\Box \rightarrow \kappa}$, $(\phi \Box \rightarrow_\kappa (\lambda \supset \phi)) \in \Delta$. So $@_\lambda \phi \in \Delta'$. By Definition B3.11, $\langle \Delta', c_\lambda \rangle \in A$.

W. Let $\langle \Delta, c_\kappa \rangle \in A$ and $\phi \in [A]$ (so $@_\kappa \phi \in \Delta$). Suppose $\phi \Box \rightarrow_\kappa (\kappa \supset \psi) \in \Delta$. By $\text{MP}_{\Box \rightarrow}$ and Ded , $\phi, \phi \Box \rightarrow_\kappa (\kappa \supset \psi), \kappa \Vdash \psi$. By $\text{Gen}_{@}$ and Ref , $@_\kappa \phi, \phi \Box \rightarrow_\kappa (\kappa \supset \psi) \Vdash @_\kappa \psi$. Hence, $@_\kappa \psi \in \Delta$. By Definition B3.13, $\langle \Delta, c_\kappa \rangle \in f_\Gamma(A, \Delta, c_\kappa)$.

C. Suppose $\langle \Delta, c_\kappa \rangle \in A$. Let $\phi \in [A]$. Thus, $@_\kappa \phi \in \Delta$. Reasoning as above, we have $@_\kappa \phi \Vdash (\phi \Box \rightarrow_\kappa (\kappa \supset \psi)) \equiv @_\kappa \psi$. So if $(\phi \Box \rightarrow_\kappa (\kappa \supset \psi)) \in \Delta$, then $@_\kappa \psi \in \Delta$, meaning $\langle \Delta, c_\kappa \rangle \in f_\Gamma(A, \Delta, c_\kappa)$. Moreover, let $\langle \Delta', c_\lambda \rangle \in f_\Gamma(A, \Delta, c_\kappa)$. So for all $\psi \in \mathcal{L}_{\Box \rightarrow}^{H+}$, if $(\phi \Box \rightarrow_\kappa (\lambda \supset \psi)) \in \Delta$, then $@_\lambda \psi \in \Delta'$. Now, by Cen , $i, \phi \Vdash (\phi \Box \rightarrow i)$. By Gen_{\downarrow} and Vac_{\downarrow} , $\phi \Vdash \downarrow i.(\phi \Box \rightarrow i)$. By $\text{Gen}_{@}$ and $\text{DA}_{@}$, $|\kappa|_1, @_\kappa \phi \Vdash @_\kappa (\phi \Box \rightarrow \kappa)$. Since $|\kappa|_1 \in \Delta$, that means $(\phi \Box \rightarrow_\kappa \kappa) \in \Delta$. By $\text{RK}_{\Box \rightarrow \kappa}$, $(\phi \Box \rightarrow_\kappa (\lambda \supset \kappa)) \in \Delta$. So $@_\lambda \kappa \in \Delta'$. By Rigid and Corollary A3.27 , $|\kappa|_1 \in \Delta'$. By $\text{Intro}_{=}$, $(\kappa = \lambda) \in \Delta'$. By Lemma A3.31, $c_\kappa = c_\lambda$. We now show $\Delta' = \Delta$. We'll just show $\Delta \subseteq \Delta'$ to illustrate. Let $\psi \in \Delta$. By $\text{Intro}_{@}$, $\text{Elim}_{@}$, and Red , $@_\kappa @_{i_\Delta} \psi \in \Delta$. Thus, $\phi \Box \rightarrow_\kappa (\kappa \supset @_{i_\Delta} \psi) \in \Delta$. So $@_\kappa @_{i_\Delta} \psi \in \Delta'$ since $\langle \Delta', c_\kappa \rangle \in f_\Gamma(A, \Delta, c_\kappa)$. So by Red , Rigid , $\text{Intro}_{@}$, and $\text{Elim}_{@}$, $\psi \in \Delta'$.

Stal. Suppose $\langle \Delta_1, c_\lambda \rangle, \langle \Delta_2, c_\mu \rangle \in f(A, \Delta, c_\kappa)$. Let $\phi \in [A]$. Thus, for all $\psi \in \mathcal{L}_{\Box \rightarrow}^{H+}$:

$$\begin{aligned} (\phi \Box \rightarrow_\kappa (\lambda \supset \psi)) \in \Delta &\Rightarrow @_\lambda \psi \in \Delta_1, \\ (\phi \Box \rightarrow_\kappa (\mu \supset \psi)) \in \Delta &\Rightarrow @_\mu \psi \in \Delta_2. \end{aligned}$$

Suppose $(\phi \Box \rightarrow_\kappa \sim \lambda) \in \Delta$. Thus, $(\phi \Box \rightarrow_\kappa (\lambda \supset \sim \lambda)) \in \Delta$ by $\text{RK}_{\Box \rightarrow \kappa}$, and so, $@_\lambda \sim \lambda \in \Delta_1, \not\in$. Hence, $(\phi \Box \rightarrow_\kappa \sim \lambda) \notin \Delta$. By CEM , $(\phi \Box \rightarrow_\kappa \lambda) \in \Delta$. By $\text{RK}_{\Box \rightarrow \kappa}$, $(\phi \Box \rightarrow_\kappa (\mu \supset \lambda)) \in \Delta$, and so, $@_\mu \lambda \in \Delta_2$. By Rigid and $\text{Intro}_{=}$, $(\lambda = \mu) \in \Gamma$ since $|\lambda|_1, |\mu|_1 \in \Gamma$. By Lemma A3.31 then, $c_\lambda = c_\mu$. Further, $(\lambda = \mu) \in \Delta_1 \cap \Delta \cap \Delta_2$ by Rigid .

We now show that $\Delta_1 \subseteq \Delta_2$ (the proof that $\Delta_2 \subseteq \Delta_1$ is symmetric). Suppose $\psi \in \Delta_1$. By $\text{Intro}_{@}$ and $\text{Elim}_{@}$, $@_{i_\Delta} \neg \psi \notin \Delta_1$. By Red , $@_\lambda @_{i_\Delta} \neg \psi \notin \Delta_1$. Since $\langle \Delta_1, c_\lambda \rangle \in f_\Gamma(A, \Delta, c_\kappa)$, $(\phi \Box \rightarrow_\kappa (\lambda \supset @_{i_\Delta} \neg \psi)) \notin \Delta$. By SubId , since $(\lambda = \mu) \in \Delta$, $(\phi \Box \rightarrow_\kappa (\mu \supset @_{i_\Delta} \neg \psi)) \notin \Delta$. By CEM , $(\phi \Box \rightarrow_\kappa \sim (\mu \supset @_{i_\Delta} \neg \psi)) \in \Delta$. Since $(\phi \Box \rightarrow_\kappa \mu) \in \Delta$, we have $(\phi \Box \rightarrow_\kappa (\mu \supset \sim @_{i_\Delta} \neg \psi)) \in \Delta$ by $\text{RK}_{\Box \rightarrow \kappa}$. Since $\langle \Delta_2, c_\mu \rangle \in f_\Gamma(A, \Delta, c_\kappa)$, we have $\neg @_{i_\Delta} \neg \psi \in \Delta_2$ by Bool . So by $\text{Dist}_{@}$, $\text{Intro}_{@}$, and $\text{Elim}_{@}$, $\psi \in \Delta_2$.

Vac. Let $A(c_\kappa) = \emptyset$. Suppose for reductio $f_\Gamma(A, \Delta, c_\kappa) \neq \emptyset$. Let $\langle \Delta', c_\lambda \rangle \in f_\Gamma(A, \Delta, c_\kappa)$ and let $\phi \in [A]$. By Corollary A3.27 and $\text{Dist}_{@}$, $@_\kappa \sim \diamond \phi \in \Delta$. By Vac and $\text{Gen}_{@}$, $\phi \Box \rightarrow_\kappa (\lambda \supset \perp) \in \Delta$. By Definition B3.13, $@_\lambda \perp \in \Delta', \not\in$.

NC, NEC, SIC. We just do NC, since NEC and SIC are similar. It's left as an exercise to the reader to show that the two versions of the relevant axiom are coderivable. Let $\phi \in [A]$ and let $\langle \Delta', c_\lambda \rangle \in f_\Gamma(A, \Delta, c_\kappa)$. So for all ψ , if $(\phi \Box \rightarrow_\kappa (\lambda \supset \psi)) \in \Delta$, then $@_\lambda \psi \in \Delta'$. By Rigid and Bool , $@_\kappa \blacksquare \kappa \in \Delta$. By NC and $\text{RK}_{\Box \rightarrow \kappa}$, $\phi \Box \rightarrow_\kappa (\lambda \supset \kappa) \in \Delta$. Hence, $@_\lambda \kappa \in \Delta'$. By Rigid , $\text{Intro}_{@}$, and $\text{Elim}_{@}$, $(\kappa = \lambda) \in \Delta'$. Thus, $c_\lambda = c_\kappa$.

R_o . We revise the definition of a canonical hyperconvention as we did for Theorem B2.2 so that $\pi_{c_\kappa} = \{X \mid [X]_\kappa \neq \emptyset\}$. Given this, let $\langle \Delta', c_\lambda \rangle \in f(A, \Delta, c_\kappa)$ and let $\alpha \in [A]$. Thus, for all χ , if $\alpha \Box \rightarrow \kappa \chi \in \Delta$, then $@_\lambda \chi \in \Delta'$. We will just show the \neg -case, i.e., that $\neg_{c_\kappa} = \neg_{c_\lambda}$, since the others are similar.

First, observe that $\pi_{c_\kappa} = \pi_{c_\lambda}$, since, e.g., if $\phi \in [X]_\lambda$, then $@_\lambda \phi \in [X]_\kappa$ by Red. So let $X \in \pi_{c_\lambda}$ and let $\phi \in [X]_\lambda$. By R_o and $\text{Gen}_@$, $@_\kappa \downarrow i. \Box_\alpha \downarrow j. (\neg \phi = @_i \neg @_j \phi) \in \Delta$. By $\text{DA}_@$, $@_\kappa \Box_\alpha \downarrow j. (\neg \phi = @_\kappa \neg @_j \phi) \in \Delta$. Hence, $@_\lambda \downarrow j. (\neg \phi = @_\kappa \neg @_j \phi) \in \Delta'$. By $\text{DA}_@$, $@_\lambda (\neg \phi = @_\kappa \neg @_j \phi) \in \Delta'$. By $\text{Dist}_@$ and Red, $(@_\lambda \neg \phi = @_\kappa \neg @_\lambda \phi) \in \Delta'$. By Lemma B5.17 (since $@_\lambda \phi \in [X]_\kappa$), $\neg_{c_\kappa} X = \neg_{c_\lambda} X$.

Acknowledgement. I am grateful to two anonymous referees for their helpful feedback.

BIBLIOGRAPHY

[1] Alechina, N., Logan, B., & Whitsey, M. (2004). A complete and decidable logic for resource-bounded agents. In Jennings, N. R., Sierra, C., Sonenberg, L., and Tambe, M., editors. *Proceedings of the Third International Joint Conference on Autonomous Agents and Multi-Agent Systems (AAMAS 2004)*. New York: ACM Press, pp. 606–613.

[2] Baltag, A. & Smets, S. (2006). Conditional doxastic models: A qualitative approach to dynamic belief revision. *Electronic Notes in Theoretical Computer Science*, **165**, 5–21.

[3] Bennett, J. F. (2003). *A Philosophical Guide to Conditionals*. Oxford: Oxford University Press.

[4] Bernstein, S. (2016). Omission impossible. *Philosophical Studies*, **173**(10), 2575–2589.

[5] Berto, F. (2010). Impossible worlds and propositions: Against the parity thesis. *The Philosophical Quarterly*, **60**(240), 471–486.

[6] Berto, F., French, R., Priest, G., & Ripley, D. (2018). Williamson on counterpossibles. *Journal of Philosophical Logic*, **47**, 693–713.

[7] Berto, F. & Nolan, D. (2021). Hyperintensionality. In Zalta, E. N., editor. *The Stanford Encyclopedia of Philosophy*. Palo Alto, CA: Metaphysics Research Lab, Stanford University. Available from: <https://plato.stanford.edu/entries/hyperintensionality/>.

[8] Bjerring, J. C. (2013). Impossible worlds and logical omniscience: An impossibility result. *Synthese*, **190**, 2505–2524.

[9] Bjerring, J. C. & Schwarz, W. (2017). Granularity problems. *The Philosophical Quarterly*, **67**(266), 22–37.

[10] Bjerring, J. C. & Skipper, M. (2019). A dynamic solution to the problem of logical omniscience. *Journal of Philosophical Logic*, **48**, 501–521.

[11] Boutilier, C. E. (1992). Conditional Logics for Default Reasoning and Belief Revision. Ph.D. Thesis, University of Toronto.

[12] Brogaard, B. & Salerno, J. (2013). Remarks on counterpossibles. *Synthese*, **190**(4), 639–660.

[13] Clarke-Doane, J. (2019). Modal objectivity. *Noûs*, **53**(2), 266–295.

[14] Cohen, D. H. (1987). The problem of counterpossibles. *Notre Dame Journal of Formal Logic*, **29**(1), 91–101.

[15] ———. (1990). On what cannot be. In Jon Michael Dunn and Anil Gupta editors, *Truth or Consequences*. Dordrecht: Springer, pp. 123–132.

[16] Duc, H. N. (1997). Reasoning about rational, but not logically omniscient, agents. *Journal of Logic and Computation*, **7**(5), 633–648.

[17] Elga, A. & Rayo, A. (2021). Fragmentation and logical omniscience. *Noûs*, **56**, 716–741.

[18] Emery, N. & Hill, C. S. (2017). Impossible worlds and metaphysical explanation: Comments on Kment's modality and explanatory reasoning. *Analysis*, **77**(1), 134–148.

[19] Fine, K. (1970). Propositional quantifiers in modal logic. *Theoria*, **36**(3), 336–346.

[20] French, R., Girard, P., & Ripley, D. (2020). Classical counterpossibles. *Review of Symbolic Logic*, **15**, 259–275.

[21] Friedman, N. & Halpern, J. Y. (1997). Modeling belief in dynamic systems, part I: Foundations. *Artificial Intelligence*, **95**, 257–316.

[22] Goodman, J. (2004). An extended Lewis/Stalnaker semantics and the new problem of counterpossibles. *Philosophical Papers*, **33**(1), 35–66.

[23] Hawke, P., Özgün, A., & Berto, F. (2019). The fundamental problem of logical omniscience. *Journal of Philosophical Logic*, **49**, 727–766.

[24] Hintikka, J. (1975). Impossible possible worlds vindicated. *Journal of Philosophical Logic*, **4**, 475–484.

[25] Hoek, D. (2022). Questions in action. *Journal of Philosophy*, **119**, 113–143.

[26] Jago, M. (2007). Hintikka and Cresswell on logical omniscience. *Logic and Logical Philosophy*, **15**, 325–354.

[27] ———. (2014). *The Impossible*. Oxford: Oxford University Press.

[28] ———. (2015). Hyperintensional propositions. *Synthese*, **192**, 585–601.

[29] Jenny, M. (2018). Counterpossibles in science: The case of relative computability. *Noûs*, **52**(3), 530–560.

[30] Kim, S. & Maslen, C. (2006). Counterfactuals as short stories. *Philosophical Studies*, **129**(1), 81–117.

[31] Kment, B. (2014). *Modality and Explanatory Reasoning*. Oxford: Oxford University Press.

[32] Kocurek, A. W. (2021a). Counterpossibles. *Philosophy Compass*, **16**(11), e12787.

[33] ———. (2021b). Logic talk. *Synthese*, **199**(5–6), 13661–13688.

[34] ———. (2022). The logic of hyperlogic. Part A: Foundations. *Review of Symbolic Logic*, 1–28. DOI: [10.1017/S1755020322000193](https://doi.org/10.1017/S1755020322000193)

[35] Kocurek, A. W. & Jerzak, E. J. (2021). Counterlogicals as counterconventionals. *Journal of Philosophical Logic*, **50**(4), 673–704.

[36] Krakauer, B. (2012). Counterpossibles. Ph.D. Thesis, University of Massachusetts, Amherst.

[37] Kratzer, A. (1979). Conditional necessity and possibility. In Bäuerle, R., Egli, U., and von Stechow, A., editors. *Semantics from Different Points of View*. Berlin–Heidelberg–New York: Springer, pp. 117–147.

[38] Lamarre, P. & Shoham, Y. (1994). Knowledge, certainty, belief and conditionalisation. In Doyle, J., Sandewall, E., and Torasso, P., editors. *Principles of Knowledge Representation and Reasoning: Proceedings 4th International Conference (KR'94)*. San Francisco: Elsevier, pp. 415–424.

- [39] Lewis, D. K. (1973). *Counterfactuals*. Cambridge: Harvard University Press.
- [40] Mares, E. D. (1997). Who's afraid of impossible worlds? *Notre Dame Journal of Formal Logic*, **38**(4), 516–526.
- [41] Merricks, T. (2001). *Objects and Persons*. Oxford: Oxford University Press.
- [42] Moses, Y. & Shoham, Y. (1993). Belief as defeasible knowledge. *Artificial Intelligence*, **64**, 299–321.
- [43] Nolan, D. (1997). Impossible worlds: A modest approach. *Notre Dame Journal of Formal Logic*, **38**(4), 535–572.
- [44] Ripley, D. (2012). Structures and circumstances: Two ways to fine-grain propositions. *Synthese*, **189**, 97–118.
- [45] Sedlár, I. (2015). Substructural epistemic logics. *Journal of Applied Non-Classical Logics*, **25**, 256–285.
- [46] Skipper, M. & Bjerring, J. C. (2020). Hyperintensional semantics: A Fregean approach. *Synthese*, **197**, 3535–3558.
- [47] Soysal, Z. (2022). A metalinguistic and computational approach to the problem of mathematical omniscience. *Philosophy and Phenomenological Research*, 1–20. DOI: [10.1111/phpr.12864](https://doi.org/10.1111/phpr.12864)
- [48] Stalnaker, R. C. (1968). A theory of conditionals. In Rescher, N., editor. *Studies in Logical Theory*. American Philosophical Quarterly Monographs, Vol. 2. Oxford: Basil Blackwell, pp. 98–112.
- [49] ———. (1976a). Possible worlds. *Noûs*, **10**(1), 65–75.
- [50] ———. (1976b). Propositions. In MacKay, A. F. and Merrill, D. D., editors. *Issues in Philosophy of Language*. New Haven: Yale University Press, pp. 79–91.
- [51] ———. (1984). *Inquiry*. Cambridge: MIT Press.
- [52] ———. (1996). Impossibilities. *Philosophical Topics*, **24**(1), 193–204.
- [53] Tan, P. (2019). Counterpossible non-vacuity in scientific practice. *Journal of Philosophy*, **116**(1), 32–60.
- [54] van Benthem, J. (2007). Dynamic logic for belief revision. *Journal of Applied Non-Classical Logics*, **17**(2), 129–155.
- [55] van Ditmarsch, H. P. (2005). Prolegomena to dynamic logic for belief revision. *Synthese*, **147**, 229–275.
- [56] Vander Laan, D. A. (2004). Counterpossibles and similarity. In Jackson, F. and Priest, G., editors. *Lewisian Themes*. Oxford: Oxford University Press, pp. 258–275.
- [57] Williamson, T. (2007). *The Philosophy of Philosophy*. Malden, MA: Blackwell Publishers.
- [58] ———. (2017) Counterpossibles in semantics and metaphysics. *Argumenta*, **2**(2), 195–226.
- [59] Yalcin, S. (2018). Belief as question-sensitive. *Philosophy and Phenomenological Research*, **97**(1), 23–47.
- [60] Zagzebski, L. T. (1990). What if the impossible had been actual? In Beatty, M., editor. *Christian Theism and the Problems of Philosophy*. Notre Dame: University of Notre Dame Press, pp. 165–183.

SAGE SCHOOL OF PHILOSOPHY
 CORNELL UNIVERSITY
 ITHACA, NY 14850, USA
 E-mail: awk78@cornell.edu