





# A parametrization of sheets of conjugacy classes in bad characteristic

Filippo Ambrosio , Giovanna Carnovale , and Francesco Esposito

*Abstract.* Let  $G$  be a simple algebraic group of adjoint type over an algebraically closed field  $k$  of bad characteristic. We show that its sheets of conjugacy classes are parametrized by  $G$ -conjugacy classes of pairs  $(M, \mathcal{O})$  where  $M$  is the identity component of the centralizer of a semisimple element in  $G$  and  $\mathcal{O}$  is a rigid unipotent conjugacy class in  $M$ , in analogy with the good characteristic case.

## 1 Introduction

Sheets in a reductive algebraic group  $G$  are the irreducible components of the locally closed subsets of  $G$  consisting of conjugacy classes of the same dimension. They occur also as irreducible components of the strata in the partition of  $G$ , defined in [10] in terms of Springer representations with trivial local system (see [2, 3]). One of the most fascinating features of strata is that they are parametrized by a family of irreducible representations of the Weyl group which depends on the root system of  $G$  and not on the characteristic of the base field. It is therefore of interest to figure out the behavior of the irreducible components of strata when the characteristic of the base field varies.

A description of sheets in good characteristic, and a parametrization of sheets in terms of  $G$ -conjugacy classes of triples  $(M, Z(M)^\circ, \mathcal{O})$  where  $M$  is the identity component of the centralizer of a semisimple element in  $G$ ,  $Z(M)^\circ$  is a suitable coset in the component group  $Z(M)/Z(M)^\circ$ , and  $\mathcal{O}$  is a rigid unipotent conjugacy class in  $M$  was given in [4] in good characteristic, and extended to the case of bad characteristic in [14]. A refinement of this parametrization in terms of pairs  $(M, \mathcal{O})$  where  $M$  and  $\mathcal{O}$  are as above was given in [3] under the assumption that  $G$  is simple of adjoint type and the characteristic of the base field is good for  $G$ . The present paper answers a question by G. Lusztig on the extension to arbitrary characteristic of this parametrization of sheets.

Observe that, even if the formulation of the statement is the same, the collection of possible centralizers of a semisimple element in  $G$  varies with the characteristic of the base field, as well as the collection of unipotent conjugacy classes. Centralizers of semisimple elements are fewer in bad characteristic than in good characteristic

---

Received by the editors September 20, 2022; accepted January 12, 2023.

Published online on Cambridge Core January 26, 2023.

The authors acknowledge support by: DOR2207212/22 “Algebra di Nichols, algebra di Hopf e gruppi algebrici” and BIRD203834 “Grassmannians, flag varieties and their generalizations” funded by the University of Padova. They are members of the INdAM group GNSAGA.

AMS subject classification: 20G15, 20E45.

Keywords: Reductive algebraic groups, conjugacy classes, sheets, bad characteristic.



(see [5–8]), whereas the number of unipotent conjugacy classes may increase when passing from good to bad characteristic.

We therefore elaborate upon results in [5, 6, 8] in the spirit of [13] in order to provide a combinatorial description of the root systems of connected centralizers of semisimple elements, which will allow us to retrieve most ingredients that were necessary for the proof of [3, Theorem 4.1]. As an example, we show that the number of sheets in type  $G_2$  is the same in good characteristic and in characteristic 3, but it is smaller in characteristic 2, showing a difference in the behavior of sheets as opposed to strata.

## 2 Notation

Let  $G$  be a connected reductive algebraic group defined over an algebraically closed field  $k$  of characteristic exponent  $p$ . Let  $\Phi$  be the root system of  $G$ , and let  $\Delta = \{\alpha_1, \dots, \alpha_n\}$  be a fixed base of  $\Phi$ . If  $\Phi$  is irreducible, the numbering of simple roots will be as in [1] and we will denote by  $\alpha_0$  the opposite of the highest root in  $\Phi$  and by  $d_i$  the coefficient of  $\alpha_i$  in the expression of  $-\alpha_0$ . We set  $d_0 := 1$  and  $\tilde{\Delta} = \Delta \cup \{\alpha_0\}$ . For a subset  $S \subset \tilde{\Delta}$ , we define  $d_S := \gcd(d_i \mid \alpha_i \notin S)$ . In particular,  $d_S = 1$  if  $S \subset \Delta$ .

The group acts on itself by conjugation and we denote  $g \cdot h = ghg^{-1}$  for  $g, h \in G$  and  $G \cdot g$  the  $G$ -conjugacy class of  $g \in G$ . For a closed subgroup  $H \leq G$ , the identity component will be denoted by  $H^\circ$ , and for  $g \in G$  the centralizer will be denoted by  $G_g$ . We will call  $G_g^\circ$  the connected centralizer of  $g$ . The Jordan decomposition of an element  $g \in G$  will be usually denoted by  $g = su$ .

For  $m \in \mathbb{N}$ , we set  $G_{(m)} := \{g \in G \mid \dim(G \cdot g) = m\}$ . These sets are locally closed and their irreducible components are called the *sheets* of the  $G$ -action. For  $Z \subset G$ , we also define  $m_Z := \max\{m \in \mathbb{N} \mid G_{(m)} \cap Z \neq \emptyset\}$  and  $Z^{reg} := Z \cap G_{(m_Z)}$ , and  $C_G(Z)$  will indicate the centralizer of  $Z$  in  $G$ .

If we fix a maximal torus  $T$  of  $G$  and  $\Phi$  is the root system of  $G$  with respect to  $T$ , then for  $\alpha \in \Phi$ , we indicate by  $X_\alpha$  the corresponding root subgroup. For a closed subset  $\Psi \subset \Phi$  (see [1, Chapter VI, n. 1.7, Définition 4]), and for  $s \in T$ , we set  $G_\Psi := \langle T, X_\alpha \mid \alpha \in \Psi \rangle$ .

### 2.1 Construction of sheets and a first parametrization

It was observed in [9, Section 3] that  $G$  has a partition into finitely many, locally closed, smooth, irreducible,  $G$ -stable sets, which we call Jordan classes, each contained in some  $G_{(m)}$ . As a set, the class containing  $g = su$  is

$$J(su) = G \cdot ((Z(G_s^\circ)^\circ s)^{reg} u).$$

In other words, a  $G$ -conjugacy class lies in  $J(su)$  if and only if it contains an element with Jordan decomposition  $s'u$  with  $G_s^\circ = G_{s'}^\circ$  and  $s' \in Z(G_s^\circ)^\circ s$ . The closure of a Jordan class is a union of Jordan classes (see [9, Section 3]); hence, the same holds for the regular locus of the closure of a Jordan class. This gives a partial order on the set of Jordan classes given by  $J_1 \leq J_2$  if and only if  $J_1 \subset \overline{J_2}^{reg}$ . The sheets in  $G$  are the locally closed sets of the form  $\overline{J}^{reg}$  where  $J$  is maximal with respect to  $\leq$ . Hence,

the set of sheets in  $G$  is in bijection with the set  $\mathcal{J}$  consisting of maximal Jordan classes (see [4, Proposition 5.1], [14, Section 3]).

If  $J = J(su)$  as above, then [3, Lemma 2.1] and [4, Proposition 4.8] give

$$\overline{J(su)}^{reg} = \bigcup_{z \in Z(G_s^\circ)^\circ s} G \cdot (z \text{Ind}_{G_s^\circ}^{G_s^\circ}(G_s^\circ \cdot u)),$$

where  $\text{Ind}_{G_s^\circ}^{G_s^\circ}(G_s^\circ \cdot u)$  is Lusztig–Spaltenstein’s induced unipotent conjugacy class (see [11]). The maximal Jordan classes are precisely those for which the class of  $u$  is *rigid* in  $G_s^\circ$ , i. e., it is not induced from any unipotent class in a proper Levi subgroup of a parabolic subgroup of  $G_s^\circ$  (see [4, Proposition 5.3], [14, Lemma 2.4]).

Sheets are then parametrized as follows.

**Theorem 2.1** ([4, Theorem 5.6], [14, Theorem 3.1]) *The assignment  $J = J(su) \mapsto (G_s^\circ, Z(G_s^\circ)^\circ s, G_s^\circ \cdot u)$  induces a bijection between  $\mathcal{J}$  and the set of  $G$ -orbits of triples  $(M, Z(M)^\circ r, \mathcal{O})$  where  $M$  is the connected centralizer of a semisimple element of  $G$ ;  $Z(M)^\circ r$  is a coset in  $Z(M)/Z(M)^\circ$  satisfying  $C_G(Z(M)^\circ r)^\circ = M$  and  $\mathcal{O}$  is a rigid unipotent conjugacy class in  $M$ .*

We aim at a simpler parametrization for  $G$  simple and of adjoint type.

## 2.2 Connected centralizers of semisimple elements

In this subsection,  $G$  is quasisimple. Identity components of centralizers of semisimple elements have been studied in [5–8, 13]. In the spirit of the latter, we give a combinatorial characterization of the root subsystem of such subgroups when the base field is an arbitrary algebraically closed field.

**Proposition 2.2** *Let  $G$  be quasisimple, let  $T$  be a maximal torus in  $G$ , and let  $\Psi$  be a closed subset of  $\Phi$ . Then  $G_\Psi$  is the connected centralizer of an element in  $T$  if and only if  $\Psi$  is conjugate to a root subsystem  $\Psi'$  admitting a base  $\Delta_{\Psi'} \subset \tilde{\Delta}$  and such that  $\gcd(p, d_i \mid i \in \tilde{\Delta} \setminus \Delta_{\Psi'}) = 1$  by an element in the normalizer  $N(T)$  of  $T$ .*

**Proof** If  $\Psi$  is  $N(T)$ -conjugate to a root subsystem  $\Psi'$  admitting a base  $\Delta_{\Psi'} \subset \tilde{\Delta}$  and such that  $\gcd(p, d_i \mid i \in \tilde{\Delta} \setminus \Delta_{\Psi'}) = 1$ , then there exists an  $\alpha_{\tilde{i}} \in \tilde{\Delta} \setminus \Delta_{\Psi'}$  such that  $p \nmid d_{\tilde{i}}$ . Replacing  $a_1$  with  $d_{\tilde{i}}$  in the proof of [13, Proposition 32], we obtain an element  $s \in Z(G_\Psi)$  such that  $G_s^\circ = G_{\Psi'}$ .

Assume now that  $G_\Psi = G_s^\circ$  for some  $s \in T$ . By [13, Proposition 30], the subgroup  $G_\Psi$  is  $G$ -conjugate to some  $G_{\Psi'}$  where  $\Psi'$  admits a base  $\Delta_{\Psi'} \subset \tilde{\Delta}$ . Conjugacy of maximal tori in  $G_{\Psi'}$  ensures that conjugation of the two subgroups, and of the corresponding root systems can be obtained using an element in  $N(T)$ . We show that  $\gcd(p, d_{\Delta_{\Psi'}}) = 1$ . If the characteristic exponent  $p = 1$ , then there is nothing to prove. Assume for a contradiction that  $p > 1$  and divides  $d_i$  for every  $\alpha_i \in \tilde{\Delta} \setminus \Delta_{\Psi'}$ . Since  $d_0 = 1$ , in this situation  $\alpha_0 \in \Delta_{\Psi'}$  and  $G_{\Psi'}$  is never the Levi subgroup of a parabolic subgroup of  $G$ .

Let  $\{\omega_j^\vee, j = 1, \dots, n\}$  be the basis for the cocharacters of  $T$  dual to  $\Delta$ , and let  $t \in T$  be such that  $G_{\Psi'} = G_{\Psi'}^\circ$ . Since  $\alpha_i(t) = 1$ , for every  $\alpha_i \in \Delta_{\Psi'}$ , we have  $t = \prod_{\substack{j=1, \dots, n \\ \alpha_j \notin \Delta_{\Psi'}}} \omega_j^\vee(\zeta_j)$ , and  $\alpha_0(t) = 1$  gives

$$(2.1) \quad \prod_{j \geq 1, \alpha_j \notin \Delta_{\Psi'}} \zeta_j^{d_j} = 1, \text{ so also } \prod_{j \geq 1, \alpha_j \notin \Delta_{\Psi'}} \zeta_j^{d_j/p^l} = 1 \text{ if } p^l | d_{\Delta_{\Psi'}}.$$

By [1, Section 1 of Chapter VI, n. 1.7, Paragraphe 1.7, Proposition 24], the system  $\Psi'$  is not  $\mathbb{Q}$ -closed, i.e., there exists an  $\alpha \in (\mathbb{Q}\Delta_{\Psi'} \cap \Phi) \setminus \Psi'$ . In other words, there exist  $a, b, a_i, b_i \in \mathbb{Z}$ , for  $i = 1, \dots, n$  such that  $\alpha_i \notin \Delta_{\Psi'}$ , with  $b, b_i \neq 0$  and  $z_j \in \mathbb{Z}$  for  $j = 1, \dots, n$  such that

$$\alpha = \frac{a}{b} \alpha_0 + \sum_{\alpha_i \in \Delta_{\Psi'} \cap \Delta} \frac{a_i}{b_i} \alpha_i = \sum_{j=1}^n z_j \alpha_j \in \Phi \setminus \Psi'.$$

Hence,  $b | d_i$  for all  $\alpha_i \notin \Delta_{\Psi'}$ , that is,  $b | d_{\Delta_{\Psi'}}$ . Observe that  $d_{\Delta_{\Psi'}}$  is either  $p^l$  for some  $l \geq 1$  or else  $d_{\Delta_{\Psi'}} = 6, \Phi = E_8, p = 2$  or  $3$ , and  $\Delta_{\Psi'} = \widetilde{\Delta} \setminus \{\alpha_4\}$ . In the first case, (2.1) gives  $\alpha(t) = 1$ , a contradiction. In the second case,  $t = \omega_4^\vee(\zeta_4)$  and (2.1) becomes  $\zeta_4^{6/p} = 1$ . Let  $\beta \in \Phi \setminus \Psi'$  be such that its coefficient of  $\alpha_4$  in its expression as a sum of simple roots is  $6/p$ . Then again  $\beta(t) = 1$ , a contradiction. ■

**Remark 2.3** The coprimality condition in Proposition 2.2 says that the centralizers we lose when passing from good characteristic to bad characteristic  $p$  are precisely those subgroups that in good characteristic are connected centralizers of elements of order divisible by  $p$ . When  $k$  is the algebraic closure of a finite field the statement could also be extracted from [7, Section 2].

**Proposition 2.4** Let  $G$  be simple of adjoint type, let  $T$  be a maximal torus in  $G$ , and let  $\Psi \subset \Phi$  be a closed subsystem such that  $G_\Psi = G_s^\circ$  for some  $s \in T$ . Assume in addition that  $\Psi$  admits a base  $\Delta_\Psi \subset \widetilde{\Delta}$ . Then:

- (a) The torsion subgroup of  $\mathbb{Z}\Phi/\mathbb{Z}\Psi$  is  $\mathbb{Z}/d_{\Delta_\Psi}\mathbb{Z}$ .
- (b)  $Z(G_\Psi)/Z(G_\Psi)^\circ$  is cyclic of order  $d_{\Delta_\Psi}$ .
- (c) For  $t \in Z(G_\Psi)$ , we have  $C_G(Z(G_\Psi)^\circ t) = G_\Psi$  if and only if  $Z(G_\Psi)^\circ t$  is a generator of  $Z(G_\Psi)/Z(G_\Psi)^\circ$ .

**Proof** (a) This is observed in [15, Section 2].  
 (b) The order of the torsion subgroup of  $\mathbb{Z}\Phi/\mathbb{Z}\Psi$  is coprime with  $p$  by Proposition 2.2. Hence, we are in a position to use the argument in [13, Lemma 33] and [15, Section 2.1], that we sketch for completeness. By construction and [12, Proposition 3.8] from which we borrow notation,  $Z(G_\Psi) = (\mathbb{Z}\Psi)^\perp, Z(G_\Psi)^\perp = \mathbb{Z}\Psi$ , and the character group  $X(Z(G_\Psi))$  is  $\mathbb{Z}\Phi/\mathbb{Z}\Psi$ . Then

$$X(Z(G_\Psi)/Z(G_\Psi)^\circ) \simeq \{\chi \in X(Z(G_\Psi)) \mid \chi(z) = 1, \forall z \in Z(G_\Psi)^\circ\}$$

consists of those torsion elements in  $\mathbb{Z}\Phi/\mathbb{Z}\Psi$  whose order is coprime with  $p$ , that is,  $\mathbb{Z}/d_{\Delta_\Psi}\mathbb{Z}$ .

- (c) In good characteristic, this is, [13, Lemma 34(1)], whose proof relies on the fact that every character of  $Z(G_\Psi)/Z(G_\Psi)^\circ$  can be represented by an element in  $\Phi$ . The proof of this statement depends on considerations on root systems which are characteristic-free and on a natural isomorphism between  $X(Z(G_\Psi)/Z(G_\Psi)^\circ)$  and  $\mathbb{Z}(\mathbb{Q}\Psi \cap \Phi)/\mathbb{Z}\Psi$ , which remains valid because  $X(Z(G_\Psi)/Z(G_\Psi)^\circ)$  is the torsion subgroup of  $\mathbb{Z}\Phi/\mathbb{Z}\Psi$  also in our situation. ■

### 2.3 The parametrization

In this subsection,  $G$  is simple of adjoint type. We are now in a position to prove the refinement of the parametrization of sheets of  $G$ . The general case can be readily deduced by standard arguments.

*Theorem 2.5* *Let  $G$  be simple and of adjoint type. The sheets in  $G$  are in bijection with the  $G$ -conjugacy classes of pairs  $(M, \mathcal{O})$  where  $M$  is the connected centralizer of a semisimple element in  $G$  and  $\mathcal{O}$  is a rigid unipotent conjugacy class in  $M$ .*

**Proof** In good characteristic, this is [2, Theorem 4.1], so we assume that  $p$  is bad for  $G$ . Sheets are parametrized by triples  $(M, Z(M)^\circ s, \mathcal{O})$  as in Theorem 2.1. The assignment  $(M, Z(M)^\circ s, \mathcal{O}) \mapsto (M, \mathcal{O})$  induces a well-defined and surjective map between the set of  $G$ -conjugacy classes of triples and the set of  $G$ -conjugacy classes of pairs as above. We show injectivity of this map. If  $G$  is classical, then  $M$  is a Levi subgroup of a parabolic subgroup of  $G$ , for any pair  $(M, \mathcal{O})$ ; hence,  $Z(M) = Z(M)^\circ$  and there is nothing to prove, so we assume that  $G$  is of exceptional type.

Let  $(M, Z(M)^\circ s, \mathcal{O})$  and  $(M, Z(M)^\circ r, \mathcal{O})$  be two triples inducing the same image. Without loss of generality  $s \in T$ ,  $M = G_\Psi$  where  $\Psi$  has base  $\Delta_\Psi \subset \tilde{\Delta}$ , and  $Z(M)^\circ s, Z(M)^\circ r \subset T$ . If  $d_{\Delta_\Psi} \leq 2$ , then necessarily  $Z(M)^\circ r = Z(M)^\circ s$ , so we assume that  $d_{\Delta_\Psi} \geq 3$ .

By [15, Proposition 7], there is a  $w$  in the stabilizer  $N_W(\Delta_\Psi)$  of  $\Delta_\Psi$  in  $W$  whose action on  $\mathbb{Z}\Phi/\mathbb{Z}\Psi$  generates the automorphism group of the torsion subgroup of  $\mathbb{Z}\Phi/\mathbb{Z}\Psi$ , which is isomorphic to  $Z(M)/Z(M)^\circ$  by Proposition 2.4(a). We claim that any representative of  $w$  in  $N(T)$  preserves  $\mathcal{O}$ . Since rigid unipotent classes in type  $A$  are trivial, it is enough to consider only the case in which  $\Delta_\Psi$  contains a (necessarily unique) component of type different from  $A$ , and  $\mathcal{O}$  is nontrivial in the corresponding subgroup. Such a component is always preserved by the action of  $w$ . The list of  $\Delta_\Psi$  with  $d_{\Delta_\Psi} \geq 3$  in the proof of [15, Proposition 7] shows that we only need to consider two cases for  $G$  of type  $E_8$ , namely  $\Delta_\Psi = A_3 + D_5$  which may occur only when  $p = 3, 5$ , and  $\Delta_\Psi = A_2 + E_6$ . Unipotent conjugacy classes in type  $D$  for  $p = 3$  and  $5$  are characteristic unless their Jordan form corresponds to a very even partition, i.e., a partition with only even terms, each occurring an even number of times. Such partitions never occur in  $D_n$  for  $n$  odd.

Rigid unipotent conjugacy classes in  $E_6$  in arbitrary characteristic can be deduced from [16, Chapitre II. Appendice] and they are characteristic for dimensional reasons. ■

Fixing a maximal torus  $T$  in  $G$ , by standard arguments, we retrieve a parametrization of sheets by orbits of the Weyl group. We set  $\mathcal{T}$  to be the set of pairs  $(M, \mathcal{O})$  where

$M$  is the connected centralizer of an element in  $T$  and  $\mathcal{O}$  is a rigid unipotent class in  $M$ , so  $N(T)$  and  $W = N(T)/T$  naturally act on  $\mathcal{T}$ .

**Corollary 2.6** *Let  $G$  be simple and of adjoint type. The sheets in  $G$  are parametrized by elements in  $\mathcal{T}/W$ .*

**Remark 2.7** If the characteristic of the base field is good, replacing “rigid” by “distinguished” in the parametrization in Theorem 2.5 one obtains the parametrization of conjugacy classes in the component group of the centraliser of unipotent conjugacy classes in  $G$  in [13, 15]. In bad characteristic, the most naive generalization of this parametrization fails to hold, as there are too few connected centralizers of semisimple elements, and it would be interesting to detect a suitable analog.

## 2.4 On the number of sheets in $G$

It was observed in [14, Remark 3.3] that for  $G$  of type  $B_2$ , the number of sheets is independent of the characteristic and it was suggested this to hold in general for  $G$  connected and simply connected. This fails in general because there exist sheets that are obtained from one another by multiplication by a central element, and such central element might no longer exist in bad characteristic: for example, in  $G = SL_2(k)$ , the sheets are:  $\{\text{id}\}$ ,  $\{-\text{id}\}$ , and  $G^{reg}$  for  $p \neq 2$ , and  $\{\text{id}\}$  and  $G^{reg}$  for  $p = 2$ . The following remark shows that the number of sheets depends on  $p$  also for  $G$  simple of adjoint type.

**Remark 2.8** Let  $\Phi = G_2$ . We use the parametrization in Theorem 2.5. The semisimple parts of the connected centralizers for  $p$  good are of type:  $G_2, A_2, A_1 + \tilde{A}_1, A_1, \tilde{A}_1$  or conjugate to  $T$ . In type  $A$ , all rigid unipotent classes are trivial, and there are three rigid unipotent classes in  $G$  (see [16, Chapitre II. Appendice]). Hence, there are eight sheets for  $p$  good.

According to Proposition A.1, the semisimple parts of the connected centralizers for  $p = 2$  are of type:  $G_2, A_2, A_1, \tilde{A}_1$  or conjugate to  $T$ , and there are three rigid unipotent classes in  $G$  (see [16, Chapitre II. Appendice]). Hence, there are seven sheets for  $p = 2$ .

According to Proposition A.1, the semisimple parts of the connected centralizers for  $p = 3$  are of type:  $G_2, A_1 + \tilde{A}_1, A_1, \tilde{A}_1$  or conjugate to  $T$ , and there is an extra unipotent conjugacy class in  $G_2$  which is rigid, as it can be deduced from the list of induced classes in [16, Chapitre II. Appendice]. Hence, the number of sheets for  $p = 3$  equals the number of sheets for  $p$  good.

## A Appendix

For the reader’s convenience, we list the possible connected centralizers in bad characteristic. When  $k$  is the closure of a finite field this list appeared in [7, pp. 25–27]. For writing the Table A.1, we made use of the analysis of  $W$ -conjugacy classes of subsets of  $\tilde{\Delta}$  in [15, Section 2.2]. In most cases, these classes are determined by their isomorphism type and the root lengths. For  $\Phi = E_7$  and  $E_8$ , we remove ambiguities

Table A.1. Root subsystems to be discarded.

$G$	$p$	$\Psi$
$E_6$	2	$4A_1, A_3 + 2A_1, A_5 + A_1$
$E_6$	3	$3A_2$
$E_7$	2	$(4A_1)', (A_3 + 2A_1)', 5A_1, (A_5 + A_1)', A_3 + 3A_1, 2A_3, D_4 + 2A_1, A_7, D_6 + A_1, 2A_3 + A_1$
$E_7$	3	$3A_2, A_2 + A_5$
$E_8$	2	$(4A_1)'', (A_3 + 2A_1)'', 5A_1, (2A_3)'', (A_1 + A_5)'', 3A_1 + A_3, 2A_1 + D_4, 4A_1 + A_2, A_7'', 2A_1 + A_5, A_3 + D_4, A_1 + D_6, A_1 + 2A_3, 2A_1 + A_2 + A_3, 2A_1 + D_5, D_8, A_1 + A_7, A_1 + A_2 + A_5, A_3 + D_5, A_1 + E_7$
$E_8$	3	$3A_2, A_2 + A_5, 3A_2 + A_1, A_8, A_1 + A_2 + A_5, A_2 + E_6$
$E_8$	5	$2A_4$
$F_4$	2	$2A_1, 2A_1 + \tilde{A}_1, A_1 + B_2, A_3, A_1 + C_3, A_3 + \tilde{A}_1, B_4$
$F_4$	3	$A_2 + \tilde{A}_2$
$G_2$	2	$A_1 + \tilde{A}_1$
$G_2$	3	$A_2$

adopting, as in *loc. cit.*, Dynkin’s convention. Namely, for  $n = 7, 8$ , we decorate with one prime the root subsystems which can be embedded in the subsystem of type  $A_n$  within  $E_n$ , whereas we decorate with two primes the root subsystems with the same label which cannot be embedded in  $A_n \subset E_n$ .

**Proposition A.1** *Let  $G$  be quasisimple with  $p$  bad for  $G$ . Let  $T$  be a maximal torus in  $G$  and  $\Psi \subset \Phi$  be a closed subsystem with base  $\Delta_\Psi \subset \tilde{\Delta}$ .*

*If  $\Phi$  is of classical type, then  $G_\Psi$  is the connected centralizer of an element in  $T$  if and only if it is the Levi subgroup of a parabolic subgroup of  $G$  (see [6]).*

*If  $\Phi$  is of exceptional type, then  $G_\Psi$  is the connected centralizer of an element in  $T$  unless  $G, p$  and  $\Psi$  occur in Table A.1.*

In other words, the subsystems to be discarded are those containing the subsystem whose base consists of all elements in  $\tilde{\Delta}$  whose coefficient in the expression of  $-\alpha_0$  is not divisible by  $p$ .

**Acknowledgment** We wish to thank Prof. G. Lusztig for asking the question leading to this short note, Prof. J. Michel for pointing us reference [7], and the referee for his/her remark on centralizers and the appendix table.

**References**

[1] N. Bourbaki, *Groupes et Algèbres de lie, Chapitres 4,5, et 6, Éléments de Mathématique*, Masson, Paris, 1981.  
 [2] G. Carnovale, *Lusztig’s partition and sheets (with an appendix by M. Bulois)*. Math. Res. Lett. 22(2015), 645–664.  
 [3] G. Carnovale, *Lusztig’s strata are locally closed*. Arch. Math. 115(2020), 23–26.  
 [4] G. Carnovale and F. Esposito, *On sheets of conjugacy classes in good characteristic*. Int. Math. Res. Not. 4(2012), 810–828.

- [5] R. W. Carter, *Centralizers of semisimple elements in finite groups of lie type*. Proc. Lond. Math. Soc. 37(1978), no. 3, 491–507.
- [6] R. W. Carter, *Centralizers of semisimple elements in the finite classical groups*. Proc. Lond. Math. Soc. 42(1981), 1–41.
- [7] D. I. Deriziotis, *The Brauer complex and its applications to the Chevalley groups*. Ph.D. thesis, University of Warwick, 1977.
- [8] D. I. Deriziotis, *Centralizers of semisimple elements in a Chevalley group*. Comm. Algebra 9(1981), no. 19, 1997–2014.
- [9] G. Lusztig, *Intersection cohomology complexes on a reductive group*. Invent. Math. 75(1984), 205–272.
- [10] G. Lusztig, *On conjugacy classes in a reductive group*. In: M. Nevens and P. Trapa (eds.), Representations of reductive groups, Progress in Mathematics, 312, Springer, Cham, 2015, pp. 333–363.
- [11] G. Lusztig and N. Spaltenstein, *Induced unipotent classes*. J. Lond. Math. Soc. (2) 19(1979), 41–52.
- [12] G. Malle and D. Testerman, *Linear algebraic groups and finite groups of lie type*, Cambridge Studies in Advanced Mathematics, 133, Cambridge University Press, Cambridge, 2011.
- [13] G. McNinch and E. Sommers, *Component groups of unipotent centralizers in good characteristic*. J. Algebra 270(2003), no. 1, 288–306.
- [14] I. Simion, *Sheets of conjugacy classes in simple algebraic groups*. Mathematica 61(2019), no. 84, 183–190.
- [15] E. Sommers, *A generalization of the Bala–Carter theorem for nilpotent orbits*. Int. Math. Res. Not. 11(1998), 539–562.
- [16] N. Spaltenstein, *Classes unipotentes et sous-groupes de Borel*, Lecture Notes in Mathematics, 946, Springer, Cham, 1982.

*Dipartimento di Matematica “Tullio Levi-Civita”, Università di Padova, via Trieste 63, 35121 Padova, Italy*  
e-mail: [ambrosio@math.unipd.it](mailto:ambrosio@math.unipd.it) [carvalho@math.unipd.it](mailto:carvalho@math.unipd.it) [esposito@math.unipd.it](mailto:esposito@math.unipd.it)