


H₂O masers and host environments of FU Orionis and EX Lupi type low-mass eruptive YSOs

Zsófia Marianna Szabó^{1,2,3,4} , Yan Gong¹, Wenjin Yang¹,
Karl M. Menten¹, Olga S. Bayandina⁵, Claudia J. Cyganowski²,
Ágnes Kóspál^{3,4,6,7}, Péter Ábrahám^{3,4,6}, Arnaud Belloche¹ and
Friedrich Wyrowski¹

¹Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany.
email: zszabo@mpifr-bonn.mpg.de

²Scottish Universities Physics Alliance (SUPA), School of Physics and Astronomy, University of St Andrews, North Haugh, St Andrews, KY16 9SS, UK

³Konkoly Observatory, Research Centre for Astronomy and Earth Sciences, Eötvös Loránd Research Network (ELKH), Konkoly-Thege Miklós út 15-17, 1121 Budapest, Hungary

⁴CSFK, MTA Centre of Excellence, Budapest, Konkoly Thege Miklós út 15-17., H-1121, Hungary

⁵INAF - Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125 Firenze, Italy

⁶ELTE Eötvös Loránd University, Institute of Physics, Pázmány Péter sétány 1/A, H-1117 Budapest, Hungary

⁷Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany

Abstract. The FU Orionis (FUor) and EX Lupi (EXor) type objects are rare pre-main sequence low-mass stars undergoing accretion outbursts. Maser emission is widespread and is a powerful probe of mass accretion and ejection on small scales in star forming region. However, very little is known about the overall prevalence of water masers towards FUors/Exors. We present results from our survey using the Effelsberg 100-m telescope to observe the largest sample of FUors and EXors, plus additional Gaia alerted sources (with the potential nature of being eruptive stars), a total of 51 targets, observing the 22.2 GHz H₂O maser, while simultaneously covering the NH₃ 23 GHz.

Keywords. Stars: pre-main sequence, Stars: low-mass, Masers

1. Introduction

Low-mass young stellar objects in their early stellar evolution can undergo accretion-driven episodic outbursts. By studying this phenomena we are able to gather crucial information on the formation and the evolution of Sun-like stars. The member of both FUor and EXor classes experience major increase in their brightness observed in the optical and near-infrared wavelengths. FUors can brighten up to 5-6 magnitudes in the optical and stay in a high-accretion state for decades, but likely centuries (e.g., [Fischer *et al.* 2022](#), and references therein), while EXors brighten between 1-5 magnitudes and remain in a bright state for months or years, and the outbursts are recurring (e.g., [Audard *et al.* 2014](#); [Cruz-Sáenz de Miera *et al.* 2022](#)). Masers have been substantially used to probe low- and high-mass star formation regions (e.g., [Abraham *et al.* 1981](#); [Omodaka *et al.* 1999](#);




Figure 1. Detected H₂O masers towards HH 354 IRS.

Hirota *et al.* 2011; Furuya *et al.* 2003, 2001), currently still little information exists on masers in FUors/EXors.

2. Results

2.1. H₂O masers associated with eruptive stars

We detected H₂O masers towards 5 sources, but only 3 of the detections are likely associated with FUors/EXors, which include one EXor: V512 Per (Class I source) commonly known as SVS 13A, and two FUors: HH 354 IRS (Class 0/I) and Z CMA (Class I). The maser towards HH 354 IRS is the first to be reported. The maser component in Z CMA detected in our survey is a new component yet to be reported (see Fig. 1, Szabó *et al.* 2023b).

2.2. Serendipitous detections towards Class 0 protostars

Towards V512 Per (SVS 13A), multiple variable maser features were detected arising from a nearby source H₂O(B) (Class 0 object, known for H₂O masers, Haschick *et al.* 1980), which is within the beam. The source was in an active flare, contaminating the spectra of V512 Per. The peak of the emission was found to be associated with H₂O(B), but one maser feature at >11 km s⁻¹ is likely to be associated with V512 Per (see Fig. 2). Water masers were also detected towards the FUor binary RNO 1B/1C, but they are most likely arising from the molecular outflow of IRAS 00338+6312, located 4'' from the FUors (see e.g., Fiebig 1995; Fiebig *et al.* 1996).

2.3. Discussion and conclusions

The detection rate of our survey of FUors/Exors is only 6%, surprising in light of the close connections between H₂O maser emission and mass accretion. Possible explanations include:

1. Evolutionary effect: H₂O maser detection rate generally decreases from Class 0 to Class II sources (e.g., Furuya *et al.* 2003).
2. Low luminosities in low-mass star formation regions: low bolometric luminosities result in lower flux densities (e.g., Urquhart *et al.* 2011).
3. Rapid time variation: time variability is known and evident from our study, however masers can be in quiescence for ~ 5 years (Claussen *et al.* 1996). Many of our targets




Figure 2. Pointed H₂O spectra towards V512 Per and H₂O(B) observed in 2022 February.

during the survey might have been inactive, despite showing maser emission in the past (see also Szabó *et al.* 2023b).

References

- Abraham, Z., Cohen, N. L., Opher, R., *et al.* 1981, *A&A*, 100, L10
 Audard, M., Ábrahám, P., *et al.* 2014, *Protostars and Planets VI*, 387
 Claussen, M. J., Wilking, B. A., Benson, P. J., *et al.* 1996, *ApJS*, 106, 111
 Cruz-Sáenz de Miera, F., Kóspál, Á., Ábrahám, P., *et al.* 2022, *ApJ*, 927, 125
 Fiebig, D., 1995, *A&A*, 298, 207
 Fiebig, D., Duschl, W. J., Menten, K. M., Tscharnuter, W. M., 1996, *A&A*, 310, 199
 Fischer, W. J., Hillenbrand, L. A., *et al.* 2022, *Protostars and Planets VII*, eprint arXiv:2203.11257
 Furuya, R. S., Kitamura, Y., Wootten, H., *et al.* 2001, *ApJ*, 559, L143
 Furuya, R. S., Kitamura, Y., Wootten, A., *et al.* 2003, *ApJS*, 144, 71
 Haschick, A. D., Moran, J. M., *et al.* 1980, *ApJ*, 237, 26
 Hirota, T., Tsuboi, M., Fujisawa, K., *et al.* 2011, *ApJ*, 739, L59
 Omodaka, T., Maeda, T., Miyoshi, M., *et al.* 1999, *PASJ*, 51, 333
 Urquhart, J. S., Morgan, L. K., Figura, C. C., *et al.* 2011, *MNRAS*, 418, 1689
 Szabó, Zs. M., Gong, Y., Yang, W., *et al.* 2023, *A&A*, 672, A158
 Szabó, Zs. M., Gong, Y., Yang, W., *et al.* 2023, *A&A*, 674, A202