

Image:




Image 2:





Image 3:



Conclusions: While comparing percentages of three different monocyte subsets, clear differences in their distribution among the control and patient groups were appreciated. After comparing the subset frequencies between active patients (AD) and patients who were in remission (NAD), significant differences among the subsets were found although without reaching values of the HC, indicating that even patients in remission show an activated monocyte profile.

Disclosure of Interest: None Declared

EPV0809

Psychiatric manifestations of Susac Syndrome: a case report

M. Gomes^{1*}, P. Veloso¹, E. Monteiro¹, R. Faria¹ and E. Freitas²

¹Psychiatry and ²Neurology, Hospital de Braga, Braga, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.2113

Introduction: Susac Syndrome (SS) is an immune-mediated endotheliopathy that mainly affects young women. It is characterized by the typical triad: subacute encephalopathy, retinal vaso-occlusive disease, and hearing loss. Encephalopathy symptoms are varied and include memory loss, psychiatric disturbances, cranial nerve disorders, seizures, and dementia. The syndrome is considered a rare but important differential diagnosis in various neurological, psychiatric, ophthalmological, and ear-nose-throat disorders.

Objectives: Report a clinical case of SS to reflect on the relationship between psychiatric and neurological symptoms and on immune-mediated psychiatric symptoms.

Methods: Collection of clinical information from the patient described below. Review of the literature about SS.

Results: A 21-year-old woman presented to the Psychiatry Emergency Department in November 2021 for complaints of sadness, anhedonia and emotional lability, with one month of evolution. She also had insomnia and confusional periods, so she stopped driving and quit her work as a storekeeper. She was given sertraline 50mg/day and trazodone 50mg/day. In the past two weeks, the patient had episodes of headache and vomiting, with 8 kg weight loss. She started a fever (38.5°C) two days before observation. The patient had a prior history of depressive symptomatology four years earlier related to her father's grief and her medical and surgical history was unremarkable. She was brought in a wheelchair by her mother and was using diapers because she was confined to bed for the past week. Objectively, the patient was somnolent, tearful and confused, with scarce speech and psychomotor slowing. No focal signs were found on neurological examination. Collaboration of Neurology was asked. Routine laboratory studies showed a slight increase in leucocyte count (12 500/mm³) and CRP (17mg/dL). Cerebrospinal fluid analysis showed 15 cells/uL and protein of 2.64 g/L. Cerebral MRI showed multiple striatocapsular periventricular lesions involving the thalamus, the left midbrain, and the medulla oblongata, as well as focal bilateral hemispherical and cerebellar subcortical lesions. The lesions presented high signal in T2 and showed restriction in the diffusion study. She was admitted to the Neurology inpatient department and treated with pulse methylprednisolone 1000 mg/day for five consecutive days, after which cognitive function improved. Ophthalmology observation found cotton-wool exudate and arteriolar interruption in the right eye, supporting the diagnosis of SS.

Conclusions: This syndrome represents the importance of diligent cooperation among different medical specialties to improve diagnosis-making, treatment and recovery. Psychiatric symptoms are frequent in neurological syndromes, so a high degree of suspicion is needed.

Disclosure of Interest: None Declared

EPV0810

Features of the immune status in patients with functional disorders of the gastrointestinal tract with non-psychotic mental disorders.

N. P. Chernus^{1*}, A. S. Sivkov², R. V. Gorenkov^{3,4}, S. I. Sivkov⁵ and ? . Золотовицкая¹

¹the Outpatient Care Department, the I.M. Sechenov First Moscow State Medical University, Москва; ²the Department of Clinical Pharmacology and Internal Diseases Propaedeutics, Первый МГМУ им. И.М. Сеченова: Москва, Россия; ³Scientific substantiation of health preservation of the population of the Russian Federation, The N. A. Semashko National Research Institute of Public Health; ⁴Higher Healthcare Organization Management, the I.M. Sechenov First Moscow State Medical University, Москва and ⁵the Department of Clinical Pharmacology and Internal Diseases Propaedeutics, the I.M. Sechenov First Moscow State Medical University, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.2114

Introduction: Актуальность исследования накопления данных о исключении иммунологического статуса у больных с

выраженными расстройствами желудочно-кишечного тракта (ФЖКТ), характерных не психотических заболеваний легких.

Objectives: изучение взаимосвязи и появление признаков иммунологического и психовегетативного статуса больных у ФЖИР

Methods: Проанализированы и рассчитаны результаты ПВ-А-обследования (тест ММПИ) с иммунологическим статусом 57 больных (средний возраст 35,36 ± 14,43 года, мужчины-18, женщины-39) с ФАД в динамике с группой здоровых лиц 34 человека, среднего возраста 28,63±9,54 года

Results: У больных с тревожной депрессией [средний профиль: пик по шкале 2 + ☒ 7-8 и ☒ 9 шкал] оценочное увеличение относительного значения Т-хелперов CD3+CD4+ (58,0 и 55,0% p = 0,015) при снижении относительного количества Т-цитотоксических лимфоцитов CD3+CD8+ (0,312 и 0,372, p=0,023; 14,0 и 19,0, p=0,001) соответственно, увеличение иммунорегуляторного индекса: Т-хелперы/Т-цитотоксические CD3+ CD4+/CD3+ CD8+ (4,143 и 2,600, p=0,001); снижение относительного количества НК-клеток CD3-CD16+CD56+ (6% и 8%, p=0,01).

У больных с преобладанием соматизированной анестезии [ведущий пик по шкале 1]: основные выявлены в относительно сниженном CD3+ Т-лимфоцитов (53,0 и 60,0, p=0,001); снижение Т-хелперов: CD3+CD4+ (28,0% и 31,0%, p=0,001); повышение ЭКГ: CD3-CD16+CD56+ (34% и 26,0%, p=0,01; 1153 и 470, p=0,01) по сравнению со здоровыми людьми. Выявленные изменения показателей иммунитета выявлялись в период обострения заболевания и носили транзиторный характер.

Conclusions: Анализ данных с позиций клинко-динамического выявления вклада иммунобиологических регуляторов организма в клиническое затяжное течение и участие иммунной системы в механизме реализации психоэмоционального контроля стресса. .

Disclosure of Interest: None Declared

EPV0811

Anti NMDAR encephalitis masked by symptoms of postpartum depression

S. Milanovic Hromin^{1*}, A. Jelcic², R. Perincic³, M. Komso¹, B. Barun⁴, M. Grah⁵, S. Zecevic Penic⁶, S. Biloglav⁷, I. Perincic⁷, M. Juresko⁷ and B. Skific⁷

¹GENERAL HOSPITAL ZADAR, Zadar; ²Ginekologija Dana d.o.o. for gynecology, obstetrics and consulting, Zagreb; ³Private psychiatric office dr Robert Perincic, Zadar; ⁴Department for neuroimmunology of CNS, Clinical hospital center Zagreb; ⁵Clinic for psychiatry Sveti Ivan, Zagreb, Croatia; ⁶Neurological outpatient clinic, Clinic for psychiatry Sveti Ivan, Zagreb and ⁷Department for psychiatry, GENERAL HOSPITAL ZADAR, Zadar, Croatia

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.2115

Introduction: Anti NMDAR encephalitis is a relatively common autoimmune encephalitis characterized by complex neuropsychiatric features and the presence of Immunoglobulin G antibodies against the NR1 subunit of the NMDA receptors in the central nervous system. It causes psychiatric features, confusion, memory