

# CERTAIN METHOD FOR GENERATING A SERIES OF LOGICS

SATOSHI MIURA and SHÛRÔ NAGATA

**Introduction.** At first, we define three relations  $\supseteq$ ,  $=$ , and  $\supset$  in connection with a pair of logics  $L$  and  $L^*$  as follows:

$L \supseteq L^*$ , if and only if every proposition provable in  $L^*$  is also provable in  $L$ ;

$L = L^*$ , if and only if  $L \supseteq L^*$  and  $L^* \supseteq L$ ;

$L \supset L^*$ , if and only if  $L \supseteq L^*$  but not  $L^* \supseteq L$ .

Next, for a logic  $L$ , we denote by  $L[A]$  the fortified logic of  $L$  by regarding a proposition  $A$  as a new axiom scheme.

By  $LOQ$ , we denote the logic obtained by adjoining Peirce's rule,

$$P \equiv ((a \rightarrow b) \rightarrow a) \rightarrow a,$$

to the primitive logic  $LO$  (cf. Ono [6], [7]). According to Nagata [4], we can obtain a *descending sequence*,  $L_1, L_2, \dots$ , from  $LOQ$  toward  $LO$  (i.e.  $LOQ = L_1 \supset L_2 \supset \dots \supset L_i \supset \dots \supset LO$ ) by the following method. A series of propositions  $P_i$  is defined recursively as follows:

$$\begin{cases} P_1 \equiv P, \\ P_{i+1} \equiv ((p_i \rightarrow P_i) \rightarrow p_i) \rightarrow p_i, \quad (i = 1, 2, \dots), \end{cases}$$

where  $p_i$ 's are mutually distinct proposition-variables not occurring in  $P$ . For the series  $P_1, P_2, \dots$ , we can assert that

$$LOQ = LO[P_1] \supset LO[P_2] \supset \dots \supset LO[P_i] \supset \dots \supset LO.$$

We have noticed that, by making use of the same method, existence of descending sequences from  $K$ -series logics ( $LQ, LN, LK$ ) toward their corresponding  $J$ -series logics ( $LP, LM, LJ$ ) (cf. Ono [6]) can be proved. We have also noticed that existence of a descending sequence from  $LN$  toward  $LD = LM[a \vee \neg a]$  (cf. Curry [1]) can be proved similarly.

---

Received September 6, 1966.

Discussing with us our recent studies on the subject, Prof. T. Tugué pointed out that we would have, in a similar manner, a descending sequence toward a logic  $L$  by starting from any proposition  $A$ , not provable in  $L$ , instead of starting from Peirce's rule. Guided by his valuable suggestion, we obtained the following conclusion.

For any proposition  $A$ , a series of propositions  $A_i$  is defined recursively as follows:

$$\begin{cases} A_1 \equiv A, \\ A_{i+1} \equiv ((p_i \rightarrow A_i) \rightarrow p_i) \rightarrow p_i, \quad (i = 1, 2, \dots), \end{cases}$$

where  $p_i$ 's are mutually distinct proposition-variables not occurring in  $A$ . The proposition  $A$  is called *kernel*. Taking Peirce's rule  $P$  as the kernel  $A$ , we can produce the descending sequences described before. If we take  $a \vee \neg a$  (law of the excluded middle) as the kernel  $A$ , we can produce a descending sequence from  $LD$  toward  $LM$ . Along this line, we would be able to give other examples as many as we like. To show these facts, we shall use certain truth-table, called  $(n, r)$ -evaluation. The  $(n, r)$ -evaluation is a slight refining of the truth-table appearing in Gödel [2]<sup>1)</sup>. The refinement lies on the evaluation of negation defined as follows:

$a$	0	1	...	$r-1$	$r$	...	$n$
$\neg a$	$r$	$r$	...	$r$	0	...	0

The main purpose of this paper is to show that, for a logic  $L$  and a kernel  $A$ , we can generate a descending sequence from  $L[A]$  toward  $L$  under certain conditions. We wish to express our thanks to Profs. K. Ono and T. Tugué for their kind guidances.

**DEFINITION.** For integers  $n$  and  $r$  such that  $1 \leq r \leq n$ , any evaluation having the following truth-value properties is called  $(n, r)$ -evaluation:

$$a \rightarrow b = \begin{cases} 0 & \text{if } a \geq b, \\ b & \text{if } a < b, \end{cases}$$

<sup>1)</sup> In Gödel [2], it is discussed that there is "eine monoton abnehmende Folge von Systemen" between  $LK$  and  $LJ$  by considering the formula  $F_n \equiv \bigvee_{1 \leq i < k \leq n} (a_i \equiv a_k)$  with respect to a many-valued evaluation. This fact enables us to do the same discussion between  $LK$  and a logic, in which every provable proposition is  $(n, r)$ -true (cf. Definition) for any  $n$  and some  $r$ . Moreover, in Umezawa [8]—[10] and Nishimura [5], there are detailed discussions on intermediate logics between  $LK$  and  $LJ$ .

$$\begin{aligned}
 a \vee b &= \text{Min}(a, b), \\
 a \wedge b &= \text{Max}(a, b), \\
 \neg a &= \begin{cases} r & \text{if } a < r, \\ 0 & \text{if } a \geq r; \end{cases}
 \end{aligned}$$

where the truth-values of propositions  $a, b$ , denoted simply by  $a, b$ , respectively, runs over the set  $\{0, 1, \dots, n\}$ . If we take the logical constant  $\lambda$  (contradiction) whose truth-value is defined by  $r$ , and define  $\neg a$  by  $a \rightarrow \lambda$ , then the above truth-value property for negation is obtained. For the predicate logics, we take a domain of  $k$  individual objects  $\{\xi_1, \xi_2, \dots, \xi_k\}$  and define the truth-value of

$$\begin{aligned}
 (\xi) a(\xi) &\text{ by } a(\xi_1) \wedge a(\xi_2) \wedge \dots \wedge a(\xi_k), \\
 (\exists \xi) a(\xi) &\text{ by } a(\xi_1) \vee a(\xi_2) \vee \dots \vee a(\xi_k).
 \end{aligned}$$

Any proposition whose truth-value is always 0 is called  $(n, r)$ -true.

For any  $n$  and  $r$ , all the axiom schemes<sup>2)</sup> of **LM** are  $(n, r)$ -true, and all the inference rules of **LM** deduce  $(n, r)$ -true conclusions, whenever their assumptions are all  $(n, r)$ -true.  $\lambda \rightarrow a$  is  $(n, n)$ -true,  $a \vee \neg a$  is  $(n, 1)$ -true, and  $(\lambda \rightarrow a) \vee b \vee \neg b$  (cf. Example) is both  $(n, 1)$ - and  $(n, n)$ -true for all  $n, r$ .

Before stating the theorem, the following two lemmas are remarkable.

LEMMA 1. *If the kernel  $A \leq n - j$  ( $0 \leq j \leq n - 1$ ) for the  $(n, r)$ -evaluation, then,  $A_i \leq n - j - i + 1$  ( $1 \leq i \leq n - j + 1$ ).*

LEMMA 2. *If the kernel  $A$  takes the truth-value  $n - j$  ( $0 \leq j \leq n - 1$ ) for the  $(n, r)$ -evaluation, then,  $A_i$  takes  $n - j - i + 1$  ( $1 \leq i \leq n - j + 1$ ).*

THEOREM. *Let  $L$  be a logic such that  $LK \supset L \supseteq LO$ . Assume that there exists a function  $r = r(n)$  ( $r = 1, 2, \dots, n$ ) satisfying the following conditions.*

- (1) *For all  $n$ , every **L**-provable<sup>3)</sup> proposition is  $(n, r)$ -true.*
- (2) *There exists a non-negative integer  $j$  such that, for all  $n(n \geq 2)$  larger than  $j$ , a proposition  $A$  can never takes the truth-value larger than  $n - j$ , but can take certainly  $n - j$  by the  $(n, r)$ -evaluation.*

*Then, there is a descending sequence from  $L[A]$  toward  $L$ , i.e.,*

<sup>2)</sup> cf. *H* system of Curry [1].

<sup>3)</sup> In this paper, for a logic **L**, a proposition  $A$  is called to be **L**-provable when  $A$  is provable in **L**.

$$L[A] = L[A_1] \supset L[A_2] \supset \dots \supset L[A_i] \supset \dots \supset L.$$

*Proof.* (i) The cases  $j \neq 0$  or  $n \geq 3$ . If we take  $(n + j - 1, r)$ -evaluation in place of  $(n, r)$ -evaluation, then,  $A \leq n - j$  turns out to be  $A \leq n - 1$ . By Lemma 1,  $A_i \leq n - i$  holds; hence,  $A_n = 0$  always holds. Since all the  $L$ -provable propositions are  $(n + j - 1, r)$ -true by assumption, all the  $L[A_n]$ -provable propositions are always  $(n + j - 1, r)$ -true. By Lemma 2, however,  $A_i = n - i$  holds; hence,  $A_{n-1} = 1$  holds. Therefore,  $A_{n-1}$  is not  $L[A_n]$ -provable. Namely,  $L[A_{n-1}] \supset L[A_n] \supset L$ .

(ii) The case  $j = 0$  and  $n = 2$ . By assumption, there exists  $r$  such that  $A_1$  can take 2 by  $(2, r)$ -evaluation. However,  $A_2 \leq 1$  by Lemma 1. Hence,  $A_1$  is not  $L[A_2]$ -provable. Therefore,  $L[A_1] \supset L[A_2]$ . *q. e. d.*

EXAMPLE. In the following table, descending sequences from  $L[A]$  toward  $L$  are exhibited by showing their kernels  $A$  and the numbers  $r$  appearing in the assumption of the theorem. We can further substitute, in the table,  $LM[\neg a \vee \neg \neg a]$  or  $LM[(a \rightarrow b) \vee (b \rightarrow a)]$  etc. for  $LM$ . In the following table,  $A \cap B$  denotes the logics in which any proposition is provable, if and only if it is both  $A$ - and  $B$ -provable.

	$L[A]$	$L$	$A$	$r$
1	<b>LOQ</b>	<b>LO</b>	$P \equiv ((a \rightarrow b) \rightarrow a) \rightarrow a$	$1 \leq r \leq n$
2	<b>LQ</b>	<b>LP</b>	$P$	$1 \leq r \leq n$
3	<b>LN</b>	<b>LM</b>	$P$	$1 \leq r \leq n$
4	<b>LK</b>	<b>LJ</b>	$P$	$r = n$
5	<b>LN</b>	<b>LD</b>	$P$	$r = 1$
6	<b>LD</b>	<b>LM</b>	$a \vee \neg a$	<i>e. g.</i> $r = n$
7	<b>LJ</b>	<b>LM</b>	$\lambda \rightarrow a$	$1 \leq r \leq n - 1$
8	$LN_i \equiv LM[P_i]$	$LD_i \equiv LM[B_i] (B \equiv a \vee \neg a)$	$P_i$	$r = 1$
9	$LN_i$	$LJ_i \equiv LM[C_{i+1}] (C \equiv \lambda \rightarrow a)$	$P_i$	$r = n$
10	$LJ \cap LN$	<b>LM</b>	$(\lambda \rightarrow a) \vee b \vee (b \rightarrow c)$	$1 \leq r \leq n - 1$
11	<b>LN</b>	$LJ \cap LN$	$P$	$r = n$
12	$LJ \cap LD$	<b>LM</b>	$(\lambda \rightarrow a) \vee b \vee \neg b$	<i>e. g.</i> $r = n - 1$

13	<b><i>LD</i></b>	<b><i>LJ</i> ∩ <b><i>LD</i></b></b>	$a \vee \neg a$	$r = n$
14	<b><i>LJ</i></b>	<b><i>LJ</i> ∩ <b><i>LD</i></b></b>	$\wedge \rightarrow a$	$r = 1$
15	<b><i>LJ</i> ∩ <b><i>LN</i></b></b>	<b><i>LJ</i> ∩ <b><i>LD</i></b></b>	$(\wedge \rightarrow a) \vee b \vee (b \rightarrow c)$	$r = 1$

(As for the correlations of logics in the lines 10–15 under  $L[A]$  and  $L$ , see Miura [3].)

## REFERENCES

- [ 1 ] Curry, H.B., Foundations of mathematical logic (1963), New York.  
 [ 2 ] Gödel, K., Zum intuitionistischen Aussagenkalkül, Akad. Wiss. Anzeiger, vol. **69** (1932), 65–66.  
 [ 3 ] Miura, S., A remark on the intersection of two logics, Nagoya Math. J., vol. **26** (1966), 167–171.  
 [ 4 ] Nagata, S., A series of successive modifications of Peirce's rule, Proc. Japan Acad., vol. **42** (1966), 859–861.  
 [ 5 ] Nishimura, I., On formulas of one variable in intuitionistic propositional calculus, J. Symb. Logic, vol. **25** (1960), 327–331.  
 [ 6 ] Ono, K., On universal character of the primitive logic, Nagoya Math. J., vol. **27-1** (1966), 331–353.  
 [ 7 ] Ono, K., A certain kind of formal theories, Nagoya Math. J., vol. **25** (1965), 59–86.  
 [ 8 ] Umezawa, T., Über die zwischensysteme der Aussagenlogik, Nagoya Math. J., vol. **9** (1955), 181–189.  
 [ 9 ] Umezawa, T., On intermediate propositional logics, J. Symb. Logic, vol. **24** (1959), 20–36.  
 [10] Umezawa, T., On logics intermediate between intuitionistic and classical predicate logic, J. Symb. Logic, vol. **24** (1959), 141–153.

*Toyota Technical College*  
 and  
*Mathematical Institute,*  
*Nagoya University*

**Added in proof** April 28, 1967.

After this paper had been admitted, we found the fact that a series of propositions  $P_i$ , appearing in Nagata [4] and also in this paper, has been introduced by A.S. Troelstra in the slightly different form. (See [11] cited below.) A result of Nagata [4] has been already used in Troelstra [11] in order to verify one of his theorems. Moreover, there are some arguments in [11] connected with ours in the present paper.

[11] Troelstra, A.S., On intermediate propositional logics, Nederl. Akad. Wetensch. Proc. Ser. A, vol. **68** (Indag. Math., vol. **27**) (1965), 141–152.