

COCHRANE CORNER

[†]This review is an abridged version of a Cochrane Review previously published in the *Cochrane Database of Systematic Reviews*, 2014, Issue 1, doi: 10.1002/14651858.CD008298.pub3 (see www.thecochranelibrary.com for information). Cochrane Reviews are regularly updated as new evidence emerges and in response to feedback, and the Cochrane Database of Systematic Reviews should be consulted for the most recent version of the review.

doi: 10.1192/apt.21.2.74

We thank the Cochrane Review Group for their support in publishing these reviews.

See commentary on pp. 75–77, this issue.

[‡]This review is an abridged version of a Cochrane Review previously published in the *Cochrane Database of Systematic Reviews*, Issue 3, doi: 10.1002/14651858.CD008567.pub3 (see www.thecochranelibrary.com for information).

doi: 10.1192/apt.21.2.74a

Physical health care monitoring for people with serious mental illness[†]

Graeme Tosh, Andrew V. Clifton, Jun Xia & Margueritte M. White

Background

Current guidance suggests that we should monitor the physical health of people with serious mental illness, and there has been a significant financial investment over recent years to provide this.

Objectives

To assess the effectiveness of physical health monitoring, compared with standard care for people with serious mental illness.

Search methods

We searched the Cochrane Schizophrenia Group Trials Register (October 2009, update in October 2012), which is based on regular searches of CINAHL, EMBASE, MEDLINE and PsycINFO.

Selection criteria

All randomised clinical trials focusing on physical health monitoring versus standard care, or comparing i) self monitoring versus monitoring by a healthcare professional; ii) simple versus complex monitoring; iii) specific versus non-

specific checks; iv) once only versus regular checks; or v) different guidance materials.

Data collection and analysis

Initially, review authors (GT, AC, SM) independently screened the search results and identified three studies as possibly fulfilling the review's criteria. On examination, however, all three were subsequently excluded. Forty-two additional citations were identified in October 2012 and screened by two review authors (JX and MW), 11 of which underwent full screening.

Main results

No relevant randomised trials which assess the effectiveness of physical health monitoring in people with serious mental illness have been completed. We identified one ongoing study.

Authors' conclusions

There is still no evidence from randomised trials to support or refute current guidance and practice. Guidance and practice are based on expert consensus, clinical experience and good intentions rather than high quality evidence.

Assessed as up to date: 1 Dec 2012

General physical health advice for people with serious mental illness[‡]

Graeme Tosh, Andrew V. Clifton, Jun Xia & Margueritte M. White

Background

There is currently much focus on provision of general physical health advice to people with serious mental illness and there has been increasing pressure for services to take responsibility for providing this.

Objectives

To review the effects of general physical healthcare advice for people with serious mental illness.

Search methods

We searched the Cochrane Schizophrenia Group's Trials Register (last update search October 2012) which is based on regular searches of CINAHL, BIOSIS, AMED, EMBASE, PubMed, MEDLINE, PsycINFO and registries of Clinical Trials. There is no language, date, document type, or publication status limitations for inclusion of records in the register.

Selection criteria

All randomised clinical trials focusing on general physical health advice for people with serious mental illness.

Data collection and analysis

We extracted data independently. For binary outcomes, we calculated risk ratio (RR) and its 95% confidence interval (CI), on an intention-to-treat basis. For continuous data, we estimated the mean difference (MD) between groups and its 95% CI. We employed a fixed-effect model for analyses. We assessed risk of bias for included studies and created 'Summary of findings' tables using GRADE.

Main results

Seven studies are now included in this review. For the comparison of physical healthcare advice versus standard care we identified six studies (total $n=964$) of limited quality. For

measures of quality of life one trial found no difference ($n=54$, 1 RCT, MD Lehman scale 0.20, CI -0.47 to 0.87, very low quality of evidence) but another two did for the Quality of Life Medical Outcomes Scale - mental component ($n=487$, 2 RCTs, MD 3.70, CI 1.76 to 5.64). There was no difference between groups for the outcome of death ($n=487$, 2 RCTs, RR 0.98, CI 0.27 to 3.56, low quality of evidence). For service use two studies presented favourable results for health advice, uptake of ill-health prevention services was significantly greater in the advice group ($n=363$, 1 RCT, MD 36.90, CI 33.07 to 40.73) and service use: one or more primary care visit was significantly higher in the advice group ($n=80$, 1 RCT, RR 1.77, CI 1.09 to 2.85). Economic data were equivocal. Attrition was large (> 30%) but similar for both groups ($n=964$, 6 RCTs, RR 1.11, CI 0.92 to 1.35). Comparisons of one type of physical healthcare advice with another were grossly underpowered and equivocal.

Authors' conclusions

General physical health could lead to people with serious mental illness accessing more health services which, in turn, could mean they see longer-term benefits such as reduced mortality or morbidity. On the other hand, it is possible clinicians are expending much effort, time and financial resources on giving ineffective advice. The main results in this review are based on low or very low quality data. There is some limited and poor quality evidence that the provision of general physical healthcare advice can improve health-related quality of life in the mental component but not the physical component, but this evidence is based on data from one study only. This is an important area for good research reporting outcome of interest to carers and people with serious illnesses as well as researchers and fundholders.

Assessed as up to date: 26 Oct 2013