

---

*Contents—continued from outside back cover]*

|                                |   |
|--------------------------------|---|
|                                | <i>Book Reviews</i>   |
| 233                            | Emerging technologies for integrated pest management: Concepts, research, and implementation                          |
| 235                            | An introduction to ecological integrated lepidopteroid rice pest management in North Vietnam (Lepidoptera: Pyralidae) |
| <i>Erratum</i>                 | i   |
| <i>Information for authors</i> | iii–iv  |
| <i>Author checklist</i>        | v   |
| <i>Announcement</i>            | vii   |

---

Cover illustration by N. M. Komeri from a sketch by W. A. Overholt:

*Cotesia flavipes* Cameron, an important larval parasitoid of stemborers of maize and sugarcane, with borer.  
Insects not drawn to scale.

# INSECT SCIENCE AND ITS APPLICATION

The International Journal of Tropical Insect Science

Volume 20 Number 3

2000

## CONTENTS

|   |     |  |
|---|-----|--|
| M. WALE, B. JEMBERE and E. SEYOUN                                   | 171 | <i>Research Articles</i><br>Biology of the pea aphid, <i>Acyrtosiphon pisum</i> (Harris) (Homoptera: Aphididae) on cool-season legumes                                 |
| A. J. NGI-SONG, P. G. N. NJAGI,<br>B. TORTO and W. A. OVERHOLT      | 181 | Identification of behaviourally active components from maize volatiles for the stemborer parasitoid <i>Cotesia flavipes</i> Cameron (Hymenoptera: Braconidae)          |
| S. EL-GENGAIHI, N. Z. DIMETRY,<br>S. A. A. AMER and S. M. MOHAMED   | 191 | Acaricidal activity of lipoidal matter of different plant extracts against the two-spotted spider mite <i>Tetranychus urticae</i> Koch.                                |
| J. SRIKANTH, S. EASWARAMOORTHY,<br>R. KUMAR and M. SHANMUGASUNDARAM | 195 | Patterns of laboratory parasitisation rates of <i>Cotesia flavipes</i> Cameron (Hymenoptera: Braconidae) in borers of sugarcane and sorghum                            |
| T. VERMA and S. P. SINGH  | 203 | Multiple resistance in forage sorghum hybrids to the sorghum shootfly, <i>Atherigona soccata</i> (Rondani) and the spotted stemborer, <i>Chilo partellus</i> (Swinhoe) |
| A. R. SHAKOORI, K. A. MUJEEB,<br>S. MAQBOOL and S. S. ALI           | 207 | Relative activity of various esterases in six Pakistani strains of the lesser grain borer, <i>Rhyzopertha dominica</i> (Fabricius)                                     |
| G. A. OMOOGUN and<br>O. A. AKINBOADE                                | 215 | <i>Short Communications</i><br>Tsetse and bovine trypanosomosis incidence at Egbe in the derived savanna zone of Nigeria   |
| B. DONDJI, D. D. DUHLINSKA<br>and A. SAME-EKOBO                     | 221 | Species composition of the phlebotomine sandfly fauna (Diptera, Phlebotominae) in Mokolo Region, northern Cameroon   |
| F. N. BALIRAINA, E. O. OSIR,<br>S. B. OBUYA and F. J. MULAA         | 227 | Protein polymorphism in two populations of the brown ear tick, <i>Rhipicephalus appendiculatus</i> Neumann (Acari: Ixodidae)   |

[Continued on inside back cover]

Indexed/Abstracted in  
Current Contents, C. A. B. Abstracts, BIOSIS, Zoological Record, Chemical Abstracts,  
The African Book Publishing Record, TROPAG & RURAL Abstracts and IBIDS database

ISSN 0191-9040  
ISIADL 20(3) 171-235 (2000)



Published and printed by ICIPE Science Press  
P.O. Box 72913, Nairobi, Kenya

Available on-line at <http://www.bdt.org.br/bioline/tl>