

CORRIGENDUM

Schistosoma mansoni cercariae experience
influx of macromolecules during skin
penetration – CORRIGENDUM

J. THORNHILL, P. M. Z. COELHO, P. McVEIGH, A. MAULE, A. D. JURBERG
and J. R. KUSEL

doi:10.1017/S0031182009990692, Published by Cambridge University Press,

September 2009, Vol. 136 (11).

In the article by Thornhill *et al.* (2009), the reference ‘Ribeiro, A. C., Maldonado Jr, A., D’Andrea, P. S., Vieira, G. O. and Rey, L. (1998). Susceptibility of *Nectomys rattus* (Pelzen, 1883) to experimental infection with *Schistosoma mansoni* (Sambon, 1907): a potential reservoir in Brazil. *Memorias do Instituto Oswaldo Cruz* **93** (Suppl.1), 295–299’ should be replaced by ‘Ribeiro, F., Coelho, P. M., Vieira, L. Q., Powell, K. and Kusel, J. R. (1998). Membrane internalisation processes in different stages of *Schistosoma mansoni* as shown by styryl dye (Frei Mao 1–43). *Parasitology* **116**, 51–59.’

REFERENCE

Thornhill, J. et al. (2009). *Schistosoma mansoni* cercariae experience influx of macromolecules during skin penetration. *Parasitology* **136**, 1257–1267.