

Research Article

Cite this article: Santangeli A, Fozzi I, De Rosa D, Aresu M, Campus A, Muzzeddu M, Secci D, Cerri J, Berlinguer F (2024). Quantifying the regulation and cultural ecosystem services associated with Griffon Vultures *Gyps fulvus* in Sardinia, Italy. *Bird Conservation International*, **34**, e35, 1–7
<https://doi.org/10.1017/S0959270924000327>

Received: 01 May 2024

Revised: 25 August 2024

Accepted: 09 September 2024

Keywords:





Avian scavengers; Carcass disposal; Cultural ecosystem services; Human–nature relationship; Nature contribution to people

Corresponding author:

Andrea Santangeli;

Email: andrea.santangeli@gmail.com

Quantifying the regulation and cultural ecosystem services associated with Griffon Vultures *Gyps fulvus* in Sardinia, Italy

Andrea Santangeli^{1,2} , Ilaria Fozzi³ , Davide De Rosa³ , Mauro Aresu⁴, Alfonso Campus³, Marco Muzzeddu⁵, Dionigi Secci⁵, Jacopo Cerri³ and Fiammetta Berlinguer³ 

¹Animal Demography and Ecology Unit, Institute for Mediterranean Studies (IMEDEA), CSIC-UIB, 07190, Esporles, Spain; ²FitzPatrick Institute of African Ornithology, University of Cape Town, Cape Town, South Africa; ³Department of Veterinary Medicine, University of Sassari, Sassari 07100, Italy; ⁴Via Crispi 5, 08015 Macomer, Italy and ⁵Agenzia Forestale Regionale per lo Sviluppo del Territorio e l'Ambiente della Sardegna (FoReSTAS), Cagliari 09123, Italy

Summary

In the Anthropocene, recognising nature's role in human well-being is pivotal for biodiversity conservation. Despite their significance, knowledge gaps persist regarding ecosystem services, even for well-studied species like vultures. Our study focuses on the Griffon Vulture *Gyps fulvus* in Sardinia, Italy, exploring their cultural and regulating services, including carcass disposal and resulting greenhouse gas (GHG) mitigation. Through surveys of natural reserve visitors and data on carcass provision and GHG emissions, we assess public perception, economic value, and environmental impacts associated with vultures. The public perception of Griffon Vultures is predominantly positive, with a strong acknowledgment of their role in disease prevention and carcass disposal, highlighting their contribution to regulation services. Furthermore, vultures are widely recognised as a key element characterising the agropastoral landscapes of Sardinia, underscoring their cultural importance. The economic evaluation, through willingness to pay for vulture-watching and photography opportunities, indicates a significant appreciation of these birds, with almost three-quarters of respondents willing to pay an entrance fee at vulture observation sites. We also show that supplanting the disposal role of vultures at studied feeding sites (during 2017–2022) would result in the emission of 96 tons of CO₂ equivalent, which highlights the critical role of vultures in climate mitigation. This study not only sheds light on the ecological and cultural significance of Griffon Vultures in Sardinia but also underscores the economic and environmental benefits of their conservation. It emphasises the need for continued efforts in vulture conservation, integrating ecological, cultural, and economic perspectives to foster a sustainable coexistence between humans and wildlife.

Introduction

Ecosystem services (i.e. nature's contribution to people) are nowadays recognised as a main pillar for biodiversity conservation because of their profound implications for environmental management and policy (Daily and Matson 2008; Geijzenborffer et al. 2017; IPBES 2019). Typically, ecosystem services refers to the various benefits that humans obtain from ecosystems, which can be categorised into three main groups: provisioning, regulation/maintenance, and cultural (Haines-Young and Potschin-Young 2018). Provisioning services are direct products obtained from ecosystems, such as food, water, raw materials, and medicinal plants, which fulfil human needs for sustenance and economic activities. Regulation and maintenance services involve the ecological processes that contribute to the overall balance and stability of ecosystems. These services include climate regulation, water purification, pollination, and disease control, which ensure the health and resilience of ecosystems. Cultural services encompass the non-material benefits that ecosystems provide to humans, including aesthetic, spiritual, educational, and recreational values. These services contribute to the cultural identity, inspiration, and overall well-being of humans.

Birds play a vital role in maintaining ecological balance, conserving biodiversity, and enhancing human well-being (Sekercioglu et al. 2016). As such, birds are often associated with key ecosystem services, including pollination for plant reproduction, natural pest control by feeding on insects, and seed dispersal that supports biodiversity and ecosystem regeneration. Birds also hold a cultural value, symbolising freedom, beauty, and connection to nature, contributing to recreational opportunities and ecotourism (Becker et al. 2005; García-Jiménez et al. 2021; Gupta et al. 2020; Moleón et al. 2014). Their role in controlling insect-borne diseases, such as

mosquitoes, benefits both wildlife and humans (Sekercioglu et al. 2016). Additionally, the presence of birds in the environment enriches the cultural heritage through folklore, art, and traditional practices (Tidemann and Gosler 2010).

Old and New World vultures play a crucial role in maintaining the ecosystem's health (Berlinguer et al. 2021; Craig et al. 2018; Gangoso et al. 2013; Grilli et al. 2019; Morales-Reyes et al. 2018; Mateo-Tomás et al. 2017; Plaza et al. 2020; Santangeli et al. 2019). Being the only obligate scavengers among vertebrates, they efficiently remove carrion, curbing disease spread and reducing the potential for pathogen proliferation impacting both wildlife and domestic animals (Berlinguer et al. 2021; Plaza et al. 2020). Beyond disease control, vultures contribute to waste reduction by consuming animal remains, and therefore prevent the release of greenhouse gases (GHG) and aid nutrient cycling (Morales-Reyes et al. 2015). In addition to their ecological services, vultures hold cultural significance, associated with death and rebirth in some cultures, enriching the spiritual identity of communities (Van Dooren 2012). These birds are also central to ecotourism in several regions, providing economic benefits to local communities and raising awareness about vulture conservation and their broader ecological services (Aguilera-Alcalá et al. 2020; Becker et al. 2005; García-Jiménez et al. 2021). Finally, by being apex scavengers, vultures regulate population dynamics across the food chain, influencing the structure of ecological communities and contributing to ecosystem stability (Morales-Reyes et al. 2017; Ogada et al. 2012). Understanding and conserving these species is essential for maintaining the ecological balance and functioning of their habitats. In the agropastoral landscapes of Europe, including Sardinia, the Griffon Vulture *Gyps fulvus* is the most abundant avian scavenger, providing an irreplaceable scavenging service through its efficient carcass consumption (Berlinguer et al. 2021; Oliva-Vidal et al. 2022a).

Research on vultures and their associated ecosystem services has witnessed a substantial growth, with numerous studies shedding light on their vital roles in disease control, waste reduction, and cultural significance (Carucci et al. 2022; Plaza and Lambertucci 2022; van den Heever et al. 2021). However, despite this progress, significant gaps still persist in our understanding of ecosystem services associated with these species. For example, there are no studies, to the best of our knowledge, on how vultures mitigate GHG emissions on Mediterranean islands. The complex and context-dependent nature of the ecosystem services provided by vultures underscores the importance of studying them at the local to regional scale. Such studies are essential for comprehensively assessing the specific services of vultures in diverse ecosystems, enabling targeted conservation strategies, and ensuring the sustained provision of these critical services. Closing these knowledge gaps is crucial for the effective conservation and management of vulture populations and to leverage the potential of vultures in providing key services that are so fundamental for modern societies. This also applies to the agropastoral communities of Sardinia, and the rest of Europe, that live on marginal landscapes, such as steppes, whose conservation is often neglected but are crucial for sustainable development and biodiversity conservation. In such landscapes, Griffon Vultures play a role in carcass decomposition, with potential implications for disease control, saving costs to livestock farmers for artificial disposal, as well as savings in GHG emissions from the process. Moreover, local and foreign visitors to areas within the vulture range may have an interest in watching, photographing or simply contributing to the species conservation (García-Jiménez et al. 2021, 2022; Oliva-Vidal et al. 2022a). Therefore, in this study we aim to fill the above knowledge gap in the ecosystem service

provision by vultures in Sardinia. Specifically, we here aim: (1) to quantify key cultural services associated with Griffon Vultures in Sardinia, specifically focused on the perceptions of ecotourists towards vultures and their interest and willingness to pay for watching vultures and for their conservation; (2) to quantify key regulation services associated with Griffon Vultures in Sardinia, specifically focused on waste decomposition and associated climate regulation and opportunity cost services. Essentially, how much animal waste they dispose, and what are the net savings in GHG emissions and money due to the natural waste disposal by vultures compared with carcass transport and incineration.

Methods

Perceptions on vultures and associated ecosystem services

Between August 2022 and October 2023, we quantified how visitors at eight protected areas in Sardinia ($n = 116$) perceived Griffon Vultures and their associated ecosystem services, by means of a structured questionnaire survey. Questionnaires (see Supplementary material [appendix](#) and some broad details below) were completed on-site ($n = 100$) or filled in online ($n = 16$). We employed these two different approaches for the questionnaire delivery to achieve a larger sample size. While we combined data from both online and on-site-filled questionnaires, the results were largely the same when the online-filled questionnaires were excluded. The questionnaire was divided into specific sections asking questions related to: the frequency of visits to Sardinia and its natural reserves; willingness to pay for observing, photographing or conserving vultures (using pre-defined numerical amounts); the ecosystem services that were perceived as being delivered by vultures (using a five-point bipolar scale from strongly disagree to strongly agree); the demographic attributes of respondents (see SM appendix). In compliance with the General Data Protection Regulation of the EU (<https://gdpr.eu/>), informed consent was collected at the beginning of the questionnaire. Participants were informed about the scope and objectives of the study, that participation was totally voluntary, and that individual anonymity and confidentiality were ensured.

Regulation services

We calculated the amount of GHG emissions, in CO₂ equivalent, resulting from the artificial disposal of all carcasses that had been disposed and consumed by vultures at supplementary feeding stations in north-western Sardinia during 2017–2022. That is, what would have been the GHG emissions of those carcasses without their consumption by vultures. To address this question, we tracked the artificial carcass disposal process from the moment the animal dies at the farm, to the point when it is incinerated, by following the framework of Morales-Reyes et al. (2015). Namely, we separated the process into three steps: (1) carcass transport to an intermediate accumulation plant; (2) transport from the accumulation plant to the incineration plant; (3) incineration at the end of the process (Figure 1).

In step 1, we assumed that one small truck (7.5 tons) would collect a carcass at each farm which has a supplementary feeding station and assumed that if multiple carcasses become available in the same day at the same farm, these are jointly collected within a single trip. We calculated the distance (km), from each farm to the intermediate plant, located in the municipality of Bitti (central Sardinia). This distance was calculated using Google Maps. We assumed the consumption of the truck to be 0.21 l/km on a diesel



Figure 1. Schematic representation of the natural versus artificial disposal process for livestock carcasses in central-western Sardinia. Top left figure represents the landscape where the livestock dies (farm locations depicted as green circles on the map) and requires disposal, either via the natural scavenger decomposition (the Griffon Vulture *Gyps fulvus* role; green area in the figure) or through artificial disposal (grey area). The latter process involves three different steps (locations marked as red circles in the map). The carcass, as it becomes available, is collected from the farm and transported to an intermediate plant (step 1). At this intermediate plant, carcasses are stored and piled up to fill a large truck that then brings them to mainland Italy (step 2), where they are incinerated at a specific plant (step 3).

engine, with 2.7 kg CO₂ emitted per litre of diesel fuel consumed (Morales-Reyes et al. 2015). Then we used historical data on daily carcass provisions for each supplementary feeding station to obtain the total mileage of all trips, and we used the conversion factors mentioned above to estimate the total CO₂ emissions associated with carcass collection. Indeed, the total mileage was doubled, as we assumed that the truck always made a return trip on each collection journey.

In step 2, based on available evidence about carrion disposal, we estimated CO₂ emissions by assuming that biomass is moved from Sardinia to peninsular Italy via trucks (approximately 538 km) and ferry transport (approximately 313 km). In Sardinia, we assumed that carrion is moved on the road network from the intermediate centre in Bitti to the harbour of Olbia. Then the truck is shipped across the Tyrrhenian Sea, from Olbia to Livorno, on a ferry. Finally, the truck would deliver the biomass from the harbour of Livorno to the incineration plant in Bologna. For this task, usually large trucks (24 tons), operating at a full load, are used. It was considered that the carrion disposed at supplementary feeding stations in 2017–2022 amounted to 95.7 tons, we estimated that four trips would have been required to deliver the biomass from Bitti to the incinerating plant. In terms of emissions, following Morales-Reyes et al. (2015), we also assumed that a large truck with a diesel engine would consume 0.26 l/km, with 2.7 kg CO₂ emitted per litre of diesel fuel consumed. For the ferry, an average emission

factor of 25 g CO₂ per ton-km was assumed based on EU maritime transport data (www.data.europa.eu/data/datasets/co2-emissions-data). Emissions were doubled to account for the return trip of the truck to Sardinia.

We calculated CO₂ emissions from step 3, the artificial carcass disposal process, by assuming a rendering procedure, where high-temperature incineration is used, which has a conversion factor of 200 kg CO₂ equivalent per each ton of carcass burned (Plaza and Lambertucci 2022).

By summing up the emissions from each of the three steps, we quantified the total CO₂ emissions equivalent related to the artificial disposal of the carcasses in a scenario where vultures are absent from Sardinia to consume this biomass.

Next, we calculated the total amount of potential CO₂ emissions mitigated by the whole Griffon Vulture population in Sardinia, per year. We used the total population estimate from the year 2022 (316–338 individuals; Berlinguer et al. 2022), each consuming 208–305 kg biomass/year (Plaza and Lambertucci 2022), amounting to a total annual biomass consumed by the entire population as 65.6–103.1 tons. We then converted the biomass to potential CO₂ emissions assuming that all that material would be disposed of artificially, by using the ratio of emissions from the tons of carcass as calculated in the steps above (which is very close to one, that is, 1 ton of biomass would emit 1 ton of CO₂ equivalent).

Finally, we calculated the economic opportunity costs (in euros) for all carcasses consumed by vultures at supplementary feeding stations in north-western Sardinia during 2017–2022. Essentially, how much it would have cost if the above carcasses were to be disposed of artificially, as in Figure 1. For this, we used the total sum of biomass (kg) disposed of at the supplementary feeding stations as above, separated by three livestock categories: cattle, horses/donkeys, and sheep/goats. We obtained the maximum cost (in €/kg biomass) of collection, transport, and incineration of carcasses by each of the three livestock categories from the Official Bulletin of the Italian Republic (series n. 123, date: 27 May 2023) for the Sardinia region in 2023 (as an average across the region). Total maximum disposal costs were reported as: €0.97/kg for cattle, €1.07/kg for horses/donkeys, and €3.08/kg for sheep/goats. We then multiplied the total biomass (kg) for the cost of disposal per each livestock category, and then summed the three to yield the total opportunity cost for disposing of those carcasses.

Results

Socio-demographics of respondents

Overall, 116 respondents filled in the questionnaire about the perception of vultures and their associated ecosystem services. This sample was balanced across genders, with 58 male and 57 female respondents, and across various age classes: 28 respondents aged 18–30 years old, 30 aged 31–40, 25 aged 41–50, and 29 aged >51. Moreover, the majority of respondents had a high level of education, with 45 having at least a high school degree, 42 having some basic university degree, and 23 with a higher than basic university degree (e.g. MSc or PhD). As expected, most respondents ($n = 91$, 79% of the total sample) were Italian, while the remaining respondents came from Germany and France (seven for each), Czech Republic (four), and Netherlands, Austria, and Poland (two for each). Moreover, 57 respondents (50% of the 114 that replied to this question) were resident in Sardinia, while the others were visiting the island for the first time (24) or had already visited Sardinia on two (10), three or

more previous occasions (6), or visited the island every year (17). Only eight respondents stated that they have a profession somewhat related to nature (e.g. biologist, nature guide, veterinarian, naturalist), while others had different private social or government employment jobs, likely unrelated to nature and its conservation. The sample of respondents was rather balanced among those that enjoy photography (51, amounting to 45% of the total; not necessarily nature related) and those that do not practise this activity (63, 55% of the total). Among all respondents who replied to the question about the frequency with which they visited a protected area, 30 had visited for the first time, 44 visited once a year, 8 once a month, and 9 once a week.

Perceptions on vultures' ecosystem services

With regard to the respondents' perceptions towards vultures and their associated ecosystem services, the results are rather clear that respondents had a broadly positive perception of vultures and were also aware of some of their main ecosystem services (Figure 2). Specifically, most respondents (almost two out of three) believed that vultures can counteract the spread of diseases, and reduce costs associated with carcass disposal, e.g. due to carcass transport and incineration (regulation services). Moreover, most respondents agreed that vultures belong to the agropastoral traditional ecological landscape of Sardinia (cultural service). Even more clear was the respondents' perception towards vultures, whereby most (71%) of the respondents did not agree that vultures are dirty, and an even higher percentage (78%) of respondents did not agree with the statement affirming that they do not like to have vultures around, i.e. they like to have vultures in the surrounding landscape.

Cultural ecosystem services

We also attempted to quantify the willingness to pay for observing and photographing vultures, as a monetary quantification of their potential cultural ecosystem service. Specifically, to the question whether respondents would be willing to visit an observation hide




Figure 2. Perceptions of visitors to natural reserves on vultures and their associated ecosystem services. The top two bars represent regulation and maintenance services, whereas the bottom three bars represent cultural services.

for watching and photographing vultures, 92 (79%) responded positively, 10 negatively, and 14 don't know. Similarly, 85 (73%) would be willing to pay an entrance fee to access such a hide. Among the latter, the average they would pay was €13 each. However, there was a large variation in the amount respondents would be willing to pay, with 54 respondents willing to pay only €10 or less, and the rest a larger sum. The results were similar with regards to willingness to join a guided visit to such a vulture observation place (92, 80% of 115 responses), while 89 (79% of 112 responses) were willing to pay an entrance fee for such a place. Among the latter, the average they would pay is €15 each, with 46 respondents willing to pay only €10 or less, 23 respondents willing to pay €20, and 9 willing to pay €30 or more. Over half (55) of the respondents stated that they would visit the vultures observation hide only as part of an entire trip to the island, whereby 34 respondents would visit it on purpose. In more general terms, about half of the respondents (58 of 114 who replied) were willing to make a donation for the conservation of the Griffon Vulture in Sardinia, amounting to an average of €21 per person. Specifically, 29 of the 51 respondents willing to support vulture conservation would donate €10 or less, 6 would donate €20, and the remaining 16 would donate €30 or more.

Regulation services

During the period 2017–2022, a total of 30 supplementary feeding stations in north-central Sardinia provisioned 95.7 tons of livestock (mainly sheep, goats, and cattle) carcasses during 609 provisioning events (each event could include from one to seven animal carcasses). These carcasses were almost entirely consumed by the local Griffon Vulture population. The CO₂ equivalent that would have been emitted into the atmosphere from the artificial disposal of such a total amount of biomass amounts to 96.07 tons, according to the calculations and related assumptions as detailed in the methods. Out of the total CO₂ emissions from the full process of artificial disposal of the above biomass, almost three-quarters (77%, 73.9 tons) of the emissions were due to local carcass transport (from the farm to the intermediate plant), only 3% (3.0 tons) were due to long haul carcass transport (from the intermediate plant in Sardinia to the final incineration plant in mainland Italy), and one fifth (20%, 19.1 tons) of the total CO₂ was emitted during the incineration phase.

Extrapolating the above CO₂ emissions to the total annual potential biomass consumed by the entire population of Griffon Vultures in Sardinia would amount to 65.9–103.5 tons of CO₂ equivalent per year (minimum and maximum estimates based on minimum and maximum vulture food intake and population estimate).

The economic opportunity costs that are saved by the disposal service of the vultures at the supplementary feeding stations are also substantial and equate to €134,016 for the period 2017–2022 at the 30 supplementary feeding stations in north-central Sardinia.

Discussion

To the best of our knowledge, this study is the first one to explore the public perception, magnitude, and economic value of ecosystem services delivered by Griffon Vultures on a Mediterranean island. We acknowledge that the study has some limitations, including the representativeness of the pool of respondents for this study, as well as the uncertainty in the economic and GHG emission savings

associated with the vultures' scavenging services. Nevertheless, we believe that the figures provided are broadly representative of the local situation based on a few key facts: i.e. it was a local study of a restricted area, which reduced spatial heterogeneity while allowing detailed information on carcass availability, carcass disposal, and vulture population numbers.

Our study revealed a broadly positive public perception of the ecosystem services associated with Griffon Vultures in Sardinia. Most respondents acknowledged the role played by vultures in preventing disease and saving costs associated with carcass disposal. Economic aspects are highlighted by the willingness of 73% of respondents to pay for vulture-watching, with an average proposed fee of €13, indicating the potential cultural value of these birds, also in economic terms. Additionally, about half of the respondents were willing to donate money for conserving vultures, averaging €21. Moreover, from an ecological perspective, vultures play a role in GHG mitigation. Between 2017 and 2022, vultures contributed to preventing 96 tons of CO₂ equivalent that would have been emitted from artificial carcass disposal, and a potential annual CO₂ mitigation of 66–103 tons for the whole vulture population in Sardinia. During the same period, the economic savings from carcass disposal by vultures at supplementary feeding stations was substantial, reaching well over €100,000.

The positive perception of Griffon Vultures among resident and foreign visitors to natural sites in Sardinia is encouraging and aligns with recent studies on the positive cultural value associated with vultures in modern society (Craig et al. 2018; Carucci et al. 2022; García-Jiménez et al. 2022; Santangeli et al. 2016). These findings underscore the deep integration of vultures into the cultural and ecological landscape of Sardinia. The acceptance and appreciation of vultures are crucial for conservation efforts, as public perception often drives policy and funding support (Oliva-Vidal et al. 2022a).

In turn, the positive perception of vultures is reflected in the visitor's willingness to pay for observing and photographing these animals. This underscores a great ecotourism potential especially related to those local livestock farms interested in managing a supplementary feeding station that would bring a financial revenue, e.g. through entrance fees for visiting the site. Indeed such cultural services related to vultures bringing local financial revenues have been quantified, at least for a few sites (Becker et al. 2005; García-Jiménez et al. 2021). These activities not only contribute to local economies but also foster a deeper appreciation for wildlife. The monetary valuation of these services is important for conservation strategies, demonstrating that protecting vultures aligns with local economic interests. Moreover, the variation in the willingness to pay suggests a diverse range of values and interests among different visitor demographics, which could be targeted by future tourism and conservation strategies. We believe that future studies should also explore how local livestock farm owners would be available to develop activities associated with ecotourism. This would allow a better understanding of which conditions the demand and offer for observing and photographing vultures would align. In turn, this could also allow the development of vulture sighting activities that would maximise the value of this cultural service for both tourists and farmers, while minimising the potential disturbance to vultures (Donázar et al. 2023).

Aside from the above cultural services, we also show that the local population of Griffon Vultures in Sardinia can contribute to important regulation/mitigation services through the carcass disposal that, if artificially performed, would result in GHG emissions and also economic costs. To place this finding in context, the

net annual savings in CO₂ emissions resulting from the total population of Griffon Vultures in Sardinia equate to the annual CO₂ emissions of 16–34 households in Sardinia. Previous studies have quantified the role of vultures in preventing GHG emissions via their scavenging role, at a global as well as national level (Morales-Reyes et al. 2015; Plaza and Lambertucci 2022). By quantifying the CO₂ equivalent emissions prevented by vulture scavenging, we contribute another compelling case for the conservation of these birds, whose ecological role spans broad contexts of global environmental challenges (Santangeli et al. 2024). Moreover, we believe that translating reductions of GHG emissions into tangible examples, such as household CO₂ emissions, could be useful for communication initiatives targeting both residents and visitors in Sardinia, with the goal of increasing overall awareness of the important ecosystem services associated with vultures. To this end, future studies should use approaches from the social sciences to test for the effectiveness of similar messaging through, for example, information panels at protected areas (Abrams et al. 2020).

Interestingly, we show that, in the context of Sardinia, over three-quarters of the emissions of the full artificial disposal process are related to local carcass collection and transport, while only one-fifth of emissions are related to the incineration process. This finding is relevant, as it indicates that local carcass transport plays a key role in CO₂ emission mitigation scenarios related to vultures. For example, a recent study quantified the global emission mitigation relative to the incineration of all biomass that is currently consumed by all vulture populations in the world as 3–61 Tg CO₂ eq. (Plaza and Lambertucci 2022). That value did not consider the emissions related to carcass transport. Our findings indicate that global values could be up to five times higher, underscoring the importance of considering the full carcass collection, transport, and incineration process when estimating emission and cost mitigation.

Overall, the findings on perception and economic valuation of vultures' ecosystem services have important implications for conservation efforts. They underscore the potential benefits these birds could bring to local peoples and their economies. They also highlight the need for continued education and awareness programmes to maintain and enhance public support for vulture conservation. This study adds important insights into our understanding of the Griffon Vulture's role for the ecosystems of Sardinia, with far-reaching implications across other Mediterranean islands and beyond where vulture populations occur. It provides a compelling case for their conservation, not only for their ecological benefits, but also for their cultural and economic services. This holistic understanding is vital for developing effective bottom-up conservation strategies that are supported by the public and aligned with sustainable development goals. As the Griffon Vulture plays a key and irreplaceable scavenging role in European agropastoral systems (Oliva-Vidal et al. 2022b), its conservation is paramount to keeping such ecosystems in balance, in Sardinia and beyond. It is now clear that the consequences of vulture declines could be severe, for both the ecosystem and human societies (Frank and Sudarshan 2024). This further reinforces the urgency and relevance of conserving healthy vulture populations locally and globally.

Supplementary material. The supplementary material for this article can be found at <http://doi.org/10.1017/S0959270924000327>.

Acknowledgements. We thank all those who helped us with monitoring and management activities within the LIFE “Under Griffon Wings” project. The study was funded by the European Commission through the Life Under

Griffon Wings project (LIFE14/NAT/IT/000484) and the Life Safe for Vultures project (LIFE19 NAT/IT/000732). The present research was carried out within the framework of the activities of the Spanish Government through the “Maria de Maeztu Centre of Excellence” accreditation to IMEDEA (CSIC-UIB) (CEX2021-001198). AS acknowledges support from the European Commission through the Horizon 2020 Marie Skłodowska-Curie Actions individual fellowships (Grant no. 101027534). We also thank two anonymous referees for their valuable comments on an early version of the article.

References

- Abrams K.M., Leong K., Melena S. and Teel T. (2020). Encouraging safe wildlife viewing in national parks: Effects of a communication campaign on visitors' behavior. *Environmental Communication* **14**, 255–270.
- Aguilera-Alcalá N., Morales-Reyes Z., Martín-López B., Moleón M. and Sánchez-Zapata J.A. (2020). Role of scavengers in providing non-material contributions to people. *Ecological Indicators* **117**, 106643. <https://doi.org/10.1016/j.ecolind.2020.106643>
- Becker N., Inbar M., Bahat O., Choresh Y., Ben-Noon G. and Yaffe O. (2005). Estimating the economic value of viewing griffon vultures *Gyps fulvus*: A Travel Cost Model study at Gamla Nature Reserve, Israel. *Oryx* **39**, 429–434.
- Berlinguer F., Ahmed F., Tamponi C., Carta S., Scala A., Cappai M.G. et al. (2021). Help from the sky: Can vultures contribute to Cystic Echinococcosis control in endemic areas? *PLOS Neglected Tropical Diseases* **15**, e0009615. <https://doi.org/10.1371/journal.pntd.0009615>
- Berlinguer F., Campus A., De Rosa D. and Aresu M. (2022). AZIONE D.5 – Censimento Annuale del Grifone (*Gyps fulvus*) in Sardegna. Terzo Report.
- Carucci T., Whitehouse-Tedd K., Yarnell R.W., Collins A., Fitzpatrick F., Botha A. et al. (2022). Ecosystem services and disservices associated with vultures: A systematic review and evidence assessment. *Ecosystem Services* **56**, 101447. <https://doi.org/10.1016/j.ecoser.2022.101447>
- Craig C.A., Thomson R.L. and Santangeli A. (2018). Communal farmers of Namibia appreciate vultures and the ecosystem services they provide. *Ostrich* **89**, 1–10.
- Daily G.C. and Matson P.A. (2008). Ecosystem services: From theory to implementation. *Proceedings of the National Academy of Sciences of the United States of America* – PNAS **105**, 9455–9456. <https://doi.org/10.1073/pnas.0804960105>
- Donázar J.A., Cortés-Avizanda A., Arrondo E., Delgado-González A. and Ceballos O. (2023). Hidden effects of high numbers of tourists in protected areas: displacement of foraging top scavengers. *Ibis* **165**, 305–311. <https://doi.org/10.1111/ibi.13121>
- Frank E. and Sudarshan A. (2024). The social costs of keystone species collapse: Evidence from the decline of vultures in India. *American Economic Review* **114**, 3007–3040.
- Gangoso L., Agudo R., Anadón J.D., de la Riva M., Suleyman A.S., Porter R. et al. (2013). Reinventing mutualism between humans and wild fauna: insights from vultures as ecosystem services providers. *Conservation Letters* **6**, 172–179. <https://doi.org/10.1111/j.1755-263X.2012.00289.x>
- García-Jiménez R., Morales-Reyes Z., Pérez-García J.M. and Margalida A. (2021). Economic valuation of non-material contributions to people provided by avian scavengers: Harmonizing conservation and wildlife-based tourism. *Ecological Economics* **187**, 107088. <https://doi.org/10.1016/j.ecolecon.2021.107088>
- García-Jiménez R., Pérez-García J.M., Margalida A. and Morales-Reyes Z. (2022). Avian scavengers' contributions to people: The cultural dimension of wildlife-based tourism. *Science of the Total Environment* **806**, 150419. <https://doi.org/10.1016/j.scitotenv.2021.150419>
- Geijzendorffer I.R., Cohen-Shacham E., Cord A.F., Cramer W., Guerra C. and Martín-López B. (2017). Ecosystem services in global sustainability policies. *Environmental Science & Policy* **74**, 40–48. <https://doi.org/10.1016/j.envsci.2017.04.017>
- Grilli M.G., Bildstein K.L. and Lambertucci S.A. (2019). Nature's clean-up crew: Quantifying ecosystem services offered by a migratory avian scavenger on a continental scale. *Ecosystem Services* **39**, 100990. <https://doi.org/10.1016/j.ecoser.2019.100990>
- Gupta U., Qureshi Q. and Kumar N. (2020). Folk perceptions for avian scavengers in a tropical megacity: implications for biocultural conservation. SocArXiv Papers. <https://doi.org/10.31235/osf.io/tmdv4>

- Haines-Young R. and Potschin-Young M. (2018). Revision of the Common International Classification for Ecosystem Services (CICES V5.1): A policy brief. *One Ecosystem* **3**, e27108. <https://doi.org/10.3897/oneeco.3.e27108>
- Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) (2019). *Summary for Policymakers of the Global Assessment Report on Biodiversity and Ecosystem Services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*. Bonn: IPBES Secretariat.
- Mateo-Tomás P., Olea P.P., Moleón M., Selva N. and Sánchez-Zapata J.A. (2017). Both rare and common species support ecosystem services in scavenger communities. *Global Ecology and Biogeography* **26**, 1459–1470. <https://doi.org/10.1111/geb.12673>
- Moleón M., Sánchez-Zapata J.A., Margalida A., Carrete M., Owen-Smith N. and Donazar J.A. (2014). Humans and scavengers: The evolution of interactions and ecosystem services. *Bioscience* **64**, 394–403. <https://doi.org/10.1093/biosci/biu034>
- Morales-Reyes Z., Martín-López B., Moleón M., Mateo-Tomás P., Botella F., Margalida A. et al. (2018). Farmer perceptions of the ecosystem services provided by scavengers: What, who, and to whom. *Conservation Letters* **11**, e12392. <https://doi.org/10.1111/conl.12392>
- Morales-Reyes Z., Pérez-García J.M., Moleón M., Botella F., Carrete M., Lazcano C. et al. (2015). Supplanting ecosystem services provided by scavengers raises greenhouse gas emissions. *Scientific Reports* **5**, 7811. <https://doi.org/10.1038/srep07811>
- Morales-Reyes Z., Sánchez-Zapata J.A., Sebastián-González E., Botella F., Carrete M. and Moleón M. (2017). Scavenging efficiency and red fox abundance in Mediterranean mountains with and without vultures. *Acta Oecologica* **79**, 81–88. <https://doi.org/10.1016/j.actao.2016.12.012>
- Ogada D.L., Torchin M.E., Kinnaird M.F. and Ezenwa V.O. (2012). Effects of vulture declines on facultative scavengers and potential implications for mammalian disease transmission. *Conservation Biology* **26**, 453–460. <https://doi.org/10.1111/j.1523-1739.2012.01827.x>
- Oliva-Vidal P., Hernández-Matías A., García D., Colomer M.À., Real J. and Margalida A. (2022a). Griffon vultures, livestock and farmers: Unraveling a complex socio-economic ecological conflict from a conservation perspective. *Biological Conservation* **272**, 109664. <https://doi.org/10.1016/j.biocon.2022.109664>
- Oliva-Vidal P., Sebastián-González E. and Margalida A. (2022b). Scavenging in changing environments: woody encroachment shapes rural scavenger assemblages in Europe. *Oikos* **12**, e09310. <https://doi.org/10.1111/oik.09310>
- Plaza P.I., Blanco G. and Lambertucci S.A. (2020). Implications of bacterial, viral, and mycotic microorganisms in vultures for wildlife conservation, ecosystem services, and public health. *Ibis* **162**, 1109–1124. <https://doi.org/10.1111/ibi.12865>
- Plaza P.I. and Lambertucci S.A. (2022). Mitigating GHG emissions: A global ecosystem service provided by obligate scavenging birds. *Ecosystem Services* **56**, 101455. <https://doi.org/10.1016/j.ecoser.2022.101455>
- Santangeli A., Arkumarev V., Rust N. and Girardello M. (2016). Understanding, quantifying, and mapping the use of poison by commercial farmers in Namibia – Implications for scavengers' conservation and ecosystem health. *Biological Conservation* **204**, 205–211. <https://doi.org/10.1016/j.biocon.2016.10.018>
- Santangeli A., Girardello M., Buechley E., Botha A., Minin E.D. and Moilanen A. (2019). Priority areas for conservation of Old World vultures. *Conservation Biology* **33**, 1056–1065. <https://doi.org/10.1111/cobi.13282>
- Santangeli A., Lambertucci S.A., Margalida A., Carucci T., Botha A., Whitehouse-Tedd K. et al. (2024). The global contribution of vultures towards ecosystem services and sustainability: An experts' perspective. *iScience* **27**, 109925. <https://doi.org/10.1016/j.isci.2024.109925>
- Sekercioglu C.H., Wenny D.G. and Whelan C.J. (2016). *Why Birds Matter: Avian Ecological Function and Ecosystem Services*. Chicago: University of Chicago Press.
- Tidemann S.C. and Gosler A. (eds) (2010). *Ethno-Ornithology: Birds, Indigenous Peoples, Culture and Society*. London: Earthscan.
- van den Heever L., Thompson L.J., Bowerman W.W., Smit-Robinson H., Shaffer L.J., Harrell R.M. et al. (2021). Reviewing the role of vultures at the human–wildlife–livestock disease interface: An African perspective. *Journal of Raptor Research* **55**, 311–327.
- Van Dooren T. (2012). *Vulture*. London: Reaktion Books.